REVIEW

Contributions of Specialized Nurses to Medication Management for Older People in Home Care: A Mixed-Method Systematic Review

Mojtaba Vaismoradi 10¹, Patricia A Logan 10²

¹Faculty of Nursing and Health Sciences, Nord University, Bodø, Nordland, Norway; ²Faculty of Science and Health, Charles Sturt University, Bathurst, NSW, Australia

Correspondence: Mojtaba Vaismoradi, Faculty of Nursing and Health Sciences, Nord University, Bodø, Nordland, 8049, Norway, Email mojtaba.vaismoradi@nord.no

Abstract: There is a growing international interest in defining the roles, responsibilities, and contributions of specialized nurses across various aspects of home care to establish them as effective members of multidisciplinary home care teams. This study aimed to identify and describe the contributions of specialized nurses to medication management for older people in home care. This systematic review was performed using a mixed method approach. A thorough search was conducted across PubMed (including MEDLINE), Scopus, CINAHL, ProQuest, and Embase, focusing on studies published in English over the last decade, from 2014 to 2024. The convergent synthesis was used to combine and analyze qualitative and quantitative evidence in parallel to address the review question. The initial search yielded 875 studies, which were then refined to 32 studies selected for data-based convergent synthesis. The findings of the review were organized into three categories: "specialized nurses' identities", "specialized nurses' roles and responsibilities", and "impact of specialized nurses' medication management in home care". Specialized nurses played key roles in enhancing medication safety, improving older adults' quality of life, and reducing emergency visits. Their support in medication management eased the home care burden and ensured timely interventions for better symptom control and healthcare outcomes. Specialized nurses bring vital expertise to home care multidisciplinary medication team, promoting safe medication practices and helping older adults manage chronic conditions effectively at home.

Keywords: home care, medication management, medication outcome, non-physician prescriber, older people, specialized nurse, patient safety

Introduction

Specialized nursing involves bachelor's degree and master's degree nurses who pursue additional education and training to assume advanced roles and practices in a specific area of the healthcare system. This specialization encompasses various definitions influenced by differing expertise, education, and skills pertinent to specific care contexts across the globe. The International Council of Nurses (ICN) outlines two key roles within specialized or advanced practice nursing: clinical nurse specialists (CNS) and nurse practitioners (NP), differentiated by their educational and training requirements.¹ There is a lack of consensus on the terminology used for specialized nursing, educational requirements, and regulatory approaches for its effective implementation in practice nursing.² Nevertheless, Advanced Practice Nurse (APN) and NP roles have been established in a number of countries, including the USA, Canada, Australia, Ireland, and the Netherlands.³ In the UK, terms such as CNS, APN, and NP are used with variations in seniority and function.⁴ In the USA, the term Advanced Practice Registered Nurse (APRN) is commonly used, encompassing both CNS and NP roles.^{5,6} Autonomy is a characteristic of specialized nursing encompassing independent practice entry, authority and access to diagnostics, and recognition as a primary care provider.⁷ The capabilities and roles of specialized nurses facilitate coordination between aged care and general practice, address the needs of the aging population, and address the growing burden of chronic conditions leading to cost-effective, holistic primary care.^{8,9} In some cases, primary care

delivered by specialized nurses has been shown to be superior to that provided by physicians, demonstrating that they can also offer safe and effective primary care.¹⁰ Specialized nurses improve health care outcomes and behaviors in older adults in long-term care settings and enhance family satisfaction.¹¹

A comprehensive analysis of the global burden of disease for 371 diseases and injuries across 204 countries and territories covering 811 subnational locations, reveals a rise in global disability-adjusted life-years from 2.63 billion (95% uncertainty interval: 2.44–2.85) in 2010 to 2.88 billion (2.64–3.15) by 2021 due to demographic changes including ageing.¹² This trend has led to a significant increase in older adults using home care services, signaling a critical need for healthcare managers and policymakers to address the quality and safety of home care.¹³ Moreover, older people primarily wish to live in their own homes for as long as possible and therefore need improved home coping and safety.¹⁴ Home care services try to align with this emerging preference, enabling older people to "age in place" in familiar surroundings rather than receiving care for chronic health conditions or aging needs in institutional care settings.¹⁵

Older age causes chronic diseases, complex healthcare needs, and comorbidities, typically resulting in frailty and use of multiple medication use.¹⁶ Administrative records from 133 certified home healthcare agencies across 32 states in the USA revealed that, among older adults receiving home healthcare services (n=87,780), 44% received opioids, 19% were prescribed benzodiazepines, and nearly 7% were prescribed non-benzodiazepine sleep medications.¹⁷ In addition, a retrospective analysis of 100 older adults in Australia seeking community care support for medication management found that 66% used \geq 5 medications, and 48% used at least one high-risk medication, mainly opioids, anticoagulants, and insulin. Additionally, medication errors occurred in 41% of the patients, with 13% experiencing an adverse medication event requiring medical consultation or hospitalization, of which 64% were preventable.¹⁸ Polypharmacy and multimorbidity due to drug interactions affect 25.1% to 100% of older people internationally, with 30 to 388.3 drug interactions per 100 persons with higher values in home care given less controlled medication management than in nursing homes.¹⁹ A study of older people's home medications at hospital admission revealed an average of 11.6 ± 4.5 medications per person, of which 95% experienced polypharmacy, 69% hyper-polypharmacy, and 77.5% had at least one severe potential drug-drug interaction.²⁰ In a study in the Netherlands, 15% of readmissions of older adults within 30 days of discharge were medication-related, with 23% potentially preventable.²¹ Therefore, early and continuous interventions in the community are crucial for healthy aging,¹⁶ and for reducing emergency department visits and hospitalization.²²

Due to the complexity of older people's medication regimens, managing chronic medications is best handled by multidisciplinary medication team,²³ and specialized nurses have been nominated to offer crucial support in medication management and provide support to other healthcare staff.²⁴

Current questions in the international literature involve identifying and describing the roles of specialized nurses in delivering geriatric care in primary healthcare settings.²⁵ International literature lacks comprehensive insights into the integration and advancement of specialized nurses' roles in medication management within home care. This review aimed to cover this knowledge gap and provide an understanding of policies, strategies, and interventions that enhance the role of specialized nurses to medication management for older people in home care. The guiding review question was: What is the scope and impact of specialized nurses' contribution to medication management for older people to ensure patient safety in home care?

Materials and Methods

Review Design

This systematic review utilized a mixed-method approach, which helps understand how complex interventions function, for whom they are effective, and how healthcare systems adapt to their implementation.²⁶ This approach aligns with the traditional integrative review framework, which connects quantitative and qualitative data to synthesize current knowl-edge and enhance practice.²⁷ This review adhered to the PRISMA guideline to ensure structured development and reporting findings (Supplementary File 1).

Protocol and Registration

The international research team collaboratively developed a review protocol before conducting the review to promote transparency, maintain integrity, and minimize bias. The protocol was registered in PROSPERO, detailing the review objectives, methodology, and search plan (ID: CRD42024523188), accessible at the following link: crd.york.ac.uk/ PROSPERO/display_record.php?RecordID=523188.

Accordingly, the review aim was structured using the PICO framework as follows:

P (population): Specialized nurses are registered nurses (RNs) who focus on a specific area of nursing practice. With additional education and training at the bachelor's or master's level, nurses such as NPs, APNs, and CNS are prepared to deliver expert care in fields such as geriatrics, oncology, critical care, and mental health. Their advanced knowledge and skills enable them to provide targeted support and care for patients requiring monitoring and continuous care.

I (interest): The current frameworks, responsibilities, and contributions of specialized nurses in medication management defined as "to address medicines-related problems and optimize the use of medicines by providing advice on prescribing, medication monitoring, management of repeat prescribing systems and education and training on prescribing and the use of medicines".²⁸ The initial interest lies in understanding how to optimize the contribution of specialized nurses in enhancing medication safety and improving the quality of care for older people.

C (context): Home care refers to older peopel living in their own homes while receiving supervised, multiprofessional short- and long-term care provided within the community.

O (outcome): Minimizing the probability of medication errors, enhancing older people's outcomes, and overcoming challenges in medication management to improve overall patient safety.

Literature Search

A systematic search of the international literature was performed across databases, including PubMed (covering MEDLINE), Scopus, CINAHL, ProQuest, and Embase, focusing on studies published from January 1, 2014, to August 31, 2024, to capture recent research within the last decade. A pilot search on Google Scholar helped identify relevant keywords. Search strings were developed by translating Medical Subject Headings (MeSH) and thesaurus terms into database-compatible terms using Boolean logic and truncation with AND/OR operators. Consultation with a librarian ensured accuracy. Various versions of key terms related to "specialized nurse", "medication management", and "home care" were used (Supplementary File 2). Cross-references from article bibliographies were also examined to broaden search coverage.

Selection of Studies

All original research studies employing qualitative, quantitative, or mixed-methods designs were considered, with selection criteria focusing on medication management involving specialized nurses within the patient's home, adult home care, published in peer-reviewed scientific journals, in English, and in the last decade (2014–2024). The exclusion criteria were commentaries, letters, case reports, case studies, conference proceedings, and books; studies without specific details on specialized nurses' roles in medication management; studies focused on non-specialized nurses, nurse assistants, or associates; pediatric, child, or neonatal care (under 18 years); publications before 2014; and non-English language studies.

Search results were uploaded to the Rayyan platform for screening. Two authors (MV and PAL) independently screened and selected studies based on the eligibility criteria, first reviewing titles and abstracts, followed by full-text reviews. They reached a consensus on article selection through discussions and shared insights by creating tables that summarized the studies' characteristics, relevance, and suitability for inclusion. Discrepancies were resolved to finalize the inclusion and exclusion of the studies.

Quality Appraisal and Risk of Bias Assessment

This review included a comprehensive assessment of the credibility, relevance, and findings of the selected studies by two review authors (PAL and MV). The Mixed Methods Appraisal Tool (MMAT), version 2018, was applied to evaluate

studies with diverse methodologies because of its user-friendliness and structured criteria for qualitative, quantitative, and mixed-methods research, ensuring a robust quality assessment across various research designs.²⁹ To evaluate the risk of bias, RoB2 and ROBINS-I tools were used for randomized clinical trials and cohort studies, respectively. In addition, the Robvis tool was used to visualize the risk of bias assessment.³⁰ Studies were independently reviewed to ensure a detailed and accurate assessment of bias.

Ethical Considerations

Although this review study did not require ethical approval as it did not involve human samples, the authors have acknowledged all available sources and adhere to citation practices that respect intellectual property rights.

Research Synthesis

Differences in methodologies, objectives, data collection tools, and measurement outcomes across experimental and cohort studies, along with the inclusion of qualitative studies, make conducting a meta-analysis impractical. Therefore, within the mixed-method approach, a data-based convergent synthesis was employed to collate and analyze qualitative and quantitative evidence in a parallel and complementary manner to address the review question. The review results were presented at both descriptive and explanatory levels that incorporated statistical and non-statistical data. This approach aims to describe and integrate quantitative and qualitative findings, extend beyond theoretical insights, and offer practical interpretations for clinical use.^{26,31}

The data extracted from the included studies were organized in a table format for straightforward extraction, comparison, and categorization of the key characteristics. It helped to structuralise, summarize, and interpret the results of the studies, allowing for a thematic analysis that informed identifying recurring codes, similarities, and contrasts in findings and comparing them across studies led to the development of categories and subcategories for presenting findings.

Results

Outcome of the Search Process

The initial database search yielded a total of 875 studies. Following the removal of duplicates (n=115) and the screening of titles and abstracts, 78 studies remained. A thorough full-text review further refined the search to include 32 articles (<u>Supplementary File 3</u>). The primary reasons for exclusion were studies that did not focus on medication management initiatives, did not involve specialized nurses, lacked details on their roles in medication management, or were centered on acute or ambulatory healthcare settings, such as hospitals. After evaluating each article using appropriate methodological checklists, all 32 studies were included in the research synthesis (Figure 1).

Overview of Included Studies

The studies were published in English, spanning the last decade from 2014 to 2024. They originated from multiple countries, including the USA,^{32–51} Australia,^{52,53} the UK,^{54–56} Canada,^{57–59} Ireland,⁶⁰ Italy,⁶¹ South Korea,⁶² and the Netherlands.⁶³

Most studies utilized quantitative research designs, including three randomized clinical trials,^{34,38,47} 12 quasiexperimental studies,^{32,37,42,43,46,49,51–53,59,61,63} five cohort studies,^{39,40,45,57,58} four cross-sectional studies,^{44,48,54,55} and one case-control study.³⁶ Additionally, the data collection included 2 mixed-method studies,^{33,56} 5 qualitative studies.^{35,41,50,60,62} The study sample consisted of older adults with various chronic health conditions. Table 1 summarizes the selected studies and their general characteristics.

Evaluation of Methodological Quality

The results of the quality appraisal are presented in <u>Supplementary File 4</u>. All studies met the required quality criteria for their respective methodologies, demonstrating an appropriate study design, suitable data collection methods, coherent derivation of findings from the data, and alignment in data analysis, interpretation, and presentation of the results.



Figure I PRISMA diagram of search process. Adapted from Page MJ, McKenzie JE, Bossuyt P, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021; 372 :n71. Creative Commons.

Regarding risk of bias assessment, randomized clinical trials^{34,38,47} showed a low risk of bias in all dimensions, including randomization, intervention, outcome, and reporting results. Cohort studies^{39,40,45,57,58} general had a low risk of bias, except for some concerns with the participants' section, missing data, and measurement of outcome (Supplementary file 4). Therefore, all studies were considered of adequate quality and were included in the study.

Specialized Nurses in Home Care Oversee Medication Management

The review findings were grouped into the following categories: "specialized nurses' identities", "specialized nurses' roles and responsibilities", and "impact of specialized nurses' medication management in home care". The last category consisted of the following six subcategories: "quality of life enhancement", "emergency department visits", "burden of care for older adults and caregivers", "monitoring and identification of medication issues", "symptom management", and "workload and burden on the healthcare system". Table 2 summarizes the areas of concentration of the studies' findings regarding the impact of specialized nurses' involvement in medication management within home care.

Specialized Nurses' Identities

Given that specialized nursing practice encompasses a diverse range of definitions and interpretations reflecting varying expertise, education, and skills, the expertise of specialized nurses varied in our review as they fulfilled different roles in medication management within home care. They were mostly NP.^{32,34,36–43,46,48–53,57,58,63} They were also identified as APN,⁴⁷ advanced practice registered nurse,^{35,44,45} advanced nurse practitioner,⁵⁶ advanced practice nurse,^{59,62} family nurse practitioner,⁶¹ palliative care clinical nurse,³³ palliative care specialist nurse,⁵⁴ mental health nurse,⁶⁰ home care cardiovascular nurse,⁶³ and CNS.⁵⁵

Specialized Nurses' Roles and Responsibilities

Specialized nurses assumed multiple roles in medication management within home care, thus ensuring the safety of medication practices. The most highlighted role was medication prescribing.^{33,35,39,52,54,56–58,60,62,63}

$\textbf{Table I} \ \textbf{A} \ \textbf{Summary of the Selected Studies and Their General Characteristic}$

Authors and Year, Country	Focus	Research Methodology	Sample and Setting for Older People in Home Care	Intervention	Nurse Degree and Qualification	Role of Nurse in Medication Management	Measure	Implication for Patient Safety
Goldman et al, 2014, USA ³⁸	Hospital support at home for older people	Randomized controlled trial	700 older people divided to the intervention (n=306) and control (n=319) groups/own home	Peri-discharge, nurse-led intervention via telehealth along with physician, pharmacist, and nurse	Nurse practitioner	Assessing adherence to medications, answered questions about medications; adjusting medications; working with pharmacies to resolve prescription problem	Emergency department visits or readmissions	A nurse-led program involving patient education and follow-up phone calls
Hall et al, 2014, USA ⁴⁰	Cardiac surgery nurse practitioner home care for coronary artery bypass graft patients	Retrospective observational cohort	401 coronary artery bypass grafting older patients: 169 patients and 232 controls/own home	Transitional care program: Follow Your Heart	Nurse practitioner	Home visit, reconciliation of discharge medications, assessing patient treatment compliance, providing education to enhance understanding, and making and communicating necessary medication adjustments to primary care physicians, cardiologists, home care providers, and the cardiac	30-day readmission and death	Ensuring continuity of care, serving as a communication hub, and managing medications, while giving home care nurses the opportunity to familiarize themselves with the patient's specific issues, treatments, and unresolved concerns.

Chan et al. 2015. USA ⁴⁴ Care manison for older mult- lingual adults net. Care controlled trail 3/47 to the intervention and 353 to the control of mult- lingual adults net. Autoration for all to the participation of registered nurse and physicin Nurse vention Assessing absence practitioner- led for more medications, informing primary care provider about medication adjustments Need for more patient controlled rail participation and physicin Nurse medications, informing primary care provider about medication adjustments Need for more patient about medications, informing primary care provider adjustments Need for more patient about medications, informing primary care conference by nurse practitioner- led for rural paliative care Readomiting pail adjustments Need for more patient adjustments Mitchell et al, 2016, USA ⁴⁴ Nurse patienter practitioner- led for rural paliative care Plot practitioner- led for rural paliative care Plot patient sprint patients Plot patient sprint practitioner- led for rural paliative care Nurse patient sprint practitioner- led for rural paliative care Plot patient sprint practitioner- led for rural paliative care Plot patient sprint practitioner- led for rural patient sprint practitioner- led for rural practitioner- led for rural practitioner- le	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~								I
Moder multi- linguid adults in the site net Controlled rul immervation 353 to the control site site in the site in the site intervention interventinterventintervention intervention intervention intervention inter	Chan et al, 2015, USA ³⁴	Care	Randomized	685 older people:	Nurse-led hospital-	Nurse	Assessing	Discharge-related	Need for more
older multi- lingual adults intervention and intervention and prisidual singual adults intervention intervention and prisidual adults medication, informing primary adults experiences informing primary informing primary informing informing primary informing primary informing primar		transition for	controlled trial	347 to the	based care	practitioner	adherence to	patient	intensive post-
Mitchell et al, 2016, Australia ⁵² Nurse practioner- led for rural palitative care proxibiner Pilot single (2 older people) 353 to the control participation of registered nurse and physician Nurse participation practioner- led for rural palitative care proxibiner Informing primary (and cognitively mapped practioner- led for rural palitative care proxibiner Pilot proxibiner Pilot proxibiner 62 older people Nurse participation of community-based healt professionals and general physician Nurse practioner- led care with the participation of community-based healt professionals and general physician Nurse physician Reconciling physician Reconciling medication and medication and m		older multi-		intervention and	transition via		medications,	experiences	hospitalization
Mitchell et al, 2016, Australia ²⁴ Nurse protectioner- led for rural pallative care Plot intervention 62 older people/ own home Single own home Nurse multidiscipilinary case conference by nurse practitioner Nurse prescription Patient-reported intervention Initial assessments own home Moroe et al, 2016, USA ⁴⁶ Nurse Plot study 22 older people/ own home Nurse prescription Nurse prescription Patient-reported intervention Initial assessments own home Moroe et al, 2016, USA ⁴⁶ Nurse Plot study 22 older people/ own home Nurse prescription Nurse prescription Readmission advanced pallative care. improving care. imp		lingual adults		353 to the control	telehealth with the		informing primary		interventions
net and cognitively and physician about medication adjustments about medication adjustments nurse and cognitively adjustments Mitchell et al, 2016, Australia ⁵² Nurse practitioner led for rural pallative care Plot 62 older people/ intervention Single own home Nurse multidisciplinary case conference by nurse practitioner led care with the participation of community-based heath professionals and general physician Nurse practitioner led intervisioner intervention Patient-reported by Durse to uncovered urgent dinical issues patients through nurse practitioner intervision of community-based heath professionals and general physician Means practitioner intervisioner intervisioner patients through patients through patients through patients through nutidisciplinary care, corroflation and boosting patient through patients through patient through patients through patient through patients through		in the safety		groups/own home	participation of		care provider		among multilingual,
Mitchell et al. 2016, Australia ²⁴ Mitricener lef for rural pallative care intervention lef for sund pallative care Nurse Nurse Nurse Pallative care Nurse Pallative Pal		net			registered nurse		about medication		and cognitively
Mitchell et al, 2016, Australia ³⁷ Nurse Pliot 62 older poople/ Single Nurse Medication Padication Padication Practitioner Initial assessments I ed for runal palliative care intervention own home multidisciplinary practitioner practitioner practitioner practitioner practitioner outcomes, by nurse to I ed for runal palliative care intervention own home runse practitioner practitioner practitioner practitioner outcomes, by nurse to I ed or runal palliative care intervention own home runse practitioner practitioner practitioner practitioner care care care promo; cather cordination and doosting patient and carer care, improving care care conditioner maldidise pilinary practitioner madication and magement, madige pilination satisfaction home more feasible Moore et al, 2016, USA*6 Nurse practitioner edication and medication and medication and medication and medication and medication and medication and					and physician		adjustments		impaired patients
Practitioner led for rural palitative care intervention intervention own home mutidiscipinary case conference b nurse practitioner- led care with the participation of community-based health professionals and general physicain practitioner- led care with the participation of care community-based health professionals and general physicain practitioner- led care with the participation of care community-based health professionals and general physicain practitioner- led home care for congestive heart failure Pilot study Pilot study 22 older people/ own home Nurse practitioner- led home care program with home program with nurse, physical therapists, a diedician, pharmacists, social workers, home Nurse practitioner- led home care for congestive heart failure Pilot study 22 older people/ own home Nurse practitioner- led home care program with home program with nurse, physical therapists, a diedician, pharmacists, social workers, home Nurse practitioner- led home care program in therapists, a diedician, pharmacists, social workers, home Reconciling medication refili, assessing changes in medication refili, assessing changes in medications Redmission Effectiveness of home nurse-led program in hospital therapists, a diedician, pharmacists, social workers, home	Mitchell et al, 2016, Australia ⁵²	Nurse	Pilot	62 older people/	Single	Nurse	Medication	Patient-reported	Initial assessments
Norse et al, 2016, USA 46Nurse practitioner led met and congention heart failurePilot study22 older people/ own homeNurse practitioner led interdisciplinary physical therapists, a dieticain, physical therapist, a dieticain, physical therapist, a dieticain, physical therapist, a dieticain, physical therapist, a dieticain, physical therapist, a dieticain, physical therapist, a dieticain, physical		practitioner-	intervention	own home	multidisciplinary	practitioner	prescription	outcomes,	by nurse to
Moore et al, 2016, USA ⁴⁶ Nurse practitioner- led new the participation of community-based health professionals and general physician Nurse practitioner- led intervision for congestive heart failure Pilot study 22 older people/ own home Nurse practitioner- led intervision alog with nurse, physical therapists, a deticain, physical therapists, a deticai		led for rural			case conference by			depression and	uncovered urgent
Moore et al, 2016, USA ⁴⁴ Nurse Practitioner Ied home care for congestive heart failure Practicians Program with home for congestive heart failure Practicioner Ied home care for congestive Practicioner Ied home Practicioner Ied Practicioner Ied Practicione Practicioner Ied Practicione Practicioner Ied Practicione Practicioner Practicione Practicioner Pr		palliative care			nurse practitioner-			anxiety, palliative	clinical issues in
Moore et al, 2016, USA ⁴⁶ Nurse practitioner- led home care for congestive heart failure heart failure					led care with the			symptoms,	advanced palliative
Moore et al, 2016, USA ⁴⁶ Nurse practitioner- led home care for congestive heart failure led inter for congestive heart failure for congestive heart failure led inter for congestive heart failure heart fa					participation of			satisfaction	patients through
Moore et al, 2016, USA ⁴⁶ Nurse practitioner- led home care for congestive heart failure Heat failure Piot study Piot st					community-based				multidisciplinary
Moore et al, 2016, USA ⁴⁶ Nurse Pilot study Pilot stu					health professionals				care, improving
Moore et al, 2016, USA ⁴⁶ Nurse Nurse Nurse Pilot study Pilot stu					and general				care coordination
Moore et al, 2016, USA ⁴⁶ Nurse practitioner- led home care for congestive heart failure let failure					physician				and boosting
Moore et al, 2016, USA ⁴⁶ Nurse Practitioner- led home care for congestive heart failure led liner fa									patient and carer
Moore et al, 2016, USA ⁴⁶ Nurse Pilot study Pilot stu									confidence in
Moore et al, 2016, USA ⁴⁶ Nurse Pilot study Pilot stu									symptom
Moore et al. 2016, USA ⁴⁶ Nurse Pilot study Pilot stu									management,
Moore et al, 2016, USA ⁴⁶ Nurse practitioner- led home care for congestive heart failure let intervision let intervision heart failure let intervision let in									making dying at
Moore et al, 2016, USA46NursePilot study22 older people/ own homeNurse practitioner- led interdisciplinary program with home telemonitoring along with nurse, physical therapists, a dietician, pharmacists, social workers, home health aidesNurse practitioner- practitionerReadmissionEffectiveness of home nurse-led program in reducing emergency and hospital visits									home more feasible
practitioner- led home care for congestive heart failureown homeled interdisciplinary program with home telemonitoring along with nurse, physical therapists, occupational therapists, social workers, home health aidespractitionermedication and medication refill, assessing changes in medicationshome nurse-led program in reducing emergency and hospital visits	Moore et al, 2016, USA ⁴⁶	Nurse	Pilot study	22 older people/	Nurse practitioner-	Nurse	Reconciling	Readmission	Effectiveness of
Ied home care for congestive heart failureprogram with home telemonitoringmedication refill, assessing changes in medicationsprogram in reducingheart failurealong with nurse, physical therapists, occupational therapists, a dietician, pharmacists, social workers, home health aidesmedication refill, assessing changes in medicationsprogram in reducingunegret physical therapists, occupational therapists, a dietician, pharmacists, social workers, home health aidesnegret occupationnegret occupation		practitioner-		own home	led interdisciplinary	practitioner	medication and		home nurse-led
for congestive telemonitoring assessing changes in reducing heart failure along with nurse, medications emergency and physical therapists, occupational therapists, a dietician, pharmacists, social workers, home uorkers, home uorkel the aides		led home care			program with home		medication refill,		program in
heart failure heart		for congestive			telemonitoring		assessing changes in		reducing
hospital visits occupational therapists, a dietician, pharmacists, social workers, home health aides		heart failure			along with nurse,		medications		emergency and
occupational therapists, a dietician, pharmacists, social workers, home health aides					physical therapists,				hospital visits
therapists, a dietician, pharmacists, social workers, home health aides					occupational				
a dietician, pharmacists, social workers, home health aides					therapists.				
pharmacists, social workers, home health aides					a dietician,				
workers, home health aides					pharmacists, social				
health aides					workers, home				
					health aides				

Vaismoradi and Logan

Table	I	(Continued).
-------	---	--------------

Authors and Year, Country	Focus	Research Methodology	Sample and Setting for Older People in Home Care	Intervention	Nurse Degree and Qualification	Role of Nurse in Medication Management	Measure	Implication for Patient Safety
Quist et al, 2016, USA ⁴⁸	Medication management team-based care for older people	Retrospective descriptive study	152 older people	GRACE team care along with social worker, geriatrician, pharmacist, mental- health professional	Nurse practitioner	Assessment and recommendation to the team for medication management	Implementation of recommendations	Facilitating medication reconciliation, education, communication, and care coordination
Hernandez, 2017, USA ⁴¹	Medication management for older people	Narrative inquiry	15 nurse practitioners/own home	Expertise in providing care for older adults while managing the complexities of medication administration	Nurse practitioner	Managing complex medication needs of patients	Experiences and perspectives	Remaining attentive to polypharmacy and the challenges of managing complex medication regimens, while simplifying medication management and ensuring continuity of care through digital solutions, as well as empowering and educating older people
Bagchi et al, 2018, USA ³²	Telemedicine care for older people	Pilot study with a mixed methods approach	10 older people	Telemedicine application	Nurse practitioner	Medication review	Satisfaction and experiences and perspectives	Identifying needs and preferences of underserved communities

							_	
Dawson et al, 2018, UK ³³	Home-based	Retrospective	92 older people/	Home-based	Clinical Nurse	Explaining	Post-transition	Clinical stability and
	medication	analysis	own home	medication	Specialist	medication	adverse event,	tolerance due to
	transitioning			transitioning by		changes, new	feasibility, safety,	the transitional
	for patients			telehealth		dosing schedule,	effectiveness	program
	with					potential side		
	pulmonary					effects, and drug		
	arterial					interactions,		
	hypertension					advising to report		
						any adverse events,		
						communicating		
						medication issues		
						to the healthcare		
						team		
Wong et al, 2018, Canada ⁵⁹	Transitional	Pilot study	77 older people	Evaluating	Advanced	Assessing	Fluid balance	Need to establish
	care from		with transcatheter	transitional care	practice nurse	medication,	monitoring,	consistent
	hospital to		aortic valve	strategies		rectifying	medication	processes to
	home for		implantation/own			medication	management, and	support older
	older people		home			discrepancies. Self-	addressing anxiety	people during
	after					care education	and depression	vulnerable care
	transcatheter					around medication		
	aortic valve					management		
	implantation							
Edge et al, 2019, Canada ⁵⁷	Medication	Population-	25,220 older	Sociodemographic	Nurse	Medication	Prescription data	Understanding care
	prescription	based	people/own home	characteristics and	practitioner	prescription	based on	patterns for
	by nurse	descriptive		comorbidities of			demographic and	shaping primary
	practitioners	retrospective		older adults			health-related	care policy
	and family	cohort		prescribed			characteristics	
	physicians			medications by				
				nurse practitioners				
				and family				
				physicians				
Fisher et al, 2019, USA ³⁷	Remote,	Pilot	130 older people/	Remote, navigator-	Nurse	Medication	Blood pressure	Efficiency and
	nonphysician-	intervention	own home	led hypertension	practitioners	titration and	•	safety of care by
	led			innovation program		training patient's		non-physician
	hypertension			along with		navigator		models
	management			a pharmacist				
L	management			" P.iai macióc				

(Continued)

Vaismoradi and Logan

Table I (Continued).

Authors and Year, Country	Focus	Research Methodology	Sample and Setting for Older People in Home Care	Intervention	Nurse Degree and Qualification	Role of Nurse in Medication Management	Measure	Implication for Patient Safety
Leask & Tennant, 2019, UK ⁵⁶	Unscheduled care advanced nurse practitioners in primary care	Mixed- methods design	239 home visits to older people/own home	Unscheduled home visits on behalf of physician in primary care	Advanced nurse practitioner	Medication prescription	Number of referrals, experiences of nurses and physicians of services	Ensuring that nurse-led care delivery is feasible and acceptable to physicians
Possin et al, 2019, USA ⁴⁷	Collaborative dementia care via telephone and internet in dementia	Single-blind, randomized clinical trial with a pragmatic design	Older people with dementia and their caregivers: 512 dyads in the intervention and 268 in the control group/own home	Telephone-based collaborative dementia care	Advanced practice nurse within the dementia expert providers' team along with social worker and pharmacist	Helping to convey medication recommendations from pharmacist to patients	Quality of life, emergency department/ system use, readmission, caregiver depression/burden	Access to experts for issues beyond the care team's scope, like medical needs, behavioural problems, legal or financial complexities, and safety concerns.
Warren et al, 2019, USA ⁵¹	Community- based care transition	Quality improvement program	7931 older people/ own home	Community-based care transitions program	Nurse practitioner	Addressing medication questions or identifying medication-related issues involves referring patients to a pharmacist and assisting in identifying and resolving various drug-related issues	30-day all-cause readmission rate, readmission rate among the high- risk cohort, achievement of target volumes for full enrolment	Successful transitions through relying on the collaboration of healthcare providers, social workers, nurse practitioners, physicians, community pharmacists, and visiting nurses

							1	1
Antunes et al, 2020, UK ⁵⁴	Anticipatory prescribing for end-of- life care in community	Online survey	261 consisting palliative medicine consultants, general practitioners, palliative care specialist nurses (n=31), general community nurses, palliative medicine trainees, other doctors, pharmacists, other nurses, and other professional group/	Anticipatory prescribing in community end-of- life care	Palliative care specialist nurse	Prescribing and dispensing injectable medications in anticipation of clinical need	Concerning changes in anticipatory prescribing during the COVID-19 pandemic and recommendations for improvement.	Challenge to community palliative care during pandemics
			home care					
Guo et al, 2020, USA ³⁹	Diabetes mellitus care by primary care team	Cohort	306741 older people within 3524 primary care practices/own home	Different types of team primary care models	Nurse practitioner	Medication prescription	Guideline- recommended diabetes care, prescription of high-risk medications, hospitalization rate	Team-based care with nurses improved adherence to diabetes guidelines compared to physician-only practices
Kulsick et al, 2020, USA ⁴³	Medication adherence in older people at nurse practitioner clinic	Quality improvement project	319 older people/ own home	A review of medical records for patients over 52 weeks before an educational seminar and 13 weeks afterward; educating nurses on evidence-based, provider-driven strategies for medication management	Nurse practitioner	Enhancing medication adherence, prescriber consistency, provider-patient interaction, and medication reminders	Medication adherence rate and clinical biomarkers including blood pressure, lipid profile, blood glucose indicators	A need for ongoing monitoring of adherence and clinical outcomes in this specific patient population

(Continued)

Vaismoradi and Logan

456

http

Authors and Year, Country	Focus	Research Methodology	Sample and Setting for Older People in Home Care	Intervention	Nurse Degree and Qualification	Role of Nurse in Medication Management	Measure	Implication for Patient Safety
Squires et al, 2020, USA ⁵⁰	Medication complexity in home care	Qualitative secondary data analysis using directed and summative content analysis	77 home healthcare professionals including 12 nurse practitioner preceptors	Interprofessional intervention based in home care with nurse practitioner, social workers, and pharmacists and their student trainees	Nurse practitioner preceptor	Medication management	Perspectives and experiences of medication management challenge	Patient, provider, and system-level factors affecting medication complexity and management leading to communication and coordination challenges
Ernecoff et al, 2021, USA ³⁶	Home-based, nurse practitioner- led care	Retrospective case-control study	110 older people and 371 controls/ own home	A nurse practitioner-led, home-based care program for individuals with serious or complex chronic illnesses on palliative care needs and coordinating with patients' primary care and specialist providers	Nurse practitioner	Complex home support for antibiotics, screening for medication adherence	30- and 90- day emergency department visits and hospitalization, hospice conversion, and 30- and 90-day mortality	Addressing care needs and gaps early to avoid unnecessary hospitalizations and receive timely hospice services
Frain et al, 2021, Ireland ⁶⁰	Mental health nurse prescribing in home care	Qualitative thematic analysis	12 mental health nurses in home care/own home	Mental health nurse prescribing	Mental health nurse	Medication prescription	Perspectives and experiences	Nurse prescribing enhances care for individuals and improve access to services. Need for facilitating discussions about reducing or discontinuing medications

Γ								1
Levine et al, 2021, USA ⁴⁴	Medication	Retrospective	105 older people/	Advanced practice	Advanced	In-home	Medication	Multidisciplinary
	therapy	analysis	own home	registered nurse-	practice	medication	therapy problems:	teamwork to
	problems			led,	registered	reconciliations and	indication,	identify cognitively
	involving			multidisciplinary	nurse	feeding pharmacist,	effectiveness,	harmful
	cognition			home-based care		geriatrician, and	safety	medications, side
	among older			team		psychiatrist with		effects of dementia
	people					data for further		treatments, and
						actions; engaging		untreated cognitive
						patients and		conditions
						caregivers in		
						discussions about		
						medication		
						recommendations		
Misra-Hebert et al, 2021, USA ⁴⁵	Home visit	Retrospective	468 older people in	Post-discharge	Advanced	Educating patients	Hospital	Medication
	for patients	cohort using	two phases within	home visit program	practice	on the reasons for	admissions,	education and
	discharged	convergent	17 months/own	by nurse and	registered	their medications	emergency	enhancing
	from the	mixed	home	paramedics	nurse	and how to use	department visits,	understanding of
	hospital	methods				them properly	and mortality	discharge
								recommendations
								can benefit
								medically complex
								patients and
								enhance care
								outcomes
Rose et al, 2021, USA ⁴⁹	Transitional	Quality	129 older people/	Transitional care	Nurse	Medication	30-day	Identifying high-risk
	care following	improvement	own home	visit following	practitioner	assessment	readmission rate,	patients discharged
	a skilled			skilled nurse facility			identifying gaps	from the hospital
	nursing facility						and needs in care	to home for
	by nurse							potential
	practitioners							readmission
					1			

(Continued)

Table I (Continued).

Authors and Year, Country	Focus	Research Methodology	Sample and Setting for Older People in Home Care	Intervention	Nurse Degree and Qualification	Role of Nurse in Medication Management	Measure	Implication for Patient Safety
Savini et al, 2021, Italy ⁶¹	A family nurse-led intervention for patients with chronic diseases	Non- randomized before-after pilot study	70 older people/ own home	Nurse-led educational intervention	Family nurse practitioner	Educating home nurses about self- management skills, such as medication adherence and recognizing and managing signs and symptoms for patient empowerment	Rate of health services' use	Nurse-led education has the potential to reduce healthcare utilization in older patients with chronic diseases
Pond et al, 2021, Australia ⁵³	Nurse practitioner- led collaborative dementia care	Feasibility study	76 older people/ own home	Nurse practitioner- led mobile memory clinic	Nurse practitioner	Advising general physician to conduct medication review	Satisfaction and advice to physician	Early diagnosis and assessment, improvement of access to care for this at-risk population
Campling et al, 2022, USA ³³	Palliative care medication in the community	Evaluative, mixed method case studies of service delivery models	40 older patients' records/own home	A structured log documenting medication access experiences over an 8-week period.	Palliative care clinical nurse specialist prescribers	Medication prescription	Perspective and experiences, cost	Growing number of nurse prescribers, along with enhanced shared inter-professional access to electronic prescribing systems, improves continuity of care and strengthens professional relationships.

1	Advant 1	Qualitati	20	Designs of t	Advance 1	Madiatia	Function 1	Effective
Lee et al, 2022, South Korea	Advanced	Qualitative	20 nurses	ratient safety	Advanced	riedication	Experiences and	Enecuve
	practice		specialized in nome	culture in nospital-	practice nurse	prescription	perspectives of	communication and
	nurses and		care/own nome	based nome			patient safety	collaboration and
	patient safety			healthcare				involving patients,
	culture in							caregivers,
	hospital-based							colleagues, and
	home care							medical
								professionals within
								community
								partnership
van Ramshorst et al, 2022, the	Connected	Feasibility	600 older people	Hospital-at-home	Nurse	Intravenous	Patient	Safety of
Netherlands ⁶³	care for	study	with endocarditis/	model using virtual	practitioner	treatment at home	satisfaction, clinical	ambulatory
	endocarditis		own home	ward with home	and home care	and prescribing the	efficacy outcomes	medication
	and heart			care nurse and	cardiovascular	necessary doctor's		treatment at home
	failure via			cardiologist	nurse	orders		and its role in
	hospital-at-			-				reducing length of
	home care							hospital stays
Kanne et al, 2023, USA ⁴²	The	Ouality	123 older patients/	Interagency care	Nurse	Reviewing	Identification of	Linking older adults
	interagency	improvement	own home	team of geriatrics	practitioner	medications and	medication issues	with complex care
	care team for	with a single		providers and	•	patient-reported		needs to geriatric
	older people	site, multi-		community		barriers, such as		expertise and
	in primary	method design		partners		affordability and		community
	care	method design		collaborating		side effects via		services through
	Care			through electronic		telephone along		collaborations with
				consultations		with pharmacist		primary care
				consultations		with pharmatist		primary care
Sector at al. 2022 Canada ⁵⁸		Deterror	102 (() de se de sée		Numer	Madiatian	At least one	Providers Resoluting home
Scott et al, 2023, Canada		Retrospective	103 664 decedents	End of life care	INUrse	Medication	At least one	Receiving nome
	nome care by	conort study	older people/own	nome visit	practitioner	prescription for	medication claim	visits from a nurse
	nurse		home			symptom	related to pain and	near the end of life
	practitioner					management	symptom	may lead to better
	and physician						management	outcomes aligned
							occurring during	with patients'
							the last 90 days of	preferences.
25							life	
Datta et al, 2024, USA ³⁵	Decision-	Qualitative	13 advanced	Home-based	Advanced	Antibiotic	Experiences and	Managing
	making in	using	practice registered	primary care	practice	prescription	perspectives	uncertainty while
	home-based	grounded	nurses/home care	program	registered			weighing competing
	primary care	theory			nurses			factors in infection
								treatment.

Abbreviations: GRACE: Geriatric Resources for Assessment and Care of Elders; COVID-19: Coronavirus Disease 2019.

Author, Year	Impact on Medication Management										
	Quality of Life Enhancements	Emergency Department Visits	Burden of Care for Older Adults and Caregivers	Monitoring and Identification of Medication Issues	Symptom Management	Workload and Burden on the Healthcare System					
Goldman et al, 2014 ³⁸ Hall et al, 2014 ⁴⁰		×				×					
Mitchell et al, 2016 ⁵²				×	×	×					
Moore et al, 2016 ¹⁵ Hernandez, 2017 ⁴¹			×	×	×	×					
Bagchi et al, 2018 ⁵⁵ Dawson et al, 2018 ⁵⁵				× ×	~						
Edge et al, 2019^{57} Eisbor et al, 2019^{37}				×	Ŷ						
Leask & Tennant, 2019 ⁵⁶	×	×	×			×					
Warren et al, 2019^{51}					×	×					
Guo et al, 2020^{39} Kulsick et al. 2020^{43}			×		×						
Squires et al, 2020^{50} Francoff et al, 2021^{36}		×		×		×					
Frain et al. 2021^{60}				×							
Misra-Hebert et al, 2021 ⁴⁵ Quist et al, 2016 ⁴⁸				×		×					
Rose et al, 2021 ⁴⁹ Savini et al, 2021 ⁶¹			×	×		×					
Pond et al, 2021 ⁵³ Campling et al, 2022 ³³			×	× ×	×	×					
Lee et al, 2022 ⁶² van Ramshorst et al, 2022 ⁶³	×		×		×	×					
Kanne et al, 2023 ⁴² Scott et al, 2023 ⁵⁸			×	×	×						
Datta et al, 2024 ³⁵				×		×					

 Table 2 Outcome Indicators of Specialized Nurses' Medication Management Within Home Care Discussed in the Reports of 32

 Included Studies

Within the multidisciplinary medication team, their roles encompassed bridging the healthcare team members and older people by providing data to healthcare providers at various healthcare levels for further actions,^{43,44} communicating medication issues and necessary medication adjustments,^{34,40,55} conveying medication recommendations from pharmacists to older people⁴⁷ and referring older people to a pharmacist, helping to identify and resolve drug-related concerns,⁵¹ and following up older people-reported barriers such as affordability and medication side effects through telephone consultations with a pharmacist.⁴² Moreover, they advised collaboration with physicians and pharmacies to perform medication reviews, resolve prescription issues, and manage medications.^{38,48,53}

In terms of educational role, specialized nurses educated other home nurses on remote navigation for medication monitoring,³⁷ self-management skills including medication adherence, and recognizing and managing signs and symptoms.⁶¹ They were educated about the reasons for their medications and proper usage,^{36,40,45} engaged older people

and caregivers in discussions about medication recommendations and self-care medication management,^{44,59} answered their questions about medications,^{38,51} explained medication changes, new dosing schedules, potential side effects, or drug interactions, and advised older people to report any adverse events.⁵⁵

Their monitoring role was to screen older people's medication adherence,^{34,36,38,40,43} improve it through prescriber consistency and medication reminders.⁴³

Regarding the general aspects of medication management, they reviewed medications,^{32,42} undertook medication management based on older people's complex medication needs,^{41,50} assessed medications,^{34,48,49,59} rectified and adjusted discrepancies,^{38,59} reconciled medications,^{40,44,46} titrated medications, and refilled dispensing devices while assessing changes in prescriptions^{37,46} and administered intravenous medications.⁶³

Impact of Specialized Nurses' Medication Management in Home Care

They were classified into six subcategories. A summary of specialized nurses' contributions to and their impact on medication management in home care for older people is presented in Table 3.

Subcategory	Description of Impact
Quality of life enhancement	Personalized-treatment and care;
	Assessment for medication issues;
	Monitoring medications' effects and side effects;
	Promotion of adherence to medications;
	Collaboration with the pharmaceutical care team;
	Meeting older people's need and expectations
Emergency department visits	Proactive medication management;
	Prevention of complications;
	Sensitization to medications effects and side effects
Burden of care for older adults and	Medication prescription and administration;
caregivers	Improvement of mood, reduction of caregiver's being overwhelmed;
	Education and empowerment of older people and caregivers regarding polypharmacy, medication effect,
	side effect and adherence and communication with provider;
	Identification and management of polypharmacy;
	Support to access medications;
	Follow up of guidelines and the use of data for medication management
Monitoring and identification of	Detection of medication discrepancies;
medication issues	Monitoring adverse drug reactions and risk of complications;
	Ongoing assessment of medication use at home;
	Communication of medication issues with the therapeutic team;
	Use of electronic health record system;
	Medication list review to check dosage, frequency, use of herbal and over the counter drugs, listening to
	older people and families' concerns;
	Detection of duplications, drugs interactions over- and under -use and effectiveness level of medications;
	Recommendations for medication review, dose reduction and deprescription of potentially inappropriate
	medications, discharge medication counselling and detection of discrepancies;
	Updating medication list for primary care provider's follow up;
	Reduction of comorbidity; encouragement of older people to collaborate in decision making;
	Suggestion of alternative treatment options;
	Readily accessible to older people;
	Handling medications in a timely accessible manner

Table 3 A Summary of the Contributions and Impact of Specialized Nurses on Medication Management in Home Care

(Continued)

Table 3 (Continued).

Subcategory	Description of Impact
Symptom management	Assessment of reaching intended therapeutic outcomes;
	Balancing older people's needs with the necessity of medication use;
	Balancing medications' risks and benefits;
	End of life medication prescription;
	Management of polypharmacy through identification of reasons for medication use, effects and side effects;
	Deprescription of medications;
	Palliative care medication re-prescription;
	Promote symptoms' relief;
	Securing home medication storage;
	Adaptation of medication type based on availability
Workload and burden on the	Reduction of hospitalization and home visit;
healthcare system	Close medication use monitoring;
	Education of older people of effects, side effects and adherence;
	Early identification of symptoms and prompt intervention;
	Education of older people about self-management, medication verification;
	End of life care medication prescription;
	Ambulatory medication treatment

Quality of Life Enhancement

Specialized nurses play a crucial role in home-care medication management, directly impacting older adults' quality of life by ensuring safe, effective, and personalized treatment. This was the result of careful assessment, identification of potential medication issues, monitoring of side effects, and promotion of adherence to prescribed regimens, in collaboration with other healthcare professionals. Accordingly, specialized nurses' active involvement in medication management in home care enhanced older people's quality of life (B = 0.53; 95% CI: 0.25-1.30; P = 0.04).⁴⁷ Older adults also stated that home care meets their expectations and needs. They would recommend such care at home and would choose this option again.⁶³

Emergency Department Visits

The impact of specialized nurses' participation in medication management in home care was changing the pattern of emergency department visits through proactive medication management and preventing complications that led to emergency care. It has improved the outcomes of older people and helped alleviate the strain on emergency healthcare services. The results showed that specialized nurses' efforts in medication management lowered emergency department visits among older adults (B = -0.14; 95% CI: -0.29, -0.01; P = 0.04).⁴⁷ In contrast, in one study, older people were more likely to have emergency department visits at 30 days (15.1 percentage points, CI: 4.9-25.3; P = 0.004) and 90 days (27.8 percentage points, CI: 16.0-39.6; P < 0.001) of the specialized nurse' care intervention.³⁶ Similarly, in another study, specialized nurses referred 3.6% of older adults to urgent care, 1.3% to the emergency department, and 1.6% to paramedics.³⁸ The increased number of emergency department visits could be due to the early identification, referral, and sensitization of older people to medication side effects.

Burden of Care for Older Adults and Caregivers

The involvement of specialized nurses in medication management enhances care quality by reducing the burden on both older people and their caregivers in home care, as highlighted in their statements, perspectives, and statistical results. For instance, specialized nurses, through intravenous treatment and prescription at home, met older people's expectations and needs, as they expressed that they would recommend it to others.⁶³ Moreover, it reduced caregiver depression (B = -1.14; 95% CI, -2.15, -0.13; P = 0.03) and burden (B = -1.90; 95% CI: -3.89, -0.08; P = 0.046).⁴⁷

Specialized nurses empowered older adults and caregivers via health information, and played a key role in identifying and managing polypharmacy, focusing on community-level efforts by educating older adults and their

caregivers. They informed the older people's family, significant others, or close friends to be aware of and recognize if they were struggling to manage or take their medications consistently.⁴¹ They also provided older adults with information on how to obtain medications, and checked for stock availability at pharmacies.³³ The outcome was that older people in the nurse-practitioner-led team received more guideline-recommended diabetes care, such as annual eye examinations (adjusted OR, 1.04; 95% CI, 1.00–1.08) and HbA1C tests (adjusted OR: 1.11, 95% CI: 1.04–1.18), affecting the process of medication management and better adherence to clinical practice guidelines for diabetes management compared to physician-only practices.³⁹ After home visits by specialized nurses, older adults' understanding of their medications improved significantly, with 33.9% reporting significant improvement and 44.44% noting some improvement.⁴⁹ In addition, 94% of primary care team members understood the program's purpose, found the recommendations easy to implement, and believed that their older adults were more likely to remain safe at home longer because of consultation with a specialized nurse. Healthcare team members also reported feeling less overwhelmed when managing complex older adults and addressing polypharmacy.⁴² Education on medication management and interaction with providers, reviews, and reminders to older people by specialized nurses improved the adherence rate for diabetes management from 85.7% (SD = 19.58) to 94.6% (SD = 13.73, P = 0.019).⁴³

Monitoring and Identification of Medication Issues

Specialized nurses identified and monitored medication issues as detected discrepancies and monitored for adverse reactions to prevent medication errors and reduce the risk of complications. Their ongoing assessment and communication with healthcare professionals regarding medication regimens ensured the quality of care. They used electronic health records (EHR) to reduce potentially inappropriate medications (PIMs) by verifying medication lists, clarifying dosages and frequencies, and addressing older people's concerns, including over-the-counter and herbal products. This involves the identification of therapeutic duplications and potential drug interactions.⁴¹ Specialized nurses in home care practices helped identify indication medication therapy problems accounting for 34% (57 out of 166) of total medication therapy problems, with 79% (45 out of 57) related to underuse and 21% (12 out of 57) to overuse, and effectiveness medication therapy problems representing 13% of the total, and safety medication therapy problems comprising over half (52%) of all medication therapy problems, often involving benzodiazepines and anticholinergics. The specialized nurse-led program, along with a multidisciplinary medication team, identified and provided recommendations including 23% for discontinuation and 19% for dose reduction. Cognitive medication therapy problems were found in 79% of older people with undesirable effects, comprising 41% (36 out of 87) of safety medication therapy problems. Acetylcholinesterase inhibitors (AChEIs) were responsible for 67% (24 out of 36) of these occurrences, leading to recommendations for discontinuation (25%) or dose reduction (8%). Inappropriate medication adherence was identified in terms of underuse, including potentially untreated dementia (62%, 28 out of 45) and depression (38%, 17 out of 45). Dose reductions (58%) or discontinuations (28%) due to unsafe medications were applied to benzodiazepines, anticholinergics, nonbenzodiazepine receptor agonists (Z-drugs) and antipsychotics.⁴⁴ Specialized nurses monitored medication issues to make decisions regarding medication changes due to inadequate responses (92% for macitentan and 77% for ambrisentan), side effects (one person switched to macitentan due to side effects, and four transitioned from bosentan to ambrisentan for issues such as rash (n=2), headache (n=1), low oxygen (n=1), abnormal liver tests, diarrhea (n=1), and intolerance (n=1). Mild ankle edema occurred in 7 macitentans and 12 ambrisentan persons.⁵⁵ In a specialized nurse care program, nine out of ten people indicated that they were likely or very likely to recommend telemedicine services to friends and family. The benefits of specialized nursing care include consultations and medication refills, observation of medication adherence, medication reviews.³² Older people and caregivers found the assessment by the specialized nurse acceptable and useful; out of 76 older people, the specialized nurse requested the general physician for a medication review for 23 older people.53

At the time of discharge from skilled nursing facilities to home, older adults experienced conflicting medication lists from both the hospital and nursing facilities, leading to confusion. Of the older adults receiving home visits by specialized nurses, 81.25% were identified as having medication discrepancies, often because they did not pick up prescriptions, resuming discontinued medications, or confusion over changes made between the hospital and nursing

home stay.⁴⁹ Discharge medication counselling by specialized nurses led to an improvement in older adults' inquiries about medications 30 days after discharge compared to the control (92.1 vs 86.0, P = 0.02).³⁴ Specialized nurses identified areas for improvement by medication assessments in home visits, including tapering narcotics, transitioning older people off sliding-scale insulin, updating the medication list in the electronic health record before discharge, and improving primary care provider follow-up, as only 25% of discharged older people had follow-ups within seven days.⁴⁹ Of the 924 medication-related recommendations provided by specialized nurses, 96% (887) were implemented. The majority (94%) involved reviewing medications (151), updating medication lists (153), and educating patients about medication safety (132). Among 58 drug-specific recommendations, 36% (n = 21) were adopted. Three of the four suggestions for discontinuing medication were as follows: Specific recommendations included starting multivitamins (15 suggested, 3 implemented), calcium with vitamin D (20, 4 implemented), antidepressants (6, 6 implemented), analgesics (2, 2 implemented), anticholinergics (6, 4 implemented) and antipsychotics (2, 1 implemented). Seven of the nine recommendations to modify or stop PIMs were also followed, including amitriptyline and diphenhydramine.⁴⁸ In another study, specialized nurses identified various medication issues (84%) related to medications leading to dose adjustment, alerts about drug interaction, geriatric precautions, and adherence.⁴²

Older people treated by specialized nurses than those primarily seen by family physicians (1.30 vs 2.04) had lower mean comorbidity index scores.⁵⁷ Additionally, Older people value being encouraged by nurse prescribers to collaborate in decision making and explore alternative treatment options for a person-centered approach to care. They appreciated the therapeutic relationship in a flexible and unhurried manner, and provided prompt access to nurses for medication-related issues and unresolved symptoms.⁶⁰ However, coordination issues within the information exchange, disagreements, and unclear line of responsibilities, such as medication reconciliation between specialized nurses and other healthcare providers, hindered medication safety as it highlighted restrictions on their practice as a key aspect of their professional authority in home care in managing complex medication regimens.⁵⁰ Moreover, older people and families are reassured by specialized nurses' medical authority so that actions can be taken, leading to enhanced medication use.³⁵ While nurse prescribers typically transferred responsibility for repeat prescriptions to physicians, they sometimes handled repeat prescriptions themselves when they judged that it would be faster for older people, such as during symptom exacerbations or when a holiday or weekend was approaching.³³

Symptom Management

Specialized nurses helped with symptom management by accurately assessing older adults' symptoms and ensuring that medications were prescribed, administered, and monitored correctly to achieve the intended therapeutic effects, leading to improved well-being. For appropriate medication management by specialized nurses, it is crucial to recognize that not everything that older people need should be fixed using medications. Additionally, one aspect of specialized nurses' responsibilities for managing polypharmacy requires that they balance quality of life with the risks and benefits of medication prescriptions.⁴¹ For older people undergoing end-of-life care, a study showed that receiving a home visit from specialized nurses increased the odds of receiving medication for symptom management in the last three weeks of life (OR: 2.54, 95% CI: 2.33 to 2.76).⁵⁸ Specialized nurses considered the dysfunctional cycle in which the side effects of one medication were mistaken for new issues, leading to more prescriptions. Therefore, they identified older people with polypharmacy and the reasons behind the use of each medication, ensuring that they were not just treating the side effects of other medications. An example was given to older people with multiple care providers and numerous medications, in which the specialized nurse referred a person to the specialist who then prescribed a medication that led to delirium and was misattributed to dementia, but the nurses recognized medication side effects and ceased the medication.⁴¹ However, in another study, older people had a higher risk of being prescribed high-risk medications by specialized nurses (adjusted OR: 1.03, 95% CI: 1.00–1.07), such as anticholinergics, antithrombotics, alpha agonists, barbiturates, vasodilators, nonbenzodiazepine hypnotics.³⁹ Specialized nurses for palliative care prescribed 24 new medications for 12 older people and adjusted the dose of nine medications in six persons as for analgesia, including antiepileptics and antidepressants, to relieve neuropathic pain.⁵² They also used their knowledge of end-of-life medicines to source and represcribe appropriate medications in a timely manner when symptoms arose during home care.³³ Specialized nurses' medication assessment and rectification helped older people (87.2%) reach no symptom and activity limitation state, with scores ranging from 1

to 4 (mean: 1.32) after 30 days.⁵⁹ In the case of symptoms arising, specialized nurses promptly investigated and addressed them using medications along with long-term symptom management through secure medication storage at home.⁶² A study on hypertension management using amlodipine as the most frequently added drug to specialized nurses showed that 91% of 116 older adults reached their target BP at an average of 7 ± 7 weeks. Systolic BP decreased from 155 ± 18 mmHg in the clinic to 124 ± 8 mmHg at home, and diastolic BP decreased from 92 ± 13 to 74 ± 8 mmHg (P < 0.001) over 6 months, with a minimal increase (1.4–1.8 on average).³⁷ During the COVID pandemic for end-of-life care, specialized nurses adapted medications based on availability, using hyoscine patches instead of glycopyrronium. They prioritized the subcutaneous route and utilized buccal or transdermal options if subcutaneous drugs were unavailable and increased coverage on weekends and holidays.⁵⁴

Workload and Burden on the Healthcare System

Medication management by specialized nurses reduced the workload of the healthcare system, especially regarding hospitalizations and home visits. By closely monitoring older people's medication schedules, managing potential side effects, and educating older people on adherence, they improved care outcomes and prevented medication-related complications, leading to extra burden on healthcare systems. For older people with Parkinson's disease who are prone to aspiration pneumonia and recurrent hospitalizations, early treatment was prioritized if symptoms such as increased cough or mild shortness of breath increased, avoiding hospitalization, even if it risked overprescribing.³⁵ Specialized nurse-medication management reduced probability of hospitalization at 30 days (11.4 percentage points, CI: -17.7 to -5.0; P < 0.001), and instead enhanced more transition to hospice care (22.4 percentage points, CI: 11.4–33.3; P < 0.001).³⁶ In another study, the readmission rate decreased to 12.5%, representing an 11% decline from the inception program by a specialized nurse to address medication issues.⁵¹ Discharge medication management intervention by specialized nurses resulted in a significantly lower 30-day readmission or death rate of 3.85% (6 out of 156) compared to 11.54% (18 out of 156) in the usual care matched group (P = 0.023) (Hall et al, 2014). Educating specialized nurses on self-management led to a reduction in health service utilization ($\chi^2 = 28.03$, P < 0.001, OR = 0.016, Cohen's d = 0.48).⁶¹ Education on medication use by specialized nurses for older people in home visits reduced the odds ratios (OR) of 30-day readmissions: 0.886 (95% CI: 0.808–0.971, P = 0.010), 90-day (OR: 0.913; 95% CI: 0.843–0.989, P = 0.026), and 180day readmissions (OR: 0.911, 95% CI: 0.847–0.980, P = 0.013).⁴⁵ Medication prescriptions by specialized nurses in home care caused no older people to go to residential aged care facilities or hospitals after it.⁵² Medication verification by sending photos to specialized nurses with daily medical specialist consultations eliminated the need for home visits and saved 847 hospital days for older people.⁶³ Medication prescriptions for end-of-life care projects by specialized nurses resulted in 20% of older adults dving at home instead of hospitals or remaining at home longer than usual.⁵² Ambulatory intravenous antibiotic and diuretic treatment by specialized nurses reduced hospital stays by 847 and 201 days for endocarditis and heart failure, respectively, and reduced the average length of stay by 25 days for older people with endocarditis without major cardiovascular events.⁶³ A clinical pathway for medication management led by specialized nurses for cardiac older people enrolling 22 older people reduced the 30-day readmission rates to 9%, which is well below the national average of 23%.⁴⁶

Their involvement in home care reduced physicians' workload for home visits due to alarms about medication, worsening statements (n=107), and self-care advice (n=47), leading to general practitioners' satisfaction (90% average score)⁵⁶ and appreciation of nurses' advice.⁵² The identification of medication issues by specialized nurses in home care led to early intervention and 28 hospital admissions.⁵⁶ Physician satisfaction was high, averaging 9/10, with all attendees recommending the service to other practices in the city, noting reduced stress, and improved capacity. With a low referral rejection rate, specialized nurses discussed the potential to expand the service to more practices.⁵⁶ The involvement of a specialized nurse prescriber streamlined and sped up the prescription process, especially when delays caused medication shortages at older people's usual pharmacies, additional medications were needed for holidays, or prescribed treatments were ineffective. Coordinated care, such as contacting pharmacists or physicians through Email or phone calls, ensures that older people have timely access to necessary medications.³³

Discussion

This review aimed to identify and describe the contributions of specialized nurses to medication management for older people in home care. According to the review results, specialized nurses assume multiple roles in medication management within home care. In addition, their contribution to medication management in home care influenced older people's quality of life, emergency department visits, burden of care for older people and caregivers, other healthcare providers, and the healthcare system and helped with the detection and resolution of medication issues and management of symptoms. The roles and responsibilities of specialized nurses in home care services have been emphasized in the international literature. Specialized nurses play a vital role in improving safety in home care by addressing potential issues with medications used at home, leading to better adherence to the therapeutic regimen and a reduced risk of adverse events.⁶⁴ Specialized nurses are recognized as key members of the multidisciplinary medication team, offering numerous benefits to the healthcare system in terms of care quality and overall patient outcomes. For example, the new multidisciplinary team systemic anti-cancer therapy clinic for neuroendocrine tumor patients in the UK, with the inclusion of CNS as a non-physician prescriber, demonstrated the positive impact of innovative clinics leveraging the expertise of specialized nurses along with the medical team.⁶⁵ In addition, a prospective safety audit of a nurse specialized-delivered injection service in an ophthalmology clinic in New Zealand revealed low incidences of postinjection complications, such as endophthalmitis (0.07%), vitreous hemorrhage (0.07%), and elevated intraocular pressure (0.17%). These findings indicate that services delivered by specialized nurses are safe and effective.⁶⁶ However, coordination and collaboration challenges between specialized nurses and other healthcare professionals persist due to disagreements and unclear lines of responsibility within healthcare, as indicated in our review results. Internationally, the role of specialized nurses as non-physician prescribers is evolving, with factors such as heightened accountability impacting nurses' confidence and competence.⁶⁷ Nurse empowerment involves nurses taking control of their practices and careers to advance nursing professionals and has a stronger impact on communities.⁶⁸ Multidisciplinary support is essential for empowering specialized nurses to advance healthcare services.⁶⁷ There is also a need to enhance standards for certification and regulation of these nurses, as well as the interpretation of their roles and competencies, in addition to the development of health policies that recognize their specialties.⁶⁹ This calls for optimizing their roles and scopes of practice across the entire care spectrum, from healthcare facilities to home care, alongside establishing strategic alignment and accountability frameworks within healthcare systems.⁷⁰

Our review found that specialized nursing practices were associated with a higher risk of prescribing high-risk medications and an increased likelihood of emergency department visits for older people. Regarding medications, in primary care, high-risk prescribing increased significantly with age, rising from 2.5% in patients under 40 years to 20.2% in those aged 80 years and above (adjusted OR: 8.64, 95% CI: 7.69–9.71). It is also associated with the number of chronic medications; 10.6% of patients taking 0–2 chronic drugs received high-risk prescriptions, compared to 19.0% of those taking 11 or more chronic medications (adjusted OR: 1.35, 95% CI: 1.29–1.40).⁷¹ Ineffective communication with individuals outside the core therapeutic team and organizational factors, such as including trade names in hospital formularies and on-call hours' prescriptions, create latent risks for high-risk medication on prescribing for older people to enhance safe medication practices.⁷³ In addition, quality improvement strategies such as clinical decision support to identify inappropriate prescribing and guide deprescribing protocols, performance feedback, and pharmacist-led interventions have been shown to be effective in reducing prescription errors. The growing use of electronic health records in primary care presents new opportunities to enhance medication safety and coordinate services among diverse healthcare stakeholders.^{74,75}

To our knowledge, this is the first systematic review to integrate the current international knowledge of specialized nurses' contributions to medication management in home care for older people. This review excluded materials predating 2014, which may have excluded relevant research from that period, and those in other languages. There is insufficient international literature to clearly identify the roles, responsibilities, and benefits of specialized nurses in medication management within home care. Although selection bias was minimized by including both qualitative and quantitative studies, the varied quality and diverse study designs could have influenced the integration of the findings. Further studies

are needed to identify the roles and competencies of specialized nurses within interdisciplinary teams for medication management in home care and their impact on the home care outcomes of various older adults, including physical, psychological, and social indicators.

Conclusion

This review highlights the critical role of specialized nurses in medication management within home care. Their diverse expertise, education, and skills are essential in supporting older adults and their caregivers who face chronic health challenges.

Specialized nurses assume multiple roles to ensure safe and effective medication practices. Their proactive involvement not only enhances medication safety but also significantly improves the quality of life for older adults. By reducing emergency department visits, they help alleviate the strain on healthcare systems and minimize disruptions for patients and caregivers.

Addressing the unique needs of older adults, specialized nurses provide the necessary support for effective medication management to achieve therapeutic goals. Their proficiency in monitoring and resolving medication-related issues ensures timely interventions, better symptom control, and improved health outcomes. Through a holistic approach, specialized nurses address immediate medication concerns while fostering a supportive environment that empowers older people and caregivers to manage chronic conditions at home.

The positive impact of their practice underscores the need for continued investment in this field. By recognizing and promoting the contributions of specialized nurses, healthcare systems can optimize medication management, improve home care for older adults, and reduce the burden on healthcare resources. Supporting specialized nurses by clarifying their roles and responsibilities will help ensure high-quality care for older adults in their homes.

Policies are needed to translate the positive impacts of specialized nursing on medication management into sustainable outcomes for home care. Priority should be given to developing funding models that explicitly reflect the long-term cost savings achieved through specialized nurses' involvement in home care, including reductions in emergency department visits, hospitalizations, and the overall strain on the healthcare system.

Data Sharing Statement

Data supporting the findings of this review are available upon reasonable request from the corresponding author.

Funding

The authors received no financial support for the research, authorship, or publication of this manuscript.

Disclosure

The authors declare that there is no conflicts of interest in this work.

References

- 1. International Council of Nurses (ICN). Guidelines on advanced practice nursing. Available from: https://www.icn.ch/system/files/documents/2020-04/ICN_APN%20Report_EN_WEB.pdf. Accessed November 1, 2024.
- 2. Dowling M, Beauchesne M, Farrelly F, Murphy K. Advanced practice nursing: a concept analysis. Int J Nurs Pract. 2013;19(2):131-140. doi:10.1111/ijn.12050
- 3. Lee G, Hendriks J, Deaton C. Advanced nursing practice across Europe: work in progress. Eur J Cardiovasc Nurs. 2020;19(7):561–563. doi:10.1177/1474515120917626
- 4. Leary A, Maclaine K, Trevatt P, Radford M, Punshon G. Variation in job titles within the nursing workforce. J Clin Nurs. 2017;26(23-24):4945-4950. doi:10.1111/jocn.13985
- 5. Mayo AM, Ray MM, Chamblee TB, Urden LD, Moody R. The advanced practice clinical nurse specialist. Nurs Adm Q. 2017;41(1):70-76. doi:10.1097/naq.00000000000201
- Parker JM, Hill MN. A review of advanced practice nursing in the United States, Canada, Australia and Hong Kong Special Administrative Region (SAR), China. Int J Nurs Sci. 2017;4(2):196–204. doi:10.1016/j.ijnss.2017.01.002
- McCleery E, Christensen V, Peterson K, Humphrey L, Helfand M. VA evidence synthesis program reports. In: Evidence Brief: The Quality of Care Provided by Advanced Practice Nurses. Department of Veterans Affairs (US); 2014.
- Bentley M, Minstrell M, Bucher H, Sproule L, Robinson A, Stirling C. Aged care nurse practitioners working in general practice. J Clin Nurs. 2015;24(23–24):3745–3747. doi:10.1111/jocn.13066

- 9. Hako L, Turunen H, Jokiniemi K. Advanced practice nurse capabilities: a mixed methods systematic review. *Scand J Caring Sci.* 2023;37(1):3–19. doi:10.1111/scs.13134
- Swan M, Ferguson S, Chang A, Larson E, Smaldone A. Quality of primary care by advanced practice nurses: a systematic review. Int J Qual Health Care. 2015;27(5):396–404. doi:10.1093/intqhc/mzv054
- 11. Donald F, Martin-Misener R, Carter N, et al. A systematic review of the effectiveness of advanced practice nurses in long-term care. J Adv Nurs. 2013;69(10):2148–2161. doi:10.1111/jan.12140
- 12. GBD. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet*. 2024;403(10440):2133–2161. doi:10.1016/s0140-6736(24)00757-8
- 13. Chang M, Michelet M, Skirbekk V, et al. Trends in the use of home care services among Norwegians 70+ and projections towards 2050: the HUNT study 1995-2017. *Scand J Caring Sci.* 2023;37(3):752–765. doi:10.1111/scs.13158
- 14. Kivimäki T, Stolt M, Charalambous A, Suhonen R. Safety of older people at home: an integrative literature review. *Int J Older People Nurs*. 2020;15(1):e12285. doi:10.1111/opn.12285
- 15. Mah JC, Stevens SJ, Keefe JM, Rockwood K, Andrew MK. Social factors influencing utilization of home care in community-dwelling older adults: a scoping review. *BMC Geriatr.* 2021;21(1):145. doi:10.1186/s12877-021-02069-1
- 16. Duffy S, Paterson R. Exploring the benefits of structured medication reviews for frail older patients in advanced clinical practice. *Br J Nurs*. 2021;30(15):894–898. doi:10.12968/bjon.2021.30.15.894
- 17. Cotton BP, Lohman MC, Brooks JM, et al. Prevalence of and factors related to prescription opioids, benzodiazepines, and hypnotics among medicare home health recipients. *Home Healthc Now.* 2017;35(6):304–313. doi:10.1097/nhh.0000000000553
- Elliott RA, Lee CY, Beanland C, Vakil K, Goeman D. Medicines management, medication errors and adverse medication events in older people referred to a community nursing service: a retrospective observational study. *Drugs Real World Outcomes*. 2016;3(1):13–24. doi:10.1007/s40801-016-0065-6
- Sánchez-Fidalgo S, Guzmán-Ramos MI, Galván-Banqueri M, Bernabeu-Wittel M, Santos-Ramos B. Prevalence of drug interactions in elderly patients with multimorbidity in primary care. Int J Clin Pharm. 2017;39(2):343–353. doi:10.1007/s11096-017-0439-1
- 20. Sheikh-Taha M, Asmar M. Polypharmacy and severe potential drug-drug interactions among older adults with cardiovascular disease in the United States. *BMC Geriatr.* 2021;21(1):233. doi:10.1186/s12877-021-02183-0
- 21. Uitvlugt EB, Janssen MJA, Siegert CEH, et al. Patients' and providers' perspectives on medication relatedness and potential preventability of hospital readmissions within 30 days of discharge. *Health Expect.* 2020;23(1):212–219. doi:10.1111/hex.12993
- 22. Wasan T, Hayhoe B, Cicek M, et al. The effects of community interventions on unplanned healthcare use in patients with multimorbidity: a systematic review. J R Soc Med. 2024;117(1):24–35. doi:10.1177/01410768231186224
- 23. Azubike N, Moseley M, Powers JS. Opioid management in older adults: lessons learned from a geriatric patient-centered medical home. *Fed Pract.* 2021;38(4):168–173. doi:10.12788/fp.0110
- 24. Fujita T, Iwaki M, Hatono Y. The role of nurses for patients with Parkinson's disease at home: a scoping review. *BMC Nurs.* 2024;23(1):318. doi:10.1186/s12912-024-01931-y
- 25. Garrard JW, Cox NJ, Dodds RM, Roberts HC, Sayer AA. Comprehensive geriatric assessment in primary care: a systematic review. *Aging Clin Exp Res.* 2020;32(2):197–205. doi:10.1007/s40520-019-01183-w
- 26. Noyes J, Booth A, Moore G, Flemming K, Tunçalp Ö, Shakibazadeh E. Synthesising quantitative and qualitative evidence to inform guidelines on complex interventions: clarifying the purposes, designs and outlining some methods. *BMJ Glob Health*. 2019;4(Suppl 1):e000893. doi:10.1136/ bmjgh-2018-000893
- 27. Whittemore R, Knafl K. The integrative review: updated methodology. J Adv Nurs. 2005;52(5):546-553. doi:10.1111/j.1365-2648.2005.03621.x
- Royal College of Nursing. Medicines management an overview for nursing: clinical professional resource. Available from: https://www.rcn.org.uk/ Professional-Development/publications/pub-009018. Accessed November 1, 2024.
- 29. Hong QN, Pluye P, Fàbregues S, et al. Mixed Methods Appraisal Tool (MMAT), version 2018. Available from: http://mixedmethodsappraisaltool public.pbworks.com/w/file/fetch/127916259/MMAT_2018_criteria-manual_2018-08-01_ENG.pdf. Accessed November 1, 2024.
- 30. Risk of bias tools. Risk of bias tools for use in systematic reviews. Available from: www.riskofbias.info. Accessed November 1, 2024.
- 31. Hong QN, Pluye P, Bujold M, Wassef M. Convergent and sequential synthesis designs: implications for conducting and reporting systematic reviews of qualitative and quantitative evidence. *Syst Rev.* 2017;6(1):61. doi:10.1186/s13643-017-0454-2
- 32. Bagchi A, Melamed B, Yeniyurt S, Holzemer W, Reyes D. Telemedicine delivery for urban seniors with low computer literacy: a pilot study citation. *Online J Nurs Inf.* 2018;22(2). doi:10.1007/s11036-013-0456-9
- 33. Campling N, Birtwistle J, Richardson A, et al. Access to palliative care medicines in the community: an evaluation of practice and costs using case studies of service models in England. Int J Nurs Stud. 2022;132:104275. doi:10.1016/j.ijnurstu.2022.104275
- 34. Chan B, Goldman LE, Sarkar U, et al. The effect of a care transition intervention on the patient experience of older multi-lingual adults in the safety net: results of a randomized controlled trial. J Gen Intern Med. 2015;30(12):1788–1794. doi:10.1007/s11606-015-3362-y
- 35. Datta R, Kiwak E, Fried TR, et al. Diagnostic uncertainty and decision-making in home-based primary care: a qualitative study of antibiotic prescribing. J Am Geriatr Soc. 2024;72(5):1468–1475. doi:10.1111/jgs.18778
- 36. Ernecoff NC, Altieri-Dunn SC, Bilderback A, et al. Evaluation of a home-based, nurse practitioner-led advanced illness care program. J Am Med Dir Assoc. 2021;22(11):2389–2393. doi:10.1016/j.jamda.2021.05.013
- 37. Fisher NDL, Fera LE, Dunning JR, et al. Development of an entirely remote, non-physician led hypertension management program. *Clin Cardiol.* 2019;42(2):285–291. doi:10.1002/clc.23141
- 38. Goldman LE, Sarkar U, Kessell E, et al. Support from hospital to home for elders: a randomized trial. Ann Intern Med. 2014;161(7):472-481. doi:10.7326/m14-0094
- 39. Guo F, Lin YL, Raji M, Leonard B, Chou LN, Kuo YF. Processes and outcomes of diabetes mellitus care by different types of team primary care models. PLoS One. 2020;15(11):e0241516. doi:10.1371/journal.pone.0241516
- 40. Hall MH, Esposito RA, Pekmezaris R, et al. Cardiac surgery nurse practitioner home visits prevent coronary artery bypass graft readmissions. *Ann Thorac Surg.* 2014;97(5):1488–93;discussion1493–5. doi:10.1016/j.athoracsur.2013.12.049

- 41. Hernandez J. Medication management in the older adult: a narrative exploration. J Am Assoc Nurse Pract. 2017;29(4):186–194. doi:10.1002/2327-6924.12427
- 42. Kanne GE, McConnell ES, Disco ME, et al. The interagency care team: a new model to integrate social and medical care for older adults in primary care. *Geriatr Nurs*. 2023;50:72–79. doi:10.1016/j.gerinurse.2022.12.008
- 43. Kulsick C, Votta J, Wright WL, White P, Strowman S. Enhancing medication adherence in older adults at two nurse practitioner-owned clinics. J Am Assoc Nurse Pract. 2020;33(7):553–562. doi:10.1097/jxx.00000000000414
- 44. Levine AMP, Emonds EE, Smith MA, et al. Pharmacist identification of medication therapy problems involving cognition among older adults followed by a home-based care team. *Drugs Aging*. 2021;38(2):157–168. doi:10.1007/s40266-020-00821-7
- 45. Misra-Hebert AD, Rothberg MB, Fox J, et al. Healthcare utilization and patient and provider experience with a home visit program for patients discharged from the hospital at high risk for readmission. *Healthc.* 2021;9(1):100518. doi:10.1016/j.hjdsi.2020.100518
- 46. Moore JA. Evaluation of the efficacy of a nurse practitioner-led home-based congestive heart failure clinical pathway. *Home Health Care Serv Q*. 2016;35(1):39–51. doi:10.1080/01621424.2016.1175992
- 47. Possin KL, Merrilees JJ, Dulaney S, et al. Effect of Collaborative Dementia care via telephone and internet on quality of life, caregiver well-being, and health care use: the care ecosystem randomized clinical trial. JAMA Intern Med. 2019;179(12):1658–1667. doi:10.1001/jamainternmed.2019.4101
- 48. Quist KK, Counsell SR, Schubert CC, Weiner M. Medication management interventions in patients enrolled in GRACE Team Care. *Geriatr Nurs*. 2016;37(5):371–375. doi:10.1016/j.gerinurse.2016.05.001
- 49. Rose T, Frith K, Zimmer R. Transitional care following a skilled nursing facility stay: utilization of nurse practitioners to reduce readmissions in high risk older adults. *Geriatr Nurs*. 2021;42(6):1594–1596. doi:10.1016/j.gerinurse.2021.06.024
- 50. Squires A, Ridge L, Miner S, McDonald MV, Greenberg SA, Cortes T. Provider perspectives of medication complexity in home health care: a qualitative secondary data analysis. *Med Care Res Rev.* 2020;77(6):609–619. doi:10.1177/1077558719828942
- Warren C, Lemieux AA, Phoenix Bittner N. Excellence in population health: a successful community-based care transitions program model. Prof Case Manag. 2019;24(1):39–45. doi:10.1097/ncm.0000000000303
- 52. Mitchell GK, Senior HE, Bibo MP, et al. Evaluation of a pilot of nurse practitioner led, GP supported rural palliative care provision. *BMC Palliat Care*. 2016;15(1):93. doi:10.1186/s12904-016-0163-y
- Pond D, Higgins I, Mate K, Merl H, Mills D, McNeil K. Mobile memory clinic: implementing a nurse practitioner-led, collaborative dementia model of care within general practice. Aust J Prim Health. 2021;27(1):6–12. doi:10.1071/py20118
- 54. Antunes B, Bowers B, Winterburn I, et al. Anticipatory prescribing in community end-of-life care in the UK and Ireland during the COVID-19 pandemic: online survey. *BMJ Support Palliat Care*. 2020;10(3):343–349. doi:10.1136/bmjspcare-2020-002394
- 55. Dawson A, Reddecliffe S, Coghlan C, Schreiber BE, Coghlan JG. Safety and feasibility audit of a home-based drug-transitioning approach for patients with pulmonary arterial hypertension: an observational study. *Eur J Cardiovasc Nurs.* 2018;17(7):612–618. doi:10.1177/ 1474515118769534
- 56. Leask CF, Tennant H. Evaluation of an unscheduled care model delivered by advanced nurse practitioners in a primary-care setting. *J Res Nurs*. 2019;24(8):696–709. doi:10.1177/1744987119852380
- 57. Edge DS, Tranmer JE, Wei X, VanDenKerkhof EG. A population profile of older adults with prescription encounters with nurse practitioners and family physicians in Ontario: a descriptive retrospective cohort study. *CMAJ Open*. 2019;7(2):E323–e332. doi:10.9778/cmajo.20190007
- Scott MM, Ramzy A, Isenberg SR, et al. Nurse practitioner and physician end-of-life home visits and end-of-life outcomes. BMJ Support Palliat Care. 2023. doi:10.1136/spcare-2023-004392
- Wong S, Montoya L, Quinlan B. Transitional care post TAVI: a pilot initiative focused on bridging gaps and improving outcomes. *Geriatr Nurs*. 2018;39(5):548–553. doi:10.1016/j.gerinurse.2018.03.003
- 60. Frain S, Chambers L, Higgins A, Donohue G. 'Not left in limbo': service user experiences of mental health nurse prescribing in home care settings. *Issues Ment Health Nurs*. 2021;42(7):660–666. doi:10.1080/01612840.2020.1820120
- 61. Savini S, Iovino P, Monaco D, et al. A family nurse-led intervention for reducing health services' utilization in individuals with chronic diseases: the ADVICE pilot study. Int J Nurs Sci. 2021;8(3):264–270. doi:10.1016/j.ijnss.2021.05.001
- 62. Lee S, Lee JY, Kim H, Lee K, Lee T. Advanced practice nurses' experiences on patient safety culture in hospital-based home healthcare: a qualitative descriptive study. *Risk Manag Healthc Policy*. 2022;15:2297–2309. doi:10.2147/rmhp.S388902
- 63. van Ramshorst J, Duffels M, de Boer SPM, et al. Connected care for endocarditis and heart failure patients: a hospital-at-home programme. *Neth Heart J*. 2022;30(6):319–327. doi:10.1007/s12471-021-01614-2
- 64. Vaismoradi M, Rae J, Turunen H, Logan PA. Specialized nurses' role in ensuring patient safety within the context of telehealth in home care: a scoping review. *Digit Health*. 2024;10:20552076241287272. doi:10.1177/20552076241287272
- 65. Hand Née Davies PR. Non-medical prescribing of systemic anticancer therapy in a multidisciplinary team oncology clinic. Br J Nurs. 2019;28 (11):715–720. doi:10.12968/bjon.2019.28.11.715
- 66. Samalia P, Garland D, Squirrell D. Nurse specialists for the administration of anti-vascular endothelial growth factor intravitreal injections. N Z Med J. 2016;129(1438):32–38.
- 67. Osborne J, Kerr H. Role of the clinical nurse specialist as a non-medical prescriber in managing the palliative care needs of individuals with advanced lung cancer. Int J Palliat Nurs. 2021;27(4):205–212. doi:10.12968/ijpn.2021.27.4.205
- 68. Woodward KF. Individual nurse empowerment: a concept analysis. Nurs Forum. 2020;55(2):136-143. doi:10.1111/nuf.12407
- 69. Dury C, Hall C, Danan JL, et al. Specialist nurse in Europe: education, regulation and role. Int Nurs Rev. 2014;61(4):454-462. doi:10.1111/ inr.12123
- Jeffs L, Kuluski K, Law M, et al. Identifying effective nurse-led care transition interventions for older adults with complex needs using a structured expert panel. Worldviews Evid Based Nurs. 2017;14(2):136–144. doi:10.1111/wvn.12196
- 71. Byrne CJ, Cahir C, Curran C, Bennett K. High-risk prescribing in an Irish primary care population: trends and variation. *Br J Clin Pharmacol*. 2017;83(12):2821–2830. doi:10.1111/bcp.13373
- Alanazi MA, Tully MP, Lewis PJ. Prescribing errors by junior doctors- A comparison of errors with high risk medicines and non-high risk medicines. *PLoS One*. 2019;14(1):e0211270. doi:10.1371/journal.pone.0211270

- Poot B, Nelson K, Zonneveld R, Weatherall M. Potentially inappropriate medicine prescribing by nurse practitioners in New Zealand. J Am Assoc Nurse Pract. 2020;32(3):220–228. doi:10.1097/jxx.0000000000239
- 74. Dreischulte T, Guthrie B. High-risk prescribing and monitoring in primary care: how common is it, and how can it be improved? *Ther Adv Drug Saf*. 2012;3(4):175–184. doi:10.1177/2042098612444867
- 75. Rochon PA, Petrovic M, Cherubini A, et al. Polypharmacy, inappropriate prescribing, and deprescribing in older people: through a sex and gender lens. *Lancet Healthy Longev*. 2021;2(5):e290–e300. doi:10.1016/s2666-7568(21)00054-4

Risk Management and Healthcare Policy



Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/risk-management-and-healthcare-policy-journal