

# Is High-Intensity Interval Training an Option for Post-Kidney Transplant Physical Rehabilitation Programmes? A Scoping Review

Viviane Lovatto<sup>1,2,\*</sup>, Bruna da Silva Sousa<sup>1,\*</sup>, Vera Regina Fernandes da Silva Marães<sup>1,3</sup>

<sup>1</sup>Postgraduate Programme in Health Sciences and Technologies, University of Brasília, Ceilândia, Brazil; <sup>2</sup>Faculty of Physiotherapy, University of Rio Verde, Rio Verde, Brazil; <sup>3</sup>Postgraduate Programme in Biomedical Engineering, University of Brasília, Gama, Brazil

\*These authors contributed equally to this work

Correspondence: Viviane Lovatto, Faculty of Physiotherapy, University of Rio Verde, Fazenda Fontes Do Saber - Campus Universitário, Mailbox: 104 - Zip Code 75901-970, Rio Verde, Brazil, Email [vivianelovatto@unirv.edu.br](mailto:vivianelovatto@unirv.edu.br)

**Purpose:** To systematically and broadly review the literature to show the available information on high-intensity interval training for kidney transplant recipients as an adjunct to physical rehabilitation.

**Methods:** A scoping review of high-intensity interval training for post-kidney transplant patients was conducted by searching the PubMed, EMBASE (Elsevier), Scopus (Elsevier), Web of Science, and PEDro databases. Full-text records on the subject were included. Articles not published in English were excluded. The selected articles went through careful production quality analysis using the PEDro scale.

**Results:** The search identified 26 articles, 3 of which met the inclusion criteria. The material demonstrated satisfaction, confidence, improvement in resting heart rate, and absence of adverse effects from high-intensity interval training for kidney transplant recipients.

**Conclusion:** Based on this scoping review, high-intensity interval training for kidney transplant patients may be beneficial for physical and mental aspects and complement physical rehabilitation programmes, but there is a need for more studies with robust samples and long-term follow-up to confirm these benefits.

**Plain Language Summary:** Patients undergoing kidney transplantation are at increased risk of developing heart disease, which can result in a high risk of health complications and even death. Physical rehabilitation programmes, especially those that include exercise, have demonstrated physical capacity benefits for several populations, including organ transplant recipients. Researchers have recently explored high-intensity interval training as an option in rehabilitation programmes, but they still lack a clear understanding of its specific effects on kidney transplant patients. This study aimed to systematically review the available literature on high-intensity interval training for kidney transplant recipients, evaluating its potential in physical rehabilitation. The team reviewed several databases and, after analysis, identified three relevant articles. These articles showed that high-intensity interval-training can improve cardiac health, can enhance patient satisfaction and confidence, and does not cause any physical damage. In conclusion, high-intensity interval training appears to be beneficial for kidney transplant patients and can be included in rehabilitation programmes. However, additional studies with larger sample sizes and long-term follow-up are needed to confirm these results.

**Keywords:** cardiac rehabilitation, exercise, chronic renal insufficiency, organ transplantation, evidence-based practice

## Introduction

Kidney transplantation is the gold standard treatment for end-stage kidney disease. Kidney transplant recipients have a significantly increased cardiovascular mortality rate compared with the general population, although lower than patients on maintenance dialysis.<sup>1</sup> Nevertheless, short-term survival among kidney transplant recipients has improved considerably due to decreased mortality from infections and acute organ rejections.<sup>2</sup>

Cardiovascular disease continues to be the leading cause of death among individuals with functioning grafts worldwide.<sup>3–5</sup> The high prevalence of cardiovascular disease in kidney transplant recipients is partially linked to the presence of traditional cardiovascular risk factors, such as diabetes, dyslipidaemia, and hypertension.<sup>6</sup> There are also post-transplant factors that contribute to cardiovascular risk, such as new-onset diabetes,<sup>7</sup> the development of metabolic syndrome,<sup>8</sup> and a sedentary lifestyle.<sup>9</sup> Most kidney transplant recipients do not reach the levels of physical activity<sup>10</sup> recommended by the Kidney Disease: Improving Global Outcomes (KDIGO) guidelines<sup>11</sup> and societies such as the National Kidney Foundation<sup>12</sup> and the Brazilian Association of Organ Transplantation<sup>13</sup> in their usual routine. Hence, these individuals could benefit from personalised social and professional guidance and support to improve their daily physical activity.<sup>14</sup>

Exercise-based rehabilitation programmes are known to mitigate cardiovascular risk factors in the general population,<sup>15</sup> and they are also expected to have beneficial effects for kidney transplant recipients. Although the effects of such programmes have been extensively studied in heart and lung transplant recipients due to the direct influence of physical exercise on cardiac and pulmonary function,<sup>16,17</sup> uncertainties persist regarding their efficacy and safety for recipients of other solid organs.<sup>16,18</sup>

Among the various modalities of physical exercise investigated in contemporary studies, high-intensity interval training stands out. This regimen involves alternating sets comprising short or prolonged bursts (ranging from 30 seconds to 4 minutes) of high-intensity effort (> 85% maximum oxygen uptake [VO<sub>2</sub>max]), interspersed with brief or extended recovery periods (also ranging from 30 seconds to 4 minutes).<sup>19</sup> High-intensity interval training has demonstrated utility, safety, and feasibility in heart transplant recipients<sup>20</sup> and, more recently, in kidney transplant recipients.<sup>21</sup> However, there is a dearth of evidence regarding the frequency, duration, benefits, and safety of high-intensity interval training specifically for kidney transplant recipients.

Based on this knowledge gap, the aim of this scoping review was to identify studies elucidating the potential use of high-intensity interval training for kidney transplant patients within physical rehabilitation programmes. A scoping review enables a comprehensive examination of the existing literature and facilitates the exploration of knowledge gaps. Therefore, this review serves as an initial step in data collection, providing a foundation for future research endeavours.

## Materials and Methods

### Protocol and Logging

We conducted a scoping review to identify and delineate the existing evidence base, adhering to the guidelines outlined in the JBI Reviewer's Manual on Scoping Reviews.<sup>22</sup> The reporting of this review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews (PRIMAS-ScR), as outlined by Tricco et al.<sup>23</sup> To ensure reproducibility and transparency, we outlined the methodology employed in this study in a protocol that is freely accessible on the Open Science Framework (OSF).<sup>24</sup> Because our research did not involve the use of human data, submission to the Human Research Ethics Committee was unnecessary.

### Eligibility Criteria

We established the eligibility criteria following the Population, Concept, and Context (PCC) framework outlined in the JBI Manual.<sup>22</sup> The study population comprised adult kidney transplant recipients. To address the concept, we included articles employing high-intensity interval training, and we designated the context as rehabilitation centred on physical exercise. We included the following study designs: clinical trials, cross-sectional analyses, and case-control studies.

This scoping review was limited to articles published between 2018 and 2023 and written in English to increase the feasibility and quality. The exclusion criteria were: (1) articles published before 2018, (2) articles published in any language other than English, (3) articles that did not relate to adult kidney transplant recipients, and (4) articles that did not address high-intensity training as an approach to exercise-based rehabilitation.

## Sources of Information and Search Strategy

To identify potentially relevant documents, we initially searched PubMed (National Library of Medicine) using the following Medical Subject Headings (MeSH) associated with the Boolean operators OR and AND, “transplantation recipients”, “kidney”, “high-intensity interval training”, and “rehabilitation exercise”. Then, we searched EMBASE (Elsevier), Scopus (Elsevier), Web of Science, and PEDro. We conducted the literature search in September 2023 and updated it in April 2024. [Table S1](#) contains the complete search strategy.

## Article Selection

Two authors selected the titles and abstracts of the documents independently. VL performed the data mapping process and BSS reviewed it. Disagreements between the reviewers were dealt with by consensus and, if there was no agreement, a final decision was made by a third reviewer (VRSFM).

The following data were extracted: title, authors, year of publication, country of data collection, study design, sample details (eg, sample size, biological sex, age, and kidney disease-related information), the primary and secondary endpoints (where applicable), protocol details, and the results.


We did not evaluate the quality of the evidence of the included articles, because a scoping review does not aim to critically assess the risk of bias. However, we did assess the methodological quality of the articles by using the PEDro scale. This assessment tool allowed us to evaluate the methodological construction and the results presented by the authors based on the characteristics of randomised clinical trials. The PEDro scale includes 11 criteria, 10 of which be scored.<sup>25</sup>

1. the eligibility criteria were specified;
2. the subjects were randomly distributed into groups (in a crossover study, the subjects were randomly placed in groups according to the treatment received);
3. the subject allocation was concealed;
4. the groups were similar regarding the most important prognostic indicators;
5. all subjects participated blindly in the study;
6. all therapists who administered the therapy did so blindly;
7. all evaluators who measured at least one key result did so blindly;
8. the measurements of at least one key result were obtained in more than 85% of the subjects initially distributed among the groups;
9. all subjects from whom outcome measurements were presented received the treatment or control condition according to allocation or, when this was not the case, at least one of the outcomes was analysed based on “intention to treat”;
10. the results of inter-group statistical comparisons were described for at least one key outcome; and
11. the study presented both precision measures and variability measures for at least one key outcome.

The author with the greatest expertise in methodological evaluation tools performed this evaluation; in case of divergence, a second author was consulted for quality analysis.

## Results

A search of the PubMed, Embase, Web of Science, Scopus, and PEDro databases returned 26 records. Following the removal of duplicates, we reviewed the titles and abstracts of 15 articles. Of these, we excluded nine articles that addressed solid organ transplantation without specifying the transplanted organ; excluded kidney transplant patients; did not use high-intensity interval training as an exercise therapy; or referred to other types of publications, such as recommendations. Finally, we evaluated five full texts for eligibility. As shown in the PRISMA flow diagram<sup>18</sup> ([Figure 1](#)), we included three articles that met the inclusion criteria.



**Figure 1** PRISMA flow diagram illustrating the process of study selection.  
**Notes:** Adapted from PRISMA Flow Diagram. Adapted from Page MJ, McKenzie JE, Bossuyt PM et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.<sup>23</sup>

The three articles were published between 2021 and 2022. In terms of study design, one article utilised a convenience sample,<sup>21</sup> while two employed randomisation.<sup>26,27</sup> Geographically, two of the articles originated from the United Kingdom and the other was from Belgium. In one article,<sup>21</sup> the sample included recipients of various organs such as the heart, lungs, liver, and kidneys. The data extracted from the included articles are summarised in [Table 1](#) and [Table 2](#) presents the evaluation of the articles based on the PEDro scale.

**Table 1** Summary of the Literature Included

Author/Year	Local	Study Design	Transplanted Organ
Cappelle M et al 2021 <sup>21</sup>	Belgium	Single-arm intervention	Heart, lung, liver, and kidneys
Billany RE et al 2022 <sup>26</sup>	United Kingdom	Randomized	Renal only
Hutchinson GM et al 2022 <sup>27</sup>	United Kingdom	Randomized	Renal only

**Table 2** Assessment of Methodological Quality Using the PEDRO Scale

Autor	Crit. 1	Crit. 2	Crit. 3	Crit. 4	Crit. 5	Crit. 6	Crit. 7	Crit. 8	Crit. 9	Crit. 10	Crit. 11
Cappelle M et al, 2021 <sup>21</sup>	YES	NO	NO	NO	NO	NO	NO	NO	YES	YES	YES
Billany RE et al, 2022 <sup>26</sup>	YES	YES	NO	YES	NO	NO	NO	NO	NO	YES	YES
Hutchinson GM et al 2022 <sup>27</sup>	YES	YES	NO	NO	NO	NO	NO	YES	YES	YES	YES

**Abbreviation:** Crit, Criterion.  
**Source:** Own preparation, 2024.

**Table 3** Describes the Objectives and Protocols of High-Intensity Interval Training Applied by the Studies, as Well as the Primary and Secondary Outcomes

Author/Year	Cappelle M et al 2021 <sup>21</sup>	Billany RE et al 2022 <sup>26</sup>	Hutchinson GM et al 2022 <sup>27</sup>
Goal	To investigate the effects of a HIIT program on physical performance in solid organ recipients	Determine the feasibility and acceptability of HIIT in RxT	To explore the physiology and immunological impact of eight weeks of HIIT on RxT recipients
Equipment	Ergonomics for lower limbs	Ergonomics for lower limbs	Ergonomics for lower limbs
Intensity	80–95% of HRmax	80% of peak VO <sub>2</sub>	> 90% peak VO <sub>2</sub>
HIIT Protocol	3 times a week, high-intensity intervals of up to 2 minutes at 80–95% of maximum HR, training lasting 30 minutes for 6 months	Warm-up 5 minutes and cool-down 10 minutes; 2 HIIT protocols: HIIT (16 minutes with intervals of 4.2 and 1-minute duration at 80% of peak VO <sub>2</sub> for 24 sessions (8 weeks) total session time 30 minutes; HIIT B (4x4 minutes interval training 80% VO <sub>2</sub> peak, separated by 3-minute active rest at 60% of peak VO <sub>2</sub> ) Total session time 30 minutes	HIIT A (4, 2 and 1 min intervals; 80–90% VO <sub>2</sub> peak), HIIT B (4 × 4 min intervals; 80–90% VO <sub>2</sub> peak) for 24 supervised stationary bike sessions (approx. 3x/week for 8 ± 2 weeks)
Primary outcome	Increase in VO <sub>2</sub> max and decrease in resting HR	HIIT has been shown to be safe and viable in RxT, with no adverse effects.	There was no impairment in immunity in the short or long-term
Secondary outcome	HIIT is safe and may result in a beneficial effect on the physical performance of stable RxT.	HIIT was well accepted and tolerated by RxT.	HIIT did not cause immunological changes

**Abbreviations:** HIIT, high intensity interval training; HRmax, maximum heart rate; VO<sub>2</sub>, oxygen volume; RxT, kidney transplant; HR, heart rate.

As shown in Table 3, all three articles used lower limb ergometry. In one article,<sup>21</sup> the authors determined the exercise intensity based on the maximum heart rate (HRmax), while the authors of the other two articles<sup>26,27</sup> used peak VO<sub>2</sub>. The interventions occurred three times a week and each session lasted 30 minutes for 6 months<sup>21</sup> or at least 6 weeks and at most 10 weeks.<sup>26,27</sup>

The summary of the anthropometric and clinical characteristics of all the included articles is shown in Table 4. The clinical presentation of the samples was heterogeneous: two articles provided biochemical data,<sup>26,27</sup> while only one provided information on immunosuppressive drugs.<sup>21</sup>

**Table 4** Anthropometric and Clinical Characteristics of the Study Samples

Characteristics / Study	Cappelle <sup>21</sup>	Billany <sup>26</sup>		Hutchinson <sup>27</sup>	
Single Groups		HIIT A	HIIT B	HIIT A	HIIT B
Sample, n; Man/Woman	15 (11/4)	8 (3/5)	8 (6/2)	8 (3/5)	8 (6/2)
Age, years	40.7±14.8	41±14	51±11	41±14	51±11
Body mass, kg	72.0±13.1	68.5±15.6	84.1±24	68.5±15.6	84.1±24
BMI, kg/m <sup>2</sup>	23.9±3.4	25.9±5.4	28.5±7.1	25.9±5.4	28.5±7.1
Serum creatinine, mmol/L	N/A	104±27	133±60	104±27	133±60
GFR, mL/min/1.73 m <sup>2</sup>	N/A	62 ± 18	57 ± 22	62 ± 18	57 ± 22
Transplant time, months	43,2 (24 –111,6)	12±20	44±86	12±20	44±86

(Continued)

**Table 4** (Continued).

Characteristics / Study	Cappelle <sup>21</sup>	Billany <sup>26</sup>		Hutchinson <sup>27</sup>	
Single Groups		HIIT A	HIIT B	HIIT A	HIIT B
Medication, n (%)					
Azathioprine	2 (13)	N/A	N/A	N/A	N/A
Mycophenolate mofetil	12 (80)	N/A	N/A	N/A	N/A
Cyclosporine	1 (7)	N/A	N/A	N/A	N/A
Tacrolimus	12 (80)	N/A	N/A	N/A	N/A
Steroids	8 (53)	6 (75)	2 (25)	6 (75)	2 (25)
BCC	N/A	8 (100)	8 (100)	8 (100)	8 (100)
Antihypertensive	N/A	7 (88)	7 (88)	7 (88)	7 (88)
Anti-diabetes	N/A	1 (13)	3 (38)	1 (13)	3 (38)
Statin	N/A	4 (50)	5 (63)	4 (50)	5 (63)

**Notes:** Data presented as mean and standard deviation.

**Abbreviations:** BMI, body mass index; GFR, glomerular filtration rate; SBP, systolic blood pressure; DBP, diastolic blood pressure, mmHg; millimeters of mercury; N/A, not informed; BCC, calcium channel blockers.

One of the articles acknowledged heterogeneity in the sample concerning the type of transplant, a factor that could potentially influence physical performance.<sup>26</sup> This is attributed to variations in immunosuppressive drug indications and dosages, which adhere to the standard treatment protocol for each specific type of transplanted organ. It is widely recognised that steroids exert a direct catabolic effect on skeletal muscle, thereby detrimentally affecting muscle strength and exercise capacity.

## Effects of High-Intensity Interval Training on Hemodynamic Variables

The pre- and post-intervention hemodynamic variables, systolic and diastolic pressures, and heart rate were measured in two articles and are presented in Table 5. For one article, in kidney transplant recipients, resting heart rate decreased after 6 months of high-intensity interval training compared with pre-training (baseline  $83.9 \pm 17.3$  bpm, post-training  $79.5 \pm 14.3$  bpm,  $p = 0.04$ ). However, there were no changes in resting SBP or DBP.<sup>21</sup> Billany et al<sup>26</sup> reported that there were no hemodynamic changes after high-intensity interval training. Overall, only 40% of the participants achieved the required intensity during the intervention, and it took just over half the duration of the intervention to achieve this. The data suggest that a familiarisation period and a longer intervention may be beneficial. In addition, constant absences for health

**Table 5** Hemodynamic Variables of the Samples From the Pre- and Post-Intervention Studies

Study	Capelle <sup>21</sup>		Billany <sup>26</sup>			
Group	Unique		HIIT A		HIIT B	
	Pre	Post	Pre	Post	Pre	Post
SBP mmHg	130.9 $\pm$ 37.7	138.8 $\pm$ 18.5	122 $\pm$ 12	122 $\pm$ 11	131 $\pm$ 17	122 $\pm$ 13
DBP mmHg	82.9 $\pm$ 13.8	81.3 $\pm$ 15.7	84 $\pm$ 6	80 $\pm$ 4	79 $\pm$ 6	75 $\pm$ 6
HR rest	83.9 $\pm$ 17.3	79.5 $\pm$ 14.3	77 $\pm$ 4	76 $\pm$ 8	76 $\pm$ 4	67 $\pm$ 4

**Notes:** Data presented as mean and standard deviation.

**Abbreviations:** SBP, systolic blood pressure; DBP, diastolic blood pressure; mmHg, millimeters of mercury.

reasons (colds, justified by the continuous administration of immunosuppressants) may have influenced the fact that the participants did not reach the intensity and frequency of the target exercise. Finally, the authors of the third included article did not describe hemodynamic variables because they aimed to investigate changes in circulating immune and inflammatory cells during the 8-week intervention.<sup>27</sup> The authors concluded that high-intensity interval training exercises do not adversely affect the immunity of kidney transplant recipients.

## Effects of High-Intensity Interval Training on Anthropometric Variables

Two articles<sup>21,26</sup> reported that there were no changes in body mass and the body mass index before and after the intervention with high-intensity interval training.

## Discussion

In this scoping review, we identified that there are scarce data regarding high-intensity interval training as an option for physical exercise-based rehabilitation programmes for kidney transplant recipients. Nevertheless, this modality is recommended for the healthy population.<sup>28</sup> Moreover, studies have demonstrated its benefit and feasibility for some clinical populations such as heart transplant patients<sup>29</sup> and lung transplant recipients.<sup>30</sup> The guidelines indicate that kidney transplant patients should practice regular physical exercise<sup>31</sup> but do not describe the ideal time to start after transplantation, as well as the frequency, intensity, and duration. However, a unanimous view for healthy and clinical populations is the continuous and regular performance of physical exercise with a focus on better quality of life and exercise capacity and, consequently, a reduction of the cardiovascular risk.<sup>28,32</sup>

The physiological profile of kidney transplant recipients is well known: they have experienced the burden of chronic kidney disease, such as heart failure, systemic arterial hypertension, and sarcopenia before transplantation.<sup>33–35</sup> After kidney transplantation, there is a need to make lifestyle changes to minimise acute or chronic rejection. These changes include medication adherence, body mass control, adequate diet, and regular physical exercise, whether supervised or at home.<sup>32</sup> There has been limited on high-intensity interval training as a viable option within exercise-based rehabilitation programmes. This scarcity has arisen partly due to the exclusion of kidney transplant recipients from studies or their grouping with recipients of other organ types, thus impeding precise data extraction.

## Conclusion

Exercise therapy, particularly high-intensity interval training, demonstrates well-known benefits for healthy individuals and has recently emerged as a approach for kidney transplant patients. Chronic kidney disease and the post-transplant process directly influence the body's physiology, but high-intensity interval training shows potential despite these challenges. Although studies remain limited, researchers have not identified significant events that would render this option unfeasible in rehabilitation programmes based on physical exercise. Qualified rehabilitation professionals closely monitor transplant patients during and after high-intensity interval training to minimise the risk of complications. As a result, high-intensity interval training can serve as a tool in specific cases for kidney transplant patients, promoting engagement and adherence to rehabilitation programmes.

## Acknowledgments

All authors thank you to University of Brasília who has facilitated us to make this research. This study was supported by the Dean's Office for Research and Innovation (DPI) and the Central Library (BCE) of the University of Brasília (UnB) (Call for Proposals No. 001/2025 DPI/BCE/UnB).

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Totti V, Fernhall B, Di Michele R, et al. Longitudinal analysis of cardiovascular risk factors in active and sedentary kidney transplant recipients. *Medicina*. 2020;56(4):183. doi:10.3390/medicina56040183



2. Hart A, Smith JM, Skeans MA, et al. OPTN/SRTR 2016 annual data report: kidney. *Am J Transplant.* 2018;18(Suppl 1):18–113. doi:10.1111/ajt.14557
3. Awan AA, Niu J, Pan JS, et al. Trends in the causes of death among kidney transplant recipients in the United States (1996–2014). *Am J Nephrol.* 2018;48(6):472–481. doi:10.1159/000495081
4. de SSR, da FCD, Taminato M, et al. Infectious events in kidney transplant recipients from deceased expanded criteria donors: a prospective cohort. *Rev esc enferm USP.* 2021;55:e20210255. doi:10.1590/1980-220X-REEUSP-2021-0255
5. SRTR - the scientific registry of transplant recipients [Internet]. srtr.transplant.hrsa.gov. Available from: [http://srtr.transplant.hrsa.gov/annual\\_reports/Default.aspx](http://srtr.transplant.hrsa.gov/annual_reports/Default.aspx). Accessed February 2, 2025.
6. Rangaswami J, Mathew RO, Parasuraman R, et al. Cardiovascular disease in the kidney transplant recipient: epidemiology, diagnosis and management strategies. *Nephrol Dial Transplant.* 2019;34(5):760–773. doi:10.1093/ndt/gfz053
7. Martinez Cantarin MP. Diabetes in kidney transplantation. *Adv Chronic Kidney Dis.* 2021;28(6):596–605. doi:10.1053/j.ackd.2021.10.004
8. Lee YH, Song SH, Song SH, et al. Clinical implications of changes in metabolic syndrome status after kidney transplantation: a nationwide prospective cohort study. *Nephrol Dial Transplant.* 2023;38(12):2743–2753. doi:10.1093/ndt/gfad115
9. Ponticelli C, Favi E. Physical inactivity: a modifiable risk factor for morbidity and mortality in kidney transplantation. *J Pers Med.* 2021;11(9):927. doi:10.3390/jpm11090927
10. Takahashi A, Hu SL, Bostom A. Physical activity in kidney transplant recipients: a review. *Am J Kidney Dis.* 2018;72(3):433–443. doi:10.1053/j.ajkd.2017.12.005
11. Kidney Disease: Improving Global Outcomes (KDIGO) Transplant Work Group. KDIGO clinical practice guideline for the care of kidney transplant recipients. *Am J Transplant.* 2009;9(Suppl 3):S1–S155. doi:10.1111/j.1600-6143.2009.02834.x
12. Your guide to a functional body of health [Internet]. Available from: <https://nkfs.org/wp-content/uploads/2020/05/NKF-Hues-Booklet.pdf>. Accessed February 14, 2025.
13. Post-transplant period [Internet]. Available from: [https://site.abto.org.br/wp-content/uploads/2020/06/manual\\_transplante\\_pos-1.pdf](https://site.abto.org.br/wp-content/uploads/2020/06/manual_transplante_pos-1.pdf). Accessed February 14, 2025.
14. Sturgeon KM, Fisher C, McShea G, Sullivan SK, Sataloff D, Schmitz KH. Patient preference and timing for exercise in breast cancer care. *Support Care Cancer.* 2018;26(2):507–514. doi:10.1007/s00520-017-3856-8
15. Adams V, Linke A. Impact of exercise training on cardiovascular disease and risk. *Biochim Biophys Acta Mol Basis Dis.* 2019;1865(4):728–734. doi:10.1016/j.bbdis.2018.08.019
16. Janaudis-Ferreira T, Mathur S, Konidis S, Tansey CM, Beaurepaire C. Outcomes in randomized controlled trials of exercise interventions in solid organ transplant. *World J Transplant.* 2016;6(4):774–789. doi:10.5500/wjt.v6.i4.774
17. Dall CH, Snoer M, Christensen S, et al. Effect of high-intensity training versus moderate training on peak oxygen uptake and chronotropic response in heart transplant recipients: a randomized crossover trial. *Am J Transplant.* 2014;14(10):2391–2399. doi:10.1111/ajt.12873
18. Calella P, Hernández-Sánchez S, Garofalo C, Ruiz JR, Carrero JJ, Bellizzi V. Exercise training in kidney transplant recipients: a systematic review. *J Nephrol.* 2019;32(4):567–579. doi:10.1007/s40620-019-00583-5
19. Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle: part I: cardiopulmonary emphasis. *Sports Med.* 2013;43(5):313–338. doi:10.1007/s40279-013-0029-x
20. Conceição LSR, Gois CO, Fernandes RES, et al. Effect of high-intensity interval training on aerobic capacity and heart rate control of heart transplant recipients: a systematic review with meta-analysis. *Braz J Cardiovasc Surg.* 2021;36(1):86–93. doi:10.21470/1678-9741-2019-0420
21. Cappelle M, Masschelein E, Vos R, et al. High-intensity training for 6 months safely, but only temporarily, improves exercise capacity in selected solid organ transplant recipients. *Transplant Proc.* 2021;53(6):1836–1845. doi:10.1016/j.transproceed.2021.03.040
22. Peters MDJ, Marnie C, Tricco AC, et al. Updated methodological guidance for the conduct of scoping reviews. *JBIM Evid Synth.* 2020;18(10):2119–2126. doi:10.11124/JBIES-20-00167
23. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71
24. OSF, Open Science Framework. doi: 10.17605/OSF.IO/XZHN6.
25. Verhagen AP, de Vet HC, de Bie RA, et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *J Clin Epidemiol.* 1998;51(12):1235–1241. doi:10.1016/s0895-4356(98)00131-0
26. Billany RE, Smith AC, Hutchinson GM, Graham-Brown MPM, Nixon DGD, Bishop NC. Feasibility and acceptability of high-intensity interval training and moderate-intensity continuous training in kidney transplant recipients: the PACE-KD study. *Pilot Feasibility Stud.* 2022;8(1):106. doi:10.1186/s40814-022-01067-3
27. Hutchinson GM, Cooper AM, Billany RE, Nixon DGD, Bishop NC, Smith AC. Efeito do treinamento intervalado de alta intensidade e treinamento contínuo de intensidade moderada em células linfóides, mielóides e inflamatórias em receptores de transplante renal. *Exerc Immunol Rev.* 2022;28:100–115.
28. *ACSM's Guidelines for Graded Exercise Testing and Prescription.* 6th ed. Philadelphia: Lippincott Williams & Wilkins;2000.
29. Perrier-Melo RJ, Famds F, Guimarães GV, Costa MDC. High-intensity interval training in heart transplant recipients: a systematic review with meta-analysis. *Arq Bras Cardiol.* 2018;110(2):188–194. doi:10.5935/abc.20180017
30. Langer D, Burtin C, Schepers L, et al. Exercise training after lung transplantation improves participation in daily activity: a randomized controlled trial. *Am J Transplant.* 2012;12(6):1584–1592. doi:10.1111/j.1600-6143.2012.04000.x
31. Abecassis M, Bartlett ST, Collins AJ, et al. Kidney transplantation as primary therapy for end-stage renal disease: a national kidney foundation/kidney disease outcomes quality initiative (NKF/KDOQITM) conference. *Clin J Am Soc Nephrol.* 2008;3(2):471–480. doi:10.2215/CJN.05021107
32. WHO guidelines on physical activity and sedentary behaviour. World Health Organization; 2020.
33. Moon SJ, Kim TH, Yoon SY, Chung JH, Hwang HJ. Relationship between stage of chronic kidney disease and sarcopenia in Korean aged 40 years and older using the Korea national health and nutrition examination surveys (KNHANES IV-2, 3, and V-1, 2), 2008–2011. *PLoS One.* 2015;10(6):e0130740. doi:10.1371/journal.pone.0130740
34. Delgado C, Grimes BA, Glidden DV, Shlipak M, Sarnak MJ, Johansen KL. Association of frailty based on self-reported physical function with directly measured kidney function and mortality. *BMC Nephrol.* 2015;16(1):203. doi:10.1186/s12882-015-0202-6
35. Roshanravan B, Robinson-Cohen C, Patel KV, et al. Association between physical performance and all-cause mortality in CKD. *J Am Soc Nephrol.* 2013;24(5):822–830. doi:10.1681/ASN.2012070702



## Journal of Multidisciplinary Healthcare

### Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>

**Dovepress**  
Taylor & Francis Group