

## The Concizumab Pen-Injector is Easy to Use and Preferred by Hemophilia Patients and Caregivers: A Usability Study Assessing Pen-Injector Handling and Preference [Corrigendum]

Kahr Rasmussen N, Berg B, Christiansen ASL, et al. *Patient Prefer Adherence*. 2024;18:1713–1727.

Page 1719, Table 4, Range column, Total row under the Time to prepare and inject section, the text “0 min 29s–4 min 7s” should read “0 min 29s–4 min 3s”. The correct table is as follows.

**Table 4** Measurements of training time, time to administer an injection, and the correct execution of the injection

	Average Time	Range	Median
<b>Training time</b>			
Adult patients	7 min 45s	5–13 min	7 min 0s
Adolescent patients	7 min 37s	5–12 min	7 min 0s
Caregiver participants	8 min 16s	6–15 min	8 min 0s
<b>Total</b>	<b>7 min 49s</b>	<b>5–15 min</b>	<b>7 min 0s</b>
<b>Time to prepare and inject</b>			
Adult patients	1 min 19s	0 min 33s–4 min 3s	1 min 5s
Adolescent patients	1 min 13s	0 min 29s–2 min 0s	1 min 2s
Caregiver participants	1 min 40s	1 min 8s–2 min 45s	1 min 43s
<b>Total</b>	<b>1 min 21s</b>	<b>0 min 29s–4 min 3s</b>	<b>1 min 10s</b>
<b>Completed injections</b>	<b>Number of correctly executed injections (n; %)</b>		
Adult patients	43; 98%		
Adolescent patients	20; 95%		
Caregiver participants	15; 100%		
<b>Total</b>	<b>78; 98%</b>		


**Notes:** There was a total of 78 (98%) complete injections, with only 2 participants not completing a step (1 adult and 1 adolescent). In both cases the participants did not inject the full dose on the first attempt. This may be considered a test artefact.

**Abbreviation:** n, number of participants.


Page 1719, Ease-of-Learning to Use and Ease-of-Use of the Concizumab Pen-Injector section, second line, the text “(n=78, 97%)” should read “(n=78, 98%)”.

Figure 3 on page 1720 is incorrect. The correct Figure 3 is as follows.

(A)



(B)



**Figure 3** The ease of learning and ease of use of the pen-injector as assessed using the HDHPA questionnaire. **(A)** Responses from participants to HDHPA question 1 on factors associated with ease of handling and use of concizumab pen-injector. Participants were asked to select the response that most closely represented how they felt using the pen-injector. **(B)** Responses from participants to HDHPA question 2 on overall difficulty or ease of use of the concizumab pen-injector. Responses were assessed on a 5-point scale ranging from very difficult to very easy. Responses for very difficult and difficult were combined to represent negative results and the responses easy and very easy were combined to represent positive results. Data are presented as number of responses with percentages % (n).

Page 1722, Discussion section, eighth line, the text “(97%)” should read “(98%)”.

The authors apologize for these oversights.

**Patient Preference and Adherence****Dovepress**  
Taylor & Francis Group**Publish your work in this journal**

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>