ORIGINAL RESEARCH

Association Between the Time of Exposure to Electronic Screen and Anxiety and Depression After Controlling Demographics, Exercises, and Lifestyles Among University Students

Lirong Zhang¹, Shuangyin Zhao¹, Shaocong Zhao¹, Yizhen Ke¹, Weichen Yang¹, Mingxing Lei^{2,3}

Department of Physical Education, Xiamen University of Technology, Xiamen, Fujian, People's Republic of China; ²Department of Orthopaedics, Hainan Hospital of Chinse PLA General Hospital, Sanya, Hainan, People's Republic of China; ³Nursing Department, The First Medical Center of Chinese PLA General Hospital, Beijing, People's Republic of China

Correspondence: Lirong Zhang, Department of Physical Education, Xiamen University of Technology, No. 600, Ligong Road, limei District, Xiamen, Fujian, 361024, People's Republic of China, Tel +8613806063882, Email 22674481@qq.com; Mingxing Lei, Department of Orthopedic Surgery, Hainan Hospital of Chinse PLA General Hospital, No. 80 Jianglin Road, Haitang District, Sanya, Hainan, 572013, People's Republic of China, Tel +8618811772189, Email leimingxing2@sina.com

Background: The use of electronic screens has become ubiquitous in modern society, but the specific relationship between hours of electronic screen exposure and mental health outcomes needs to be further investigated. Hence, this study aims to examine the detailed relationship between electronic screen exposure time and mental health status.

Methods: A sample of 2896 university students were recruited and their baseline characteristics, including demographics, lifestyle habits, and mental health status, were collected. The association between electronic screen exposure time and mental health status was investigated using linear regression analysis. The predictive performance of electronic screen exposure time for predicting mental health status, including severe anxiety and depression was, also analyzed using the area under the curve (AUC) analysis.

Results: The time of exposure to electronic screen was significantly associated with anxiety (p < 0.001) and depression (p < 0.001). The AUC value was 0.641 (95% CI: 0.571-0.712) for severe anxiety and 0.665 (95% CI: 0.613-0.716) for severe depression. The optimal cut-off value for predicting anxiety and depression based on electronic screen exposure time was found to be both 6.5 hours. A subgroup analysis confirmed that participants with higher exposure time to electronic screens had significantly higher anxiety (p < p(0.001) and depression scores (p < (0.001)). The association between the hours of exposure to electronic screen and severe anxiety (p = (0.042) and severe depression (p < 0.001) remained significant after controlling for demographics, exercises, and other lifestyle factors. Conclusion: The findings suggest that excessive electronic screen exposure time is associated with increased risk of anxiety and depression among university students. These findings highlight the need for interventions targeting electronic screen use to improve mental health outcomes, and less than 6.5 hours per day of exposure to electronic screen are recommended among university students. Keywords: university students, exposure to electronic screen, mental health, anxiety, depression

Introduction

The mental health of university students is a serious concern worldwide, with a significant portion experiencing mental distress, such as anxiety and depression. Studies have shown that 23.6% to 41.8% university students had anxiety,¹⁻³ and 27.2% to 51.3% had depression.^{4–6} These problems can have a devastating impact on academic performance, social life, and overall well-being among university students.^{7–9} However, the exact factors contributing to the development of these problems are not fully understood.

Identifying the risk factors associated with mental distress presents an opportunity for the prevention of anxiety and depression.¹⁰ Lifestyle factors play a crucial role in this regard, as they are easily modifiable variables.^{11,12} Among the various lifestyle factors, such as exercise routines, which has been found to be a favorable measure to

by and incorporate the Creative Commons Attribution - Non Commercial (unported, v4.0) License (http://creativecommons.org/licenses/by-nc/4.0/). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

mitigate mental distress,^{11,13} the influence of hours of exposure to electronic screens on mental distress remains unclear and warrants further investigation. While some studies have explored the association between smartphone or internet addiction and mental health problems,^{14,15} a significant gap remains in the literature concerning the specific relationship between hours of electronic screen exposure and mental health outcomes, particularly among university students. For instance, further research is needed to explore the impact of screen time on mental health, with a particular emphasis on establishing clear thresholds or cut-off values that could inform evidence-based recommendations for safe screen use.

In the modern context, university students heavily rely on computers for academic work, completing assignments, and engaging in recreational activities such as watching short videos on their smartphones. However, the specific relationship between hours of exposure to electronic screens and anxiety and depression has not been extensively studied. Given that the current generation of university students spends a significant amount of time using computers for homework, entertainment, and social media,¹⁶ it was plausible to speculate that hours of exposure to electronic screens might play a role in the development of mental health problems. By exploring this relationship, researchers can potentially identify a cut-off value associated with anxiety and depression in terms of the exposure time to electronic screen, enabling personalized recommendations for daily computer usage. This approach holds significance in promoting mental well-being among university students.

Therefore, the primary objective of this study was to delve into the detailed relationship between electronic screen exposure time and mental health status specifically among university students. Additionally, the study aimed to pinpoint both risk factors and protective factors influencing severe anxiety and depression within this population. Through this investigation, valuable insights can be gained to inform preventive strategies and interventions aimed at fostering mental health resilience among university students. We hypothesize that excessive exposure to electronic screens is significantly associated with higher levels of anxiety and depression among university students.

Methods

Participants

This study conducted a comprehensive analysis of 2896 university students from three institutions in China, namely Xiamen University of Technology, Chongqing Normal University, and North University of China, during the period spanning January 2024 to February 2024. The sampling method involved collaboration with university administrative staff and faculty members. Specifically, online questionnaires were distributed to student groups through university teacher. The research protocol was designed to ensure voluntary participation, with stringent measures in place to safeguard participant anonymity and confidentiality. Prior to their involvement, all potential participants were explicitly informed about the study's objectives and procedures, and their written informed consent was obtained. Only those who voluntarily agreed to participate were granted access to the online questionnaire.

The primary objective of this investigation was to examine the complex interplay between various psychosocial factors and mental health outcomes among the university student population. The inclusion criteria were strictly limited to currently enrolled students who voluntarily completed the survey instrument. Participants who chose not to participate or failed to complete the survey were systematically excluded from the final analysis. This approach ensured the reliability and validity of the collected data while maintaining the highest ethical standards in research involving human subjects.

The research methodology strictly adhered to the ethical principles outlined in the Declaration of Helsinki, with particular emphasis on participant autonomy and data protection. In addition, this study has adhered to the STROBE checklist to ensure transparent and comprehensive reporting of the methods and findings. The study protocol received ethical approval from the Academic Committee and Ethics Board of Xiamen University of Technology. To maintain data integrity and participant confidentiality, all collected information was anonymized, ensuring that no individual could be identified through the research data.

Sample Size Calculation

Based on literature reports, the incidence rates were set at 33% for anxiety and 39% for depression. The significance level (α) was set at 0.05, and the power (1- β) was set at 0.80. The number of predictors was approximately 20, including electronic screen exposure time, demographics, exercise, and lifestyle factors. Using the commonly applied empirical formula for sample size calculation in logistic regression:

$$n = \frac{10}{p \cdot (1-p)}$$

Applying this formula, the estimated sample size for anxiety is approximately 905 participants, while for depression, it is approximately 841 participants. This study included a total of 2,896 university students, which fully meets the requirements of the sample size calculation.

Data Collection

The study collected a wide range of individual characteristics, including (1) basic demographics such as gender, age, grade, and marital status, (2) dietary habits such as low salt food, oily food, barbecue, meat, vegetables, and fruits, (3) personal addiction such as smoking and alcohol use, (4) lifestyles such as sedentary time, and physical activity. The hours of exposure to electronic screen were also collected. The hours of exposure to electronic screen per day was a self-reported metric which quantified the total time spent using electronic devices, such as computers, smartphones, or televisions, for a particular individual during a specific day. This metric was collected using self-reporting methods, where individuals recorded their own use of electronic devices. Dietary habits were assessed through self-reporting by participants based on their dietary practices, and this approach aligns with methods used in previous studies.¹⁷ Life stress events were defined as breakup, bereavement, or financial difficulties within the past three months.

Definition of Anxiety and Depression

Anxiety was evaluated using the GAD-7 scale, and depression was evaluated using the PHQ-9 scale. Participants were categorized into different severity categories based on their scores on these scales. The GAD-7 is a validated instrument designed to quantify the severity of anxiety symptoms, $^{6,18,19}_{,,0}$ utilizing a scoring range from 0 to 21. This scale categorizes symptom severity into four distinct levels: minimal (0-4), mild (5-9), moderate (10-14), and severe (15–21), based on the frequency and intensity of anxiety manifestations. Similarly, the PHQ-9 serves as a robust diagnostic tool for evaluating depressive symptoms, $^{6,18,19}_{0,10}$ employing a scoring system from 0 to 27. It classifies depression severity into analogous categories: minimal (0-4), mild (5-9), moderate (10-14), and severe (15–27). Both the GAD-7 and PHQ-9 have demonstrated strong psychometric properties, including high reliability and validity, making them clinically reliable instruments for accurately assessing and quantifying anxiety and depression symptoms in both research and clinical settings. Their widespread adoption in mental health assessment is attributed to their ability to provide precise measurements that correlate well with clinical diagnostic criteria. Notably, the GAD-7 and PHQ-9 scales were used to assess symptoms of anxiety and depression, which may overlap with stress-related symptoms. However, it should be emphasized that stress itself was not directly measured in this study, and we have not characterized stress as a disorder. This study calculated the Cronbach's alpha values for both the GAD-7 and the PHQ-9 scales, and the results indicated excellent internal consistency, with a Cronbach's alpha of 0.94 for GAD-7 and 0.93 for PHQ-9.

Statistical Analysis

The qualitative data in this research paper were presented in the form of proportion. The quantitative data were presented in the form of mean and standard deviation. In this study, linear regression analysis was used to assess the association between electronic screen exposure time and anxiety and depression scores. AUC analysis was used to evaluate the model performance for predicting anxiety and depression based on hours of exposure to electronic screens. Subgroup analysis was conducted to compare the effect of exposure time on anxiety and depression scores between different subgroups of participants. Multiple logistic regression analysis was conducted to control for demographics, exercises, and

other lifestyles while assessing the association between hours of exposure to electronic screen and severe anxiety and depression, as well as identify risk and protective factors. All analysis was performed using R programming software (R version 4.1.2), with a P value of less than 0.05 indicating statistical significance.

Results

Participant's Baseline Characteristics

The sample consisted of 2896 participants, with a majority being female (55.5%) aged around 19.5 (SD: 1.76) years old (Table 1). Most students were single (77.2%) and had relatively normal dietary habits, with a majority reporting no low salt (71.1%) and oily (69.0%) food intake. In addition, 48.9% of participants reported that they had vegetables diet, and 60.2% of participants had fruits diet. Smoking (6.8%) and drinking (22.3%) were relatively uncommon, with only a small percentage of students reporting current use. Sedentary behavior and exposure to electronic device screens were also reported, 33.8% of participants had 6 or above hours of sedentary time, and the

Fable	I	Baseline	Characteristics	Among	University	Students
--------------	---	----------	-----------------	-------	------------	----------

Characteristics	Participants
n	2896
Gender (male/female, %)	1606/1290 (55.5/44.5)
Age (years, mean [SD])	19.54 (1.76)
Grade (%)	
First	1243 (42.9)
Second	1227 (42.4)
Third	233 (8.0)
Fourth	171 (5.9)
Delayed	22 (0.8)
Marital status (%)	
Single	2235 (77.2)
Dating	642 (22.2)
Married	19 (0.7)
Low salt food (no/yes, %)	2060/836 (71.1/28.9)
Oily food (no/yes, %)	1999/897 (69.0/31.0)
Barbecue (no/yes, %)	1843/1053 (63.6/36.4)
Meat (no/yes, %)	929/1967 (32.1/67.9)
Vegetables (no/yes, %)	1481/1415 (51.1/48.9)
Fruits (no/yes, %)	1153/1743 (39.8/60.2)
Smoking (%)	
None	2608 (90.1)
Previous	91 (3.1)
Current	197 (6.8)

Characteristics	Participants
Drinking (%)	
None	2149 (74.2)
Previous	100 (3.5)
Current	647 (22.3)
Sedentary time (hours, %)	
< I hour	219 (7.6)
\geq I and <3 hours	716 (24.7)
≧3 and <6 hours	982 (33.9)
≧6 hours	979 (33.8)
Exposure time of electronic device screens per day (hours, mean [SD])	6.46 (2.67)
Life stress event (yes/no, %)	982/1914 (33.9/66.1)
Sport frequency (per week, %)	
None	617 (21.3)
I–2	1463 (50.5)
3-4	529 (18.3)
≧5	287 (9.9)
GAD-7 score (mean [SD])	3.97 (4.33)
Severity of anxiety	
None	1750 (60.4)
Mild	877 (30.3)
Moderate	191 (6.6)
Severe	78 (2.7)
PHQ-9 score (mean [SD])	4.74 (5.07)
Severity of depression	
None	1616 (55.8)
Mild	935 (32.3)
Moderate	208 (7.2)
Severe	137 (4.7)
Chronic disease (yes/no, %)	109/2787 (3.8/96.2)

Table I (Continued).

Abbreviations: SD, Standard Deviation; GAD-7, Generalized Anxiety Disorder-7; PHQ-9, Patient's Health Questionnaire-9.

mean exposure time of electronic screens was 6.46 (SD: 2.67) hours. Additionally, the majority of students reported anxiety (39.6%) and depression (44.2%), with a small percentage experiencing severe anxious (2.7%) or depressive (4.7%) symptoms.

Multiple Analysis of Risk Factors for Severe Anxiety and Depression

After controlling demographics, exercises, and other lifestyles, the hours of exposure to electronic screen were still significantly associated with severe anxiety (p = 0.042, Table 2). It also found that low salt diet (p = 0.038) and barbecue (p = 0.031) were risk factors for severe anxiety, and no life stress event (p < 0.001), sport (p = 0.001), and no chronic disease (p = 0.037) were protective factors for severe anxiety (Figure 1). In addition, after controlling demographics,

Characteristics	OR	95%	S CI	P value
		LL	UL	
(Intercept)	0.015	0.000	0.477	0.017
Gender				
Male	Ref.			
Female	0.828	0.489	1.402	0.481
Age	1.064	0.904	1.253	0.454
Grade				
First	Ref.			
Second	0.956	0.544	1.678	0.874
Third	1.581	0.664	3.766	0.301
Fourth	0.385	0.067	2.205	0.284
Delayed	2.186	0.231	20.719	0.495
Marital status				
Single	Ref.			
Dating	0.621	0.329	1.173	0.142
Married	0.570	0.061	5.356	0.623
Low salt diet				
No	Ref.			
Yes	1.774	1.032	3.050	0.038
Oily food diet				
No	Ref.			
Yes	1.134	0.669	1.921	0.641
Barbecue				
No	Ref.			
Yes	1.806	1.054	3.095	0.031
Meat				
No	Ref.			
Yes	1.464	0.802	2.671	0.214

Table 2 Multiple Logistic Regression Analysis for Variables to Predict Severe Anxiety

Characteristics	OR	95%	CI	P value
		LL	UL	
Vegetable				
No	Ref.			
Yes	1.027	0.594	1.775	0.925
Fruit				
No	Ref.			
Yes	0.789	0.453	1.373	0.402
Smoking				
None	Ref.			
Previous	1.106	0.285	4.288	0.884
Current	1.452	0.537	3.920	0.462
Drinking				
None	Ref.			
Previous	0.730	0.193	2.768	0.644
Current	0.670	0.345	1.302	0.238
Sedentary time (hours, per day)				
< I hour	Ref.			
≧I and <3 hours	1.575	0.410	6.043	0.508
≧3 and <6 hours	1.786	0.498	6.407	0.373
≧6 hours	3.356	0.981	11.484	0.054
Exposure time of electronic screen (hours, per day)				
≦7.5	Ref.			
>7.5	1.682	1.019	2.778	0.042
Life stress event				
Yes	Ref.			
No	0.186	0.108	0.319	0.000
Sport frequency (per week)				
None	Ref.			
1–2	0.407	0.235	0.706	0.001
34	0.427	0.186	0.982	0.045
≧5	0.662	0.279	1.568	0.348
Chronic disease				
Yes	Ref.			
No	0.440	0.203	0.953	0.037

Table 2 (Continued).

Abbreviations: OR, Odds Rate; CI, Confident Interval; LL, Lower Limit; UL, Upper Limit.



Figure I The forest plot of significant variables for predicting severe anxiety. Abbreviations: OR, Odds Ratio; Ref., Reference.

exercises, and other lifestyles, the hours of exposure to electronic screen were still significantly associated with severe depression (p < 0.001, Table 3). Oily food diet (p = 0.041), barbecue (p = 0.041), current smoking (p = 0.011), and a sedentary time of 6 or above hours (p = 0.017) were risk factors for severe depression, and no life stress event (p < 0.001), sport (p < 0.001), and no chronic disease (p < 0.001) were protective factors for severe depression (Figure 2).

Characteristics	OR	95%	S CI	P value
		LL	UL	
(Intercept)	0.054	0.004	0.654	0.022
Gender				
Male	Ref.			
Female	0.732	0.483	1.109	0.141
Age	1.079	0.961	1.211	0.200
Grade				
First	Ref.			
Second	0.765	0.501	1.169	0.215
Third	0.728	0.343	1.548	0.410
Fourth	0.389	0.120	1.264	0.116
Delayed	1.387	0.272	7.081	0.694

Table 3 Multiple Logistic Regression Analysis for Variables to Predict Severe Depression

Table 3 (Continued).

Characteristics	OR	95%	S CI	P value
		LL	UL	
Marital status				
Single	Ref.			
Dating	0.621	0.381	1.014	0.057
Married	1.691	0.433	6.611	0.450
Low salt diet				
No	Ref.			
Yes	1.165	0.752	1.806	0.494
Oily food diet				
No	Ref.			
Yes	1.522	1.018	2.273	0.041
Barbecue				
No	Ref.			
Yes	1.538	1.017	2.325	0.041
Meat				
No	Ref.			
Yes	0.881	0.573	1.356	0.566
Vegetable				
No	Ref.			
Yes	1.023	0.670	1.561	0.916
Fruit				
No	Ref.			
Yes	0.793	0.515	1.219	0.290
Smoking				
None	Ref.			
Previous	0.809	0.275	2.386	0.702
Current	2.453	1.233	4.881	0.011
Drinking				
None	Ref.			
Previous	1.008	0.403	2.521	0.987
Current	0.633	0.378	1.061	0.083

Characteristics	OR	95% CI		P value
		LL	UL	
Sedentary time (hours, per day)				
< I hour	Ref.			
≧I and <3 hours	1.822	0.692	4.800	0.225
≧3 and <6 hours	1.871	0.734	4.772	0.190
≧6 hours	2.991	1.212	7.378	0.017
Exposure time of electronic screen (hours, per day)				
≦8.5	Ref.			
>8.5	2.103	1.424	3.106	0.000
Life stress event				
Yes	Ref.			
No	0.167	0.110	0.253	0.000
Sport frequency (per week)				
None	Ref.			
1–2	0.417	0.269	0.647	0.000
3-4	0.591	0.326	1.069	0.082
≧5	0.709	0.363	1.387	0.315
Chronic disease				
Yes	Ref.			
No	0.322	0.178	0.582	0.000

 Table 3 (Continued).

Abbreviations: OR, Odds Rate; CI, Confident Interval; LL, Lower Limit; UL, Upper Limit.

Association Between the Hours of Exposure to Electronic Screen and Mental Health This study further investigated the relationship between electronic screen exposure time and mental health status, including anxiety and depression. Linear regression analysis revealed a significant association between electronic screen exposure time and anxiety score (R square=0.17, p < 0.001, Figure 3A) and depression score (R square=0.20, p < 0.001, Figure 3B). Figure 4 exemplifies the increasing anxiety and depression scores concomitant with increasing exposure time to electronic screens.

Analysis of the Hours of Exposure to Electronic Screen for Predicting Mental Health

This study tested the prediction performance of the hours of exposure to electronic screen to predict mental health status (Table 4). The AUC value was 0.568 (95% CI: 0.547–0.589) for predicting anxiety (Figure 5A) and 0.641 (95% CI: 0.571–0.712) for predicting severe anxiety (Figure 5B). Moreover, the AUC was 0.590 (95% CI: 0.570–0.611) for predicting depression (Figure 5C) and 0.665 (95% CI: 0.613–0.716) for predicting severe depression (Figure 5D). The optimal cut-off value (threshold) for predicting severe anxiety and depression based on electronic screen exposure time was found to be 7.5 hours and 8.5 hours, respectively. To further corroborate this finding, a subgroup analysis was conducted according to the hours of exposure to electronic screen (≤ 7.5 vs >7.5 hours) and (≤ 8.5 vs >8.5 hours). The results revealed that participants with higher exposure time to electronic screens had significantly higher anxiety (p <



Figure 2 The forest plot of significant variables for predicting severe depression. Abbreviations: OR, Odds Ratio; Ref., Reference.



Figure 3 Association between the exposure time of electronic screen and mental health status among university students. (A) GAD-7; (B) PHQ-9. Abbreviations: GAD-7, General Anxiety Disorder 7; PHQ-9, Patient's Health Questionnaire 9.

(0.001) and depression scores (p < (0.001)) (Table 5 and Table 6). These findings provide further evidence to support the relationship between electronic screen exposure time and the severity of anxiety and depression.

Discussion

Main Findings

The study found that exposure time to electronic screens was significantly associated with anxiety and depression scores among university students. The association remained significant after controlling for demographics, exercises, and other lifestyle factors. In addition, the study also identified risk factors and protective factors for severe anxiety and depression. The findings suggest that interventions targeting electronic screen use could improve mental health outcomes among



Figure 4 Dynamic changes of the exposure time of electronic screen with the score of anxiety and depression. Abbreviations: GAD-7, General Anxiety Disorder 7; PHQ-9, Patient's Health Questionnaire 9.

university students. Notably, this study revealed that the optimal cut-off value for predicting anxiety and depression based on electronic screen exposure time was found to be both 6.5 hours, suggesting that a less than 6.5 hours of exposure to electronic screen was recommended to university students to prevent mental distress.

Detailed Analysis of Screen Exposure and Mental Health

In recent years, there has been growing concern about the potential mental health effects of electronic screen use among university students, as people's dependence on electronic devices is increasing. There were several studies that reported smartphone or internet addiction was significantly associated with mental health problems.^{14,15} For example, Seki et al¹⁵ found that internet addiction was an independent variable for depression. Meanwhile, Demirci et al²⁰ revealed that the high smartphone use students had higher scores on measures of depression, anxiety, and daytime dysfunction than the low smartphone use individuals after analyzing 319 university students, and the study also pointed out that there were also positive correlations between smartphone addiction scale scores and depression, anxiety, and sleep quality scores.

Metrics	Prediction Outcome							
	Anxiety	Severe Anxiety	Depression	Severe Depression				
AUC	0.568 (0.547–0.589)	0.641 (0.571–0.712)	0.590 (0.570–0.611)	0.665 (0.613–0.716)				
Threshold	6.500	7.500	6.500	8.500				
Specificity	0.617	0.671	0.636	0.801				
Sensitivity	0.484	0.590	0.497	0.460				
Accuracy	0.565	0.669	0.574	0.785				
Precision	0.453	0.047	0.519	0.103				
Youden	1.101	1.261	1.132	1.261				

Table 4 Analysis of Exposure Time of the Electronic Screen to Predict Mental Health StatusAmong University Students

Abbreviation: AUC, Area Under the Curve.



Figure 5 The area under the curve analysis of the exposure time of electronic screen to predict mental health. (A) No anxiety vs anxiety; (B) No severe anxiety vs severe anxiety; (C) No depression vs depression; (D) No severe depression vs severe depression. Abbreviation: AUC, Area Under the Curve.

Koly et al²¹ found that university students who spent more than 6 hours a day on social media were 4.6 times more likely to develop depression. In addition, Matar Boumosleh et al¹⁴ found that personality type A, class year, younger age at first smartphone use, excessive use during a weekday, using it for entertainment rather than calling family members, and having depression or anxiety were associated with smartphone addiction after analyzing 688 undergraduate university students in a cross-sectional study. Thus, the findings of the above studies revealed a substantial association between smartphone or internet addiction and the presence of mental health problems among university students.

Our study examined the association between the hours of exposure to electronic screens and mental health status. The findings indicated that with increasing hours of exposure to electronic screens, anxiety and depression scores correspondingly increased. Specifically, individuals who spent more time using electronic screens exhibited higher levels of anxiety and depression than those who spent less time using electronic screens. Additionally, our study found that the optimal

Characteristics	Overall	Exposure Time of Elec	р	
		≦7.5	>7.5	
n	2896	1924	972	
GAD-7 (mean [SD])	3.97 (4.33)	3.53 (3.96)	4.83 (4.87)	<0.001
Severity of anxiety				<0.001
None	1750 (60.4)	1222 (63.5)	528 (54.3)	
Mild	877 (30.3)	566 (29.4)	311 (32.0)	
Moderate	191 (6.6)	104 (5.4)	87 (9.0)	
Severe	78 (2.7)	32 (1.7)	46 (4.7)	
PHQ-9 (mean [SD])	4.74 (5.07)	4.16 (4.53)	5.90 (5.82)	<0.001
Severity of depression				<0.001
None	1616 (55.8)	1158 (60.2)	458 (47.1)	
Mild	935 (32.3)	590 (30.7)	345 (35.5)	
Moderate	208 (7.2)	119 (6.2)	89 (9.2)	
Severe	137 (4.7)	57 (3.0)	80 (8.2)	

Table 5 Subgroup Analysis of Mental Health According to the Exposure Time of Electronic Screen (\leq 7.5 vs >7.5 hours)

Abbreviations: SD, Standard Deviation; GAD-7, Generalized Anxiety Disorder-7; PHQ-9, Patient's Health Questionnaire-9.

Table 6	Subgroup	Analysis o	f Mental	Health	According to	the	Exposure	Time	of	Electronic
Screen (≦8.5 vs 8.	5 hours)								

Characteristics	Overall	Exposure Time of Elec	р	
		≦8.5	>8.5	
n	2896	2285	611	
GAD-7 (mean [SD])	3.97 (4.33)	3.67 (4.06)	5.08 (5.07)	<0.001
Severity of anxiety				<0.001
None	1750 (60.4)	1433 (62.7)	317 (51.9)	
Mild	877 (30.3)	675 (29.5)	202 (33.1)	
Moderate	191 (6.6)	134 (5.9)	57 (9.3)	
Severe	78 (2.7)	43 (1.9)	35 (5.7)	
PHQ-9 (mean [SD])	4.74 (5.07)	4.31 (4.62)	6.36 (6.22)	<0.001
Severity of depression				<0.001
None	1616 (55.8)	1341 (58.7)	275 (45.0)	
Mild	935 (32.3)	722 (31.6)	213 (34.9)	
Moderate	208 (7.2)	148 (6.5)	60 (9.8)	
Severe	137 (4.7)	74 (3.2)	63 (10.3)	

Abbreviations: SD, Standard Deviation; GAD-7, Generalized Anxiety Disorder-7; PHQ-9, Patient's Health Questionnaire-9.

cut-off value for the exposure time to electronic screens to predict anxiety and depression was 6.5 hours, suggesting that a more than 6.5-hour exposure of electronic screens was more likely to develop mental health problems among university students. Furthermore, our study demonstrated that a more than 7.5-hour exposure of electronic screens tended to be linked to more severe anxiety, and 8.5 hours tended to be associated with more severe depression. This may be thought to be due to the fact that electronic screens emit blue light, which can disrupt the body's production of melatonin, a hormone that helps regulate sleep-wake cycles.^{22,23} Additionally, the use of electronic screens can lead to sedentary behavior, which can also contribute to anxiety and depression.²⁴ In a word, our study provided evidence that excessive hours of exposure to electronic screens were associated with higher levels of anxiety and depression among university students. The findings suggest that limiting the use of electronic screens, particularly those exceeding 6.5 hours, may be beneficial for improving mental health outcomes. Our study contributed to a deeper understanding of the potential impact of electronic screen use on mental health and could help establish a foundation for the development of interventions to mitigate its negative effects.

Addressing Confounding Factors and Identifying Detailed Relationship Between Screen Exposure and Mental Health

The impact of the time of exposure to electronic screens on mental health has been studied, but the detailed relationship is inconclusive. It is important to note that the above studies have not been able to control for a number of confounding factors, such as age, gender, and socioeconomic status. Our study examined the detailed relationship between electronic screen exposure time and mental health status, as well as to identify risk factors and protective factors for severe anxiety and depression. We found that after controlling for individual's demographics, eating habits, and exercises, the time of exposure to electronic screens still held significance with anxiety and depression. In addition, this study also pointed out that a healthy eating habit (such as no low salt diet, no barbecue, and no oily food diet), no smoking, more exercise, less sedentary time, no life stress event, and no chronic disease were significantly associated with less anxiety and depression scores.

Implications for Education and Future Research

Based on above findings, therefore, effective measures to address student anxiety and depression can include lifestyle interventions such as regular exercise, a balanced diet, smoking quitting, and stress reduction techniques like mindfulness meditation and cognitive behavioral therapy. Additionally, seeking professional help from counselors and therapists can be beneficial for individuals experiencing significant mental health difficulties.²⁵ Other studies also provided evidence that cognitive, behavioral, and mindfulness interventions were effective in reducing stress in university students.²⁶ Furthermore, the findings of this study have important implications for university students and policymakers. To improve the mental health of university students, interventions targeting electronic screen use should be implemented. These interventions could include limiting screen time, providing educational resources about the health effects of electronic screen use, and promoting healthy lifestyle behaviors. The findings also suggest that universities could consider implementing policies that encourage students to reduce their electronic screen use, such as mandatory breaks during classes or designated screen-free hours in residence halls. By incorporating these evidence-based interventions into university programs and student support services, universities can foster a more supportive and inclusive environment for all students.

Future research should build on these findings to further explore the relationship between electronic screen exposure and mental health outcomes. Longitudinal studies are needed to examine the causal relationships between screen use and mental health over time, helping to determine whether excessive screen use is a risk factor for the development of anxiety and depression or a consequence of pre-existing mental health conditions. Additionally, research should investigate the underlying mechanisms linking screen exposure to mental health outcomes, such as the role of disrupted sleep patterns, reduced physical activity, and increased social isolation, as understanding these pathways could inform targeted interventions. Intervention studies are also critical to design and evaluate programs aimed at reducing screen time and promoting healthy screen use habits, such as mindfulness-based programs, digital detox initiatives, or educational campaigns. Expanding the scope of research to include diverse populations, such as students from different cultural backgrounds, age groups, and academic disciplines, would

help determine whether the observed associations are generalizable across different contexts. Furthermore, future studies should examine the differential impacts of various types of screen use (eg, social media, gaming, academic work) on mental health, as this could provide insights into which forms of screen use are most detrimental and inform more tailored interventions, particularly as we navigate an increasingly digital world and the rapid integration of various artificial intelligence into daily life.²⁷ Incorporating validated stress-specific scales, such as the Perceived Stress Scale or the Depression, Anxiety, and Stress Scale-21, would also help disentangle the effects of stress from anxiety and depression symptoms. Meanwhile, it is crucial to sensitize both students and teachers to the potential mental health risks associated with excessive screen use, Finally, evaluating the effectiveness of university-level policies and environmental changes, such as screen-free zones, mandatory breaks, or wellness programs, could provide valuable evidence for reducing screen time and improving mental health outcomes. By addressing these research gaps, future studies can further elucidate the complex relationship between screen exposure and mental health, ultimately contributing to the development of more effective prevention and intervention strategies.

Strengths and Limitations

This study has several limitations. First, the study participants were recruited from university students, which limits the generalizability of the findings to other populations. Second, the study did not control for all potential confounding factors, such as socioeconomic status and family history of mental illness. Third, the study did not assess the impact of different types of electronic screen use on mental health, such as social media use or gaming use. Fourth, the study did not assess the long-term effects of electronic screen use on mental health. Lastly, while the GAD-7 and PHQ-9 provide valuable insights into anxiety and depression symptoms, they do not capture stress as a separate dimension. Considering that the GAD-7 and PHQ-9 scores reflect symptoms that may be influenced by stress, it should be cautioned against interpreting these scores as definitive indicators of clinical disorders without further evaluation. Hence, although our study shielded an important light on the association between the time of exposure to electronic screen and mental distress, further research was still needed to explore the above issues and to develop effective interventions.

Conclusions

The findings of this study suggest that prolonged exposure to electronic screens is associated with increased risk for experiencing anxiety and depression among university students. Less than 6.5 hours of exposure to electronic screen are recommended among university students to largely avoid anxiety and depression. This study highlights the need to intervene and manage effectively the use of electronic screens in order to promote mental well-being in university students. Additionally, raising awareness about the potential negative impacts of excessive electronic screen use and promoting healthier lifestyle behaviors may serve as effective tools for managing mental health challenges among university students.

Funding

This study was funded by the teaching reform research project of Xiamen University of Technology (JYCG202459) and the Study of the Mechanism and Pathways of Artificial Intelligence Empowering Sports to Prevent and Control Myopia among Chinese Adolescents (No. SKHF24029).

Disclosure

The authors declare that they have no conflict of interest.

References

^{1.} Wang Y, Wang X, Wang X, et al. Stressors in university life and anxiety symptoms among international students: a sequential mediation model. *BMC Psychiatry*. 2023;23(1):556. doi:10.1186/s12888-023-05046-7

- 2. Meeks K, Peak AS. Dreihaus A: depression, anxiety, and stress among students, faculty, and staff. J Am Coll Health. 2023;71(2):348-354. doi:10.1080/07448481.2021.1891913
- 3. Ramón-Arbués E, Gea-Caballero V, Granada-López JM, Juárez-Vela R, Pellicer-García B, Antón-Solanas I. The prevalence of depression, anxiety and stress and their associated factors in college students. *Int J Environ Res Public Health*. 2020;17(19):7001.
- 4. Rotenstein LS, Ramos MA, Torre M, et al. Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. *JAMA*. 2016;316(21):2214–2236. doi:10.1001/jama.2016.17324
- 5. Iqbal S, Gupta S, Venkatarao E. Stress, anxiety and depression among medical undergraduate students and their socio-demographic correlates. *Indian J Med Res.* 2015;141(3):354–357. doi:10.4103/0971-5916.156571
- 6. Li Y, Jin S, Li Y, et al. The effects of research activities on biomedical students' mental health: a national cross-sectional study. *Front Psychiatry*. 2022;13:796697. doi:10.3389/fpsyt.2022.796697
- Al-Khani AM, Sarhandi MI, Zaghloul MS, Ewid M, Saquib N. A cross-sectional survey on sleep quality, mental health, and academic performance among medical students in Saudi Arabia. BMC Res Notes. 2019;12(1):665. doi:10.1186/s13104-019-4713-2
- March-Amengual JM, Cambra Badii I, Casas-Baroy JC, et al. Psychological distress, burnout, and academic performance in first year college students. Int J Environ Res Public Health. 2022;19(6). doi:10.3390/ijerph19063356
- Barbosa-Camacho FJ, Romero-Limón OM, Ibarrola-Peña JC, et al. Depression, anxiety, and academic performance in COVID-19: a cross-sectional study. BMC Psychiatry. 2022;22(1):443. doi:10.1186/s12888-022-04062-3
- 10. Guo F, Yi M, Sun L, et al. A novel model to predict mental distress among medical graduate students in China. *BMC Psychiatry*. 2021;21(1):569. doi:10.1186/s12888-021-03573-9
- 11. Herbert C. Enhancing mental health, well-being and active lifestyles of university students by means of physical activity and exercise research programs. *Front Public Health*. 2022;10:849093. doi:10.3389/fpubh.2022.849093
- 12. Xu M, Yin X, Gong Y. Lifestyle factors in the association of shift work and depression and anxiety. JAMA Netw Open. 2023;6(8):e2328798. doi:10.1001/jamanetworkopen.2023.28798
- Grasdalsmoen M, Eriksen HR, Lønning KJ, Sivertsen B. Physical exercise, mental health problems, and suicide attempts in university students. BMC Psychiatry. 2020;20(1):175. doi:10.1186/s12888-020-02583-3
- Matar Boumosleh J, Jaalouk D. Depression, anxiety, and smartphone addiction in university students- A cross sectional study. *PLoS One*. 2017;12 (8):e0182239. doi:10.1371/journal.pone.0182239
- 15. Seki T, Hamazaki K, Natori T, Inadera H. Relationship between internet addiction and depression among Japanese university students. J Affect Disord. 2019;256:668–672. doi:10.1016/j.jad.2019.06.055
- Landa-Blanco M, García YR, Landa-Blanco AL, Cortés-Ramos A, Paz-Maldonado E. Social media addiction relationship with academic engagement in university students: the mediator role of self-esteem, depression, and anxiety. *Heliyon*. 2024;10(2):e24384.
- 17. Zhang L, Zhao S, Yang Z, Zheng H, Lei M. An artificial intelligence platform to stratify the risk of experiencing sleep disturbance in university students after analyzing psychological health, lifestyle, and sports: a multicenter externally validated study. *Psychol Res Behav Manag.* 2024;17:1057–1071.
- Villarreal-Zegarra D, Barrera-Begazo J, Otazú-Alfaro S, Mayo-Puchoc N, Bazo-Alvarez JC, Huarcaya-Victoria J. Sensitivity and specificity of the Patient Health Questionnaire (PHQ-9, PHQ-8, PHQ-2) and General Anxiety Disorder scale (GAD-7, GAD-2) for depression and anxiety diagnosis: a cross-sectional study in a Peruvian hospital population. *BMJ Open*. 2023;13(9):e076193. doi:10.1136/bmjopen-2023-076193
- 19. Zhang L, Zhao S, Yang Z, Zheng H, Lei M. An artificial intelligence tool to assess the risk of severe mental distress among college students in terms of demographics, eating habits, lifestyles, and sport habits: an externally validated study using machine learning. *BMC Psychiatry*. 2024;24 (1):581.
- 20. Demirci K, Akgönül M, Akpinar A. Relationship of smartphone use severity with sleep quality, depression, and anxiety in university students. *J Behav Addict*. 2015;4(2):85–92. doi:10.1556/2006.4.2015.010
- Koly KN, Sultana S, Iqbal A, Dunn JA, Ryan G, Chowdhury AB. Prevalence of depression and its correlates among public university students in Bangladesh. J Affect Disord. 2021;282:689–694. doi:10.1016/j.jad.2020.12.137
- 22. Ostrin LA. Ocular and systemic melatonin and the influence of light exposure. Clin Exp Optom. 2019;102(2):99-108. doi:10.1111/cxo.12824
- 23. Wahl S, Engelhardt M, Schaupp P, Lappe C, Ivanov IV. The inner clock-Blue light sets the human rhythm. J Biophotonics. 2019;12(12): e201900102. doi:10.1002/jbio.201900102
- 24. Wang X, Li Y, Fan H. The associations between screen time-based sedentary behavior and depression: a systematic review and meta-analysis. *BMC Public Health.* 2019;19(1):1524. doi:10.1186/s12889-019-7904-9
- 25. Ip V, Chan F, Chan JY, Lee JK, Sung C, Wilson EH. Factors influencing Chinese college students' preferences for mental health professionals. J Ment Health. 2016;25(2):142–147. doi:10.3109/09638237.2015.1057328
- 26. Regehr C, Glancy D, Pitts A. Interventions to reduce stress in university students: a review and meta-analysis. J Affect Disord. 2013;148(1):1–11. doi:10.1016/j.jad.2012.11.026
- 27. Rossettini G, Cook C, Palese A, Pillastrini P, Turolla A. Pros and cons of using artificial intelligence Chatbots for musculoskeletal rehabilitation management. J Orthop Sports Phys Ther. 2023;53(12):728-734. doi:10.2519/jospt.2023.12000

Neuropsychiatric Disease and Treatment



Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal

