REVIEW

# Exercise Prescription Training in Chronic Obstructive Pulmonary Disease: Benefits and Mechanisms

Siqi Liu<sup>1,\*</sup>, Ailin Yang<sup>1,2,\*</sup>, Yue Yu<sup>1,3</sup>, Bo Xu<sup>1</sup>, Ganggang Yu<sup>1</sup>, Haoyan Wang<sup>1</sup>

<sup>1</sup>Department of Respiratory Medicine, Beijing Friendship Hospital, Capital Medical University, Beijing, People's Republic of China; <sup>2</sup>Department of Respiratory and Critical Medicine, Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China, Chengdu, People's Republic of China; <sup>3</sup>Department of Pulmonary and Critical Care Medicine, Xinqiao Hospital of Army Medical University, Chongqing, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Ganggang Yu; Haoyan Wang, Department of Respiratory Medicine, Beijing Friendship Hospital, Capital Medical University, No. 95 Yong An Road, Xichen District, Beijing, 100050, People's Republic of China, Email gangyu.603@163.com; haoyanw@ccmu.edu.cn

**Abstract:** Exercise rehabilitation training has emerged as one of the most promising modalities for enhancing clinical outcomes and overall well-being in patients with chronic obstructive pulmonary disease (COPD). Distinct exercise prescriptions yield different clinical benefits in this population. Endurance training has been demonstrated to significantly improve exercise capacity, alleviate dyspnea, and enhance health-related quality of life metrics. High-intensity interval training offers a time-efficient approach to boosting cardiorespiratory fitness and metabolic function. Resistance training addresses progressive muscle atrophy through targeted myofiber recruitment, thereby augmenting musculoskeletal performance and translating to enhanced exercise tolerance in patients with COPD. Exercise-mediated rehabilitation attenuates COPD progression and mitigates acute exacerbation risks via multifactorial mechanisms such as mitigation of inflammatory responses, reduction of oxidative stress, and improvement of endothelial cell function. Elucidating the pathophysiological mechanisms underlying exercise-induced benefits will pave the way for precision rehabilitation protocols, ultimately advancing COPD disease management paradigms, refining patient-centered outcome measures, and achieving sustainable health optimization in this clinical cohort. **Keywords:** chronic obstructive pulmonary disease, exercise training, pulmonary rehabilitation, inflammation, oxidative stress

#### Introduction

COPD is characterized by progressive airflow limitation, airway inflammation, and alveolar destruction, leading to impaired pulmonary function and significantly reduced quality of life.<sup>1</sup> As the third leading cause of death globally, COPD imposes significant economic and social burdens worldwide.<sup>2,3</sup>

Patients with COPD can benefit from pulmonary rehabilitation (PR), which has been shown to reduce symptoms and improve activity capacity, both of which are essential for enhancing quality of life.<sup>2,3</sup> Physical activity (PA), a key component of PR, encompasses exercise assessment and training therapy. PA is traditionally defined as any skeletal muscle movement that results in energy expenditure.<sup>4</sup> Physical training is a complex behavior influenced by various subjective (such as motivation and self-efficacy) and objective factors (such as environmental conditions and measurement tools).<sup>5</sup> Notably, patients with COPD have significantly lower levels of PA compared to age-matched healthy individuals,<sup>6</sup> highlighting the need for focused attention.

While extensive research confirms that exercise improves the general condition of patients with COPD, the optimal exercise intensity and the mechanisms by which it alleviates COPD symptoms require further investigation. This review summarizes advancements in exercise methods for COPD and explores the primary mechanisms underlying their effects on patients.

# Various Forms of Exercise Training

Endurance training (ET), high-intensity interval training (HIIT), resistance training (RT), flexibility exercise training, and neuromotor exercise training each contribute to overall physical fitness and improved health, particularly for individuals with chronic respiratory conditions such as COPD. ET focuses on enhancing aerobic efficiency, increasing the anaerobic threshold, and improving cardiovascular and pulmonary function.<sup>7,8</sup> HIIT alternates intense exercise bursts with recovery periods, enhancing cardiorespiratory fitness in patients with COPD.<sup>9</sup> It also incorporates inspiratory muscle training (IMT), which improves respiratory muscle strength and tolerance, alleviating some COPD symptoms,<sup>10</sup> though it may not significantly reduce dyspnea.<sup>11</sup> IMT, utilizing a specialized inspiratory muscle trainer, such as Expand-A-Lung and Powerbreathe, can significantly improve exercise capacity, health-related quality of life, and daily physical activity (DPA) in patients with inspiratory muscle mass, enhancing quality of life,<sup>13</sup> while flexibility training improves joint mobility and reduces muscle damage.<sup>14</sup> Neuromotor training, such as yoga, Tai Chi, split-belt treadmill, and perturbation-based training, promotes balance and coordination, supporting the overall well-being of older adults.<sup>15,16</sup>

A multifaceted exercise program should be tailored to individual capabilities, incorporating a combination of these training modalities to optimize respiratory function and overall health. Regular participation in these exercises can significantly improve exercise capacity, muscle strength, flexibility, and quality of life. For patients with COPD, a comprehensive rehabilitation program integrating these exercise modalities is essential for effective disease management. A summary of the exercise types and recommendations is provided in Table 1.

Type of Exercise	Examples	Focus/Benefit	Recommendations
Endurance training (aerobic training) <sup>8</sup>	Walking, running/jogging, swimming, biking, jumping rope, dancing, climbing stairs	Improves aerobic efficiency, lung function, cardiorespiratory fitness, and arterial stiffness	60–80% max work rate, 20 -60 min/session, 3–5 times/ week; gradually increase work rate and duration; include respiratory muscle endurance training
High-intensity interval training <sup>11,17,18</sup>	Running, dancing, rowing machines, stationary bicycling, climbing stairs	Enhances cardiorespiratory fitness, exercise capacity, and lung function	≥80% <sup>19</sup> max intensity, 2–3 days/ week. Include a 5-min warm-up and 5–10 min cool-down; add high-intensity inspiratory muscle training for patients with moderate-to-severe COPD.
Resistance training	Free weights, weight machines,	Builds strength, increases	60-70% IRM, increase training
(strength/ weight training) <sup>13</sup>	medicine balls, bands, kettlebells,	muscle mass, and improves	load as tolerated; 2–4 sets/
	elastic tubing, bodyweight exercise	quality of life	session, 8–12 repetitions, 2–3 times/week for ≥8 weeks
Flexibility exercise training <sup>14,20</sup>	Static stretching, dynamic	Enhances flexibility and range	Smooth and slow movements for
	stretching	of motion; mitigates muscle	60s daily (10–30 s increments),
		damage	at least 2 to 3 days/week
Balance and coordination	Balance balls, heel-to-toe walks,	Enhances sciatic nerve	≥45 min/session, 2–3 times/
training <sup>15,16,21,22</sup>	one-leg balance exercises, Tai	regeneration; improves	week for≥ 8 weeks
	Chi, dancing, split-belt treadmill,	balance, exercise tolerance,	
	perturbation-based training,	and health-related quality of	
	Sensorimotor training	life; decreases the risk of falls	

Table I Exercise Types and Prescription Recommendations for Chronic Obstructive Pulmonary Disease

Abbreviations: IRM, One-repetition maximum; COPD, chronic obstructive pulmonary disease.

# **Biomolecular Mechanism of Exercise Training** Promoting Inflammation Resolution

Systemic and airway inflammation are two major features of COPD and disrupt muscle synthesis, leading to muscle dysfunction.<sup>17</sup> These inflammatory processes significantly influence the progression and severity of COPD, highlighting the importance of managing both systemic and airway inflammation in chronic respiratory conditions.

The effects of exercise on inflammatory markers are complex. A substantial body of evidence suggests that exercise can improve the overall inflammatory response. However, some inflammatory markers remained unaffected by exercise (Table 2).

Although the precise mechanism by which exercise mitigates inflammation in patients with COPD remains unclear, recent research suggests that exercise can regulate inflammation by altering chemerin levels.<sup>39</sup> Chemerin is a novel adipokine that plays a major role in adipogenesis and lipid metabolism.<sup>36</sup> This is particularly important in diseases such as COPD, where the chemerin/CMKLR1 axis plays a role in both inflammation and metabolism. The inhibitory effect of exercise on chemerin expression provides valuable insights into exercise-induced pulmonary rehabilitation, which shows potential for clinical applications in COPD management. Additionally, exercise reduces STAT3 activation in various cell types involved in COPD pathogenesis, including peribronchial leukocytes, parenchymal leukocytes, and airway epithelial cells, thereby alleviating inflammation in patients with COPD.<sup>37</sup> Furthermore, Bufei Yishen Formula III, a classical traditional Chinese Medicine, exerts therapeutic effects by modulating pulmonary function, reducing inflammation, and enhancing immune regulation in chronic respiratory diseases. A study demonstrated that, when combined with exercise rehabilitation, it may synergistically reduce lung inflammation by inhibiting the EGFR/MAPK pathway<sup>40</sup> (Figure 1).

# Effects of Exercise Training on Oxidative Stress

The effects of exercise training on oxidative stress in patients with COPD are complex and influenced by various factors, such as the type and intensity of the exercise and the severity of the disease. While some studies suggest that exercise may trigger oxidative stress by increasing airflow and exposure to pollutants, other research indicates that exercise has minimal effects on oxidative stress in COPD, with no significant changes observed in the ferric-reducing ability of plasma or thiobarbituric acid reactive substances after exercise.<sup>41</sup> Notably, most studies support the view that exercise positively mitigates oxidative stress in patients with COPD, as demonstrated by the findings summarized in Table 3.

The beneficial effects of exercise on oxidative stress in the diaphragm of patients with COPD involve multiple pathways. Clinical and animal studies by Lei et al<sup>10</sup> showed that sequential IMT alleviates oxidative stress by modulating the SOCS5/JAK2/STAT3 signaling pathway. Irisin, a muscle-derived hormone released during exercise, may protect against oxidative stress by activating the Nrf2 and HO-1 pathways, potentially mitigating cigarette smoke-induced emphysema in COPD<sup>45</sup> (Figure 1).

# Others

Exercise has been shown to enhance muscle strength, endurance, and overall exercise capacity in patients with COPD by modulating the expression levels of myokines, including interleukin (IL)-6, IL-15, myostatin, irisin/FNDC5, insulin-like growth factor-1, and semaphorin 3A, thereby effectively addressing skeletal muscle dysfunction.<sup>46,47</sup> Additionally, moderate-intensity exercise improves mitochondrial function in the skeletal muscles of patients with COPD by increasing PGC-1 $\alpha$  expression, enhancing ATP production through mitochondrial complex I, and regulating antioxidant capacity via the Keap1–Nrf2–ARE pathway.<sup>48</sup>

Endothelial dysfunction is another common issue in individuals with COPD and appears to be related to the severity of airflow limitation.<sup>49</sup> This dysfunction can significantly impair patients' abilities to engage in physical exercise but can be effectively mitigated through structured exercise programs. Notably, the benefits of exercise on endothelial function appear to be independent of supplemental oxygen during training, suggesting that exercising in regular room air might be even more beneficial for improving peripheral endothelial function. This effect might stem from the stimulatory impact of local hypoxic conditions, which trigger beneficial vascular adaptation. Exercise increases the production and bioavailability of nitric oxide (NO), which helps regulate shear stress and improves vascular function.<sup>50</sup> Furthermore,

Study ID	Design	Sample Size	Participants	Intervention	Biomarkers Related to Inflammation Change
F. Rodrigues et al <sup>23</sup>	Cross-sectional	21	Patients with COPD, clinically stable (past 4 weeks), exercise- capable, and able to complete health status	Treadmill/Cycle ergometer until exhaustion; low-intensity activities maintained, no high-intensity exercise in prior 48 h	No significant changes in IFN-γ, IL-1β, IL6, IL8, TNF-α, TGF-β and iNOS at rest, 1 h, and 24 h post-exercise
Paul D. Loprinzi, PhD et al <sup>24</sup>	Cross-sectional	238	questionnaires Former or current smokers with self- reported COPD and complete data on study variables	ActiGraph 7164 accelerometers worn on right hip for 7 days (LPA, MVPA, TPA)	LPA, MVPA, and TPA inversely associated with WBC and neutrophil count
Olaf holz I, Stefan roepcke et al <sup>25</sup>	Prospective clinical	46 (moderate COPD: 23; age- and sex- matched healthy controls: 23)	Current smokers with a smoking history of at least 10 pack-years	Peak work capacity and constant-load exercise	IL6, sICAMI, CRP ↑; MPO ↓; Healthy smokers: IL6, CRP ↑ > COPD; COPD: IL6, MPO, CRP, sICAMI, VWF, LTB4 changes/min > controls. Recovery: IL6 ↑ in healthy smokers only
Gilson Pires Dorneles et al <sup>26</sup>	Cross-sectional	41 (moderate:14; severe: 14; very severe: 13)	Patients with COPD	6MWT, patients walked on a 30-m track (10-m markings). Stopped for symptoms (eg, chest pain, respiratory distress, muscle pain). Distance was measured using a 20-m tape	IL6: $\uparrow$ in very severe > severe (post-test). IL4: $\downarrow$ in severe < very severe/moderate (pre-test); $\uparrow$ in moderate > severe (post- test). Very severe: IL4 pre-post $\uparrow$ effect
Ryrsø CK <sup>27</sup>	Cross-sectional	38 (RT: 15; ET: 15; healthy sedentary participants: 8)	Patients with moderate-to -severe COPD and healthy controls	RT and ET, both groups completed 35 min of supervised exercise three times a week for 8 weeks.	TNF- $\alpha$ $\uparrow;$ IL-1 $\beta,$ IL6, IL8, and IL18 unchanged
Juliana S. Uzeloto et al <sup>28</sup>	RCT	23 (PEG: 15; RPG: 8)	Participants with stable	PEG or RPG	IL-8 ↓; IL-13, TNF-α↑ in CD4 <sup>+</sup> T lymphocytes
Shehab M et al <sup>29</sup>	RCT	80 (AE group: 40; no exercise training: 40)	Patients with moderate severity of COPD	Aerobic exercise on a treadmill	TNF- $\alpha$ , IL-4, IL-6, and CRP $\downarrow$ in exercise group
Bruna S. de Alencar Silva et al <sup>30</sup>	RCT	48 (EBG: 14; ETG:18; MG:16)	Clinically stable patients with moderate to severe COPD	ETG Group: Used specially prepared chairs with elastic tubes (resistance levels 200–204); EBG Group: Utilized elastic bands with resistance categorized by color: yellow (lowest) to black (highest); MG Group: Employed traditional weight machines for upper limb exercises	IL-6: $\downarrow$ after 12 weeks, group differences and interaction; TNF- $\alpha$ : $\downarrow$ after 12 weeks, no group differences; IL-10: $\downarrow$ in EG vs MG; IL-15: $\downarrow$ after 12 weeks, no group/ interaction effects; IL-10/TNF- $\alpha$ Ratio: Higher in EG vs MG

#### Table 2 Summary of Studies on the Impact of Exercise on Inflammation in COPD

	-				
Alex R. Jenkins <sup>31</sup>	Prospective	67 (COPD, Phase 1: 40;	Patients with COPD were	8-week exercise program	Acute exercise (Phase I): Fibrinogen,
	cohort	COPD, Phase 2: 27)	recruited with no	(cardiorespiratory + muscular strength)	CRP, and neutrophil markers unchanged;
			exacerbations for		leukocytes, neutrophils, and immature
			≥4 weeks and no		neutrophils increased. Pulmonary rehab
			antibiotic or steroid		(Phase 2): Fibrinogen response increased
			treatment during this		post-exercise
			period		
Chun-Hua Wang, Pai-Chien Chou	Clinical pilot	26 (control:14; mobile	Patients with COPD	Mobile phone-guided ET (80% of the	Reduced CRP and IL8 in mobile phone
et al <sup>32</sup>		phone:12)		maximal capacity) vs control group	group; no changes in IL6 and TNF- $\alpha$
Shehab M et al <sup>33</sup>	Prospective	100 (AE: 50; RT:50)	Patients with COPD	AE vs RT (3months)	TNF-α, II-2, IL-4, IL-6, and CRP were
24	cohort				reduced in both groups
Wei Liu et al <sup>34</sup>	RCT	102 (control: 26,	Patients with mild to	TC, TBRS, TBRS-TC	IL-6, IL-8, TNF- $\alpha$ , and copeptin were
		TBRS: 25, TC:26,	severe stable COPD		statistically significant between groups.
		TBRS-TC: 25)			Greater reduction in IL8 and TNF- $\alpha$ in
27					TBRS-TC
Xishuai Wang et al <sup>35</sup>	Animal	30 (control:10;	Male ICR mice	COPD: LPS via intratracheal injection;	BALF: total cell count, neutrophils,
	experiment	COPD:10; COPD		COPD + exercise: LPS via intratracheal	macrophages, TGF- $\beta$ , IL-1 $\beta$ , TNF- $\alpha \downarrow$ ;
		+AE:10)		injection and low-intensity exercise using	IL10 and CXCL1 $\uparrow$ ; lymphocytes,
				a six-lane treadmill	eosinophils, IL17 unchanged; Serum:
24					TNF-α ↓, IL10 ↑
Hidetaka IMAGITA et al <sup>36</sup>	Animal	21 (control: 7;	9-week-old male Wistar	Treadmill running in CS and LPS-induced	No significant difference in IL-1 $\beta$ , IL6,
	experiment	pulmonary	rats	pulmonary emphysema	TNF-α, IL4, and IL10
		emphysema group: 7;			
		emphysema + exercise			
		group: 7)			
Maysa Alves Rodrigues Brandao-	Animal	64 (control:16;	Male C57Bl/6 mice	Treadmill aerobic training in CS-induced	BALF: IL10 ↑; total cells, lymphocytes,
Rangel et al <sup>37</sup>	experiment	exercise:16; COPD:16		COPD mice	neutrophils, IL-1 $\beta$ , CXCL1, IL-17, and
		COPD+Exercise:16)			TNF- $\alpha \downarrow$ ; Serum: IL10 $\uparrow$ ; total leukocytes,
					neutrophils, lymphocytes, monocytes, IL-
					1 $\beta$ , CXCLI, IL-17, and TNF- $\alpha \downarrow$
Panpan Liu, Hongchang Gao	Animal		C57BL/6 male mice	Motor-driven treadmill exercise in CS-	Reversed TNF- $\alpha$ , IL-1 $\beta$ , and IL6
et al <sup>30</sup>	experiment			induced COPD mice	expression

**Abbreviations**: COPD, Chronic obstructive pulmonary disease; RCT, randomized controlled clinical trial; LPA, light-intensity physical activity, 100–2019 counts per min; MVPA, moderate-to-vigorous physical activity,  $\geq$  2020 counts per min; TPA, total physical activity,  $\geq$  100 counts per min; slCAMI, soluble intercellular adhesion molecule 1; CRP, C-reactive protein; MPO, myeloperoxidase; VWF, von Willebrand factor; LTB4, leukotriene B4; 6MWT, 6-min walk test; RT, resistance training; ET, endurance training; AE, aerobic exercise; PEG, physical exercise group; RPG, respiratory physiotherapy group; EBG, Elastic band group; ETG, elastic tube group; MG, weight machines equipment group; TBRS, total body recumbent stepper; TC, Tai Chi; TBRS-TC, combined TBRS exercise and Tai Chi; CS, cigarette smoke; LPS, lipopolysaccharide solution.



Figure I Mechanisms of exercise-induced improvement in COPD symptoms (created using Figdraw). Abbreviation: COPD, chronic obstructive pulmonary disease.

a randomized, double-anonymized, sham-controlled study confirmed that the application of noninvasive positive pressure ventilation during high-intensity exercise acutely modulates endothelial functions in patients with COPD-related heart failure.<sup>51</sup>

Study	Participants	Exercise Type	Changes in Oxidative Stress Markers	Main Conclusion
Mayalen Valero-Breton et al <sup>42</sup> Denisse Valladares-Ide et al <sup>43</sup>	Patients with COPD Patients with COPD	12-week eccentric (ECC) and concentric (CONC) cycling training Moderate-intensity CONC (CONC-M), moderate-intensity ECC (ECC-M), high-intensity ECC (ECC-H)	<b>CONC</b> : $\uparrow$ TAC, $\downarrow$ TBARS, $\downarrow$ GPX activity; <b>ECC</b> : $\downarrow$ GPX, no significant changes in TBARS, SOD, or CAT <b>MDA</b> : $\downarrow$ after CONC-M (-28%), ECC-M (-14%), ECC-H (-17%); <b>GPX</b> : No change; <b>EBC</b> H <sub>2</sub> O <sub>2</sub> , <b>NO</b> <sup>2</sup> -, <b>pH</b> : No change	CONC training significantly improved antioxidant capacity; ECC had weaker effects on oxidative stress Exercise decreased oxidative stress in the plasma of patients with COPD at moderate stage of the disease

Table 3 Effects of Different Exercise Interventions on Oxidative Stress in COPD

(Continued)

#### Table 3 (Continued).

Study	Participants	Exercise Type	Changes in Oxidative Stress Markers	Main Conclusion
Wang et al <sup>35</sup>	LPS-induced COPD mice	Aerobic exercise	↓ MDA, MPO; ↑ SOD, GSH	Exercise reduced oxidative stress and improved antioxidant capacity
Renata Tiscoski Nesi et al <sup>44</sup>	CS-induced COPD mice	Swimming	Prevented increases in hydroxyproline content, SOD activity, CAT activity, 2-thiobarbituric acid reactive species, and carbonyl group content	Exercise mitigated CS- induced oxidative damage
Yirou Lei et al <sup>10</sup>	Patients with COPD + CS- induced COPD mice	NIPPV, IMT, and sequential NIPPV + IMT	↓ ROS, MDA; ↑ SOD, GSH	Sequential NIPPV + IMT reduced oxidative stress and improved COPD outcomes

**Abbreviations**: TAC, total antioxidant capacity; TBARS, barbituric acid reactive substances; GPX, glutathione peroxidase activity; SOD, superoxide dismutase; CAT, Catalase; MDA, malondialdehyde; EBC, exhaled breath condensate; H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide; LPS, lipopolysaccharide; GSH, glutathione; CS, Cigarette smoke; NIPPV, noninvasive positive pressure ventilation; IMT, inspiratory muscle training.

The lactate threshold—the point where lactate production surpasses clearance—occurs at lower exercise intensities in patients with COPD due to impaired oxygen delivery, leading to early lactic acid buildup, breathlessness, and fatigue. Tailored exercise training can raise this threshold, allowing patients to exercise at higher intensities before symptoms arise, improving their physical capacity and disease management.<sup>52</sup> Excess post-exercise oxygen consumption (EPOC), the elevated oxygen uptake post-exercise, is less pronounced in patients with COPD due to limited lung capacity and impaired oxygen use. Managing intensity is key to optimizing EPOC while minimizing respiratory strain. Activities like walking, cycling, and aquatic exercises are ideal, as they promote recovery while accommodating reduced respiratory function.<sup>53</sup>

Current research supports the idea that exercise directly influences pulmonary function by modulating inflammation and oxidative stress. Additionally, exercise enhances skeletal muscle and endothelial function, thereby significantly influencing the pathophysiology of COPD. This multifaceted effect underscores the importance of incorporating physical activity into therapeutic strategies to improve health outcomes in individuals with this debilitating respiratory condition.

# **Exercise Training Assessment in Clinical Practice**

## Acute Exacerbations of Chronic Obstructive Pulmonary Disease (AE-COPD)

GOLD 2023 has highlighted the potential of PR in reducing hospitalization rates among patients with recent exacerbations (within 4 weeks of their previous hospitalization). Initiating exercise training within 3 days of hospitalization likely enhances exercise capacity and improves overall quality of life.<sup>54</sup> A clinical study revealed that Medicare beneficiaries who completed appropriate exercise training within 90 days of being discharged for COPD experienced a reduction in rehospitalization rates within a year.<sup>55</sup> A retrospective study demonstrated that combined respiratory and exercise training significantly lowered the rates of acute exacerbation, rehospitalization, and mortality compared to controls.<sup>56</sup> A randomized controlled trial<sup>57</sup> showed that patients receiving rigorous supervised training had fewer readmissions and a longer time to first readmission than those receiving usual care. Additionally, RT was safer, more feasible, and more effective than ET for improving muscle function in patients with severe AECOPD. However, ET, when performed after the acute phase of AECOPD, can improve functional exercise tolerance.<sup>58</sup>

## Dyspnea

Dyspnea is a prominent symptom among individuals with COPD. The severity of dyspnea can be assessed using various tools, including the modified Medical Research Council (mMRC) scale, Borg Dyspnea Scale, Visual Analogue Scale, Numeric Rating Scale, and the Baseline Dyspnea Index/Transition Dyspnea Index. GOLD 2023 indicates that PR improves dyspnea in patients with COPD. Exercise reduces central ventilatory drive, improves dynamic ventilatory mechanics, and enhances respiratory muscle function, thereby alleviating dyspnea.<sup>59</sup> Additionally, a decrease in the diaphragm activation ratio (measured via electromyography) during maximal inspiration explains the reduction in perceived dyspnea, even under prolonged high ventilation and mechanical loading during exercise.<sup>60</sup> Upper extremity exercise and IMT alleviate dyspnea in patients with COPD,<sup>61</sup> with the ET group reporting significantly lower mMRC scores compared to controls.<sup>62</sup> Additionally, a 4-week lower limb resistance training program effectively reduced exertional dyspnea, while both elastic tube and conventional resistance training reduced mMRC scores by 30%.<sup>63</sup> However, the effectiveness of HIIT in reducing dyspnea remains debated. A study<sup>64</sup> indicated a reduction in breathlessness, while a subsequent meta-analysis<sup>11</sup> of 689 participants suggested that HIIT improves pulmonary function, exercise capacity, and quality of life without significantly affecting the sensation of breathlessness. This inconsistency is noteworthy, as the larger sample size of the meta-analysis may provide a more reliable conclusion regarding HIIT's impact. Furthermore, sensitivity analyses indicated that the effect on dyspnea could vary depending on the study design and measurement tools used. Therefore, HIIT may not be suitable for all patients with COPD, particularly those with significant dyspnea. Therefore, a more appropriate exercise method should be selected for patients with COPD who experience significant dyspnea.

# Lung Function

Spirometry is an inexpensive, noninvasive, easily accessible, and repeatable diagnostic test widely used for diagnosing and monitoring COPD. Commonly reported parameters in pulmonary function tests include forced expiratory volume in 1 s (FEV1), forced vital capacity (FVC), and the ratio of these measurements (FEV1/FVC). A meta-analysis of 20 RCTs involving 962 patients with COPD demonstrated that HIIT improved peak oxygen consumption (standardized mean difference [SMD] = 0.30, 95% confidence interval [CI]: 0.14–0.46), peak minute ventilation (SMD = 0.26, 95% CI: 0.05-0.47), and peak work rate (SMD = 0.34, 95% CI: 0.17-0.51) in these patients. However, no significant effects were observed regarding the FEV1/FVC ratio or St. George's Respiratory Questionnaire scores (P > 0.05). Additionally, no significant difference was found in the FEV1/FVC ratio between the HIIT and control groups.<sup>64</sup> A clinical study revealed that ET combined with resistance training (ERT) and ET alone increased vital capacity (VC), FVC, and forced expiratory flow at 25%–75% (FEF25%–75%), while only ET and ERT increased maximum voluntary ventilation (MVV); however, none of these interventions significantly affected FEV1 or FEV1/FVC in this study.<sup>65</sup> Notably, high-volume RT may impair lung function indices due to reduced engagement of ventilatory and respiratory muscles.<sup>66</sup> Nonetheless, a shortterm (12 weeks) high-intensity RT program for patients with moderate to severe COPD resulted in a significant 5.3% increase in FEV1 and a trend towards improved daily peak-flow measurements.<sup>67</sup> Therefore, selecting the appropriate exercise intensity and frequency for patients with varying degrees of severity is crucial. Several other indicators reflecting ventilation and gas exchange function, such as MVV, PEF, and FEF25%-75%, can also be assessed simultaneously.

## **Exercise Ability**

The Six-Minute Walk Distance (6MWD) is a widely utilized clinical tool for evaluating the functional exercise capacity of individuals with COPD. The 6MWD test provides valuable information about the severity of COPD, its impact on exercise tolerance, and the effectiveness of interventions such as pulmonary rehabilitation and pharmacotherapy. ET, RT, combined ET and RT, HIIT, and respiratory training have all been shown to improve 6MWD.<sup>56,64,68</sup> Cardiopulmonary exercise testing (CPET) is a comprehensive, noninvasive procedure that evaluates the integrated functions of the heart and lungs during physical activity. It has consistently been regarded as the gold standard for assessing exercise capacity in patients with COPD,<sup>69</sup> especially those with pulmonary artery hypertension. High-intensity interval cycle ergometer training for 6 weeks resulted in significant reductions in lactate values, minute ventilation, ventilatory equivalent for oxygen, heart rate, oxygen uptake, and carbon dioxide output.<sup>70</sup> Additionally, notable differences were observed between

active (DPA  $\geq$  30 min a day, 5 days a week) and inactive (DPA < 30 min a day, 5 days a week) patients with COPD in their cardiovascular responses to exercise, such as peak O<sub>2</sub> pulse, double product reserve, and heart rate recovery. However, no significant variations were observed in their ventilatory response or operational volumes during exercise.<sup>71</sup> Therefore, both 6MWD and CPET are convenient and practical tools for monitoring COPD. The 6MWD can be used to assess the benefits of exercise, while CPET results allow for individualized adjustments to training intensity, ultimately optimizing exercise programs for patients.

#### Health-Related Quality of Life

Various methods are available for evaluating quality of life in patients with COPD, including the St. George's Respiratory Questionnaire (SGRQ), COPD Assessment Test (CAT), Chronic Respiratory Questionnaire (CRQ), and Clinical COPD Questionnaire (CCQ). The use of ET resulted in a significantly lower CAT score compared to the control group in a study.<sup>62</sup> In another study, both the RT and ET groups significantly improved CRQ scores compared to the control group, with no notable differences between the two groups.<sup>70</sup> Conversely, conventional RT was more effective in improving CRQ scores than ET using elastic tubes.<sup>63</sup> Short-term (12 weeks) high-intensity RT significantly improved SGRQ scores in patients with moderate to severe COPD.<sup>67</sup> Additionally, HIIT significantly improved the quality of life in patients with COPD, although SGRQ scores did not significantly differ between the HIIT and control groups.<sup>64</sup> Neuromotor exercise training, specifically sensorimotor training, has significantly reduced pain compared to usual care. It is recommended that the training intensity be moderated to ensure participants do not experience excessive fatigue. Furthermore, a frequency of two to three sessions per week is suggested for optimal results. Additionally, whole-body vibration therapy has demonstrated promise in improving sleep quality and is particularly effective in reducing depressive symptoms. Aerobic exercises like brisk walking, swimming, cycling, and dancing help reduce stress, improve mood, and enhance cognitive function by boosting endorphin production, increasing blood flow to the brain, enhancing prefrontal oxygenation, and elevating circulating neurotrophins.<sup>72</sup>

# Conclusion

In conclusion, exercise interventions, including ET, HIIT, and RT, offer significant benefits for patients with COPD by improving exercise capacity, reducing dyspnea, and enhancing quality of life. These modalities not only improve physical function but also reduce inflammation and oxidative stress, potentially slowing disease progression and reducing exacerbation risk. Monitoring inflammatory markers like IL-4, IL-6, IL-8, and IL-10 can help assess intervention efficacy.

To optimize patient outcomes, individualized exercise prescriptions should be tailored to disease severity, comorbidities, and physical capability, with a focus on gradual progression in intensity and duration. Supervision is essential for patients with more severe disease, ensuring safety during exercise. Clinicians should integrate exercise with pharmacological treatments and pulmonary rehabilitation for a holistic management approach.

Future research should aim to elucidate the molecular mechanisms behind exercise-induced improvements, standardize exercise protocols, and evaluate long-term adherence and efficacy. By incorporating exercise as a core component of COPD management, clinicians can significantly improve the functional status and quality of life of their patients.

# Acknowledgments

Funding/Financial support: This study was supported by the National Natural Science Foundation of China (Grant numbers: 81870029, 81700038, 82000042, and 82000043) and the Key Clinical Specialty Construction Program of Beijing (Grant number: 2020–2022).

## Disclosure

The authors report no conflicts of interest in this work.

# References

- 1. Decramer M, Janssens W, Miravitlles M. Chronic obstructive pulmonary disease. *The Lancet*. 2012;379(9823):1341–1351. doi:10.1016/S0140-6736(11)60968-9
- Nici L, Donner C, Wouters E, et al. American Thoracic Society/European Respiratory Society Statement on Pulmonary Rehabilitation. Am J Respir Crit Care Med. 2006;173(12):1390–1413. doi:10.1164/rccm.200508-1211ST
- Higashimoto Y, Ando M, Sano A, et al. Effect of pulmonary rehabilitation programs including lower limb endurance training on dyspnea in stable COPD: a systematic review and meta-analysis. *Respir Investig.* 2020;58(5):355–366. doi:10.1016/j.resinv.2020.05.010
- 4. Caspersen CJ, Powell KE, Christenson GM. Physical Activity, Exercise, and Physical Fitness: definitions and Distinctions for Health-Related Research. *Public Health Reports*. 1985;100(2):126.
- Silfee VJ. Objective measurement of physical activity outcomes in lifestyle interventions among adults\_ A systematic review. Prev Med Rep. 2018. doi:10.1016/j.pmedr.2018.05.003
- 6. Spruit MA, Pitta F, McAuley E, ZuWallack RL, Nici L. Pulmonary Rehabilitation and Physical Activity in Patients with Chronic Obstructive Pulmonary Disease. Am J Respir Crit Care Med. 2015;192(8):924–933. doi:10.1164/rccm.201505-0929CI
- 7. Jones AM, Carter H. The Effect of Endurance Training on Parameters of Aerobic Fitness. Sports Med. 2000;29(6):373-386. doi:10.2165/00007256-200029060-00001
- Vivodtzev I, Minet C, Wuyam B, et al. Significant Improvement in Arterial Stiffness After Endurance Training in Patients With COPD. Chest. 2010;137(3):585–592. doi:10.1378/chest.09-1437
- 9. Vogiatzis I. Dynamic hyperinflation and tolerance to interval exercise in patients with advanced COPD. Eur Respir J. 2004;24(3):385–390. doi:10.1183/09031936.04.00128903
- Lei Y, He J, Hu F, et al. Sequential inspiratory muscle exercise-noninvasive positive pressure ventilation alleviates oxidative stress in COPD by mediating SOCS5/JAK2/STAT3 pathway. BMC Pulm Med. 2023;23(1):385. doi:10.1186/s12890-023-02656-5
- Gao M, Huang Y, Wang Q, Liu K, Sun G. Effects of High-Intensity Interval Training on Pulmonary Function and Exercise Capacity in Individuals with Chronic Obstructive Pulmonary Disease: a Meta-Analysis and Systematic Review. *Adv Ther*. 2022;39(1):94–116. doi:10.1007/s12325-021-01920-6
- 12. Kkp M, Lr N, Pr A, Jc P, Lft S. A Review on Respiratory Muscle Training Devices. J Pulm Respir Med. 2018;08(02). doi:10.4172/2161-105X.1000451
- 13. Liao W, Chen J, Chen X, et al. Impact of Resistance Training in Subjects With COPD: a Systematic Review and Meta-Analysis. *Respir Care*. 2015;60(8):1130–1145. doi:10.4187/respcare.03598
- 14. Ketenci A. What is the benefits flexibility exercise training for adults with fibromyalgia? A Cochrane review summary with commentary. *Turk J Phys Med Rehabil.* 2020;66(4):468–472. doi:10.5606/tftrd.2020.0508
- 15. Gerards M, Marcellis R, Senden R, et al. The effect of perturbation-based balance training on balance control and fear of falling in older adults: a single-blind randomised controlled trial. *BMC Geriatr.* 2023;23(1):305. doi:10.1186/s12877-023-03988-x
- Conradsson D, Hinton DC, Paquette C. The effects of dual-tasking on temporal gait adaptation and de-adaptation to the split-belt treadmill in older adults. Exp Gerontol. 2019;125:110655. doi:10.1016/j.exger.2019.110655
- 17. Aakerøy L, Nørstebø EA, Thomas KM, et al. High-intensity interval training and pulmonary hemodynamics in COPD with hypoxemia. *Eur Clin Respir J.* 2021;8(1):1984642. doi:10.1080/20018525.2021.1984642
- 18. Hill K. High-intensity inspiratory muscle training in COPD. Eur Respir J. 2006;27(6):1119-1128. doi:10.1183/09031936.06.00105205
- 19. Sawyer A, Cavalheri V, Hill K. Effects of high intensity interval training on exercise capacity in people with chronic pulmonary conditions: a narrative review. *BMC Sports Sci Med Rehabil.* 2020;12(1):22. doi:10.1186/s13102-020-00167-y
- Chen CH, Nosaka K, Chen HL, Lin MJ, Tseng KW, Chen TC. Effects of Flexibility Training on Eccentric Exercise-Induced Muscle Damage. Med Sci Sports Exerc. 2011;43(3):491–500. doi:10.1249/MSS.0b013e3181f315ad
- Bonetti LV, Korb A, Da Silva SA, et al. Balance and coordination training after sciatic nerve injury. *Muscle Nerve*. 2011;44(1):55–62. doi:10.1002/ mus.21996
- 22. Dunsky A. The Effect of Balance and Coordination Exercises on Quality of Life in Older Adults: a Mini-Review. *Front Aging Neurosci*. 2019;11:318. doi:10.3389/fnagi.2019.00318
- Rodrigues F, Papoila AL, Ligeiro D, Gomes MJM, Trindade H. Acute exercise amplifies inflammation in obese patients with COPD. Rev Port Pneumol Engl Ed. 2016;22(6):315–322. doi:10.1016/j.rppnen.2016.05.005
- Loprinzi PD, Walker JF, Lee H. Association between Physical Activity and Inflammatory Markers among U.S. Adults with Chronic Obstructive Pulmonary Disease. Am J Health Promot. 2014;29(2):81–88. doi:10.4278/ajhp.130510-QUAN-235
- Holz O, Roepcke S, Watz H, Tegtbur U, Lahu G, Hohlfeld J. Constant-load exercise decreases the serum concentration of myeloperoxidase in healthy smokers and smokers with COPD. Int J Chron Obstruct Pulmon Dis. 2015;1393. doi:10.2147/COPD.S83269
- 26. Dorneles GP, Vianna P, Del Duca Lima D, et al. Cytokine response to the 6-min walk test in individuals with different degrees of COPD. *Clin Respir* J. 2016;10(3):326–332. doi:10.1111/crj.12221
- Ryrsø CK, Thaning P, Siebenmann C, et al. Effect of endurance versus resistance training on local muscle and systemic inflammation and oxidative stress in COPD. Scand J Med Sci Sports. 2018;28(11):2339–2348. doi:10.1111/sms.13227
- 28. Uzeloto JS, de Toledo-Arruda AC, Silva BSA, et al. Effect of physical training on cytokine expression in CD4+ T lymphocytes in subjects with stable COPD. *Ther Adv Respir Dis.* 2022;16:17534666221091179. doi:10.1177/17534666221091179
- El-Kade SM A, Al-Jiffri OH. Exercise alleviates depression related systemic inflammation in chronic obstructive pulmonary disease patients. *Afr Health Sci.* 2017;16(4):1078. doi:10.4314/ahs.v16i4.25
- Silva BSDA, Lira FS, Rossi FE, et al. Inflammatory and Metabolic Responses to Different Resistance Training on Chronic Obstructive Pulmonary Disease: a Randomized Control Trial. Front Physiol. 2018;9:262. doi:10.3389/fphys.2018.00262
- Jenkins AR, Holden NS, Jones AW. Inflammatory responses to acute exercise during pulmonary rehabilitation in patients with COPD. Eur J Appl Physiol. 2020;120(10):2301–2309. doi:10.1007/s00421-020-04452-z
- 32. Wang CH, Chou PC, Joa WC, et al. Mobile-phone-based home exercise training program decreases systemic inflammation in COPD: a pilot study. *BMC Pulm Med.* 2014;14(1):142. doi:10.1186/1471-2466-14-142

- El-Kader SM A, Al-Jiffri OH, Al-Shreef FM. Plasma inflammatory biomarkers response to aerobic versus resisted exercise training for chronic obstructive pulmonary disease patients. *Afr Health Sci.* 2016;16(2):507–515. doi:10.4314/ahs.v16i2.19
- 34. Liu W, Liu XM, Huang YL, et al. Tai Chi as a complementary exercise for pulmonary rehabilitation in chronic obstructive pulmonary disease: a randomised controlled trial. *Complement Ther Med.* 2023;78:102977. doi:10.1016/j.ctim.2023.102977
- Wang X, Wang Z, Tang D. Aerobic Exercise Alleviates Inflammation, Oxidative Stress, and Apoptosis in Mice with Chronic Obstructive Pulmonary Disease. Int J Chron Obstruct Pulmon Dis. 2021;16:1369–1379. doi:10.2147/COPD.S309041
- 36. Tan L, Lu X, Danser AHJ, Verdonk K. The Role of Chemerin in Metabolic and Cardiovascular Disease: a Literature Review of Its Physiology and Pathology from a Nutritional Perspective. Nutrients. 2023;15(13):2878. doi:10.3390/nu15132878
- Rodrigues Brandao-Rangel MA, Bachi ALL, Oliveira-Junior MC, et al. Exercise Inhibits the Effects of Smoke-Induced COPD Involving Modulation of STAT3. Oxid Med Cell Longev. 2017;2017:6572714. doi:10.1155/2017/6572714
- 38. Liu P, Gao H, Wang Y, Li Y, Zhao L. LncRNA H19 Contributes to Smoke-Related Chronic Obstructive Pulmonary Disease by Targeting miR-181/ PDCD4 Axis. COPD J Chronic Obstr Pulm Dis. 2023;20(1):119–125. doi:10.1080/15412555.2023.2165906
- 39. Li J, Lu Y, Li N, et al. Chemerin: a Potential Regulator of Inflammation and Metabolism for Chronic Obstructive Pulmonary Disease and Pulmonary Rehabilitation. *BioMed Res Int.* 2020;2020:1–20. doi:10.1155/2020/4574509
- 40. Xu K, Ma J, Lu R, et al. Effective-compound combination of Bufei Yishen formula III combined with ER suppress airway mucus hypersecretion in COPD rats: via EGFR/MAPK signaling. *Biosci Rep.* 2023;43(11):BSR20222669. doi:10.1042/BSR20222669
- 41. Domaszewska K, Górna S, Pietrzak M, Podgórski T. Oxidative Stress and Total Phenolics Concentration in COPD Patients—The Effect of Exercises: a Randomized Controlled Trial. *Nutrients*. 2022;14(9):1947. doi:10.3390/nu14091947
- Valero-Breton M, Valladares-Ide D, Álvarez C, Peñailillo RS, Peñailillo L. Changes in Blood Markers of Oxidative Stress, Inflammation and Cardiometabolic Patients with COPD after Eccentric and Concentric Cycling Training. *Nutrients*. 2023;15(4):908. doi:10.3390/nu15040908
- 43. Valladares-Ide D, Bravo MJ, Carvajal A, et al. Changes in pulmonary and plasma oxidative stress and inflammation following eccentric and concentric cycling in stable COPD patients. Eur J Appl Physiol. 2021;121(6):1677–1688. doi:10.1007/s00421-021-04652-1
- 44. Nesi RT, de Souza PS, Dos Santos GP, et al. Physical exercise is effective in preventing cigarette smoke-induced pulmonary oxidative response in mice. Int J Chron Obstruct Pulmon Dis. 2016;11:603–610. doi:10.2147/COPD.S93958
- 45. Kubo H, Asai K, Kojima K, et al. Exercise Ameliorates Emphysema Of Cigarette Smoke-Induced COPD In Mice Through The Exercise-Irisin-Nrf2 Axis. Int J Chron Obstruct Pulmon Dis. 2019;14:2507–2516. doi:10.2147/COPD.S226623
- 46. Arabzadeh E, Shirvani H, Masjedi MR, Ghanei M, Hofmeister M, Rostamkhani F. Treadmill exercise with nanoselenium supplementation affects the expression of Irisin/FNDC5 and semaphorin 3A in rats exposed to cigarette smoke extract. 3 Biotech. 2024;14(1):4. doi:10.1007/s13205-023-03849-9
- Han L, Li P, He Q, et al. Revisiting Skeletal Muscle Dysfunction and Exercise in Chronic Obstructive Pulmonary Disease: emerging Significance of Myokines. *Aging Dis.* 2023. doi:10.14336/AD.2023.1125
- Wang Y, Li P, Cao Y, Liu C, Wang J, Wu W. Skeletal Muscle Mitochondrial Dysfunction in Chronic Obstructive Pulmonary Disease: underlying Mechanisms and Physical Therapy Perspectives. *Aging Dis.* 2023;14(1):33–45. doi:10.14336/AD.2022.0603
- 49. Vivodtzev I, Tamisier R, Baguet JP, Borel JC, Levy P, Pépin JL. Arterial Stiffness in COPD. Chest. 2014;145(4):861-875. doi:10.1378/chest.13-1809
- 50. Francescomarino SD, Sciartilli A, Valerio VD, Baldassarre AD, Gallina S. The Effect of Physical Exercise on Endothelial Function. *Sports Med.* 2009;2009:1.
- Da luz Goulart C, Caruso FR, De Araújo AS G, et al. Non-invasive ventilation improves exercise tolerance and peripheral vascular function after high-intensity exercise in COPD-HF patients. *Respir Med.* 2020;173:106173. doi:10.1016/j.rmed.2020.106173
- 52. Spruit MA, Burtin C, De Boever P, et al. COPD and exercise: does it make a difference? *Breathe*. 2016;12(2):e38-e49. doi:10.1183/20734735.003916
- Brsheim E, Bahr R. Effect of Exercise Intensity, Duration and Mode on Post-Exercise Oxygen Consumption. Sports Med. 2003;33(14):1037–1060. doi:10.2165/00007256-200333140-00002
- 54. Zhang D, Zhang H, Li X, et al. Pulmonary Rehabilitation Programmes Within Three Days of Hospitalization for Acute Exacerbation of Chronic Obstructive Pulmonary Disease: a Systematic Review and Meta-Analysis. Int J Chron Obstruct Pulmon Dis. 2021;16:3525–3538. doi:10.2147/ COPD.S338074
- 55. Stefan MS, Pekow PS, Priya A, et al. Association between Initiation of Pulmonary Rehabilitation and Rehospitalizations in Patients Hospitalized with Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med.* 2021;204(9):1015–1023. doi:10.1164/rccm.202012-4389OC
- Wang L, Wu X. Clinical Effects of Exercise Combined with Respiratory Training in the Rehabilitation Treatment of Patients with Chronic Obstructive Pulmonary Disease. *Altern Ther.* 2024;30:188–194.
- 57. Ko FW, Tam W, Siu EHS, et al. Effect of short-course exercise training on the frequency of exacerbations and physical activity in patients with COPD: a randomized controlled trial. *Respirology*. 2021;26(1):72–79. doi:10.1111/resp.13872
- 58. Wuyts M, Hermans F, Breuls S, et al. Development and feasibility of an exercise training program in primary care for patients with COPD experiencing an acute exacerbation. *Physiotherapy*. 2024;123:81–90. doi:10.1016/j.physio.2023.09.003
- 59. Sheel AW, Foster GE, Romer LM. Exercise and its impact on dyspnea. Curr Opin Pharmacol. 2011;11(3):195-203. doi:10.1016/j. coph.2011.04.004
- 60. Langer D, Ciavaglia C, Faisal A, et al. Inspiratory muscle training reduces diaphragm activation and dyspnea during exercise in COPD. J Appl Physiol. 2018;125(2):381–392. doi:10.1152/japplphysiol.01078.2017
- 61. Pan L, Guo YZ, Yan JH, Zhang WX, Sun J, Li BW. Does upper extremity exercise improve dyspnea in patients with COPD? A meta-analysis. *Respir Med.* 2012;106(11):1517–1525. doi:10.1016/j.rmed.2012.08.002
- 62. Li Y, Wu W, Wang X, Chen L. Effect of Endurance Training in COPD Patients Undergoing Pulmonary Rehabilitation: a Meta-Analysis. Comput Math Methods Med. 2022;2022:4671419. doi:10.1155/2022/4671419
- 63. de Silva BS, Gobbo A, Freire LA, Trevisan IB, Silva IG, Ramos EMC. Effects of a Resistance Training with Elastic Tubing in Strength, Quality of Life and Dypsnea in Patients with Chronic Obstructive Pulmonary Disease. J Phys Educ. 2016;27:e2722. doi:10.4025/jphyseduc.v27i1.2722
- 64. Wang H, Liu Q, Liu L, Cao J, Liang Q, Zhang X. High-intensity interval training improves the outcomes of patients with chronic obstructive pulmonary disease: a meta-analysis of randomized controlled trials. *Respir Med.* 2023;208:107128. doi:10.1016/j.rmed.2023.107128

- 65. Khosravi M, Tayebi SM, Safari H. Single and Concurrent Effects of Endurance and Resistance Training on Pulmonary Function. *Iran J Basic Med Sci.* 2013;16(4):628–634.
- 66. Hackett D. Acute effects of high-volume compared to low-volume resistance exercise on lung function. J Exerc Rehabil. 2020;16(5):450–457. doi:10.12965/jer.2040656.328
- Wright P, Heck H, Langenkamp H. Effects of a resistance training on pulmonary function and performance measurements in patients with chronic obstructive pulmonary disease. Eur J Sport Sci. 2003;3(3):1–10. doi:10.1080/17461390300073304
- Li P, Li J, Wang Y, Xia J, Liu X. Effects of Exercise Intervention on Peripheral Skeletal Muscle in Stable Patients With COPD: a Systematic Review and Meta-Analysis. Front Med. 2021;8. doi:10.3389/fmed.2021.766841.
- 69. Chen L, Su J. Pulmonary Rehabilitation Exercise Assessments and Training Methods for Patients with COPD: a Literature Review. *J Rehabil Ther.* 2021;3(1):1.
- 70. Casaburi R, Patessio A, Ioli F, Zanaboni S, Donner CF, Wasserman K. Reductions in Exercise Lactic Acidosis and Ventilation as a Result of Exercise Training in Patients with Obstructive Lung Disease. Am Rev Respir Dis. 1991;143(1):9–18. doi:10.1164/ajrccm/143.1.9
- 71. Aiello M, Frizzelli A, Pisi R, et al. Effects of Daily Physical Activity on Exercise Capacity in Chronic Obstructive Pulmonary Disease. *Medicina* (*Mex*). 2024;60(7):1026. doi:10.3390/medicina60071026
- 72. Mador MJ, Bozkanat E, Aggarwal A, Shaffer M, Kufel TJ. Endurance and Strength Training in Patients with COPD. Cardiopulm Phys Ther J. 2004;15(4):22. doi:10.1097/01823246-200415040-00007

International Journal of Chronic Obstructive Pulmonary Disease



Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www. dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal

1082 🖪 💥 in 🗖