ORIGINAL RESEARCH

Establishing Sitafloxacin Epidemiological Cut-off Values (ECOFFs) for Clinical Bacterial Isolates

Jingjia Zhang^{1,2,*}, Xue Li^{1,*}, Xinmiao Jia^{3,*}, Yun Li⁴, Bo Zheng⁴, Yunsong Yu⁵, Fupin Hu⁶, Ziyong Sun⁷, Zhongju Chen⁷, Ge Zhang¹, Wei Kang¹, Tong Wang¹, Jin Li¹, Haotian Gao¹, Qiaolian Yi¹, Wei Yu^{1,8}, Xiaobing Chu^{1,8}, Yingchun Xu¹, Qiwen Yang^{1,9}

¹Department of Clinical Laboratory, State Key Laboratory of Complex Severe and Rare Diseases, Peking Union Medical College Hospital, Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, People's Republic of China; ²Clinical Biobank, Biomedical Engineering Facility of National Infrastructures for Translational Medicine, Chinese Academy of Medical Sciences, Peking Union Medical College Hospital, Beijing, Peoples Republic of China; ³Department of Medical Research Center, State Key Laboratory of Complex Severe & Rare Diseases, Peking Union Medical College Hospital, Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, People's Republic of China; ⁴Institute of Clinical Pharmacology, Peking University, Beijing, People's Republic of China; ⁵Department of Clinical Infectious Diseases, Zhejiang Provincial People's Hospital, Hangzhou, People's Republic of China; ⁶Huashan Hospital of Fudan University, Shanghai, People's Republic of China; ⁷Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, People's Republic of China; ⁸Graduate School, Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, People's Republic of China; ⁹Key Laboratory of Pathogen Infection Prevention and control (Peking Union Medical College), Ministry of Education, Beijing, People's Republic of China

*These authors contributed equally to this work

Correspondence: Qiwen Yang, Department of Clinical Laboratory, Peking Union Medical College Hospital, Email yangqiwen81@vip.163.com

Objective: To establish the epidemiological cut-off values (ECOFFs) of sitafloxacin against *Escherichia coli, Klebsiella pneumoniae, Proteus mirabilis, Pseudomonas aeruginosa, Acinetobacter baumannii, Staphylococcus aureus, Enterococcus faecalis, Enterococcus faecium,* and *Streptococcus pneumoniae.*

Methods: We collected 2264 clinical isolates from five different labs located in four cities in China. The minimum inhibitory concentrations (MICs) and inhibition zone diameters of sitafloxacin for all isolates were determined by using the broth microdilution method (BMD) and the disk diffusion method according to European Committee on Antimicrobial Susceptibility Testing (EUCAST) guidelines. MIC ECOFFs were determined using ECOFFinder software, with the 99% calculated ECOFF selected as the initial value. Zone diameter ECOFFs were determined via the visual estimation method. Whole-genome sequencing was performed on *E. coli* strains exhibiting overlapping MICs between wild-type (WT) and non-wild-type (NWT) groups to analyze resistance mechanisms.

Results: Sitafloxacin MICs ranged from 0.002 to 64 mg/L, while inhibition zone diameters ranged from 6 to 45 mm across the nine species. MIC ECOFFs were determined as 0.032, 0.064, 0.125, 0.5, 0.064, 0.125, 0.5, 0.25, and 0.125 mg/L for *E. coli, K. pneumoniae, P. mirabilis* (tentative ECOFF), *P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecium*, and *S. pneumoniae*, respectively. Except for *S. pneumoniae*, MICs of the other eight species showed a high correlation with zone diameters $(|\mathbf{r}| > 0.8, P < 0.0001)$. Consequently, the zone diameter ECOFFs were established as 26, 25, 24, 24, 25, 26, 21, and 22 for *E. coli, K. pneumoniae, P. mirabilis, P. aeruginosa, A. baumannii, S. aureus, E. faecium*, respectively.

Conclusion: We established MIC and zone diameter ECOFFs for sitafloxacin against the nine species listed above. The MIC ECOFF for *P. mirabilis* was classified as tentative. For *S. pneumoniae*, the correlation between zone diameters and MICs was insufficient to establish a zone diameter ECOFF.

Keywords: sitafloxacin, ECOFF, MIC, zone diameter, BMD, disk diffusion

Introduction

The escalating prevalence of antibiotic-resistant bacterial strains constitutes a critical global health threat, driving clinical and diagnostic laboratories to prioritize the development of novel antimicrobial therapies.^{1,2} Among fluoroquinolones, antibiotics such as levofloxacin remain widely prescribed in clinical practice. Nevertheless, their extensive application has been accompanied by a concerning rise in resistance phenotypes. This investigation aims to establish the

Infection and Drug Resistance 2025:18 1993-2004

1993

Graphical Abstract



epidemiological cutoff values (ECOFFs) for sitafloxacin - a third-generation fluoroquinolone - through systematic analysis of susceptibility data from multicenter clinical isolates.

Sitafloxacin, a synthetic broad-spectrum 8-chloro-fluoroquinolone, exhibits in vitro activity against both Grampositive and Gram-negative bacteria.³ It inhibits DNA gyrase and topoisomerase IV, enzymes essential for bacterial DNA replication and transcription.⁴ The N-1-cyclopropyl substitution in sitafloxacin enhances its antibacterial activity,⁵ resulting in superior bacteriostatic effects compared to other quinolones such as levofloxacin, moxifloxacin and ciprofloxacin.^{6–9}

First developed in Japan in 1988,¹⁰ sitafloxacin was approved for clinical use in 2008 and later introduced to the Chinese market in 2019. It is indicated for infections caused by *Escherichia coli, Klebsiella pneumoniae, Proteus mirabilis, Pseudomonas aeruginosa, Acinetobacter baumannii, Staphylococcus aureus, Enterococcus faecalis, Enterococcus faecium, Streptococcus pneumoniae, Mycobacterium, and Helicobacter pylori.^{6,11–17} However, clinical evidence for its efficacy in treating pneumonia and urinary tract infections remains limited,^{18–20} and no clinical breakpoint or ECOFFs have been established in China.*

In 2002, the European Committee on Antimicrobial Susceptibility Testing (EUCAST) defined ECOFFs as the highest minimum inhibitory concentration (MIC) for bacteria or fungi lacking phenotypically detectable acquired resistance mechanisms.²¹ This threshold distinguishes wild-type (WT) from non-wild-type (NWT) populations and serves as a provisional guideline for clinical treatment until clinical breakpoint are validated.²² To date, ECOFFs have been established for numerous antibiotics and pathogens in Europe.^{23–25}

Despite increasing clinical use of sitafloxacin, neither its ECOFFs nor clinical breakpoints have been defined. Consequently, physicians currently rely on extrapolated data from other quinolones with established breakpoints and limited clinical reports to guide empirical use – a practice that lacks precision. This study aims to address this gap by establishing ECOFFs for sitafloxacin against common clinical strains.

The broth microdilution method (BMD), recommended by both the Clinical and Laboratory Standard Institute (CLSI) and EUCAST for antimicrobial susceptibility testing (AST), remains the gold standard. However, the disk diffusion method offers greater practicality in routine clinical settings. Validating the correlation between these two methods is therefore critical to ensure the reliability of disk diffusion results for sitafloxacin.

To our knowledge, neither EUCAST nor CLSI has established clinical breakpoints or ECOFFs for sitafloxacin. This study therefore seek to determine ECOFFs for sitafloxacin against prevalent Gram-negative and Gram-positive bacterial species.

Methods

Bacterial Isolates and Quality Control Strains

This multicenter study analyzed 2,264 non-duplicate clinical isolates collected from five tertiary hospitals across four Chinese cities between 2017 and 2020. The participating hospitals included Peking Union Medical College Hospital (PU, Beijing; central laboratory), Peking University First Hospital (BD, Beijing), Sir Run Run Shaw Hospital of Zhejiang University (YF, Hangzhou), Huashan Hospital of Fudan University (HS, Shanghai), and Tongji Hospital of Tongji Medical College (WH, Wuhan).

All isolates were obtained from hospitalized patients across various clinical specimens, primarily including sputum, urine, blood, bile, bronchoalveolar lavage fluid (BALF), drainage fluid, abscess, and others. Species identification was performed using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS), with the following distribution: 250 *A. baumannii*, 253 *E. faecalis*, 205 *E. faecium*, 240 *E. coli*, 250 *K. pneumoniae*, 206 *P. mirabilis*, 353 *P. aeruginosa*, 250 *S. aureus*, and 257 *S. pneumoniae* (Supplementary material, Table S1).

For quality control, 27 external quality control (EQC) isolates (three per species with stratified MIC values) were analyzed as blind samples. Reference strains (*E. coli* ATCC 25922, *P. aeruginosa* ATCC 27853, *S. aureus* ATCC 29213, *E. faecium* ATCC 29212, and *S. pneumoniae* ATCC 49619) were included. MICs for all blind samples and control strains were determined in triplicate over three consecutive days across five independent laboratories. Results were validated if $\geq 2/3$ measurements fell within one two-fold dilution of the central laboratory's values.²¹

Antimicrobial Susceptibility Testing

The BMD concentration range of sitafloxacin was from 0.002 to 64 mg/L. The content of sitafloxacin disk (Daiichi Sankyo Company Limited) was 5 μ g. In accordance with the EUCAST SOP 11.0,²⁶ all 2264 isolates were detected by BMD and disk diffusion method in five separate laboratories. Levofloxacin was used as control agent to perform experiments on BMD and disk diffusion.

Whole-Genome Sequencing

Because of the overlap of WT and NWT MIC distribution of sitafloxacin against *E.coli* strains, we performed wholegenome sequencing to deeply clarify the resistance mechanism distribution on the *E.coli* strains with MICs of 0.032 and 0.064 mg/L. Genomic DNA was extracted with the SDS method.²⁷ The harvested DNA was detected by the agarose gel electrophoresis and quantified by Qubit[®] 2.0 Fluorometer (Thermo Scientific). A total amount of 1µg DNA per sample was used as input material for the DNA sample preparations. Sequencing libraries were generated using NEBNext[®] Ultra[™] DNA Library Prep Kit for Illumina (NEB, USA) following manufacturer's recommendations and index codes were added to attribute sequences to each sample. The whole genome of clinical isolates were sequenced using Illumina NovaSeq PE150. The raw data obtained by sequencing (Raw Data) had a certain proportion of low-quality data. In order to ensure the accuracy and reliability of the subsequent information analysis results, the original data must be filtered to obtain valid data (Clean Data). The specific processing steps for genome assembly with Clean Data.

Data Analysis

MICs and inhibition zone diameters from five laboratories were centrally analyzed. According to EUCAST Standard Operating Procedure,²⁸ MIC ECOFFs of nine species were calculated and determined by ECOFFinder software (<u>www.eucast.org/mic_distributions and_ecoffs/</u>) and visual estimation in central laboratory, the two sets of data should be consistent or within one dilution difference. Zone diameter ECOFFs were determined by the visual estimation. When it had high correlation between MICs and zone diameters, we can set zone diameter ECOFFs for species.²⁹

To explore whether there is a correlation between inhibition zone diameter and MIC ECOFFs, we used PRISM 8.0 software (GraphPad, La Jolla, CA, USA) to analyze the correlation of MICs and zone diameters, linear regression, calculate the P value and R square.

After the whole genome assembly was completed, Resfinder and PointFinder software were used to predict the resistance genes and point mutations of the strain, respectively.^{30,31}

Results

MIC Distributions and ECOFFs Establishment

All results of the EQC samples and QC strains were within one dilution step and satisfied the criteria.

MICs against the nine species were distributed in the concentration range of 0.002–64 mg/L (<u>Table S1</u>). The modal MICs of the centers for the same strain were almost always within a dilution factor. The maximum value of MIC of *S. pneumoniae* was 1 mg/L, which was much smaller than the value of the other 8 species. *K. pneumoniae* has the widest MIC distribution, ranging from 0.004 to 64 mg/L. Only the modal MIC of *E. coli* and *E. faecium* produced by the WH lab had a large deviation (Table 1). The modal MIC for all *E. coli* at the five centers was 0.008 mg/L, compared to 1 mg/L for the WH lab. Similarly, WH lab had an *E. faecium* modal MIC of 2 mg/L, but an overall *E. faecium* modal MIC was 0.064 mg/L.

We selected 99% calculated ECOFF as the initially formulated ECOFF value, these were consistent or within one dilution difference with visual estimate ECOFFs (Figure 1). The calculated ECOFF values were relatively small, with the largest being 0.25 mg/L for *P. aeruginosa* and *E. faecalis*.

Among them, we believe that the data of *P. mirabilis* was not amenable to analysis, according to EUCAST SOP10.1,²⁸ the species was only formulated tentative ECOFF (TECOFF). We analyzed the *P. mirabilis* data of the five centers one by one, and made the MIC distribution (Table 1). Except for YF, the data of the other 4 centers peaked at 0.032 mg/L. While YF was relatively concentrated in the range of 0.032 to 0.5 mg/L, these five concentrations had 8, 7, 6, 5, and 5 MIC value distributions, respectively. Therefore, the data of *P. mirabilis* from YF lab were considered substandard data, and only 4 centers were included in the end.

In the end, we determined that the MIC ECOFFs were 0.032, 0.064, 0.125, 0.5, 0.064, 0.125, 0.5, 0.25, and 0.125 mg/ L for *E. coli, K. pneumoniae, P. mirabilis* (TECOFF), *P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecium*, and *S. pneumoniae*, respectively.

At the same time, the MIC ECOFFs of levofloxacin were calculated using the ECOFFinder software. Combined with the visual method, we determined the MIC ECOFFs of levofloxacin to be 0.125, 0.25, 0.25, 2, 0.5, 1, 4, 8, and 2 mg/L for *E. coli, K. pneumoniae, P. mirabilis, P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecium*, and *S. pneumoniae* (Table S2), which is consistent with the data published on the EUCAST website.

Correlation Between MICs and Inhibition Zone Diameters

The distribution of MICs and inhibition zone diameters for the 9 species was shown in Figure 2. Except for *S. pneumoniae*, the other 8 species had Pearson correlation coefficients between MIC values and inhibition zone diameters were R > 0.8 (*P*<0.0001) (Table 2), with very strong correlation between two methods. However, the R value of *S. pneumoniae* was 0.46, which meant that the two methods were poorly correlated. Therefore, ECOFF of the *S. pneumoniae* zone diameter cannot be established.

Species/Lab	Number of Isolates With MIC (mg/L)											Total	Range	Modal MICs	Median	IQRs	99%					
	0.002	0.004	0.008	0.016	0.032	0.064	0.125	0.25	0.5	I	2	4	8	16	32	64	number	(mg/L)	(mg/L)	(mg/L)	(mg/L)	calculated ECOFFs (mg/L)
E. coli																						
PU	0	1 I	24	7	3	7	2	2	0	3	3	4	Т	0	0	0	57	0.004–8	0.008	0.016	0.045	
BD	0	0	7	3	3	0	3	6	0	3	3	2	0	0	0	0	30	0.008–4	0.008	0.125	0.984	
YF	1	6	12	1	13	5	0	1	2	6	3	3	Т	1	0	0	55	0.002-16	0.032	0.032	0.92	
нs	1	3	5	7	2	5	4	1	3	4	6	2	0	0	0	0	43	0.002–4	0.016	0.064	0.984	
₩Н	0	0	9	8	4	6	3	5	1	10	7	1	0	1	0	0	55	0.008-16	T	0.125	0.984	
Total	2	10	57	26	25	23	12	15	6	26	22	12	2	2	0	0	240	0.002-16	0.008	0.064	0.992	0.032
K. pneumoniae																						
PU	0	0	0	10	30	3	0	0	0	0	1	4	0	2	0	0	50	0.016-16	0.032	0.032	0	
BD	0	0	2	13	13	4	0	7	1	0	4	5	0	0	1	0	50	0.008–32	0.016; 0.032	0.032	0.234	
YF	0	0	0	27	7	3	0	2	2	0	0	0	0	2	6	1	50	0.016-64	0.016	0.016	0.234	
HS	0	1	3	14	4	I.	6	6	0	2	3	1	4	5	0	0	50	0.004–16	0.016	0.125	1.984	
₩Н	0	0	0	14	17	I.	3	2	2	Т	6	3	Т	0	0	0	50	0.016-8	0.032	0.032	0.484	
Total	0	1	5	78	71	12	9	17	5	3	14	13	5	9	7	I.	250	0.004–64	0.016	0.032	0.234	0.064
P. mirabilis																						
PU	0	0	0	6	20	4	3	5	2	6	2	3	0	0	0	0	51	0.016-4	0.032	0.032	0.468	
BD	0	0	0	2	11	9	0	4	2	7	1	1	0	0	0	0	37	0.016-4	0.032	0.064	0.468	
YF	0	0	2	I	8	7	6	5	5	3	0	1	0	0	0	0	38	0.008–4	0.032	0.25	0.218	
HS	0	0	0	0	10	T	T	4	8	4	3	3	2	0	3	0	39	0.032–32	0.032	0.5	1.968	
WН	0	0	0	4	15	2	3	5	4	5	2	0	Т	0	0	0	41	0.016-8	0.032	0.064	0.468	
Total	0	0	2	13	64	23	13	23	21	25	8	8	3	0	3	0	206	0.008–32	0.032	0.125	0.468	0.125 ^a
P. aeruginosa																						
PU	0	0	0	0	0	T	16	22	9	9	1	0	0	0	0	0	58	0.064–2	0.25	0.25	0.375	
BD	0	0	0	0	1 I	4	23	8	4	3	5	2	0	0	0	0	50	0.032-4	0.125	0.125	0.375	
YF	0	0	1 I	0	4	8	43	6	Т	2	3	3	4	0	0	0	75	0.008–8	0.125	0.125	0.125	
HS	0	0	0	0	1	13	21	7	10	3	4	4	0	0	0	1	64	0.032–64	0.125	0.125	0.375	
WН	0	0	0	0	1	13	59	21	2	3	1	3	Т	2	0	0	106	0.032-16	0.125	0.125	0.125	
Total	0	0	1	0	7	39	162	64	26	20	14	12	5	2	0	1	353	0.008–64	0.125	0.125	0.125	0.5
A. baumannii																						
PU	0	0	0	22	14	3	0	0	0	0	0	3	3	5	0	0	50	0.016-16	0.016	0.032	0.016	
BD	0	0	0	13	17	5	T	0	0	2	9	3	0	0	0	0	50	0.016-4	0.032	0.032	0.984	
YF	0	0	9	22	2	2	0	0	T	2	8	3	Т	0	0	0	50	0.008–8	0.016	0.016	0.984	
HS	0	0	2	16	9	7	0	0	0	Т	П	4	0	0	0	0	50	0.008–4	0.016	0.032	1.984	
₩Н	0	0	0	7	22	9	0	0	0	3	8	Т	0	0	0	0	50	0.016-4	0.032	0.032	0.032	
Total	0	0	11	80	64	26	1	0	Т	8	36	14	4	5	0	0	250	0.008–16	0.016	0.032	0.984	0.064

Table I MIC Distributions of Five Labs and (T)ECOFFs for Sitafloxacin Against Nine Species

(Continued)

Zhang et al

Table I	(Continued).
---------	--------------

Species/Lab	Number of Isolates With MIC (mg/L)												Total	Range	Modal MICs	Median	IQRs	99 %				
	0.002	0.004	0.008	0.016	0.032	0.064	0.125	0.25	0.5	I	2	4	8	16	32	64	number	(mg/L)	(mg/L)	(mg/L)	(mg/L)	calculated ECOFFs (mg/L)
S. aureus																						
PU	0	0	0	11	19	7	2	0	0	5	4	0	Т	1	0	0	50	0.016-16	0.032	0.032	0.093	
BD	0	0	0	1 I	19	13	6	0	0	4	5	1	0	0	1	0	50	0.016-32	0.032	0.064	0.093	
YF	0	0	1	9	18	7	0	1	3	5	4	1	0	1	0	0	50	0.008–16	0.032	0.032	0.468	
HS	0	0	1 I	12	23	2	0	0	4	6	0	0	2	0	0	0	50	0.008–8	0.032	0.032	0.048	
WH	0	0	0	1 I	16	21	T	0	2	5	2	2	0	0	0	0	50	0.016-4	0.064	0.064	0.032	
Total	0	0	2	34	95	50	9	1	9	25	15	4	3	2	Т	0	250	0.008–32	0.032	0.032	0.093	0.125
E. faecalis																						
PU	0	0	0	I	I	13	17	6	Т	7	4	0	0	0	0	0	50	0.016–2	0.125	0.125	0.186	
BD	0	0	0	0	0	0	22	14	2	4	6	2	0	0	0	0	50	0.125-4	0.125	0.25	0.375	
YF	0	0	0	0	I	П	21	3	1	5	8	0	0	0	0	0	50	0.032–2	0.125	0.125	0.875	
HS	0	0	0	0	0	10	26	4	0	4	7	2	0	0	0	0	53	0.064_4	0.125	0.125	0.125	
WH	0	0	0	0	0	8	25	4	Т	2	8	2	0	0	0	0	50	0.064_4	0.125	0.125	0.375	
Total	0	0	0	I	2	42	111	31	5	22	33	6	0	0	0	0	253	0.016-4	0.125	0.125	0.375	0.5
E. faecium																						
PU	I	T	1	0	2	10	4	3	10	6	2	Т	Т	0	0	0	42	0.002–8	0.064; 0.5	0.25	0.436	
BD	0	0	0	0	4	П	4	3	3	Т	4	8	Т	0	0	0	39	0.032–8	0.064	0.25	1.936	
YF	0	0	1	8	13	10	3	0	1	3	6	2	Т	0	0	0	48	0.008–8	0.064	0.064	0.468	
HS	0	0	0	I.	4	15	5	3	0	2	8	4	0	1	0	0	43	0.016-16	0.064	0.125	1.936	
WH	0	0	0	0	4	6	4	4	0	2	7	- I	Т	4	0	0	33	0.032-16	2	0.25	1.936	
Total	I	I	2	9	27	52	20	13	14	14	27	16	4	5	0	0	205	0.002-16	0.064	0.125	1.936	0.25
S. pneumoniae																						
PU	0	0	0	0	10	35	4	- I	0	0	0	0	0	0	0	0	50	0.032-0.25	0.064	0.064	0	
BD	0	0	0	0	20	30	0	0	0	0	0	0	0	0	0	0	50	0.032-0.064	0.064	0.064	0.032	
YF	0	0	1	3	39	3	0	3	1	0	0	0	0	0	0	0	50	0.008–0.5	0.032	0.032	0	
HS	0	I	0	4	42	7	2	0	0	I	0	0	0	0	0	0	57	0.004–1	0.032	0.032	0	
WH	0	0	0	2	25	22	I	0	0	0	0	0	0	0	0	0	50	0.016-0.125	0.032	0.064	0.032	
Total	0	I	1	9	136	97	7	4	1	I	0	0	0	0	0	0	257	0.004–1	0.032	0.032	0.032	0.125

Notes: a: Proteus mirabilis - We thought the data of YF was not amenable to analysis, resulting in only 4 distributions, and only a tentative ECOFF.



Figure I (A–I): The MIC distributions of E. coli, K. pneumoniae, P. mirabilis, P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecium, and S. pneumoniae, respectively. The dotted lines represent estimated ECOFFs of each species using ECOFFinder.

Inhibition Zone Diameter Distributions and ECOFFs Establishment

As shown in Figure 2, we visually estimated the zone diameter for 8 species except *S. pneumoniae*. Zone diameter ECOFFs were that *E. coli* and *S. aureus* was 26 cm, *K. pneumoniae* and *A. baumannii* was 25 cm, *P. mirabilis* and *P. aeruginosa* was 24 cm, *E. faecium* was 22 cm, *E. faecalis* was 21 cm.

Resistance Mechanisms Distribution in E. Coli Strains with Overlapped MICs Between WT and NWT Group

The combination of ECOFF inder and visual inspection led to the determination of a MIC ECOFF value of 0.032 mg/L for *E. coli*. As can be seen from Figure 1A, the number of *E. coli* with MIC values of 0.032 and 0.064 mg/L (close to calculated ECOFF) were very similar (25 and 23 strains, respectively), which indicate an overlap between WT and NWT strains in these two MIC distributions. To deeply investigate the resistance mechanisms distribution, these two groups of strains were selected for whole-genome sequencing to detect sitafloxacin resistance-related genes (*qnr*) or mutations (*gyrA/parC/parE*).



Figure 2 (A–I): Distribution of inhibition zone diameters and MICs (mg/L) of E. coli, K. pneumoniae, P. mirabilis, P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecalis, E. faecalis, and S. pneumoniae, respectively.

Twelve-point mutations associated with quinolone resistance in whole genomes were detected from 52.1% (25/48) of *E. coli* strains by using Pointfinder software. The mutations identified were *gyrA p.S83L*, *gyrA p.D87N*, *gyrA p.D87A*, *parC p.S57T*, *parC p.S80I*, *parC p.E84V*, *parE p.I529L*, *parE p.S458A*, *parE p.S458T*, *parE p.I355T*, *parE p.L416F*, and *parE p.P439S*. Among the bacteria with a MIC of 0.032 mg/L, 48.0% (12/25) of the strains had at least one mutation, with *gyrA p.S83L* being the most common mutation detected in up to 40.0% (10/25) of strains. In addition, *gyrA p.D87N* and *parC p.S80I* were also commonly detected at frequencies of 16.0% (4/25) and 12.0% (3/25), respectively. Among the

Species	Total Number	R	P value	Liner Regression Equation ^a
E. coli	240	-0.93	<0.0001	Y = -0.39X + 7.18
K. pneumoniae	250	-0.95	<0.0001	Y = -0.40X + 6.62
P. mirabilis	206	-0.81	<0.0001	Y = -0.32X + 6.91
P. aeruginosa	353	-0.89	<0.0001	Y = -0.24X + 4.65
A. baumannii	250	-0.94	<0.0001	Y = -0.44X + 8.49
S. aureus	250	-0.88	<0.0001	Y = -0.31X + 5.59
E. faecalis	253	-0.83	<0.0001	Y = -0.33X + 5.73
E. faecium	205	-0.92	<0.0001	Y = -0.39X + 6.80
S. pneumoniae	257	-0.46	<0.0001	Y = −0.13X - 0.28

 Table 2 The Correlation of MIC Values and Inhibition Zone Diameters for 9

 Species

Notes: a: Y is log2 (MIC), X is the inhibition diameter (mm).



Figure 3 (A) Distribution of resistance mechanisms of E. coli with MIC of 0.032 and 0.064 mg/L in 48 strains. (B) ECOFFinder calculated the ECOFF value of E. coli.

bacteria with a MIC of 0.064 mg/L, mutation sites were detected in 65.2% (15/23) of strains, with *gyrA p.S83L, gyrA p.* D87N, and *parC p.S80I* being the most common mutations detected at frequencies of 56.5% (13/23), 8.7% (2/23), and 13.0% (3/23), respectively (Figure 3A).

Resfinder software detected two sitafloxacin resistance-related genes, *qnrS1* and *qnrB19*, from the whole genomes of 20.8% (10/48) of *E. coli*. Among the 25 strains with a MIC of 0.032 mg/L, 4 strains carried *qnrS1* gene and 1 strain carried *qnrB19* gene. In 23 strains with a MIC of 0.064 mg/L, 4 *qnrS1* gene and 1 *qnrB19* gene-carried isolates were also detected (Figure 3A).

Overall, 17 strains of *E. coli* with MIC of 0.032 mg/L carried resistance mechanisms, and 8 strains did not carry any resistance mechanisms (neither resistance genes nor point mutations). Among *E. coli* with MIC of 0.064 mg/L, 20 strains carried resistance mechanism, and only 3 strains did not carry any quinolone–resistance determinants. The resistance mechanism distribution exhibited statistical difference between the two groups (P = 0.003).

Discussion

Nowadays, the widespread use of antibiotics has led to a large increase in the number of various drug-resistant strains, which has increased the difficulty of clinical treatment. As a result, new antibiotics are constantly being developed to combat drug-resistant strains with their better antimicrobial activity. Sitafloxacin is a new fluoroquinolone drug, which has good antibacterial activity against gram-negative bacteria and positive bacteria.³²

From the data on the EUCAST website, the MIC (T)ECOFF of levofloxacin for these 9 strains was 0.125, 0.25, 0.25, 2, 0.5, 1, 4, 8, and 2 mg/L for *E. coli, K. pneumoniae, P. mirabilis, P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecalis, and S. pneumoniae*, respectively.³³ In comparison, the MIC (T)ECOFF of levofloxacin are larger than the sitafloxacin established in this study. This demonstrates that sitafloxacin exhibits more potent antibacterial activity compared to levofloxacin. The introduction of a methoxy group ($-OCH_3$) at the C-8 position in sitafloxacin significantly enhances its affinity and stability for DNA gyrase. This modification enables the drug to bind more securely to the β -subunit, reducing the dissociation rate of the enzyme and thereby prolonging the duration of inhibition. Compared to the shorter side chain ($-NHCOCH_3$) in levofloxacin, sitafloxacin features a longer C-7 substituent ($-CF_3-CH_2-N(CH_3)_2$), which may enhance hydrophobicity, promote penetration of the bacterial outer membrane lipid bilayer, and increase intracellular concentrations.³⁴

In recent years, some domestic reports have also confirmed that sitafloxacin has better efficacy. Li et al evaluated the efficacy and safety of sitafloxacin and levofloxacin in the treatment of complicated urinary tract infections in adults in China. The clinical cure rate and microbiological success rate of sitafloxacin in the treatment of complicated UTI were better than levofloxacin, especially bacterial eradication rate, which was 93.3% (14/15) versus 63.6% (7/11).⁷ Similarly, the pharmacokinetic/pharmacodynamic studies of multidrug-resistant bacteria in the model of sitafloxacin in dynamic urinary tract infection showed significantly lower MIC values than levofloxacin for drug-resistant *Escherichia coli*, *Pseudomonas aeruginosa*, and *Enterococcus faecium*.³⁵ Shi et al studied the comparison of the activity of sitafloxacin, levofloxacin and 1–2 concentration gradients smaller than moxifloxacin.⁹ So sitafloxacin exhibited strong bactericidal activity against organisms resistant to other fluoroquinolones. Because sitafloxacin had not yet developed a clinical breakpoint, it was important to establish ECOFF first.

The distribution of sitafloxacin MIC and zone diameter was relatively wide, ranging from 0.002 to 64 mg/L and from 6 to 45 mm, respectively. The 99% calculated ECOFFs was one concentration gradient smaller than the verified ECOFFs, with *P. mirabilis, P. aeruginosa*, and *E. faecalis*. Among *P. aeruginosa* and *E. faecalis*, we may believe that visual results of visual estimate method were more inclined to 0.5 mg/L than 0.25 mg/L.

Regarding *P. mirabilis*, the MIC ECOFF ECOFF inder calculated was 0.064 mg/L. However, we thought that the data of YF was not amenable to analysis about *P. mirabilis*. In addition to YF, the *P. mirabilis* data of the other four labs had obvious peak. However, the YF data did not show obvious quantitative changes from 0.032–0.5 mg/L concentration gradient, and the changes of YF enrollment *P. mirabilis* could not be judged. This may be related to the resistance mechanism of the enrolled strains, or it may be that laboratory staff readings were biased.

Among the 9 species in this study, the results of *P. aeruginosa* were the most ideal from the experimental data obtained. Its MIC distribution was symmetrical with fewer NWT strains. The presence of an obvious resistance mechanism may not be evident from the MIC distribution.

As shown in Figure 1, there were 2 peaks of different sizes in the distribution of *Escherichia coli* MIC. It may be because *Escherichia coli* enrolled showed 2 different resistance mechanisms. The molecular mechanisms of fluoroquinolone resistance, including plasmid-mediated quinolone resistance (PMQR) along with mutation in the quinolone-resistance determining regions (QRDRs) of *gyrA*, *gyrB*, *parC* and *parE*.^{36–38} The clinically common extended-spectrum β -lactamase-producing (ESBL) *Enterobacteriaceae* also have more than one-third resistance to fluoroquinolones.³⁹

Furthermore, the analysis results of whole genome drug resistance information showed a similar situation of carrying drug resistance genes and point mutations for the two groups of *E. coli* with MIC values of 0.032 and 0.064 mg/L. The strains with MIC values less than or equal to the ECOFF value were WT strains and did not carry any resistance mechanism. In this study, only 8 out of the 25 *E. coli* strains with MIC values of 0.032 mg/L did not carry drug resistance mechanisms. It was possible that some of the low-tolerance strains in the NWT strains crossed with the WT strains near the ECOFF value (Figure 3B). We followed EUCAST SOP 10.2 as described, the WT distribution is visually assessed to be a log-normal distribution.

Except for *S. pneumoniae*, all other species showed a high correlation between MIC and zone diameter (R > 0.8, P < 0.0001). Because it was more convenient to operate the disk diffusion method in the microbiology laboratories, we recommend the disk diffusion method to test the resistance for 8 strains other than *S. pneumoniae* to sitafloxacin. Most of the MIC values obtained by our enrolled *S. pneumoniae* distributed in concentrations of 0.032 and 0.064 mg/L, and the largest MIC value was 1 mg/L. The ECOFF of *S. pneumoniae* MIC were 0.125 mg/L, so the number of NWT in the enrolled strains was very small, and it was not good to analyze the reasons for the poor consistency of the results of the two methods of MIC and disk diffusion method from the level of resistance genes. In conclusion, we did not recommend the disk diffusion method in clinical microbiology laboratories to detect the tolerance of *S. pneumoniae* to sitafloxacin.

In summary, we established MIC and zone diameter ECOFFs for sitafloxacin against *E. coli, K. pneumoniae, P. mirabilis, P. aeruginosa, A. baumannii, S. aureus, E. faecalis, E. faecium*, and *S. pneumoniae*. The MIC ECOFF of the above nine strains were 0.032, 0.064, 0.125, 0.5, 0.064, 0.125, 0.5, 0.25, and 0.125 mg/L, respectively. The zone diameter ECOFF of the above nine strains were 26, 25, 24, 24, 25, 26, 21.22, and NA. The MIC ECOFF for *P. mirabilis* remains tentative pending further validation. For *S. pneumoniae*, MICs and zone diameters showed weak correlation, preventing the establishment of zone diameter ECOFFs.

Data Sharing Statement

The WGS data of 48 *E.coli* to are available in the China National Center for Bioinformation (<u>https://ngdc.cncb.ac.cn</u>) under project number PRJCA023379.

Ethical Approval

This study complies with the Declaration of Helsinki and was approved by the Health Research Ethics Board of Peking Union Medical College Hospital (No. S-K1240).

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

The authors acknowledge the National High-Level Hospital Clinical Research Funding Grant (2022-PUMCH-B-074), (2022-PUMCH-B-028), (2022-PUMCH-C-060), the National Natural Science Foundation of China (82272380, 82072318) and the National Key Research and Development Program of China (2021YFC2301002) for financial support.

Disclosure

The authors declare no conflict of interest.

References

- 1. Hussen NH, Hamid SJ, Sabir MN. et al Novel penicillin derivatives against selected multiple-drug resistant bacterial strains: design, synthesis, structural analysis, in silico and in vitro studies. *Curr Org Synth.* 2024;21:684–703. doi:10.2174/1570179420666230510104319
- 2. Thiazole-pyrazoline hybrids as potential antimicrobial agent: synthesis, biological evaluation, molecular docking, DFT studies and POM analysis. *J mol Struct.* 2023;1282. doi:10.1016/j.molstruc.2023.135191
- 3. Chen CK, Cheng IL, Chen YH, Lai CC. Efficacy and safety of sitafloxacin in the treatment of acute bacterial infection: a meta-analysis of randomized controlled trials. *Antibiotics*. 2020;9:106. doi:10.3390/antibiotics9030106
- 4. Okumura R, Hirata T, Onodera Y, et al. Dual-targeting properties of the 3-aminopyrrolidyl quinolones, DC-159a and sitafloxacin, against DNA gyrase and topoisomerase IV: contribution to reducing in vitro emergence of quinolone-resistant Streptococcus pneumoniae. J Antimicrob Chemother. 2008;62:98–104. doi:10.1093/jac/dkn136
- 5. Jolley A, Andrews JM, Brenwald N, Wise R. The in-vitro activity of a new highly active quinolone, DU-6859a. J Antimicrob Chemother. 1993;32:757-763. doi:10.1093/jac/32.5.757
- 6. Keating GM. Sitafloxacin: in bacterial infections. Drugs. 2011;71:731-744. doi:10.2165/11207380-00000000-00000
- Li Y, Yin Y, Peng X, et al. A randomized, active-controlled, multicentre clinical trial to evaluate the efficacy and safety of oral sitafloxacin versus levofloxacin in Chinese adults with acute uncomplicated or complicated urinary tract infection. Ann Med. 2021;53:217–226. doi:10.1080/ 07853890.2020.1861322
- 8. Li Y, Zhu D, Peng Y, et al. A randomized, controlled, multicenter clinical trial to evaluate the efficacy and safety of oral sitafloxacin versus moxifloxacin in adult patients with community-acquired pneumonia. *Curr Med Res Opin*. 2021;37:693–701. doi:10.1080/03007995.2021.1885362
- 9. Shi G, Chen X, Wang H, et al. Activity of sitafloxacin against extracellular and intracellular Staphylococcus aureus in vitro and in vivo: comparison with levofloxacin and moxifloxacin. *J Antibiot*. 2012;65:229–236. doi:10.1038/ja.2012.7
- Sato K, Hoshino K, Tanaka M, Hayakawa I, Osada Y. Antimicrobial activity of DU-6859, a new potent fluoroquinolone, against clinical isolates. *Antimicrob Agents Chemother*. 1992;36:1491–1498. doi:10.1128/AAC.36.7.1491
- 11. Nishizawa T, Munkjargal M, Ebinuma H, Toyoshima O, Suzuki H. Sitafloxacin for third-line helicobacter pylori eradication: a systematic review. *J Clin Med.* 2021;10:2722. doi:10.3390/jcm10122722
- 12. He S, Guo Q, Zhao L, et al. Sitafloxacin expresses potent anti-mycobacterium abscessus activity. Front Microbiol. 2021;12:779531. doi:10.3389/ fmicb.2021.779531
- 13. Miyazaki T, Nakamura S, Hashiguchi K, et al. The efficacy and safety of sitafloxacin and garenoxacin for the treatment of pneumonia in elderly patients: a randomized, multicenter, open-label trial. J Infect Chemother. 2019;25:886–893. doi:10.1016/j.jiac.2019.04.019
- 14. Xu N, Wang G, Leng Y, et al. Sulbactam enhances the in vitro activity of sitafloxacin against extensively-drug resistant Acinetobacter baumannii. Exp Ther Med. 2018;16:3485–3491. doi:10.3892/etm.2018.6630
- 15. Leechawengwongs M, Prammananan T, Jaitrong S, et al. In vitro activity and mic of sitafloxacin against multidrug-resistant and extensively drug-resistant mycobacterium tuberculosis isolated in Thailand. *Antimicrob Agents Chemother*. 2018;62. doi:10.1128/AAC.00825-17
- 16. Yi L, Aono A, Chikamatsu K, et al. In vitro activity of sitafloxacin against Mycobacterium tuberculosis with gyrA/B mutations isolated in Japan. J Med Microbiol. 2017;66:770–776. doi:10.1099/jmm.0.000493

- Huang YS, Wang JT, Sheng WH, Chuang YC, Chang SC. Comparative in vitro activity of sitafloxacin against bacteremic isolates of carbapenem resistant Acinetobacter baumannii complex. J Microbiol Immunol Infect. 2015;48:545–551. doi:10.1016/j.jmii.2014.02.002
- Manosuthi W, Wiboonchutikul S. Treatment outcomes of oral sitafloxacin in acute complicated urinary tract infection and pyelonephritis. Springerplus. 2016;5:410. doi:10.1186/s40064-016-2044-5
- Malaisri C, Phuphuakrat A, Wibulpolprasert A, Santanirand P, Kiertiburanakul S. A randomized controlled trial of sitafloxacin vs. ertapenem as a switch therapy after treatment for acute pyelonephritis caused by extended-spectrum beta-lactamase-producing Escherichia coli: a pilot study. J Infect Chemother. 2017;23:556–562. doi:10.1016/j.jiac.2017.05.005
- 20. Paiboonvong T, Nosoongnoen W, Sathirakul K, et al. Pharmacokinetics and penetration of sitafloxacin into alveolar epithelial lining fluid in critically III Thai patients with pneumonia. *Antimicrob Agents Chemother*. 2019;63. doi:10.1128/AAC.00800-19
- 21. Yang Q, Li X, Jia P, et al. Determination of norvancomycin epidemiological cut-off values (ECOFFs) for Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus haemolyticus and Staphylococcus hominis. J Antimicrob Chemother. 2021;76:152–159. doi:10.1093/jac/dkaa414
- 22. Vading M, S. BH, Sundsfjord AS, Giske CG, Giske CG. Comparison of disk diffusion, Etest and VITEK2 for detection of carbapenemase-producing Klebsiella pneumoniae with the EUCAST and CLSI breakpoint systems. *Clin Microbiol Infect.* 2011;17:668–674. doi:10.1111/j.1469-0691.2010.03299.x
- 23. Froberg G, Maurer FP, Chryssanthou E, et al. Towards clinical breakpoints for non-tuberculous mycobacteria determination of epidemiological cut off values for the mycobacterium avium complex and Mycobacterium abscessus using broth microdilution. *Clin Microbiol Infect.* 2023;29:758–764. doi:10.1016/j.cmi.2023.02.007
- 24. Schon T, Werngren J, Machado D, et al. Multicentre testing of the EUCAST broth microdilution reference method for MIC determination on Mycobacterium tuberculosis. *Clin Microbiol Infect.* 2021;27:288e281–288e284. doi:10.1016/j.cmi.2020.10.019
- 25. Tack B, Phoba M-F, Thong P, et al. Epidemiological cut-off value and antibiotic susceptibility test methods for azithromycin in a collection of multi-country invasive non-typhoidal Salmonella. *Clin Microbiol Infect*. 2022;28:1615–1623. doi:10.1016/j.cmi.2022.06.009
- 26. Procedure for optimizing disk contents (potencies) for disk diffusion testing of antimicrobial agents using harmonized CLSI and EUCAST criteria. *Standard Operating Procedure EUCAST SOP.* 2020;11.
- 27. Lim HJ, Lee EH, Yoon Y, Chua B, Son A. Portable lysis apparatus for rapid single-step DNA extraction of bacillus subtilis. *J Appl Microbiol*. 2016;120:379–387. doi:10.1111/jam.13011
- 28. Testing TECOAS. MIC distributions and epidemiological cut-off value (ECOFF) setting. EUCAST SOP. 2019;10(1).
- 29. Karatuna O, Dance DAB, Matuschek E, et al. Burkholderia pseudomallei multi-centre study to establish EUCAST MIC and zone diameter distributions and epidemiological cut-off values. *Clin Microbiol Infect.* 2020. doi:10.1016/j.cmi.2020.07.001
- Bortolaia V, Kaas RS, Ruppe E, et al. ResFinder 4.0 for predictions of phenotypes from genotypes. J Antimicrob Chemother. 2020;75:3491–3500. doi:10.1093/jac/dkaa345
- 31. Zankari E, Allesøe R, Joensen KG, et al. PointFinder: a novel web tool for WGS-based detection of antimicrobial resistance associated with chromosomal point mutations in bacterial pathogens. J Antimicrob Chemother. 2017;72:2764–2768. doi:10.1093/jac/dkx217
- 32. Tateda K, Ohno A, Ishii Y, et al. Investigation of the susceptibility trends in Japan to fluoroquinolones and other antimicrobial agents in a nationwide collection of clinical isolates: a longitudinal analysis from 1994 to 2016. J Infect Chemother. 2019;25:594–604. doi:10.1016/j. jiac.2019.03.008
- 33. EUCAST. MIC and zone diameter distributions and ECOFFs. Available from: https://www.eucast.org/mic_distributions_and_ecoffs/. Accessed April 14, 2025
- 34. Kuhn EMA, Sominsky LA, Chittò M, Schwarz EM, Moriarty TF. Antibacterial mechanisms and clinical impact of sitafloxacin. *Pharmaceuticals*. 2024;17:1537. doi:10.3390/ph17111537
- 35. Guo S, Li X, Li Y, et al. Sitafloxacin pharmacokinetics/pharmacodynamics against multidrug-resistant bacteria in a dynamic urinary tract infection in vitro model. *J Antimicrob Chemother*. 2022;78:141–149. doi:10.1093/jac/dkac365
- 36. Piekarska K, Zacharczuk K, Wołkowicz T, et al. The molecular mechanisms of fluoroquinolone resistance found in rectal swab isolates of enterobacterales from men undergoing a transrectal prostate biopsy: the rationale for targeted prophylaxis. *Ann Clin Microbiol Antimicrob*. 2021;20:81. doi:10.1186/s12941-021-00487-y
- Touyama M, Higa F, Nakasone C, et al. In vitro activity of sitafloxacin against clinical strains of Streptococcus pneumoniae with defined amino acid substitutions in QRDRs of gyrase A and topoisomerase IV. J Antimicrob Chemother. 2006;58:1279–1282. doi:10.1093/jac/dkl427
- 38. Mayumi Tanaka TW, Onodera Y, Uchida Y, Sato K, Sato K. Mechanism of quinolone resistance in staphylococcus aureus. J Infect Chemother. 2000;6:131–139. doi:10.1007/s101560070010
- Dalhoff A. Resistance surveillance studies: a multifaceted problem--the fluoroquinolone example. *Infection*. 2012;40:239–262. doi:10.1007/s15010-012-0257-2

Infection and Drug Resistance



Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/infection-and-drug-resistance-journal

2004 🖪 💥 in 🔼