

# Sex Differences in Clinical Features and Outcomes in Patients with Acute Coronary Syndrome Treated with Bifurcation Stenting Using the Double-Kissing Culotte and Culotte Technique – 1-year Follow-up

Piotr Włodarczak<sup>1</sup>, Mateusz Barycki<sup>2</sup>, Szymon Włodarczak<sup>1</sup>, Łukasz Furtan<sup>2</sup>, Andrzej Giniewicz<sup>3</sup>, Anita Chudaś<sup>1</sup>, Waldemar Banasiak<sup>4,5</sup>, Adrian Doroszko<sup>4,5</sup>, Maciej Lesiak<sup>6</sup>, Adrian Włodarczak<sup>1,4</sup>, Piotr Rola<sup>2,4</sup>

<sup>1</sup>Department of Cardiology, The Copper Health Centre (MCZ), Lubin, 59-301, Poland; <sup>2</sup>Department of Cardiology, Provincial Specialized Hospital, Legnica, 59-220, Poland; <sup>3</sup>Faculty of Pure and Applied Mathematics, Wrocław University of Science and Technology, Wrocław, 50-981, Poland; <sup>4</sup>Faculty of Medicine, Wrocław University of Science and Technology, Wrocław, 50-981, Poland; <sup>5</sup>Department of Cardiology, Center for Heart Diseases, 4th Military Hospital, Wrocław, 50-981, Poland; <sup>6</sup>1st Department of Cardiology, University of Medical Sciences, Poznań, 61-848, Poland

Correspondence: Piotr Rola, Faculty of Medicine, Wrocław University of Science and Technology, Wrocław, 50-981, Poland, Tel +0048767211442, Email Piotr.rola@gmail.com

**Purpose:** Sex differences in the clinical course of coronary artery disease (CAD) particularly in Acute Coronary Syndrome (ACS), have been extensively hypothesized. Although coronary bifurcations account for approximately 20% of ACS cases, data concerning potential sex disparities in the outcomes of percutaneous interventions (PCI) remain scarce. Several data suggest the double kiss (DK) Culotte may provide advantages in bifurcation PCI. Therefore, we evaluate potential sex differences in relation to the Culotte technique.

**Patients and Methods:** This study retrospectively analyzes sex disparities following PCI interventions in ACS patients using the DK-Culotte or Culotte technique for bifurcation lesions. The primary endpoint was Target Lesion Failure (TLF), a composite of cardiovascular death, target vessel myocardial infarction, or clinically driven target lesion revascularization (TLR). The secondary endpoint included major adverse cardiac events MACE (myocardial infarction, cardiac death, and target lesion revascularization).

**Results:** There were no significant differences between sexes regarding TLF (DK-Culotte: Females 10.3% vs Males 5.7%;  $p=0.401$ ; Culotte: Females 16.2% vs Males 11.8%;  $p=0.481$ ) or MACE (DK-Culotte: Females 13.8% vs Males 12.5%;  $p=0.771$ ; Culotte: Females 24.3% vs Males 17.6%;  $p=0.370$ ) after a 1-year follow-up for both bifurcation techniques.

**Conclusion:** The study found no significant differences in clinical outcomes between sexes following PCI for bifurcation lesions in the ACS cohort, regardless of whether the two-stent techniques (DK- Culotte or Culotte) were used.

**Keywords:** sex differences, gender differences, acute coronary syndrome, coronary bifurcation, double-kiss culotte, culotte technique, percutaneous coronary intervention

## Introduction

The average age of patients undergoing cardiac revascularization during an acute coronary syndrome (ACS) event is on the rise. This is in line with a broader demographic shift in modern societies. As a result, patients undergoing percutaneous coronary intervention (PCI) for ACS are more likely to have an increased prevalence of independent adverse events prognostic factors (diabetes and renal insufficiency) after percutaneous revascularization.<sup>1–3</sup> Furthermore, both phenomena have been demonstrated to increase the probability of the occurrence of complex lesions, which emphasizes the necessity to pay close attention to the most appropriate PCI technique.

Interestingly, these demographic changes have erased the traditional differences between sexes in the prevalence of coronary artery disease.<sup>4</sup> However, females still tend to be underrepresented in many contemporary clinical trials and are

more likely to receive conservative therapy in the course of acute coronary syndrome.<sup>5</sup> Furthermore, even if they receive percutaneous revascularization, the comprehensiveness of treatment along with long-term outcomes is less favorable in comparison to males.<sup>6,7</sup> The precise mechanisms underlying these discrepancies remain elusive, despite the undeniable advancements in interventional techniques and armamentarium observed in recent times in the field of percutaneous revascularization.

Coronary bifurcation lesions are a prevalent phenomenon, representing approximately 20%<sup>8</sup> of percutaneous coronary intervention (PCI) cases, and continuing to present a significant challenge for interventional cardiologists, particularly in the context of acute coronary syndrome (ACS) conditions. Although the recently presented expert consensus considers the provisional single-stent technique to be the most convenient for the majority of bifurcation lesions,<sup>9</sup> Nevertheless, in select cases, including those with significant anatomical complexity where diseases affect critical side branches, two-stent techniques may offer clinical advantages when compared with other options.<sup>10</sup>

An increasing body of evidence indicates that the double kiss (DK)-crush technique may offer benefits in the context of left main bifurcation, particularly in comparison with the classic crush technique.<sup>11</sup> A minor modification to the conventional crush methodology, which involved the incorporation of an additional kissing balloon (KB), led to a notable enhancement in the long-term outcomes of the DK crush.<sup>12</sup> On the other hand, since the introduction of the Culotte technique into clinical practice,<sup>13</sup> this bifurcation stenting technique has become one of the most commonly used procedures. It has been demonstrated to be safe and efficient<sup>14</sup> and its use is now widespread. A review of recent pre-clinical trials and a small number of pilot studies suggests that additional kissing dilatation before main branch (MB) stenting may improve the culotte technique in a manner similar to what was observed in the context of the crush technique.<sup>15–17</sup>

In light of the data presented, this study was designed to evaluate the difference between sexes in the short-term clinical outcome (one year) in ACS patients with culprit true bifurcation lesions treated with one of two stent techniques (Culotte vs DK-Culotte).

## Materials and Methods

### Study Population

The complete trial design, along with a full list of inclusion and exclusion criteria, was previously reported (the study was registered in ClinicalTrials.gov: NCT06284057).<sup>18</sup> In brief, the study population is constituted by all subjects included in the Lower Silesia Culotte Bifurcation Registry (LSCBR). This registry comprises clinical data from two cooperating high-volume cardiac centers of subjects with ACS who underwent percutaneous coronary intervention (PCI) of a culprit bifurcation lesion using two different two-stent techniques, the DK Culotte and the Culotte. The decision to perform PCI was based on clinical indications, in accordance with the European Society of Cardiology recommendations. The decision to perform PCI with the selected two-stent technique was left to the discretion of the operators. All PCI procedures were preformed with second-generation drug-eluting stents - Orsiro (Biotronik, Berlin, Germany), Xience (Abbott Chicago, USA), Onyx (Medtronic, Dublin, Republic of Ireland) Cruz (SMT, Mumbai, India) Biofreedom (Biosensor, Singapore), Synergy (Boston Scientific, Marlborough, USA). Prior to undergoing percutaneous coronary intervention (PCI), all patients were provided with comprehensive information regarding the procedure and its associated risks. This information was presented in a written consent form, which the patient was required to sign before the procedure could be performed. The Bioethics Committee of the Lower Silesian Medical Association (Poland), which oversees research centers in accordance with local legal restrictions, has approved this study (01/BO/2023). Furthermore, the study is in accordance with the Helsinki Declaration.

### Study Endpoints

The study's main points and clinical outcomes were in line with general consensus<sup>19</sup> The study had composed of Primary Endpoint- Target Lesion Failure (TLF)), which consisted of cardiovascular death (CVD), target vessel myocardial infarction (TV-MI), or clinically driven target lesion revascularization (TLR). Furthermore, the study revealed the prevalence of several secondary endpoints, including major adverse cardiac events (MACE), which encompasses myocardial infarction, cardiac death, and target lesion revascularization (TLR), along with TLR, overall mortality, and selected procedural factors (radiation dose and contrast use).

## Statistical Analysis

Continuous variables were evaluated using nonparametric two-sample Mann–Whitney *U*-tests, whereas categorical variables were analyzed using Fisher's exact tests. Kaplan-Meier curves were employed to illustrate the cumulative one-year rates of target lesion failure, major adverse cardiovascular events, all-cause death, and target lesion revascularization in the study groups. Log rank tests were used to assess the significance of the observed differences. The statistical cutoff point for significance was established at 0.05. The statistical analyses were conducted by an experienced medical statistician, who used the R programming language to perform analysis.

## Results

### Patient Characteristic

The Culotte cohort comprised 85 males and 37 females, thus the DK-Culotte arm consisted of 88 males and 29 females. With regard to the basic clinical characteristics of the Culotte cohort, a significantly lower mean age was observed in the male group compared to the female group ( $64.6 \pm 8.7$  vs  $70.7 \pm 8.7$ ;  $p = 0.006$ ). Furthermore, a lower prevalence of type 2 diabetes mellitus was observed in the male subjects compared to the female subjects (35 (41.2%) vs 23 (62.2%),  $p = 0.048$ ). Conversely, the initial creatinine levels were found to be significantly higher in the male subjects of this study arm compared to the female subjects ( $88.4$  [75–102.5] vs  $79.1$  [63.6–90];  $p = 0.013$ ). On the other hand, except for a similar trend in creatinine levels in the DK-Culotte group ( $83.2$  [72.2–99.8] vs  $73$  [63.5–92.4];  $p = 0.033$ ), no significant differences in terms of basic clinical features were observed. All basic clinical and demographic data are summarized in [Table 1](#).

### Lesions and Procedural Features

In terms of anatomical and procedural characteristics in the Culotte arm we could observe statistically lower anatomical complexity of CAD measured by Syntax I Score in the male cohort compared to females (15 [12–22] vs 18 [14–29];  $p = 0.046$ ). At the same time in DK - Culotte group males compared to females were more prone to receive radial access point during index PCI (89.8% vs 69%;  $p = 0.015$ ). On the other hand, men in this arm received higher contrast ( $235.6 \pm 75.4$  vs  $205.3 \pm 63.5$ ;  $p = 0.039$ ) and radiation doses ( $1950.5$  [1324.2–3016.5] vs  $1604$  [693–2658];  $p = 0.027$ ). [Table 2](#) contains all lesions and procedural features.

**Table 1** Baseline Clinical Characteristics of Both Study Arms

	Culotte Group			DK Culotte Group		
	Male N-85	Female N-37	p-value	Male N-88	Female N-29	p-value
Age [years]	$64.6 \pm 8.7$	$70.7 \pm 8.7$	$p = 0.006$	$67.4 \pm 9.0$	$65.4 \pm 10.5$	$p = 0.377$
Unstable angina	33 (38.8%)	10 (27%)	$p = 0.010$	43 (48.9%)	13 (44.8%)	$p = 0.958$
NSTEMI	31 (36.5%)	24 (64.9%)	$p = 0.010$	30 (34.1%)	11 (37.9%)	$p = 0.958$
STEMI	21 (24.7%)	3 (8.1%)	$p = 0.010$	15 (17%)	5 (17.2%)	$p = 0.958$
Diabetes mellitus type 2	35 (41.2%)	23 (62.2%)	$p = 0.048$	25 (28.4%)	11 (37.9%)	$p = 0.359$
Oral anti-diabetic treatment	30 (35.3%)	16 (43.2%)	$p = 0.043$	20 (22.7%)	11 (37.9%)	$p = 0.117$
Insulin	5 (5.9%)	7 (18.9%)	$p = 0.043$	5 (5.7%)	5 (17.2%)	$p = 0.117$
Hypertension	70 (82.4%)	33 (89.2%)	$p = 0.423$	70 (79.5%)	24 (82.8%)	$p = 0.793$
Hyperlipidemia	59 (69.4%)	32 (86.5%)	$p = 0.069$	75 (85.2%)	23 (79.3%)	$p = 0.561$
Atrial Fibrillation	19 (22.4%)	8 (21.6%)	$p = 1$	10 (11.4%)	3 (10.3%)	$p = 1$
Post PCI status	25 (29.4%)	17 (45.9%)	$p = 0.098$	28 (31.8%)	9 (31%)	$p = 1$
Primary Diagnosis of MI	19 (22.4%)	11 (29.7%)	$p = 0.493$	24 (27.3%)	10 (34.5%)	$p = 0.485$
LVEF	55 [43–60]	45 [35–60]	$p = 0.193$	58 [45–63]	55 [50–60]	$p = 0.514$
Total Cholesterol (mmol/L)	$5 \pm 1.6$	$4.8 \pm 1.4$	$p = 0.579$	$4.7 \pm 1.3$	$4.5 \pm 0.9$	$p = 0.490$
LDL (mmol/L)	$2.5$ [1.9–3.6]	$2.8$ [1.9–3.6]	$p = 0.923$	$2.7 \pm 1.2$	$2.6 \pm 0.9$	$p = 0.573$
HDL (mmol/L)	$1.2$ [1–1.5]	$1.3$ [1.1–1.6]	$p = 1$	$1.2$ [1–1.4]	$1.3$ [1.1–1.8]	$p = 0.08$
Creatine (μmol/l)	$88.4$ [75–102.5]	$79.1$ [63.6–90]	$p = 0.013$	$83.2$ [72.2–99.8]	$73$ [63.5–92.4]	$p = 0.033$
Maximal TnI peak	$186.1$ [62–1359.5]	$431$ [108.8–1363.8]	$p = 0.342$	$155.1$ [33.2–1067.6]	$354.1$ [113.9–702.6]	$p = 0.295$

**Abbreviations:** NSTEMI, no ST-Elevation Myocardial Infarction; STEMI ST-Elevation Myocardial Infarction; PCI, percutaneous coronary intervention; MI, Myocardial Infarction; LVEF, Left ventricle ejection fraction.

**Table 2** Procedural Characteristics of Both Study Arms

Procedural Characteristic	Culotte Group			DK Culotte Group		
	Male N-85	Female N-37	p-value	Male N-88	Female N-29	p-value
SYNTAX Score I	15 [12–22]	18 [14–29]	p = 0.046	15.5 [11.8–22.2]	13 [9–16]	p = 0.107
Logistic SYNTAX Score	3.1 [1.4–9.6]	7.1 [3–23.2]	p = 0.004	2.9 [1.4–5.9]	2.9 [2.1–5.2]	p = 0.682
PCI SYNTAX Score II	31.3 ± 13	43.4 ± 14.5	p < 0.001	28.5 [21.8–37.4]	33.1 [24.6–43.7]	p = 0.233
Bifurcation location: LM	29 (34.1%)	12 (32.4%)	p = 1	35 (39.8%)	12 (41.4%)	p = 1
Non-LM (LAD/D)	28 (32.9%)	15 (40.5%)	p = 0.419	32 (36.4%)	16 (55.2%)	p = 0.085
Non-LM (Cx/OM)	24 (28.2%)	7 (18.9%)	p = 0.367	16 (18.2%)	1 (3.4%)	p = 0.067
Non-LM (RCA/PLA/PDA)	4 (4.7%)	3 (8.1%)	p = 0.432	5 (5.7%)	0 (0%)	p = 0.331
Femoral access	10 (11.8%)	10 (27%)	p = 0.060	9 (10.2%)	9 (31%)	p = 0.015
Radial access	74 (87.1%)	28 (75.7%)	p = 0.182	79 (89.8%)	20 (69%)	p = 0.015
Bail out two stent strategy	7 (8.2%)	2 (5.4%)	p = 0.721	7 (8%)	2 (6.9%)	p = 1
Side branch stent diameter, mm	3 [2.5–3.5]	2.8 [2.5–3.5]	p = 0.173	3 [2.8–3.5]	2.8 [2.5–3]	p = 0.062
Side branch stent length, mm	22 [18–28]	22 [18–26]	p = 0.849	20 [18–26]	18 [18–22]	p = 0.303
Main branch stent diameter, mm	3.5 [3–3.5]	3 [3–3.5]	p = 0.232	3.5 [3–3.5]	3.5 [3–3.5]	p = 0.386
Main branch stent length, mm	26 [18–30]	26 [18–28]	p = 0.418	26 [18–34.2]	18 [18–26]	p = 0.006
Stent to the side branch first	72 (84.7%)	31 (83.8%)	p = 1	74 (84.1%)	25 (86.2%)	p = 1
Side branch predilatation	0 (0%)	0 (0%)	p = 1	88 (100%)	29 (100%)	p = 1
Main branch predilatation	75 (88.2%)	31 (83.8%)	p = 0.563	63 (71.6%)	22 (75.9%)	p = 0.811
Pre POT	38 (44.7%)	17 (45.9%)	p = 1	78 (88.6%)	27 (93.1%)	p = 0.727
KB after the first stent implantation	0 (0%)	0 (0%)	p = 1	88 (100%)	29 (100%)	p = 1
KB after the second stent implantation	83 (97.6%)	36 (97.3%)	p = 1	88 (100%)	29 (100%)	p = 1
Final POT	70 (82.4%)	28 (75.7%)	p = 0.459	83 (94.3%)	28 (96.6%)	p = 1
IVUS/OCT imaging	6 (7.1%)	0 (0%)	p = 0.176	14 (15.9%)	3 (10.3%)	p = 0.558
Rotational Atherectomy	4 (4.7%)	2 (5.4%)	p = 1	3 (3.4%)	2 (6.9%)	p = 0.596
Intravascular lithotripsy	0 (0%)	1 (2.7%)	p = 0.303	2 (2.3%)	0 (0%)	p = 1
GP IIb/IIIa use	8 (9.4%)	3 (8.1%)	p = 1	2 (2.3%)	0 (0%)	p = 1
Radiation dose (mGy)	2212 [1459–3330]	1771 [1108–2752]	p = 0.061	1950.5 [1324.2–3016.5]	1604 [693–2658]	p = 0.027
Contrast media amount (mL)	240 [200–270]	220 [180–270]	p = 0.263	235.6 ± 75.4	205.3 ± 63.5	p = 0.039
ASA	85 (100%)	37 (100%)	p = 1	88 (100%)	29 (100%)	p = 1
Clopidogrel	55 (64.7%)	28 (75.7%)	p = 0.293	50 (56.8%)	18 (62.1%)	p = 0.669
Ticagrelor	29 (34.1%)	9 (24.3%)	p = 0.395	33 (37.5%)	10 (34.5%)	p = 0.827
Prasugrel	0 (0%)	0 (0%)	p = 1	4 (4.5%)	0 (0%)	p = 0.571

**Abbreviations:** LM, Left main; OCT, optical coherence tomography; PCI, percutaneous coronary intervention; ASA, acetylsalicylic acid.

## Clinical Outcome

The one-year study results revealed no statistically significant differences between the two sexes in the two primary study outcomes at the one-year follow-up and the index procedure. However, a noticeable unfavorable trend for female subjects in terms of higher rates of primary outcome was observed in both study groups (Culotte 16.2% vs 11.8%;  $p=0.481$ ; DK-Culotte 10.3% vs 5.7%;  $p=0.401$ ). It is also noteworthy that women in the DK-Culotte arm suffered from restenosis in the target lesion more often than men (10.3% vs 1.1%;  $p=0.018$ ). Clinical outcomes were presented in [Table 3](#). Moreover, [Figure 1](#) (TLF) and [Figure 2](#) (MACE) demonstrate Kaplan–Meier curves, indicating 12-month survival rates for both study cohorts, respectively.

The data presented in [Table 4](#) indicates that there were no statistically significant differences between the female Culotte and DK Culotte arms in terms of TLF and MACE. However, the female Culotte group exhibited higher rates of target lesion failures and major adverse cardiac events (MACE) compared to the DK Culotte group (primary outcome: 16.2% vs 10.3%,  $p = 0.401$ ; and principal secondary outcome: 24.3% vs 13.8%,  $p = 0.114$ ). The 1-year Survival-Free Kaplan–Meier curves for both composite outcomes are presented in [Figure 3](#).

**Table 3** Clinical Outcomes in Both Study Arms

Clinical outcomes	Culotte Group			DK Culotte Group		
	Male N-85	Female N-37	p-value	Male N-88	Female N-29	p-value
I-Year Follow up						
Primary outcome: Target lesion failure (cardiac death, target vessel myocardial infarct, target lesion revascularisation)	10 (11.8%)	6 (16.2%)	$p = 0.481$	5 (5.7%)	3 (10.3%)	$p = 0.401$
Principal secondary outcome: MACE (myocardial infarct, cardiac death, target lesion revascularization)	15 (17.6%)	9 (24.3%)	$p = 0.370$	11 (12.5%)	4 (13.8%)	$p = 0.771$
Target lesion-revascularisation	8 (9.4%)	3 (8.1%)	$p = 0.878$	3 (3.4%)	3 (10.3%)	$p = 0.149$
All - cause mortality	4 (4.7%)	5 (13.5%)	$p = 0.090$	4 (4.5%)	2 (6.9%)	$p = 0.632$
Stent thrombosis	4 (4.7%)	1 (2.7%)	$p = 0.620$	3 (3.4%)	0 (0%)	$p = 0.318$
Stent restenosis	6 (7.1%)	2 (5.4%)	$p = 0.792$	1 (1.1%)	3 (10.3%)	$p = 0.018$


**Abbreviation:** MACE, major adverse cardiac events.

## Discussion

To the best of our knowledge presented study is among the first to examine the effect of sex on the results of percutaneous coronary intervention (PCI) of bifurcation lesions treated with the novel two-stent technique – DK Culotte– in the ACS subset in routine clinical practice.


The study results demonstrated that there was no statistically significant difference between the two genders in terms of the one-year follow-up outcomes of the two investigated two-stent techniques. However, a trend towards a lower incidence of one-year TLF and MACEs was observed in male subjects when the two investigated bifurcation techniques were considered. Furthermore, a similar reduction in composite outcomes was observed in the DK culotte group compared with the classic culotte group in both genders, although this did not reach statistical significance.

There are well-established sex differences with respect to the pathophysiology, clinical presentation, and prognosis of CAD. Although women are less likely to have extensive CAD and more favorable plaque characteristics than men,<sup>20,21</sup> the long-term prognosis and outcomes of revascularization, particularly percutaneous, remain less favorable in females. Several factors have been postulated to be involved in phenomena. Females due to slower progression of CAD tend to be older and have more comorbid conditions at the time of initial diagnosis. Furthermore, the less typical clinical manifestations of the disease result in longer delays in diagnosis and treatment, particularly with regard to revascularization procedures in the acute subset.<sup>22,23</sup> The present study's findings offer partial confirmation of the hypothesis that the



**Figure 1** Target lesion failure I-year Survival-Free Kaplan–Meier curves.

**Abbreviations:** F, Female; M, Male; DK, Double Kiss.



**Figure 2** MACE I-year Survival-Free Kaplan–Meier curves.  
**Abbreviations:** F, Female; M Male.

female population who undergo PCI are generally of more advanced age. A statistically significant disparity in age between males and females was identified in the Culotte cohort, with the female population exhibiting a higher mean age than their male counterparts ( $70.7 \pm 8.7$  vs  $64.6 \pm 8.7$ ;  $p = 0.006$ ). Still, this observation was not confirmed in the DK-Culotte arm, and a reverse trend was observed, noteworthy without statistical significance. We can generally confirm the findings of previous studies regarding a higher burden of comorbidities at the time of hospital admission for primary PCI,<sup>24,25</sup> although women undergoing coronary intervention in our study cohort only had a statistically higher prevalence of diabetes mellitus (DM) type 2, still notable trend was observed for other cardiovascular risk factors.

Despite the underrepresentation of females in studies conducted so far, the available evidence suggests that the clinical outcomes following PCI are less favorable, particularly concerning an increased incidence of periprocedural complications, predominantly related to an elevated risk of bleeding,<sup>5,26,27</sup> yet outcomes are not focused on patients with bifurcation lesions.


Generally, it can be assumed that the technical and clinical complexities associated with bifurcation lesions have led to less favorable outcomes following PCI than those observed in non-bifurcation lesions. Furthermore, women with ACS are less likely to receive evidence-based therapies and undergo cardiac catheterization and complete revascularization.<sup>28,29</sup> Notwithstanding the fact that, in accordance with good clinical practice, all physicians involved in the therapeutic process in our study were encouraged to achieve complete revascularization, this aspect of CAD management was not addressed at the

**Table 4** Clinical Outcomes in the Female Subgroup in Both Study Arms

Clinical outcomes	Culotte Female (n=37)	DK-Culotte Female (n=29)	p-value
I Year Follow up			
Primary outcome: Target lesion failure (cardiac death, target vessel myocardial infarct, target lesion revascularisation)	6 (16.2%)	3 (10.3%)	$p = 0.401$
Principal secondary outcome: MACE (myocardial infarct, cardiac death, target lesion revascularization)	9 (24.3%)	4 (13.8%)	$p = 0.114$
Target lesion-revascularisation	3 (8.1%)	3 (10.3%)	$p = 0.932$
All - cause mortality	5 (13.5%)	2 (6.9%)	$p = 0.343$
Stent thrombosis	1 (2.7%)	0 (0%)	$p = 0.341$
Stent restenosis	2 (5.4%)	3 (10.3%)	$p = 0.578$

**Abbreviation:** MACE, major adverse cardiac events.





**Figure 3** Females cohort 1-year Survival-Free Kaplan–Meier curves of Primary and Principal secondary Outcome.  
**Abbreviations:** DK, Double Kiss.

time of discharge. This appears to be a significant study limitation, given the potential prognostic significance in terms of future unfavorable clinical outcomes.

Despite, data on potential sex differences between different two-stent bifurcation techniques are limited surprisingly the vast majority of available studies suggest no significant differences in clinical outcomes.<sup>30–32</sup> The results of the current study did not demonstrate significant disparities between males and females in the clinical outcomes of the investigated bifurcation stenting technique. However, the increased rate of composite unfavorable outcomes was notable and is partially consistent with the few studies published to date.<sup>33,34</sup> Surprisingly, despite the lower complexity of CAD progression as measured by the Syntax Score in our study cohort, we observed a trend towards a higher rate of stent thrombosis in men compared to women. Although this trend did not reach statistical significance, potential mechanisms underling under this observation remain unclear. In the present study, all subjects were treated with novel second-generation drug-eluting stents (DES). No significant differences were observed in terms of dual antiplatelet therapy (DAPT) agents between sexes still, females tend to more often receive clopidogrel instead of novel antiplatelet agents (ticagrelor/prasugrel). Furthermore, females tended to have a lower average stent diameter implanted in the main and side branches. In the theoretical domain, considering the fact that females tend to have lower compliance in terms of pharmacological treatment after PCI<sup>35,36</sup> this should result in a higher probability of thrombotic events in the female cohort. Surprisingly, the opposite relationship was observed in our study. It is even more confusing when we analyze the basic demographic characteristics between the study cohorts, women in both study arms had higher rates of diabetes, hypertension, and lower Left ventricular ejection fraction (LVEF). Conversely, no substantial disparities were observed between the sexes with regard to peak TnI levels. This marker has been documented as a well-established independent risk factor for future cardiovascular incidents.<sup>37</sup>

Although the above observation did not reach statistical significance, it was observed in both study arms (Culotte and DK-Cullote) and further large scale studies are needed to fully evaluate this issue.

It is interesting to note that, despite the growing body of evidence supporting the mandatory use of proximal optimization technique (POT) and kissing balloons inflation (KBI) in terms of bifurcation lesions, the numbers reported for these two optimization techniques from everyday clinical practice still show utility of approximately 20–40% of all bifurcation PCI performed, with this figure rising to 60–70% in two-stent techniques.<sup>30,38</sup> In this context, data from our real-life cohort (POT: 76%–97%, final KBI: 97%–100%) demonstrates good compliance with the recommendations of the best medical practices, thereby confirming the high value of the preceding data and the relatively low prevalence of intravascular imaging in the study population.

Despite the fact that data focused on the impact of KBI on clinical outcomes might be inconsistent<sup>39,40</sup> Previous reports from bench tests<sup>15,41</sup> suggest a potential significant role for additional KBI during the performance of the classic

Culotte technique. Nevertheless, the data from the study cohort did not confirm a significant impact on one-year outcomes. Nonetheless, a favorable trend has been observed in terms of the clinical outcomes of the ‘Double Kiss-Culotte’ technique in comparison to ‘classical’ Culotte techniques for both sexes. Furthermore, this observation has been already recently postulated.<sup>16,18,42–44</sup>

## Limitations

This study has several limitations. Firstly, data were collected retrospectively with a relatively brief observation period (1-year follow-up). Secondly, the relatively small group size resulted in the underrepresentation of females in both study cohorts. Thirdly, prevalence of IVUS/OCT imaging use is low in the study cohort, and might have an impact on clinical outcomes. Although all operators were encouraged to achieve complete revascularisation, particularly concerning multi-vessel disease, no data were collected on residual syntax score at discharge. Finally, it should be noted that data from angiography had not been subject to external core lab evaluation, particularly in the case of quantitative coronary angiography.

## Conclusion

The present study has revealed no significant differences in clinical outcomes between genders in terms of percutaneous coronary intervention in bifurcation lesions in the ACS cohort after utilization of two different two-stent techniques (Culotte and DK Culotte). Still future randomized, large-scale studies are necessary to fully evaluate potential sex disappearance in clinical outcomes of both stenting techniques.

## Data Sharing Statement

Deidentified data may be released to investigators whose proposed use of the data has been approved by an independent review committee appointed for this purpose. The data-sharing process should be conducted in line with local legal restrictions. The proposal should be submitted to the corresponding author.

## Disclosure

Prof. Dr. Maciej Lesiak reports personal fees from Abbott Vascular, Boston Scientific, Medtronic, and Terumo Medical, outside the submitted work. The authors report no other conflicts of interest in this work.

## References

- Jarrah M, Alrabadi N, Al-Nusair N, et al.. clinical outcomes and cardiovascular-related events in young diabetic patients with acute coronary syndrome. *Vasc Health Risk Manag.* 2022;18:55–60. doi:10.2147/VHRM.S349209
- Jankowski P, Topór-Mądry R, Gąsior M, et al.. Management and predictors of clinical events in 75 686 patients with acute myocardial infarction. *Kardiol Pol.* 2022;80(4):468–475. doi:10.33963/KP.a2022.0058
- Li L, Li G, Chen H, et al.. Role of diabetes mellitus in acute coronary syndrome patients with heart failure and midrange ejection fraction who have undergone percutaneous coronary intervention: a 3-year case-series follow-up retrospective study. *Diabetes Metab Syndr Obes.* 2021;14:4931–4944. doi:10.2147/DMSO.S339209
- Satish P, Avenatti E, Patel J, Agarwala A. Understanding the spectrum of cardiovascular risk in women - A primer for prevention. *Prog Cardiovasc Dis.* 2024;66:1–6. doi:10.1016/j.pcad.2024.05.003
- Alder MR, Adamek KE, Lowenstein A, Raj LM, Lindley KJ, Sutton NR. Acute coronary syndrome in women: an update. *Curr Cardiol Rep.* 2024;26(5):293–301. doi:10.1007/s11886-024-02033-6
- Iwańczyk S, Dit Al Hakim SA, Skrzypińska M, et al.. Sex-based differences in complex percutaneous coronary intervention: insights from the COMPLEX registry. *Hellenic J Cardiol.* 2024;65:1–8. doi:10.1016/j.hjc.2024.04.005
- Yagel O, Shadafny N, Eliaz R, et al.. Long-term prognosis in young patients with acute coronary syndrome treated with percutaneous coronary intervention. *Vasc Health Risk Manag.* 2021;17:153–159. doi:10.2147/VHRM.S298436
- Finch W, Lee MS. Percutaneous coronary intervention for coronary bifurcation lesions. *Rev Cardiovasc Med.* 2017;18(2):59–66. doi:10.3909/ricm0868
- Burzotta F, Louvard Y, Lassen JF, et al. Percutaneous coronary intervention for bifurcation coronary lesions using optimized angiographic guidance: the 18th consensus document from the European bifurcation club. *EuroIntervention.* 2024;19:1–8. doi:10.4244/EIJ-D-24-00160
- Zhang JJ, Ye F, Xu K, et al.. Multicenter, randomized comparison of two-stent and provisional stenting techniques in patients with complex coronary bifurcation lesions: the DEFINITION II trial. *Eur Heart J.* 2020;41(27):2523–2536. doi:10.1093/eurheartj/ehaa543
- Chen SL, Zhang JJ, Han Y, et al. Double kissing crush versus provisional stenting for left main distal bifurcation lesions: DKCRUSH-V randomized trial. *J Am Coll Cardiol.* 2017;70(21):2605–2617. doi:10.1016/j.jacc.2017.09.1066



12. Bujak K, Verardi FM, Arevalos V, et al. Clinical outcomes following different stenting techniques for coronary bifurcation lesions: a systematic review and network meta-analysis of randomized controlled trials. *EuroIntervention*. 2023;19(8):664–675. doi:10.4244/EIJ-D-23-00013
13. Chevalier B, Glatt B, Royer T, Guyon P. Placement of coronary stents in bifurcation lesions by the “culotte” technique. *Am J Cardiol*. 1998;82(8):943–949. doi:10.1016/s0002-9149(98)00510-4
14. Ferenc M, Gick M, Comberg T, et al.. Culotte stenting vs. TAP stenting for treatment of de novo coronary bifurcation lesions with the need for side-branch stenting: the Bifurcations Bad Krozingen (BBK) II angiographic trial. *Eur Heart J*. 2016;37(45):3399–3405. doi:10.1093/eurheartj/ehw345
15. Toth GG, Sasi V, Franco D, et al.. Double-kissing culotte technique for coronary bifurcation. *EuroIntervention*. 2020;16:e724–e733. doi:10.4244/EIJ-D-20-00130
16. Tu S, Zhang L, Tian Q, Hu F, Wang Y, Chen L. Five-year outcomes of double kissing mini-culotte stenting vs mini-culotte stenting using drug-eluting stents for the treatment of true coronary bifurcation lesions. *Front Cardiovasc Med*. 2024;11:1336750. doi:10.3389/fcvm.2024.1336750
17. Barycki M, Włodarczak A, Rola P, Włodarczak S, Jatrzebski A, Lesiak M. Left main calcified lesion treated with orbital atherectomy and two-stent double-kissing Culotte technique in a very elderly patient with frailty syndrome. *Postępy Kardiologii Interwencyjnej*. 2023;19(4):380–381. doi:10.5114/aic.2023.133253
18. Barycki M, Włodarczak A, Włodarczak S, et al.. Comparative short-term outcomes of double-kissing culotte and culotte techniques in acute coronary syndrome from the lower silesia culotte bifurcation registry. *J Clin Med*. 2024;13(23):7392. doi:10.3390/jcm13237392
19. Garcia-Garcia HM, McFadden EP, Farb A, et al. Standardized end point definitions for coronary intervention trials: the academic research consortium-2 consensus document. *Eur Heart J*. 2018;39(23):2192–2207. doi:10.1093/eurheartj/ehy223
20. Ten Haaf ME, Rijndertse M, Cheng JM, et al. Sex differences in plaque characteristics by intravascular imaging in patients with coronary artery disease. *EuroIntervention*. 2017;13:320–328. doi:10.4244/EIJ-D-16-00361
21. Song BY, Chen C, Xu WH, et al. Gender differences in the correlations between immune cells and organ damage indexes of acute myocardial infarction patients. *Vasc Health Risk Manag*. 2022;18:839–850. doi:10.2147/VHRM.S374157
22. Park S, Ahn JM, Park H, et al.. Comparison of long-term outcomes following coronary revascularization in men vs women with unprotected left main disease. *Am J Cardiol*. 2021;153:9–19. doi:10.1016/j.amjcard.2021.05
23. Sakkers TR, Mokry M, Civelek M, et al.. Sex differences in the genetic and molecular mechanisms of coronary artery disease. *Atherosclerosis*. 2023;384:117279. doi:10.1016/j.atherosclerosis.2023.117279
24. Birkemeyer R, Schneider H, Rillig A, et al.. Do gender differences in primary PCI mortality represent a different adherence to guideline recommended therapy? A multicenter observation. *BMC Cardiovasc Disord*. 2014;14:71. doi:10.1186/1471-2261-14-71
25. Fu WX, Zhou TN, Wang XZ, et al.. Sex-related differences in short- and long-term outcomes among young and middle-aged patients with ST-segment elevation myocardial infarction who underwent percutaneous coronary intervention. *Chin Med J*. 2018;131(12):1420–1429. doi:10.4103/0366-6999.233965
26. Pham P, Pham PT, Pham P, et al.. Implications of bleeding in acute coronary syndrome and percutaneous coronary intervention. *Vasc Health Risk Manag*. 2011;7:551–567. doi:10.2147/VHRM.S23862
27. Sabatowski K, Malinowski KP, Siudak Z, et al.. Sex-related differences and rotational atherectomy: analysis of 5177 percutaneous coronary interventions based on a large national registry from 2014 to 2020. *Kardiologia Pol*. 2021;79(12):1320–1327. doi:10.33963/KP.a2021.0131
28. Akhter N, Milford-Beland S, Roe MT, et al.. Gender differences among patients with acute coronary syndromes undergoing percutaneous coronary intervention in the American College of Cardiology-National Cardiovascular Data Registry (ACC-NCDR). *Am Heart J*. 2009;157(1):141–148. doi:10.1016/j.ahj.2008.08.012
29. Epps K, Goel R, Mehran R, et al.. Influence of race/ethnicity and sex on coronary stent outcomes in diabetic patients. *J Soc Cardiovasc Angiogr Interv*. 2023;2(5):101053. doi:10.1016/j.jscai.2023.101053
30. Doolub G, Iannaccone M, Rab T, et al.. Sex-based treatment and outcomes for coronary bifurcation stenting: a report from the e-ULTIMASTER registry. *Catheter Cardiovasc Interv*. 2023;102(3):430–439. doi:10.1002/ccd.30770
31. Ahn HJ, Bruno F, Kang J, et al.. Sex differences in procedural characteristics and clinical outcomes among patients undergoing bifurcation PCI. *Korean Circ J*. 2024. doi:10.4070/kcj.2024.0172
32. Hosseinpour A, Attar A, Agrawal A, et al.. Sex-specific differences in clinical outcomes in patients with coronary bifurcation lesions. *Cardiol Rev*. 2023. doi:10.1097/CRD.0000000000000629
33. Osman M, Ghaffar YA, Osman K, et al.. Gender-based outcomes of coronary bifurcation stenting: a report from the national readmission database. *Catheter Cardiovasc Interv*. 2022;99(2):433–439. doi:10.1002/ccd.29704
34. Schamroth Pravda N, Perl L, Greenberg G, et al. Impact of sex on outcomes of bifurcation lesion percutaneous coronary intervention: results from a single-center prospective registry. *Coron Artery Dis*. 2022;31(1):31–36. doi:10.1097/MCA.0000000000001039
35. Ifthikhar MF, Rana MOR, Naeem A, et al.. Sex disparity of DAPT noncompliance in patients with left main stem PCI with DES. *Medicine*. 2024;103(26):e38724. doi:10.1097/MD.00000000000038724
36. Malik AH, Yandrapalli S, Shetty SS, Aronow WS, Cooper HA, Panza JA. Meta-analysis of dual antiplatelet therapy versus monotherapy with P2Y12 inhibitors in patients after percutaneous coronary intervention. *Am J Cardiol*. 2020;127:25–29. doi:10.1016/j.amjcard.2020.04.027
37. Jia X, Sun W, Hoogeveen RC, et al.. High-sensitivity troponin i and incident coronary events, stroke, heart failure hospitalization, and mortality in the ARIC study. *Circulation*. 2019;139(23):2642–2653. doi:10.1161/CIRCULATIONAHA.118.038772
38. Lee JM, Lee SH, Kim J, et al.. Ten-year trends in coronary bifurcation percutaneous coronary intervention: prognostic effects of patient and lesion characteristics, devices, and techniques. *J Am Heart Assoc*. 2021;10(18):e021632. doi:10.1161/JAHA.121.021632
39. Kini AS, Dangas GD, Baber U, et al.. Influence of final kissing balloon inflation on long-term outcomes after PCI of distal left main bifurcation lesions in the EXCEL trial. *EuroIntervention*. 2020;16(3):218–224. doi:10.4244/EIJ-D-19-00851
40. Liu G, Ke X, Huang ZB, et al.. Final kissing balloon inflation for coronary bifurcation lesions treated with single-stent technique: a meta-analysis. *Herz*. 2019;44(4):354–362. doi:10.1007/s00059-017-4647-1
41. Hu F, Tu S, Cai W, et al.. Double kissing mini-culotte versus mini-culotte stenting: insights from micro-computed tomographic imaging of bench testing. *EuroIntervention*. 2019;15(5):465–472. doi:10.4244/EIJ-D-18-00688
42. Barycki M, Rola P, Włodarczak A, et al.. Evaluation of small vessel bifurcation stenting using the double-kissing culotte and culotte technique in acute coronary syndrome: 12-month clinical outcomes. *Clin Cardiol*. 2024;47(11):e70043. doi:10.1002/clc.70043

43. Fan L, Chen L, Luo Y, et al.. DK mini-culotte stenting in the treatment of true coronary bifurcation lesions: a propensity score matching comparison with T-provisional stenting. *Heart Vessels*. 2016;31(3):308–321. doi:10.1007/s00380-014-0611-7
44. Barycki M, Rola P, Włodarczak A, et al.. One-year comparative outcomes of DK culotte and culotte techniques in left main bifurcation in acute coronary syndrome: sub-study of lower silesia culotte bifurcation registry. *Kardiol Pol*. 2024;82(10):999–1002. doi:10.33963/v.phj.102075

## Vascular Health and Risk Management

### Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/vascular-health-and-risk-management-journal>

**Dovepress**  
Taylor & Francis Group