

Modifying tetramethyl–nitrophenyl–imidazoline with amino acids: design, synthesis, and 3D-QSAR for improving inflammatory pain therapy

Xueyun Jiang¹
Yuji Wang¹
Haimei Zhu¹
Yaonan Wang¹
Ming Zhao^{1,2}
Shurui Zhao¹
Jianhui Wu¹
Shan Li¹
Shiqi Peng¹

¹Beijing Area Major Laboratory of Peptide and Small Molecular Drugs, Engineering Research Center of Endogenous Prophylactic of Ministry of Education of China, Beijing Laboratory of Biomedical Materials, College of Pharmaceutical Sciences, Capital Medical University, Beijing, People's Republic of China; ²Faculty of Biomedical Science and Environmental Biology, Kaohsiung Medical University, Kaohsiung, Taiwan

Correspondence: Shiqi Peng/Ming Zhao
College of Pharmaceutical Sciences,
Capital University of Medical Sciences,
No.10 Xitoutiao, You An Men, Beijing
100069, People's Republic of China
Tel +86 10 8391 1528
Fax: +86 10 8280 2482
Email shiqipeng@163.com/ sqpeng@
bjmu.edu.cn; mingzhao@bjmu.edu.cn

Abstract: With the help of pharmacophore analysis and docking investigation, 15 novel 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)-oxyacetyl-L-amino acids (6a–o) were designed, synthesized, and assayed. On tail-flick and xylene-induced ear edema models, 10 $\mu\text{mol/kg}$ 6a–o exhibited excellent oral anti-inflammation and analgesic activity. The dose-dependent assay of their representative 6f indicates that the effective dose should be 3.3 $\mu\text{mol/kg}$. The correlation of the three-dimensional quantitative structure–activity relationship with the docking analysis provides a basis for the rational design of drugs to treat inflammatory pain.

Keywords: tetramethylimidazoline, analgesic, anti-inflammatory, 3D-QSAR

Introduction

A variety of etiologies such as inflammation and neural destruction can complicate chronic pain. Though neuropathic pain is simply accompanied by hypersensitivity to mechanical or thermal stimulus,^{1–4} inflammatory pain is accompanied by various painful responses to injury of peripheral tissue, trauma, infection, surgery, burns, or related diseases.^{5–8} Treating inflammatory pain represents a medical need, and a lot of agents such as pyridazin derivatives,⁹ pyrazine N-acylhydrazone derivatives,¹⁰ oxadiazole derivatives,¹¹ pyrazolone–pyridazine conjugates,¹² and triazine derivatives¹³ were reported to possess anti-inflammation and analgesic activities. Besides, 1-hydroxyl- and 1-oxyl-tetramethylimidazolines were found to be analgesic pharmacophores, of which 1-oxyl-2-(3'-nitrophen-1'-yl)-4,4,5,5-tetramethylimidazoline and 1-hydroxyl-2-(3'-nitrophen-1'-yl)-4,4,5,5-tetramethylimidazoline had the highest efficacy; ie, if the 2-position of 4,4,5,5-tetramethylimidazoline was 3-nitro-phenyl, the pharmacophore will possess the highest analgesic activity.^{14,15} The merge of the characteristics of the structures in the blue box and 2-(3'-nitrophen-1'-yl)-4,4,5,5-tetramethylimidazoline had the highest analgesic activity this paper designed 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)-oxyacetyl-L-amino acids (TMNPIZA) as the therapeutic agents for inflammatory pain, docked them into the active site of cyclooxygenase-2 (COX-2) enzyme, found them properly fitting the active site, and observed the important interactions of TMNPIZA with Ser530, Phe518, Arg513, and Ser353 in particular (Figure 1).

In this context, here we present the preparation and evaluation of the oral analgesic and anti-inflammatory activities of 15 novel TMNPIZA on tail-flick and xylene-induced ear edema mouse models, respectively, as well as analyze their structure–activity relationship with Discovery Studio 4.0 three-dimensional quantitative structure–activity relationship (3D-QSAR) module.




Figure 1 Design diagram: **(A)** the merge of the known structures of clinical anti-inflammatory pain agents in the blue box and tetramethylimidazolines been analgesic pharmacophore resulting in novel structure of TMNPZA; **(B)** TMNPZA fitting the active site of COX-2 enzyme and having a desirable docking feature; **(C)** estimating TMNPZA being able to treat inflammatory pain, the feature of TMNPZA in the active site of COX-2 is highlighted with a yellow box and interactions of TMNPZA with the amino acid residues of the active site is clarified by amplifying the yellow box.

Abbreviations: AA, amino acid; TMNPZA, 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)-oxyacetyl-L-amino acids; COX-2, cyclooxygenase-2.

Materials and methods

Chemical synthesis

General method

All the reactions were carried out under nitrogen (1 bar). ^1H (300 and 500 MHz) and ^{13}C (75 and 125 MHz) nuclear magnetic resonance (NMR) spectra were recorded on Bruker Avance II-300 and Avance II-500 spectrometers for solution $\text{DMSO}-d_6$ or CDCl_3 with tetramethylsilane as internal standard. Infrared (IR) spectra were recorded with a Perkin-Elmer 983 instrument. Electrospray ionization mass spectrometry (ESI/MS) was determined on Waters Micromass Quattro Micro ZQ2000. Melting points were measured on an XT5 hot stage microscope (Beijing key electro-optic factory). All L-amino acids were purchased from China Biochemical Corp. Thin-layer chromatography (TLC) was made with Qingdao silica gel GF₂₅₄. Chromatography was performed with Qingdao silica gel H₆₀ or Sephadex-LH₂₀. The purities of the intermediates were measured on TLC (Merck silica gel plates of type 60 F₂₅₄, 0.25 mm layer thickness, three systems of mixed solvents) and showed a single component, while the purities of the products were measured on high-performance liquid chromatography (Waters, C₁₈ column 4.6×150 mm, three elution systems of mixed solvents) and more than 96%. All solvents were distilled and dried before use according to literature procedures. Optical rotations were determined with a Jasco P-1020 Polarimeter at 20°C. The statistical analysis of all the biological data was carried out by use of analysis of variance (ANOVA) test. $P < 0.05$ is considered significant.

Synthesis

2,3-dimethyl-2,3-dinitrobutane

At -5°C , to a solution of 34.5 g (0.39 mol) of 2-nitropropane in 65 mL of aqueous NaOH (6.0 M) 10 mL (0.19 mol) of Br_2 was added dropwise within 1 hour. With stirring, 128 mL of ethanol was added, and this solution was then stirred at

84°C for 3 hours. The hot reaction mixture was transferred into 400 mL of ice water. The formed colorless crystals were collected by filtration to yield 25 g (73%) of the title compound. Mp 110°C – 112°C .

2,3-dihydroxylamino-2,3-dimethylbutane

To a solution of 17.6 g of 2,3-dimethyl-2,3-dinitrobutane (0.1 mol) in 300 mL of tetrahydrofuran (THF) and 50 mL of water of 9°C 27 g of Zn powder was added to get a slurry-like mixture. To this stirring mixture a solution of 43 g of NH_4Cl (0.8 mol) in 150 mL of H_2O was added dropwise within 2 hours and was then stirred at 10°C for 1 hour and stored at 4°C for 16 hours. The slurry-like mixture was filtered, the collected precipitates were washed with THF (100 mL ×4), and diethyl ether (50 mL ×3) was added to provide 59 g of precipitates. The solution was evaporated under vacuum to remove THF. Then, the solution was protected from air, to which 50 g of sodium carbonate and 30 g of sodium chloride were added with cooling. The solids were continuously extracted with 400 mL of chloroform over 18 hours to provide 9.4 g (63%) of the title compound as a colorless powder. Mp 182°C .

1,3-dihydroxyl-2-(3-nitrophenyl)-4,4,5,5-tetramethylimidazolidine (1)

A solution of 296 mg (2 mmol) of 2,3-bis(hydroxylamino)-2,3-dimethylbutane and 302 mg (2 mmol) of 3-nitrobenzaldehyde in 3 mL of methanol was stirred at room temperature for 16 hours. The title compound was collected by filtration to provide 421 mg (75%) of the title compound. Mp 179°C – 81°C . ESI-MS (m/z) 281 $[\text{M}+\text{H}]^+$. IR (KBr) 3,315, 1,530, 1,360, 1,600, 875, 790, 685 cm^{-1} . ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ /ppm=8.03 (s, 2H), 7.77 (s, 1H), 7.57 (d, $J=7.5$ Hz, 1H), 7.34 (d, $J=6.0$ Hz, 1H), 7.04 (dd, $J=8.7$ Hz, $J=8.1$ Hz, 1H), 4.19 (s, 1H), 0.52 (s, 12H).

1-hydroxyl-2-(3-nitrophen-1-yl)-4,4,5,5-tetramethylimidazoline (2)

A suspension of 281 mg (1 mmol) of (1) and 360 mg of anhydrous MgSO_4 in 40 mL of anhydrous THF was stirred at 80°C for 24 hours, and TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}$, 20:1) indicated the complete disappearance of (1). The reaction mixture was cooled to room temperature. After removal of MgSO_4 by filtration, the filtrate was evaporated to dryness. The residue was purified on chromatography column of silica gel ($\text{CHCl}_3/\text{CH}_3\text{OH}$, 30:1) to give 174 mg (66%) of the title compound as yellow powder. Mp 97°C – 99°C . EI-MS (m/z) 263 $[\text{M}]^+$. IR (KBr) 3,345, 1,602, 1,513, 1,455, 1,370, 811, 790, 680 cm^{-1} . ^1H NMR (300 MHz, CDCl_3 -d) δ/ppm =8.55 (s, 1H), 8.22 (d, J =7.20 Hz, 1H), 7.99 (d, J =7.21 Hz, 1H), 7.55 (t, J =7.21 Hz, 1H), 2.65 (s, 1H), 1.15 (s, 6H), 1.13 (s, 6H).

Ethyl 2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetate (3)

At 0°C , to a solution of 500 mg (1.77 mmol) of (2) in 40 mL of THF and 10 mL of 80 mg 2.12 mmol of NaH was added and stirred for 30 minutes. To this suspension, 0.3 mL (2.12 mmol) of $\text{BrCH}_2\text{COOC}_2\text{H}_5$ was added and stirred at room temperature for 5 hours, and TLC ($\text{CHCl}_3/\text{CH}_3\text{OH}$ =30:1) indicated the complete disappearance of (2). The reaction mixture was filtered, the filtrate was evaporated under vacuum to remove THF, and the residue was dissolved in 200 mL of ethyl acetate. This solution was washed with saturated aqueous NaHCO_3 (50 mL \times 3) and saturated aqueous NaCl (50 mL \times 2). The ethyl acetate phase was dried with anhydrous Na_2SO_4 over light and filtered, the filtrate was evaporated under vacuum, and the residue was purified with column chromatography (EtOAC:petroleum=1:3) to provide 529 mg (82%) of the title compound as yellow syrup. ESI/MS (m/z) 350 $[\text{M}+\text{H}]^+$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.41 (s, 1H), 8.34 (m, 1H), 8.09 (d, J =7.8 Hz, 1H), 7.74 (t, J =7.8 Hz, 1H), 4.70 (s, 2H), 4.11 (q, J =7.2 Hz, 2H), 1.22 (s, 6H), 1.17 (t, J =7.2 Hz, 3H), 1.13 (s, 6H). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$): δ =168.66, 162.45, 147.85, 134.79, 132.80, 130.43, 125.21, 123.15, 74.08, 73.02, 68.40, 61.00, 23.88, 19.74, 14.41. Anal. Calcd for $\text{C}_{17}\text{H}_{23}\text{N}_3\text{O}_5$: C, 58.44; H, 6.64; N, 12.03; O, 22.90.

2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetic acid (4)

At 0°C , to a solution of 2.0 g (5.7 mmol) of (3) in 20 mL of methanol 4 mL of aqueous NaOH (4 M) was added. The reaction mixture was stirred at 0°C for 2 hours, and TLC (EtOAC: petroleum=1:3) indicated the complete disappearance of (3).

The reaction mixture was adjusted to pH 2 and evaporated to remove methanol. The residue was extracted with ethyl acetate (50 mL \times 3), the ethyl acetate phase was dried with anhydrous Na_2SO_4 over light and filtered, and the filtrate was evaporated under vacuum to provide 1.6 g (87%) of the title compound as yellow powder. Mp 66°C – 69°C ; ESI/MS (m/z) 320 $[\text{M}-\text{H}]$. $[\alpha]_{20}^{\text{D}}=-7.5$ (c =1.2, CH_3OH). ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.42 (s, 1H), 8.32 (m, 1H), 8.10 (d, J =7.8 Hz, 1H), 7.74 (t, J =7.8 Hz, 1H), 4.71 (s, 2H), 1.20 (s, 6H), 1.12 (s, 6H). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$): δ =169.14, 163.10, 148.52, 134.95, 132.63, 131.25, 124.87, 122.87, 76.12, 66.40, 58.75, 23.60, 20.20, 14.97. Anal. Calcd for $\text{C}_{15}\text{H}_{19}\text{N}_3\text{O}_5$: C, 56.07; H, 5.96; N, 13.08. Found C, 56.30; H, 6.15; N, 13.30.

General procedure of preparing 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetyl-L-amino acid benzylesters (5a–o)

At 0°C , 642 mg (2.0 mmol) of (4), 270 mg (2.0 mmol) of HOBt, and 500 mg (2.4 mmol) of Dicyclohexylcarbodiimide (DCC) was successively dissolved in 50 mL of anhydrous THF, and this solution was stirred at 0°C for 30 minutes. To this solution 2 mmol of L-amino acid benzylester was added and was adjusted to pH 8. The reaction mixture was stirred at room temperature for 6 hours, and TLC indicated the complete disappearance of (4). The reaction mixture was filtered and the filtrate was evaporated under vacuum to remove THF. The residue was dissolved in 200 mL of ethyl acetate. This solution was washed with saturated aqueous NaHCO_3 (50 mL \times 3) and saturated aqueous NaCl (50 mL \times 2). The ethyl acetate phase was dried with anhydrous Na_2SO_4 over light and filtered, the filtrate was evaporated under vacuum, and the residue was purified with column chromatography (EtOAC:petroleum=1:1) to provide 5a–o.

Benzyl (2S,3S)-3-methyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoate (5a)

Yield: 925 mg (88%). ESI/MS (m/z) 525 $[\text{M}+\text{H}]^+$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.43 (t, J =3.0 Hz, 1H), 8.33 (m, 1H), 8.18 (d, J =8.1 Hz, 1H), 8.12 (d, J =8.1 Hz, 1H), 7.71 (t, J =8.1 Hz, 1H), 7.35 (m, 5H), 5.12 (m, 2H), 4.53 (s, 2H), 4.28 (dd, J =8.1 Hz, J =6.3 Hz, 1H), 1.78 (m, 1H), 1.32 (m, 2H), 1.20 (s, 6H), 1.12 (s, 6H), 0.991 (m, 6H). ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$): δ/ppm =171.49, 167.78, 162.49, 147.97, 136.21, 134.75, 132.87, 130.42, 128.87, 128.60, 128.55, 125.20, 123.12, 75.42, 72.91, 68.34, 66.47, 56.49, 36.79, 25.10, 23.90, 15.78, 11.48. Anal. Calcd for

$C_{28}H_{36}N_4O_6$; C, 64.10; H, 6.92; N, 10.68. Found: C, 64.29; H, 6.78; N, 10.90.

Benzyl (2S)-3-methyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoate (5b)

Yield: 650 mg (71%). ESI/MS (m/z) 511 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.47 (m, 1H), 8.35 (dd, *J*=8.4 Hz, *J*=2.1 Hz, 1H), 8.21 (d, *J*=8.1 Hz, 1H), 8.15 (d, *J*=7.8 Hz, 1H), 7.74 (t, *J*=7.8 Hz, 1H), 7.36 (m, 5H), 5.13 (s, 2H), 4.58 (s, 2H), 4.23 (dd, *J*=8.1 Hz, *J*=6.3 Hz, 1H), 2.06 (m, 1H), 1.23 (s, 6H), 1.11 (s, 6H), 0.83 (d, *J*=6.6 Hz, 3H), 0.82 (d, *J*=6.6 Hz, 3H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=171.48, 167.72, 162.58, 147.99, 136.23, 134.91, 130.50, 128.88, 128.60, 128.54, 123.34, 75.57, 73.24, 68.05, 66.49, 57.56, 30.29, 23.72, 19.78, 19.30, 18.48. Anal. Calcd for $C_{27}H_{34}N_4O_6$; C, 63.51; H, 6.71; N, 10.97. Found: C, 63.30; H, 6.55; N, 10.76.

Benzyl (2S)-4-methyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoate (5c)

Yield: 500 mg (65%). ESI/MS (m/z) 525 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.42 (s, 1H), 8.32 (d, *J*=8.4 Hz, 1H), 8.25 (d, *J*=7.5 Hz, 1H), 8.11 (d, *J*=7.5 Hz, 1H), 7.71 (t, *J*=7.8 Hz, 1H), 7.34 (m, 5H), 5.10 (s, 2H), 4.49 (s, 2H), 4.35 (m, 1H), 1.56 (m, 3H), 1.20 (s, 6H), 1.12 (s, 6H), 0.85 (d, *J*=3.9 Hz, 3H), 0.80 (d, *J*=4.2 Hz, 3H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=172.38, 167.67, 162.44, 147.99, 136.33, 134.74, 132.89, 130.41, 128.87, 128.51, 128.25, 125.17, 123.11, 75.54, 72.92, 68.35, 66.45, 50.52, 24.64, 23.88, 23.12, 21.65, 19.83. Anal. Calcd for $C_{28}H_{36}N_4O_6$; C, 64.10; H, 6.92; N, 10.68. Found: C, 63.89; H, 6.77; N, 10.47.

Benzyl (2S)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)propanoate (5d)

Yield: 425 mg (44%). ESI/MS (m/z) 483 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.42 (t, *J*=1.8 Hz, 1H), 8.32 (d, *J*=6.6 Hz, 2H), 8.11 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=8.1 Hz, 1H), 7.34 (m, 5H), 5.11 (s, 2H), 4.50 (s, 2H), 4.33 (t, *J*=7.2 Hz, 1H), 1.28 (d, *J*=7.2 Hz, 3H), 1.20 (s, 6H), 1.12 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=172.47, 167.35, 162.42, 147.98, 136.37, 134.73, 132.80, 130.46, 128.87, 128.48, 128.17, 125.22, 123.10, 75.64, 72.92, 68.33, 66.43, 47.83, 23.90, 17.31. Anal. Calcd for $C_{25}H_{30}N_4O_6$; C, 62.23; H, 6.27; N, 11.61. Found: C, 62.04; H, 6.11; N, 11.83.

Benzyl (2S)-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)acetate (5e)

Yield: 810 mg (87%). ESI/MS (m/z) 469 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.41 (m, 1H), 8.33 (m, 1H), 8.23 (t, *J*=7.5 Hz, 1H), 8.10 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=7.8 Hz, 1H), 7.36 (m, 5H), 5.11 (s, 2H), 4.53 (s, 2H), 3.89 (d, *J*=6.0 Hz, 2H), 1.22 (s, 6H), 1.13 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=169.82, 167.88, 162.37, 148.05, 136.29, 134.64, 132.78, 130.49, 128.86, 128.53, 128.39, 125.22, 123.09, 75.06, 72.95, 68.35, 66.38, 23.88, 19.87. Anal. Calcd for $C_{24}H_{28}N_4O_6$; C, 61.53; H, 6.02; N, 11.96. Found: C, 61.74; H, 6.18; N, 12.17.

Benzyl (2S)-3-(4-hydroxyphenyl)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)propanoate (5f)

Yield: 952 mg (83%). ESI/MS (m/z) 573 [M-H]⁻. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=9.22 (s, 1H), 8.42 (t, *J*=1.9 Hz, 1H), 8.32 (dd, *J*=8.4 Hz, *J*=2.1 Hz, 1H), 8.14 (d, *J*=8.1 Hz, 1H), 8.07 (d, *J*=7.5 Hz, 1H), 7.69 (t, *J*=8.1 Hz, 1H), 7.34 (m, 3H), 7.29 (m, 2H), 6.93 (d, *J*=8.4 Hz, 2H), 6.62 (d, *J*=6.5 Hz, 2H), 5.08 (m, 2H), 4.52 (m, 1H), 4.45 (s, 2H), 2.93 (dd, *J*=14.0 Hz, *J*=6.0 Hz, 1H), 2.84 (dd, *J*=13.5 Hz, *J*=8.4 Hz, 1H), 1.15 (s, 6H), 1.12 (m, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=171.51, 167.43, 162.40, 156.59, 148.02, 136.15, 134.63, 132.80, 130.45, 128.83, 128.51, 128.34, 127.08, 125.19, 123.09, 115.55, 75.45, 72.92, 68.32, 66.55, 53.94, 36.26, 23.87, 14.55. Anal. Calcd for $C_{31}H_{34}N_4O_7$; C, 64.80; H, 5.96; N, 9.75. Found: C, 64.61; H, 5.80; N, 9.52.

Benzyl (2S)-dibenzyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanedioate (5g)

Yield: 600 mg (69%). ESI/MS (m/z) 631 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.42 (s, 1H), 8.30 (m, 2H), 8.09 (d, *J*=7.2 Hz, 1H), 7.68 (t, *J*=7.8 Hz, 1H), 7.34 (m, 10H), 5.11 (s, 2H), 5.07 (s, 2H), 4.49 (s, 2H), 4.39 (m, 1H), 2.39 (t, *J*=7.5 Hz, 2H), 2.07 (m, 1H), 1.91 (m, 1H), 1.19 (s, 6H), 1.12 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=172.37, 171.53, 167.82, 162.39, 148.00, 136.54, 136.26, 134.69, 132.85, 130.39, 128.66, 128.52, 128.47, 128.37, 128.26, 125.16, 123.12, 75.58, 72.92, 68.33, 66.60, 66.01, 51.40, 30.20, 26.37, 23.87, 19.80. Anal. Calcd for $C_{34}H_{38}N_4O_8$; C, 64.75; H, 6.07; N, 8.88. Found: C, 64.54; H, 5.91; N, 8.67.

Benzyl (2S)-6-(tert-butoxycarbonyl)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)hexanoate (5h)

Yield: 1,100 mg (86%). ESI/MS (m/z) 640 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.43 (t, *J*=4.8 Hz, 1H), 8.32 (dd, *J*=7.5 Hz, *J*=1.5 Hz, 1H), 8.23 (d, *J*=7.5 Hz, 1H), 8.10 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=8.1 Hz, 1H), 7.36 (m, 5H), 6.73 (t, *J*=5.1 Hz, 1H), 5.11 (s, 2H), 5.00 (s, 2H), 4.28 (m, 1H), 2.85 (dd, *J*=12.6 Hz, *J*=6.3 Hz, 2H), 1.78 (m, 2H), 1.68 (m, 2H), 1.36 (s, 9H), 1.31 (m, 2H), 1.21 (s, 6H), 1.23 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=172.02, 167.70, 162.43, 156.01, 148.01, 136.33, 134.71, 132.87, 130.42, 128.86, 128.50, 128.25, 125.18, 123.12, 77.79, 75.61, 72.93, 68.34, 66.44, 52.19, 30.90, 29.48, 28.71, 23.03, 21.21. Anal. Calcd for C₃₃H₄₅N₅O₈: C, 61.96; H, 7.09; N, 10.95. Found: C, 61.76; H, 7.02; N, 10.78.

Benzyl (2S)-4-amino-4-oxo-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoate (5i)

Yield: 852 mg (74%). ESI/MS (m/z) 526 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.41 (m, 1H), 8.31 (m, 1H), 8.18 (d, *J*=8.1 Hz, 1H), 8.08 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=8.1 Hz, 1H), 7.42 (s, 1H), 7.33 (m, 5H), 6.95 (s, 1H), 5.09 (s, 2H), 4.71 (dd, *J*=13.72 Hz, *J*=5.4 Hz, 1H), 4.51 (s, 2H), 2.59 (t, *J*=4.5 Hz, 2H), 1.19 (s, 6H), 1.12 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=171.54, 171.23, 167.32, 162.44, 148.04, 136.36, 134.58, 132.75, 130.52, 128.79, 128.36, 128.00, 125.23, 123.03, 75.72, 72.94, 68.34, 66.52, 48.80, 36.65, 25.76, 24.92, 23.88, 15.62. Anal. Calcd for C₂₆H₃₁N₅O₇: C, 59.42; H, 5.95; N, 13.33. Found: C, 59.20; H, 5.70; N, 13.11.

Benzyl (2S)-5-amino-5-oxo-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoate (5j)

Yield: 832 mg (85%). ESI/MS (m/z) 540 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.42 (m, 1H), 8.37 (m, 2H), 8.01 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=8.1 Hz, 1H), 7.35 (m, 5H), 7.24 (s, 1H), 6.77 (s, 1H), 5.11 (s, 2H), 4.49 (s, 2H), 4.32 (m, 1H), 2.09 (m, 2H), 1.96 (m, 1H), 1.81 (m, 1H), 1.21 (s, 6H), 1.09 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=173.65, 171.83, 167.74, 162.41, 148.02, 136.33, 134.69, 132.85, 130.43, 128.86, 128.49, 128.24, 125.18, 123.13, 75.62, 72.94, 68.33, 66.48, 52.00, 31.44, 26.89, 23.91, 19.84. Anal. Calcd for C₂₇H₃₃N₅O₇: C, 60.10; H, 6.16; N, 12.98. Found: C, 60.29; H, 6.00; N, 12.75.

Benzyl (2S)-3-hydroxy-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoate (5k)

Yield: 700 mg (70%). ESI/MS (m/z) 513 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.43 (m, 1H), 8.32 (dd, *J*=8.1 Hz, *J*=1.5 Hz, 1H), 8.12 (d, *J*=7.8 Hz, 1H), 7.82 (d, *J*=7.6 Hz, 1H), 7.72 (t, *J*=7.8 Hz, 1H), 7.36 (m, 5H), 5.13 (m, 2H), 4.62 (s, 2H), 4.37 (dd, *J*=8.1 Hz, *J*=3.9 Hz, 1H), 4.17 (m, 1H), 1.21 (s, 6H), 1.13 (s, 6H), 1.02 (d, *J*=6.3 Hz, 3H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=170.62, 168.07, 162.51, 148.03, 136.33, 134.63, 132.77, 130.52, 128.83, 128.44, 128.16, 125.24, 123.06, 75.55, 72.95, 68.37, 66.55, 57.82, 23.91, 20.64. Anal. Calcd for C₂₆H₃₂N₄O₇: C, 60.93; H, 6.29; N, 10.93. Found: C, 60.70; H, 6.12; N, 10.71.

Benzyl (2S)-3-(tert-butylthio)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)propanoate (5l)

Yield: 470 mg (42%). ESI/MS (m/z) 569 [M-H]⁻. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.42 (t, *J*=1.8 Hz, 1H), 8.32 (dd, *J*=8.1 Hz, *J*=1.5 Hz, 1H), 8.24 (d, *J*=8.1 Hz, 1H), 8.11 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=7.8 Hz, 1H), 7.36 (m, 5H), 5.12 (s, 2H), 4.53 (m, 1H), 4.52 (s, 2H), 2.87 (m, 1H), 2.76 (m, 1H), 1.26 (s, 6H), 1.22 (s, 9H), 1.13 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=170.44, 167.53, 162.44, 148.03, 136.13, 134.67, 132.84, 130.49, 128.84, 128.54, 128.31, 128.27, 125.21, 123.06, 75.49, 72.96, 68.38, 66.80, 52.77, 42.73, 31.14, 29.68, 23.90, 19.86. Anal. Calcd for C₂₉H₃₈N₄O₆S: C, 61.03; H, 6.71; N, 9.82. Found: C, 61.24; H, 6.88; N, 9.97.

Benzyl (2S)-4-(methylthio)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoate (5m)

Yield: 842 mg (78%). ESI/MS (m/z) 543 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.43 (m, 1H), 8.32 (d, *J*=7.2 Hz, 2H), 8.11 (d, *J*=8.1 Hz, 1H), 7.71 (t, *J*=8.1 Hz, 1H), 7.35 (m, 5H), 5.12 (s, 2H), 4.55 (s, 2H), 4.47 (m, 1H), 2.42 (m, 2H), 1.99 (s, 3H), 1.93 (m, 2H), 1.21 (s, 6H), 1.13 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=171.68, 167.81, 162.41, 148.00, 136.30, 134.72, 132.86, 130.41, 128.87, 128.54, 128.30, 125.18, 123.12, 75.60, 72.93, 68.34, 66.59, 51.21, 30.75, 29.88, 23.89, 14.94. Anal. Calcd for C₂₇H₃₄N₄O₆S: C, 59.76; H, 6.32; N, 10.32. Found: C, 59.55; H, 6.16; N, 10.10.

Benzyl (2S)-5-(3-nitroguanidino)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoate (5n)

Yield: 1,102 mg (90%). ESI/MS (m/z) 613 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.48 (m, 1H), 8.43 (m, 1H), 8.33 (m, 2H), 8.11 (d, *J*=7.8 Hz, 1H), 7.96 (m, 2H), 7.71 (t, *J*=7.8 Hz, 1H), 7.35 (m, 5H), 5.11 (s, 2H), 4.51 (s, 2H), 4.34 (m, 1H), 3.12 (dd, *J*=12.3 Hz, *J*=6.3 Hz, 2H), 1.76 (m, 1H), 1.64 (m, 1H), 1.47 (m, 2H), 1.20 (s, 6H), 1.12 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=171.86, 167.76, 162.41, 148.02, 136.27, 134.69, 132.85, 130.43, 128.87, 128.52, 128.27, 125.18, 123.12, 79.64, 75.61, 72.91, 68.34, 66.55, 51.95, 28.43, 23.90, 19.84. Anal. Calcd for C₂₈H₃₆N₈O₈: C, 54.89; H, 5.92; N, 18.29. Found: C, 54.67; H, 5.75; N, 18.07.

Benzyl (2S)-1-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetyl)pyrrolidine-2-carboxylate (5o)

Yield: 500 mg (53%). ESI/MS (m/z) 509 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.43 (m, 1H), 8.32 (d, *J*=8.1 Hz, 1H), 8.11 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=8.1 Hz, 1H), 7.34 (m, 5H), 5.11 (m, 2H), 4.73 (s, 2H), 4.35 (m, 1H), 3.50 (m, 2H), 2.16 (m, 1H), 1.89 (m, 3H), 1.21 (s, 6H), 1.09 (s, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=172.07, 166.23, 162.49, 148.00, 136.42, 134.69, 132.85, 132.73, 130.44, 128.94, 128.72, 128.45, 125.19, 123.16, 75.31, 73.05, 68.30, 66.28, 58.93, 46.06, 28.90, 24.90, 23.92. Anal. Calcd for C₂₇H₃₂N₄O₆: C, 63.77; H, 6.34; N, 11.02. Found: C, 63.54; H, 6.20; N, 10.81.

General procedure of preparing 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetyl-L-amino acids (6a–o)

At 0°C, to a solution of 1 mmol of 5a–o in 20 mL of methanol 2 mL of aqueous NaOH (4 M) was added. The reaction mixture was stirred for 1 hour, and TLC (EtOAc:petroleum=1:1) indicated the complete disappearance of 5a–o. The reaction mixture was adjusted to pH 2 and evaporated under vacuum to remove methanol and water. The residue was purified with column chromatography (CH₂Cl₂:CH₃OH=5:1) to provide 6a–o.

(2S,3S)-3-methyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoic acid (6a)

Yield: 320 mg (72%). Mp 75°C–76°C; ESI/MS (m/z) 433 [M–H][–]. [α]₂₀^D=0.3 (c=1.1, CH₃OH). IR (KBr): 3,413, 2,971, 2,905, 2,877, 1,682, 1,533, 1,451, 1,349, 1,214, 1,158, 1,062,

902, 823, 718, 573 cm^{–1}. ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.43 (m, 1H), 8.32 (dd, *J*=8.1 Hz, *J*=1.5 Hz, 1H), 8.12 (d, *J*=7.8 Hz, 1H), 7.81 (d, *J*=8.4 Hz, 1H), 7.73 (d, *J*=8.1 Hz, 1H), 4.53 (s, 2H), 4.17 (m, 1H), 1.75 (m, 1H), 1.40 (m, 2H), 1.22 (s, 6H), 1.13 (s, 6H), 0.81 (m, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=173.11, 167.22, 162.53, 148.02, 134.72, 132.89, 130.46, 128.47, 125.17, 123.08, 75.62, 72.94, 68.37, 56.59, 37.11, 25.10, 23.89, 15.92, 11.77. Anal. Calcd for C₂₁H₃₀N₄O₆: C, 58.05; H, 6.96; N, 12.89. Found: C, 57.84; H, 6.82; N, 13.12.

(2S)-3-methyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoic acid (6b)
Yield: 450 mg (78%). Mp 90°C–92°C; ESI/MS (m/z) 419 [M–H][–]. [α]₂₀^D=–18.7 (c=1.2, CH₃OH). IR (KBr): 3,210, 2,970, 2,925, 2,870, 2,699, 1,734, 1,684, 1,536, 1,460, 1,352, 1,205, 1,109, 1,011, 887, 829, 700 cm^{–1}; ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=12.90 (s, 1H), 8.81 (m, 1H), 8.50 (dd, *J*=8.4 Hz, *J*=1.5 Hz, 1H), 8.42 (d, *J*=8.1 Hz, 1H), 8.38 (d, *J*=8.2 Hz, 1H), 7.93 (t, *J*=8.1 Hz, 1H), 4.84 (s, 2H), 4.09 (dd, *J*=5.7 Hz, *J*=8.4 Hz, 1H), 2.03 (m, 1H), 1.47 (s, 6H), 1.42 (s, 6H), 0.85 (d, *J*=6.6 Hz, 6H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=172.82, 166.16, 163.29, 147.95, 136.45, 131.11, 128.90, 127.04, 125.37, 76.77, 75.91, 65.55, 63.29, 57.45, 30.23, 22.12, 19.44, 18.32, 15.63. Anal. Calcd for C₂₀H₂₈N₄O₆: C, 57.13; H, 6.71; N, 13.33. Found: C, 57.35; H, 6.88; N, 13.56.

(2S)-4-methyl-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoic acid (6c)

Yield: 401 mg (82%). Mp 112°C–120°C; ESI/MS (m/z) 433 [M–H][–]. [α]₂₀^D=–44.9 (c=1.2, CH₃OH). IR (KBr): 3,800, 3,216, 2,959, 2,680, 2,694, 1,682, 1,624, 1,585, 1,448, 1,351, 1,202, 1,160, 1,103, 902, 825, 700 cm^{–1}; ¹H NMR (300 MHz, DMSO-*d*₆): δ/ppm=8.61 (s, 1H), 8.45 (d, *J*=7.8 Hz, 1H), 8.27 (t, *J*=7.5 Hz, 2H), 7.82 (t, *J*=7.8 Hz, 1H), 4.62 (s, 2H), 4.20 (m, 1H), 1.50 (m, 3H), 1.35 (s, 6H), 1.27 (s, 6H), 0.88 (d, *J*=3.9 Hz, 3H), 0.81 (d, *J*=4.2 Hz, 3H). ¹³C NMR (75 MHz, DMSO-*d*₆): δ/ppm=173.93, 166.69, 162.88, 147.97, 135.58, 130.78, 128.47, 126.99, 124.22, 76.20, 74.40, 66.99, 65.38, 50.42, 42.71, 23.24, 23.01, 21.69, 19.62, 15.64. Anal. Calcd for C₂₁H₃₀N₄O₆: C, 58.05; H, 6.96; N, 12.89. Found: C, 58.24; H, 6.81; N, 12.65.

(2S)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)propanoic acid (6d)

Yield: 270 mg (20%). Mp 80°C–90°C; ESI/MS (m/z) 391 [M–H][–]. [α]₂₀^D=–2.7 (c=1.3, CH₃OH). IR (KBr): 3,404,

2,980, 2,972, 1,672, 1,612, 1,532, 1,452, 1,351, 1,158, 1,059, 905, 823, 717, 573 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$): δ/ppm =8.39 (s, 1H), 8.23 (d, J =6.3 Hz, 1H), 8.19 (d, J =5.7 Hz, 1H), 7.73 (m, 2H), 4.48 (s, 2H), 3.94 (m, 1H), 1.23 (m, 6H), 1.22 (s, 3H), 1.13 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO-}d_6$): δ/ppm =166.23, 162.52, 148.01, 134.64, 132.72, 130.59, 125.26, 122.99, 76.06, 72.91, 68.34, 65.39, 23.9, 15.63. Anal. Calcd for $\text{C}_{18}\text{H}_{24}\text{N}_4\text{O}_6$: C, 55.09; H, 6.16; N, 14.28. Found: C, 55.28; H, 6.33; N, 14.51.

(2S)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)acetic acid (6e)

Yield: 400 mg (54%). Mp 172°C–176°C; ESI/MS (m/z) 377 $[\text{M-H}]^-$. $[\alpha]_{20}^{\text{D}}=-5.7$ ($c=1.2$, CH_3OH). IR (KBr): 3,431, 3,086, 2,983, 2,960, 1,726, 1,682, 1,606, 1,535, 1,444, 1,348, 1,223, 1,164, 1,120, 1,064, 888, 822, 712, 555 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$): δ/ppm =8.41 (m, 1H), 8.32 (d, J =8.1 Hz, 1H), 8.11 (d, J =7.8 Hz, 1H), 7.86 (m, 1H), 7.73 (t, J =7.8 Hz, 1H), 4.50 (s, 2H), 3.68 (d, J =5.4 Hz, 2H), 1.23 (s, 6H), 1.15 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO-}d_6$): δ/ppm =171.50, 167.32, 162.42, 148.04, 134.66, 132.76, 130.52, 125.23, 123.07, 75.79, 72.95, 68.34, 23.89, 21.58, 19.90. Anal. Calcd for $\text{C}_{17}\text{H}_{22}\text{N}_4\text{O}_6$: C, 53.96; H, 5.86; N, 14.81. Found: C, 53.74; H, 5.70; N, 14.59.

(2S)-3-(4-hydroxyphenyl)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)propanoic acid (6f)

Yield: 327 mg (69%). Mp 79°C–82°C; ESI/MS (m/z) 483 $[\text{M-H}]^-$. $[\alpha]_{20}^{\text{D}}=-0.6$ ($c=1.2$, CH_3OH). IR (KBr): 3,399, 2,980, 2,932, 1,672, 1,619, 1,532, 1,445, 1,351, 1,231, 1,166, 1,113, 830, 709, 539 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$): δ/ppm =12.74 (s, 1H), 9.18 (s, 1H), 8.42 (s, 1H), 8.32 (d, J =7.8 Hz, 1H), 8.07 (d, J =7.5 Hz, 1H), 7.86 (d, J =7.5 Hz, 1H), 7.70 (t, J =7.8 Hz, 1H), 6.93 (d, J =8.1 Hz, 2H), 6.61 (d, J =8.1 Hz, 2H), 4.45 (s, 2H), 4.39 (m, 1H), 2.93 (dd, J =13.5 Hz, J =4.5 Hz, 1H), 2.78 (m, 1H), 1.15 (m, 12H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO-}d_6$): δ/ppm =173.00, 167.12, 162.48, 156.46, 148.02, 134.65, 130.44, 127.52, 125.28, 123.13, 115.45, 75.53, 73.01, 68.25, 65.76, 53.06, 36.24, 23.84, 15.62. Anal. Calcd for $\text{C}_{24}\text{H}_{28}\text{N}_4\text{O}_7$: C, 59.50; H, 5.83; N, 11.56. Found: C, 59.72; H, 5.99; N, 11.78.

(2S)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanedioic acid (6g)

Yield: 230 mg (25%). Mp 54°C–58°C; ESI/MS (m/z) 449 $[\text{M-H}]^-$. $[\alpha]_{20}^{\text{D}}=-20.4$ ($c=1.2$, CH_3OH). IR (KBr): 3,330, 3,112, 2,984, 2,865, 1,740, 1,541, 1,357, 1,250, 1,026, 906,

824, 701, 636, 515 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$): δ/ppm =12.24 (m, 2H), 8.67 (s, 1H), 8.48 (d, J =7.5 Hz, 1H), 8.38 (m, 1H), 8.30 (m, 1H), 7.86 (m, 1H), 4.69 (s, 2H), 4.20 (m, 1H), 2.23 (m, 2H), 1.96 (m, 1H), 1.77 (m, 1H), 1.38 (s, 6H), 1.31 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO-}d_6$): δ/ppm =174.03, 173.11, 172.40, 166.59, 163.05, 148.00, 135.82, 130.93, 124.57, 76.42, 74.92, 66.56, 65.39, 51.44, 30.45, 26.64, 22.78, 21.58, 19.56, 15.66. Anal. Calcd for $\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}_8$: C, 53.33; H, 5.82; N, 12.44. Found: C, 53.54; H, 5.97; N, 12.68.

(2S)-6-amino-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)hexanoic acid (6h)

Yield: 372 mg (80%). Mp 64°C–66°C; ESI/MS (m/z) 448 $[\text{M-H}]^-$. $[\alpha]_{20}^{\text{D}}=-21.2$ ($c=1.2$, CH_3OH). IR (KBr): 3,392, 3,220, 2,982, 2,723, 1,734, 1,684, 1,535, 1,352, 1,206, 1,159, 1,107, 1,020, 883, 826, 698, 507 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$): δ/ppm =8.82 (m, 1H), 8.62 (m, 2H), 8.43 (d, J =8.1 Hz, 1H), 8.14 (m, 2H), 7.94 (t, J =7.8 Hz, 1H), 4.78 (s, 2H), 4.11 (m, 1H), 2.71 (m, 2H), 1.65 (m, 1H), 1.56 (m, 2H), 1.44 (m, 12H), 1.34 (m, 3H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO-}d_6$): δ/ppm =173.27, 165.97, 163.31, 147.97, 136.48, 131.17, 128.99, 125.43, 123.78, 76.86, 76.01, 65.52, 60.21, 52.16, 30.52, 26.83, 22.72, 22.12, 21.54, 21.22, 19.38, 14.55. Anal. Calcd for $\text{C}_{21}\text{H}_{31}\text{N}_5\text{O}_6$: C, 56.11; H, 6.95; N, 15.58. Found: C, 55.85; H, 6.87; N, 15.72.

(2S)-4-amino-4-oxo-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoic acid (6i)

Yield: 326 mg (51%). Mp 89°C–92°C; ESI/MS (m/z) 436 $[\text{M+H}]^+$. $[\alpha]_{20}^{\text{D}}=0.6$ ($c=1.3$, CH_3OH). IR (KBr): 3,370, 3,230, 3,111, 2,982, 2,850, 1,737, 1,678, 1,540, 1,450, 1,357, 1,244, 1,170, 1,028, 906, 829, 702, 638, 517 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$): δ/ppm =9.48 (s, 1H), 8.77 (s, 1H), 8.61 (d, J =8.1 Hz, 1H), 8.32 (t, J =6.9 Hz, 2H), 7.96 (t, J =7.8 Hz, 1H), 7.38 (s, 1H), 6.90 (s, 1H), 4.80 (s, 2H), 4.50 (m, 1H), 2.51 (m, 2H), 1.48 (s, 6H), 1.41 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO-}d_6$): δ/ppm =172.64, 171.62, 165.56, 163.83, 148.11, 136.13, 131.39, 129.16, 125.13, 123.58, 77.08, 76.24, 65.33, 48.82, 36.61, 22.14, 19.28, 15.61. Anal. Calcd for $\text{C}_{19}\text{H}_{25}\text{N}_5\text{O}_7$: C, 52.41; H, 5.79; N, 16.08. Found: C, 52.20; H, 5.63; N, 15.84.

(2S)-5-amino-5-oxo-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoic acid (6j)

Yield: 378 mg (74%). Mp 61°C–63°C; ESI/MS (m/z) 448 $[\text{M-H}]^-$. $[\alpha]_{20}^{\text{D}}=-13.7$ ($c=1.1$, CH_3OH). IR (KBr): 3,413,

3,301, 2,950, 2,875, 1,739, 1,630, 1,541, 1,357, 1,285, 1,243, 1,169, 1,028, 637, 516 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ/ppm =10.74 (m, 1H), 8.77 (s, 1H), 8.61 (d, J =8.1 Hz, 1H), 8.44 (d, J =7.2 Hz, 1H), 8.34 (d, J =7.5 Hz, 1H), 7.96 (t, J =8.1 Hz, 1H), 7.23 (s, 1H), 6.80 (s, 1H), 4.77 (s, 2H), 4.14 (m, 1H), 2.08 (m, 2H), 1.94 (m, 1H), 1.76 (m, 1H), 1.49 (s, 6H), 1.41 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO}-d_6$): δ/ppm =173.79, 173.20, 165.84, 163.71, 148.07, 136.21, 131.31, 129.16, 125.20, 123.62, 76.94, 76.28, 65.47, 51.90, 31.57, 21.07, 22.11, 19.31, 15.61. Anal. Calcd for $\text{C}_{20}\text{H}_{27}\text{N}_5\text{O}_7$: C, 53.45; H, 6.05; N, 15.58. Found: C, 53.22; H, 6.23; N, 15.33.

(2S,3R)-3-hydroxy-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoic acid (6k)

Yield: 280 mg (62%). Mp 121°C – 123°C ; ESI/MS (m/z) 421 $[\text{M}-\text{H}]^-$. $[\alpha]_{20}^{\text{D}}=-2.0$ ($c=1.2$, CH_3OH). IR (KBr): 3,406, 2,979, 2,920, 1,668, 1,611, 1,532, 1,408, 1,351, 1,159, 1,080, 811, 716, 579 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.40 (m, 1H), 8.32 (d, J =8.1 Hz, 1H), 8.09 (d, J =7.5 Hz, 1H), 7.74 (t, J =8.1 Hz, 1H), 7.43 (m, 1H), 4.51 (s, 2H), 3.89 (m, 2H), 1.23 (s, 6H), 1.13 (s, 6H), 0.84 (m, 3H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO}-d_6$): δ/ppm =166.40, 162.53, 148.04, 134.56, 132.68, 130.63, 125.27, 122.96, 76.03, 72.59, 68.32, 65.39, 23.88, 19.23, 15.64. Anal. Calcd for $\text{C}_{19}\text{H}_{26}\text{N}_4\text{O}_7$: C, 54.02; H, 6.20; N, 13.26. Found: C, 53.80; H, 6.13; N, 13.02.

(2S)-3-mercapto-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)propanoic acid (6l)

Yield: 204 mg (24%). Mp 117°C – 118°C ; ESI/MS (m/z) 425 $[\text{M}+\text{H}]^+$. $[\alpha]_{20}^{\text{D}}=-52.8$ ($c=1.3$, CH_3OH). IR (KBr): 3,381, 3,240, 3,105, 2,944, 1,743, 1,668, 1,540, 1,451, 1,356, 1,289, 1,240, 1,170, 1,031, 829, 703, 637, 516 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ/ppm =12.10 (m, 1H), 8.77 (s, 1H), 8.60 (d, J =8.1 Hz, 1H), 8.48 (d, J =7.8 Hz, 1H), 8.32 (d, J =7.8 Hz, 1H), 7.93 (m, 1H), 4.81 (s, 2H), 4.40 (m, 1H), 3.06 (m, 1H), 2.84 (m, 1H), 1.48 (s, 6H), 1.41 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO}-d_6$): δ/ppm =171.82, 165.97, 163.76, 148.09, 136.14, 131.33, 129.11, 125.15, 123.71, 76.88, 76.24, 65.55, 65.36, 22.18, 19.34, 15.62. Anal. Calcd for $\text{C}_{18}\text{H}_{24}\text{N}_4\text{O}_6\text{S}$: C, 50.93; H, 5.70; N, 13.20. Found: C, 50.70; H, 5.54; N, 13.01.

(2S)-4-(methylthio)-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)butanoic acid (6m)

Yield: 393 mg (73%). Mp 59°C – 63°C ; ESI/MS (m/z) 451 $[\text{M}-\text{H}]^-$. $[\alpha]_{20}^{\text{D}}=1.0$ ($c=1.1$, CH_3OH). IR (KBr): 3,406, 2,978,

2,923, 1,679, 1,533, 1,442, 1,350, 1,228, 1,163, 1,117, 1,063, 714, 570 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.43 (m, 1H), 8.33 (d, J =6.9 Hz, 1H), 8.12 (d, J =7.5 Hz, 1H), 8.09 (m, 1H), 7.73 (t, J =8.1 Hz, 1H), 4.51 (s, 2H), 4.29 (m, 1H), 2.40 (m, 2H), 2.00 (s, 3H), 1.96 (m, 1H), 1.84 (m, 1H), 1.22 (s, 6H), 1.13 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO}-d_6$): δ/ppm =173.35, 167.46, 162.49, 147.98, 134.73, 132.81, 130.46, 125.22, 123.11, 79.65, 75.69, 72.93, 68.33, 51.31, 31.16, 30.03, 23.90, 14.96. Anal. Calcd for $\text{C}_{20}\text{H}_{28}\text{N}_4\text{O}_6\text{S}$: C, 53.08; H, 6.24; N, 12.38. Found: C, 53.30; H, 6.39; N, 12.62.

(2S)-5-guanidino-2-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetamido)pentanoic acid (6n)

Yield: 300 mg (64%). Mp 68°C – 75°C ; ESI/MS (m/z) 478 $[\text{M}+\text{H}]^+$. $[\alpha]_{20}^{\text{D}}=-11.2$ ($c=1.2$, CH_3OH). IR (KBr): 3,384, 3,450, 2,983, 1,671, 1,535, 1,354, 1,267, 1,173, 1,037, 641, 517 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.43 (s, 1H), 8.32 (d, J =6.9 Hz, 1H), 8.11 (d, J =6.6 Hz, 1H), 7.97 (d, J =5.7 Hz, 1H), 7.73 (m, 2H), 7.11 (m, 4H), 4.51 (s, 2H), 4.18 (m, 1H), 3.07 (m, 2H), 1.73 (m, 1H), 1.58 (m, 1H), 1.43 (m, 2H), 1.23 (s, 6H), 1.13 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO}-d_6$): δ/ppm =173.70, 167.27, 162.46, 157.21, 148.03, 134.69, 132.83, 130.46, 125.20, 123.09, 75.75, 72.95, 68.36, 51.97, 49.05, 28.94, 25.52, 23.90, 19.84. Anal. Calcd for $\text{C}_{21}\text{H}_{31}\text{N}_7\text{O}_6$: C, 52.82; H, 6.54; N, 20.53. Found: C, 52.61; H, 6.37; N, 20.30.

(2S)-1-(2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yloxy)acetyl)pyrrolidine-2-carboxylic acid (6o)

Yield: 220 mg (23%). Mp 112°C – 115°C ; ESI/MS (m/z) 417 $[\text{M}-\text{H}]^-$. $[\alpha]_{20}^{\text{D}}=-64.9$ ($c=1.2$, CH_3OH). IR (KBr): 3,420, 3,087, 2,978, 2,850, 2,780, 1,727, 1,656, 1,532, 1,445, 1,349, 1,268, 1,162, 1,082, 822, 718 cm^{-1} ; $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ/ppm =8.43 (s, 1H), 8.33 (m, 1H), 8.12 (d, J =7.2 Hz, 1H), 7.73 (t, J =8.1 Hz, 2H), 4.70 (m, 2H), 4.21 (m, 1H), 3.45 (m, 2H), 2.07 (m, 2H), 1.86 (m, 2H), 1.24 (s, 6H), 1.12 (s, 6H). $^{13}\text{C NMR}$ (75 MHz, $\text{DMSO}-d_6$): δ/ppm =166.23, 165.87, 162.51, 147.98, 134.72, 132.84, 130.48, 125.18, 123.23, 75.42, 73.06, 68.27, 65.39, 46.49, 45.95, 29.03, 24.77, 23.93, 15.63. Anal. Calcd for $\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}_6$: C, 57.41; H, 6.26; N, 13.39. Found: C, 57.20; H, 6.11; N, 13.16.

Pain threshold in vivo assay

Male ICR mice weighing 25 ± 2 g were housed in a 12/12 light/dark cycle at a room temperature of $21^\circ\text{C}\pm 2^\circ\text{C}$ for

2 days before being used. Food and tap water were supplied ad libitum. The ethical guidelines described in the National Institutes of Health Guide for the Care and Use of Laboratory Animals were followed throughout the experiments.

The mice were randomly divided into three groups of 12 mice, named the test group, vehicle control group, and positive control group. The mice in the vehicle control group were orally given a single dose of normal saline (NS), in the positive group they were orally given a single dose of 167 $\mu\text{mol/kg}$ of aspirin, while in the test group they were orally given a single dose of 10 $\mu\text{mol/kg}$, 3.3 $\mu\text{mol/kg}$, or 1 $\mu\text{mol/kg}$ of 6a–o in NS. The analgesic effects of 6a–o were evaluated by the tail-flick test. The basic pain threshold value of each mouse was measured four times.

Thirty minutes after administration, the pain thresholds were measured at 30-minute intervals. This measurement was carried out for 180 minutes. The potency of analgesia was indicated by pain threshold variation. The values were calculated according to the equation $\text{PTV} = \text{AAPT}/\text{BPT}$, where PTV = pain threshold variation, BPT = basic pain threshold, and AAPT = pain threshold after administration – basic pain threshold. All values of the pain threshold variation for each animal were averaged and constituted one sample.

Anti-inflammatory assay in vivo

Male ICR mice weighing 25 ± 2 g were housed in a 12/12 light/dark cycle at a room temperature of $21^\circ\text{C} \pm 2^\circ\text{C}$ for 2 days before being used. Food and tap water were supplied ad libitum. The ethical guidelines described in the National Institutes of Health Guide for the Care and Use of Laboratory Animals were followed throughout the experiments.

The mice were randomly divided into three groups of 12 mice, named the test group, vehicle control group, and positive control group. Each of the mice in the vehicle control group, positive control group, and test group was orally given a single dose of NS, 167 $\mu\text{mol/kg}$ of aspirin in NS, and 10 $\mu\text{mol/kg}$, 3.3 $\mu\text{mol/kg}$, or 1 $\mu\text{mol/kg}$ of 6a–s in NS, respectively. Thirty minutes later, 0.03 mL of xylene was applied to both the anterior and posterior surfaces of the right ear. The left ear was considered as a control. Two hours after xylene application, mice were sacrificed and both ears were removed. Using a cork borer with a diameter of 7 mm, several circular sections were taken and weighed. The increase in weight caused by the irritant was measured through subtracting the weight of the untreated left ear section from that of the treated right ear section. The statistical analysis of the data was carried out by use of ANOVA test. $P < 0.05$ was considered significant.

3D-QSAR

The 3D-QSAR of 15 anti-inflammatory agents was analyzed with Discovery Studio 4.0. A subset of 12 anti-inflammatory agents was utilized as a training set for QSAR modeling. To assess the predictive power of the resulting QSAR models, the remaining three antithrombotic agents were employed as a testing set. The Diverse Molecules protocol in Discovery Studio 4.0 was used to build the training and test sets. The inhibition values of the 15 anti-inflammatory agents were represented with % inhibition, and used for QSAR analyses as the response variable. The CHARMM force field was selected for the agents. The electrostatic potential and the van der Waals potential were treated as separate terms: a +1e point charge is used as the electrostatic potential probe, and distance-dependent dielectric constant is used to mimic the solvation effect; a carbon atom with a 1.73 Å radius is used as a probe for the van der Waals potential. The truncation for both the steric and the electrostatic energies was set to 30 kcal/mol. The standard parameters implemented in Discovery Studio 4.0 were used. A partial least-squares model was built using energy grids as descriptors. QSAR models were built using the create 3D-QSAR protocol of Discovery Studio 4.0. The inhibition values for the 15 agents are 6a=25.56%, 6b=17.07%, 6c=35.95%, 6d=20.55%, 6e=20.75%, 6f=63.98%, 6g=39.37%, 6h=35.45%, 6i=11.35%, 6j=20.92%, 6k=-4.10%, 6l=42.93%, 6m=32.60%, 6n=36.57%, 6o=28.87%.

Molecular docking

In the molecular docking, the structure of COX-2 enzyme was treated as rigid and prepared by AutoDockTools 1.5L; ie, merging nonpolar hydrogens and assigning gasteiger charges and autodock elements. Then, the energy-minimized 3D conformations of 15 anti-inflammatory agents were treated as flexible and prepared by AutoDockTools 1.5; ie, merging nonpolar hydrogens, assigning gasteiger charges, finding root and aromatic carbons, detecting rotatable bonds, and setting torsions. The grid box dimensions were set to $22.5 \text{ \AA} \times 30 \text{ \AA} \times 30 \text{ \AA}$ using a grid spacing of 0.375 Å for two average structures.

To determine the probable binding conformations, AutoDock4, a very popular docking program with a high success rate, and the energy-minimized 3D conformations of 15 anti-inflammatory agents were used to dock them into the active site of COX-2 enzyme.

Lamarckian genetic algorithm was used to find the appropriate binding positions, orientations, and conformations of 15 anti-inflammatory agents in the binding site of

the active site of COX-2 enzyme. The global optimization was started with parameters of a population of 100 randomly positioned individuals. The maximum number of energy evaluations was increased to 2.5×10^7 and the maximum number of generations in the Lamarckian genetic algorithm was increased to 2.7×10^5 . A Solis and Wets local search was performed with a maximum number of 3,000. During each simulation experiment, 256 runs were carried out for each antithrombotic agent. The resulting 256 conformations of 15 anti-inflammatory agents were scored by the lowest binding energy and clustered using a root mean squared tolerance of 2.0 Å.

Results and discussion

Preparation of TMNPIZA

1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)oxyacetyl-L-amino acids (TMNPIZA, 6a-o) were synthesized with the synthetic route of Figure 2. In brief, after the bromination of 2-nitropropane, the formed 2,3-dimethyl-2,3-dinitrobutane (73% yield) was reduced to 2,3-dihydroxylamino-2,3-dimethylbutane in 63% yield, which was condensed with 3-nitrobenzaldehyde to provide 1,3-dihydroxyl-2-(3-nitrophenyl)-4,4,5,5-tetramethylimidazolidine ([1], 75% yield). The dehydrolysis of

(1) formed 1-hydroxyl-2-(3-nitrophenyl)-4,4,5,5-tetramethylimidazoline (2) in 66% yield. The alkylation of (2) with $\text{BrCH}_2\text{CO}_2\text{C}_2\text{H}_5$ provided ethyl 2-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)-1-oxyacetate (3) in 82% yield. The saponification of (3) formed (4) (87% yield), which was coupled with L-amino acid benzylesters to form 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)oxyacetyl-L-amino acid benzylesters (5a-o) in 42%–90% yields. The saponification of 5a-o provided TMNPIZA (6a-o) in 20%–87% yields. The common reaction condition and the acceptable yield suggest that this synthetic route is suitable for preparing TMNPIZA.

Analgesic activities of TMNPIZA

The in vivo oral analgesic evaluation was performed on a tail-flick mouse model for a single dose of 10 µmol/kg of TMNPIZA, a single dose of 167 µmol/kg of aspirin (positive control), or a single dose of 10 mL/kg of NS (negative control). The values of basic pain threshold (BPT) and the pain threshold after administration (PTAD) of the mice were tested at 30-minutes intervals for 180 minutes. Using BPT, PTAD, and AAPT (PTAD minus BPT) values, the pain threshold variation (PTV) was calculated based on $\text{PTV} = \text{AAPT}/\text{BPT}$. The data are listed in Figure 3.




Figure 2 Synthetic route of TMNPIZA.

Notes: Reagents used in the reactions: Br₂, NaOH, ethanol; Zn, NH₄Cl, THF; OHC-C₆H₄-NO₂-m, methanol; MgSO₄, NaH, BrCH₂CO₂C₂H₅, NaOH in methanol (4 M), DCC, HOBT, and NMM. (i) Br₂, NaOH, ethanol; (ii) Zn, NH₄Cl, THF; (iii) OHC-C₆H₄-NO₂-m, methanol; (iv) anhydrous MgSO₄, anhydrous THF; (v) NaH, BrCH₂CO₂C₂H₅, anhydrous THF; (vi) NaOH in methanol (4 M); (vii) DCC, HOBT, NMM, anhydrous THF. In 5a and 6a AA=L-Ile; 5b and 6b AA=L-Val; 5c and 6c AA=L-Leu; 5d and 6d AA=L-Ala; 5e and 6e AA=Gly; 5f and 6f AA=L-Tyr; 5g and 6g AA=L-Glu; 5h and 6h AA=L-Lys; 5i and 6i AA=L-Asn; 5j and 6j AA=L-Gln; 5k and 6k AA=L-Thr; 5l and 6l AA=L-Cys; 5m and 6m AA=L-Met; 5n and 6n AA=L-Arg; 5o and 6o AA=L-Pro residues.

Abbreviations: AA, amino acid; TMNPIZA, 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)oxyacetyl-L-amino acids; Bzl, benzyl.




Figure 3 Effect of TMNPZA (6a–o) on the PTV of mice, $n=12$. The analgesic action of 6c, 6e, 6f, and 6n; 6h and 6k; and 6a, 6d, 6i, 6j, and 6l continued for 180 minutes, 150 minutes, and 120 minutes, respectively (compared with NS, $P<0.05$ or $P<0.01$), 6o exhibited no analgesic action at the 30-minute point, 6g exhibited no analgesic action at the 30- and 150-minute points (compared with NS, $P>0.05$), and the analgesic action of 6m continued for 90 minutes (compared with NS, $P<0.05$ or $P<0.01$).

Abbreviations: SD, standard deviation; TMNPZA, 1-(4,4,5,5-tetra-methyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)-oxyacetyl-L-amino acids; PTV, pain threshold variation; NS, normal saline; Asp, aspirin.

It was found that the analgesic action of 6c, 6e, 6f, and 6n; 6h and 6k; and 6a, 6d, 6i, 6j, and 6l continued 180 minutes, 150 minutes, and 120 minutes, respectively (being significantly higher than NS). Of 15 compounds, 6o exhibited no analgesic action at the 30-minute point (being equal to that of NS), 6g exhibited no analgesic action at the 30-minute and 150-minute points (being equal to that of NS), and the

analgesic action of 6m continued only for 90 minutes (being significantly higher than NS). These data emphasized that TMNPZA exhibited analgesic action in a rapid and long-acting manner. On the other hand, at different time points, the TMNPZA offered different activities. At the 60-minute and 90-minute points, 6f possessed the highest activity. This means that the side chain of the amino acid residue significantly affects the analgesic activity, and when the side chain contains the group $-\text{CH}_2\text{C}_6\text{H}_4-\text{OH}-\text{p}$, the analgesic agent will have desirable activity.

Dose-dependent action of 6f in analgesic action

To reveal the dose dependence of the analgesic action, 6f was selected as a representative of TMNPZA. Its activities were tested at 10, 3.3, and 1 $\mu\text{mol}/\text{kg}$ doses, and the data are shown in Figure 4. At a 10 $\mu\text{mol}/\text{kg}$ dose, the analgesic action continued for 180 minutes. At a 3.3 $\mu\text{mol}/\text{kg}$ dose, the analgesic action was observed at 60-, 90-, and 120-minute points only, while 1 $\mu\text{mol}/\text{kg}$ of dose gave no observable analgesic action. These data demonstrate that at 60-, 90-, and 120-minute points, 6f increased the pain threshold in a dose-dependent manner.

Anti-inflammatory activities of TMNPZA

To explore the utility of TMNPZA in treating inflammatory pain, 6a–o were assayed on xylene-induced ear edema of mice *in vivo*. The data are shown in Figure 5. It was




Figure 4 Effect of 6f dose on the PTV of mice, $n=12$.

Abbreviations: SD, standard deviation; PTV, pain threshold variation.




Figure 5 Anti-inflammatory activities of 6a–o on xylene-induced ear edema model, n=12; NS=vehicle; doses of aspirin-1 and -2 were 167 $\mu\text{mol/kg}$ and 16.7 $\mu\text{mol/kg}$, respectively; dose of 6a–o was 10 $\mu\text{mol/kg}$.

Abbreviations: SD, standard deviation; NS, normal saline.

noted that at 10 $\mu\text{mol/kg}$ of dose, except for 6i and 6k, all the others exhibited anti-inflammatory action. The activities of 6b ($P < 0.05$) and 6a, 6c–h, 6j, and 6l–o ($P < 0.01$) were significantly higher than those of NS. The activities of 6g, 6l, and 6n were significantly higher than those of 6d, 6e, and 6j ($P < 0.05$). Besides, the activity of 10 $\mu\text{mol/kg}$ (6f) was equal to that of 167 $\mu\text{mol/kg}$ aspirin, suggesting that the effective dose of 6f was 16.7-fold lower than that of aspirin.

Of 15 anti-inflammatory agents, 6f possessed the highest activity. This means that the side chain of the amino acid residue significantly affects the activity, and when the side chain contains the group $-\text{CH}_2\text{C}_6\text{H}_4-\text{OH}-p$, the anti-inflammatory agent will have desirable activity.

Dose-dependent action of 6f in anti-inflammatory action

To reveal the dose dependence of the anti-inflammatory action, 6f was selected as a representative of TMNPIZA. Its activities were tested at 10, 3.3, and 1 $\mu\text{mol/kg}$ doses, and the data are shown in Figure 6. On a xylene-induced ear edema model, the ear edema of mice in 10 $\mu\text{mol/kg}$ group was significantly lower than that of mice in the 3.3 $\mu\text{mol/kg}$ group ($P < 0.01$), and the ear edema of mice in the 1 $\mu\text{mol/kg}$ group was significantly higher than that of mice in the 3.3 $\mu\text{mol/kg}$ group ($P < 0.01$) and was significantly lower than that of mice in the NS group ($P < 0.01$). Thus, on the xylene-induced ear

edema model, 6f gave a dose-dependent anti-inflammatory response, and its effective dose was 1 $\mu\text{mol/kg}$.

3D-QSAR of TMNPIZA inhibiting inflammation

To rationally design the inhibitors of inflammatory pain afterwards, the 3D-QSAR of the 15 anti-inflammatory agents was analyzed, for which a training set of 12 agents (6b–g, 6i, and 6k–o) and a test set of three agents (6a, 6h, and 6j) was set. Their % inhibition values (from -4.10% to 63.98%)




Figure 6 Effect of various doses of 6f on xylene-induced mouse ear edema, n=12. **Abbreviations:** SD, standard deviation; NS, normal saline.




Figure 7 Graphical relationship of experimental and predicted inhibitions of 12 agents in the training set and three agents in the test set.

in inhibiting xylene-induced ear edema of mice were used, and the 3D-QSAR was analyzed according to a standard method. In the analysis, the comparative molecular field analysis (CoMFA) field descriptors were used as independent variables, and the % inhibition values were used as the dependent variables. The values of n , r , and r^2 were 12, 0.994, and 0.989, respectively. The relationship of the experimental and predicted inhibitions is shown in Figure 7.

With the relationship of Figure 7, the inhibitions of the test set (6a, 6h, and 6j) were calculated and the relationship of their experimental inhibitions of 6a-, 6h-, and 6j-treated mice (25.56%, 35.45%, and 20.92%) and the calculated inhibitions of 6a-, 6h-, and 6j-treated mice (24.66%, 35.63%, and 22.86%) are also shown in Figure 7. The prediction errors were -3.5% ,

0.5% , and 9.3% , respectively, suggesting the the relationship of Figure 7 had good validity. The visual 3D-QSAR contains steric fields, the green region mirroring a bulky group increases the activity and the yellow region mirroring bulky group decreases the activity (Figure 8A), and electrostatic fields, the blue region mirroring electron-donating group increases the activity and the red region mirroring an electron-withdrawing group decreases the activity (Figure 8B).

According to Figure 8B, all side chains of the amino acid residues of TMNPIZA is located in the blue region, and the electron-donating side chains will enhance the anti-inflammatory activity.

As mentioned, of 15 agents, 6f and 6n had the highest analgesic activity at 60- and 90-minute points, and possessed




Figure 8 Graphical 3D-QSAR correlating to the steric field (A) and electrostatic field (B) with the chemical environment of 15 anti-inflammatory agents in the active site of COX-2 enzyme.

Abbreviations: 3D-QSAR, three-dimensional quantitative structure–activity relationship; COX-2, cyclooxygenase-2.

the highest anti-inflammation activity. Based on Figure 8B, it is hypothesized that the p-electron-containing group is the electron-donating group, and when it is located in the blue region, the agent will exhibit good analgesic and anti-inflammation activities. Due to the side chain of the amino acid, residue of 6f contains $-\text{CH}_2\text{C}_6\text{H}_4-\text{OH}-\text{p}$, which is p-electron-containing groups, and is located in the blue region of Figure 8B. 6f not only possesses good analgesic activity but also possesses good anti-inflammatory activity.

Conclusion

The docking investigation, the in vivo anti-inflammation/analgesic evaluations, and the 3D-QSAR analysis provided evidence that the 15 novel 1-(4,4,5,5-tetramethyl-2-(3-nitrophenyl)-4,5-dihydroimidazol-1-yl)oxyacetyl-L-amino acids should be promising candidates for developing drugs to improve inflammatory pain therapy.

Acknowledgments

This work was supported by the Beijing Municipal Science and Technology Commission (Z141100002114049), the Project of Construction of Innovative Teams and Teacher Career Development for Universities and Colleges Under Beijing Municipality, the NSFC (81172930, 81270046, 81373265 and 81202412), and Beijing Natural Science Foundation (7132032).

Disclosure

The authors report no conflicts of interest in this work.

References

- Li X, Bantel C, Conklin D, Conklin D, Childers SR, Eisenach JC. Repeated dosing with oral allosteric modulator of adenosine A1 receptor produces tolerance in rats with neuropathic pain. *Anesthesiology*. 2004; 100(4):956–961.
- Grim TW, Ghosh S, Hsu KL, Cravatt BF, Kinsey SG, Lichtman AH. Combined inhibition of FAAH and COX produces enhanced anti-allodynic effects in mouse neuropathic and inflammatory pain models. *Pharmacol Biochem Behav*. 2014;124:405–411.
- Kam YL, Back SK, Kang B, et al. HYP-1, a novel diamide compound, relieves inflammatory and neuropathic pain in rats. *Pharmacol Biochem Behav*. 2012;103(1):33–42.
- Paulino N, Rodrigues NC, Pardi PC, et al. Evaluation of anti-inflammatory effect of synthetic 1,5-bis(4-acetoxy-3-methoxyphenyl)-1,4-pentadien-3-one, HB2. *Bioorg Med Chem*. 2009;17(13):4290–4295.
- Buvanendran A, Reuben SS, Kroin JS. Recent advances in nonopioid analgesics for acute pain management. *Tech Reg Anesth Pain Manag*. 2007;11(1):19–26.
- García-Martínez C, Fernández-Carvajal A, Valenzuela B, et al. Design and characterization of a noncompetitive antagonist of the transient receptor potential vanilloid subunit 1 channel with in vivo analgesic and anti-inflammatory activity. *Pain*. 2006;7(10):735–746.
- Sobczak M, Pilarczyk A, Jonakowski M, Lipkowski AW, Fichna J. Anti-inflammatory and antinociceptive action of the dimeric dimeric-kephalin peptide biphalin in the mouse model of colitis: new potential treatment of abdominal pain associated with inflammatory inflammatory bowel diseases. *Peptides*. 2014;60:102–106.
- Cao D, Zhang Z, Xie RG, Jiang BC1, Ji RR, Gao YJ. Chemokine CXCL1 enhances inflammatory pain and increases NMDA receptor activity and COX-2 expression in spinal cord neurons via activation of CXCR2. *Exp Neurol*. 2014;261:328–336.
- Kono M, Matsumoto T, Imaeda T, et al. Design, synthesis, and biological evaluation of a series of piperazine ureas as fatty acid amide hydrolase inhibitors. *Bioorg Med Chem*. 2014;22(4):1468–1478.
- Da Silva YKC, Augusto CV, de Castro Barbosa ML, et al. Synthesis and pharmacological evaluation of pyrazine N-acylhydrazone derivatives designed as novel analgesic and anti-inflammatory drug candidates. *Bioorg Med Chem*. 2010;18(14):5007–5015.
- Palkar MB, Singhai AS, Ronad PM, et al. Synthesis, pharmacological screening and in silico studies of new class of diclofenac analogues as a promising anti-inflammatory agents. *Bioorg Med Chem*. 2014;22(10):2855–2866.
- Khalil NA, Ahmed EM, Mohamed KO, Nissan YM, Zaitone SA. Synthesis and biological evaluation of new pyrazolone–pyridazine conjugates as anti-inflammatory and analgesic agents. *Bioorg Med Chem*. 2014;22(7):2080–2089.
- Irannejad H, Kebriaieezadeh A, Zarghi A, et al. Synthesis, docking simulation, biological evaluations and 3D-QSAR study of 5-Aryl-6-(4-methylsulfonyl)-3-(methylthio)-1,2,4-triazine as selective cyclooxygenase-2 inhibitors. *Bioorg Med Chem*. 2014;22(2):865–873.
- Zhao M, Li Z, Peng L, et al. A new class of analgesic agents toward prostacyclin receptor inhibition: Synthesis, biological studies and QSAR analysis of 1-hydroxyl-2-substituted phenyl-4,4,5,5-tetramethylimidazolines. *Eur J Med Chem*. 2008;43(5):1048–1058.
- Zhao M, Li Z, Peng L, Tang YR, et al. Novel 1-oxyl-2-substituted phenyl-4,4,5,5-tetramethylimidazolines: synthesis, selectively analgesic action, and QSAR analysis. *Bioorg Med Chem*. 2007;15(8):2815–2826.

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>

Dovepress

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.