

Cognitive dysfunction and poor health literacy are common in veterans presenting with acute coronary syndrome: insights from the MEDICATION study

Lucas N Marzec¹
Evan P Carey¹
Anne C Lambert-Kerzner¹
Eric J Del Giacco²
Stephanie D Melnyk³
Chris L Bryson⁴
Ibrahim E Fahdi²
Hayden B Bosworth³
Fran Fiocchi⁵
P Michael Ho¹

¹Division of Cardiology, Denver VA Medical Center, Denver, CO, USA;

²Department of Medicine, Little Rock VA Medical Center, Little Rock, AR, USA; ³Department of Medicine, Durham VA Medical Center, Durham, NC, USA; ⁴Department of Medicine, Puget Sound VA Medical Center, Seattle, WA, USA; ⁵American College of Cardiology, Washington, DC, USA

Background: Patient nonadherence to cardiac medications following acute coronary syndrome (ACS) is associated with increased risk of recurrent events. However, the prevalence of cognitive dysfunction and poor health literacy among ACS patients and their association with medication nonadherence are poorly understood.

Methods: We assessed rates of cognitive dysfunction and poor health literacy among participants of a clinical trial that tested the effectiveness of an intervention to improve medication adherence in patients hospitalized with ACS. Of 254 patients, 249 completed the Rapid Estimate of Adult Literacy in Medicine, Revised (REALM-R) survey, an assessment of risk for poor literacy, and the St Louis University Mental Status (SLUMS) exam, a tool assessing for neurocognitive deficits, during ACS hospitalization. We assessed if SLUMS or REALM-R scores were associated with medication adherence.

Results: Based on SLUMS score, 14% of patients were categorized as having dementia, and 52% with mild neurocognitive disorder (MNCD). Based on REALM-R score of ≤ 6 , 34% of patients were categorized as at risk for poor health literacy. There was no association between poor health literacy and medication nonadherence. Of those with MNCD, 35.5% were nonadherent, compared to 17.5% with normal cognitive function and 6.7% with dementia. In multivariable analysis, cognitive dysfunction was associated with medication nonadherence ($P=0.007$), mainly due to an association between MNCD and nonadherence (odds ratio =12.2, 95% confidence interval =1.9 to 243; $P=0.007$). Cognitive status was not associated with adherence in patients randomized to the intervention.

Conclusion: Cognitive dysfunction and risk for poor health literacy are common in patients hospitalized with ACS. We found an association between MNCD and medication nonadherence in the usual care group but not in the intervention group. These findings suggest efforts to screen for MNCD are needed during ACS hospitalization to identify patients at risk for nonadherence and who may benefit from an adherence intervention.

Keyword: medication adherence

Introduction

Mild cognitive dysfunction is present in 15% of those over the age of 70,¹ affecting an estimated 5.4 million people in the United States.² Cognitive dysfunction may have implications following acute coronary syndrome (ACS) as its presence may impact patients' ability to understand and adhere to secondary prevention medication regimens that potentially consist of several new medicines with differing dosing schedules.³ Up to 12% of patients with ACS have cognitive dysfunction;⁴ however, the relationship

Correspondence: Lucas N Marzec
Academic Office I, Room #7104, 12631
E 17th Avenue, Campus Box B-130,
Aurora, CO 80045, USA
Tel +1 303 724 2105
Fax +1 303 724 2094
Email lucas.marzec@ucdenver.edu

between cognitive dysfunction at time of presentation of ACS and subsequent medication nonadherence is poorly defined.

Poor health literacy is present in roughly 30% of ambulatory patients and is associated with cognitive dysfunction.⁵ Independent of cognitive dysfunction, level of health literacy is an important predictor of medication adherence,^{6,7} and patients with low levels of literacy are at higher risk for clinically relevant medication errors.⁸ Limited data exist regarding the prevalence and impact of poor health literacy in those presenting with ACS.⁴ Further, the prevalence of coincident poor health literacy and cognitive dysfunction in the setting of ACS, and their relationship with medication nonadherence, is unknown.

To assess the prevalence of cognitive dysfunction and poor health literacy, and their associations with medication nonadherence, we analyzed data from the Multifaceted Intervention to Improve Cardiac Medication Adherence and Secondary Prevention Measures (MEDICATION) study. This was a randomized clinical trial that assessed the effect of a multifaceted intervention on medication adherence in veterans presenting with ACS.⁹ In the study, a greater proportion of patients randomized to the intervention were adherent compared to the usual care patients (89% versus 74%; $P=0.03$). The objectives of the current analysis were to describe the prevalence of cognitive dysfunction and poor health literacy in patients presenting with ACS. In addition, we sought to evaluate if either cognitive dysfunction or health literacy were independently associated with medication nonadherence in the year following ACS hospital discharge among patients randomized to usual care. Then, we assessed the extent to which these associations would be modified among patients randomized to the intervention group given the positive effects of the intervention demonstrated in the main study.

Methods

Patient population

The design of the randomized trial has been described in detail previously.¹⁰ In brief, this multicenter, prospective clinical trial enrolled veterans at one of four Veterans Affairs (VA) medical centers with ACS as the primary reason for hospital admission. All patients admitted with ACS (defined as ST-elevation myocardial infarction, non-ST elevation myocardial infarction, or unstable angina) who used the VA as their usual source of healthcare were screened for eligibility to participate. Patients were ineligible if they were admitted for a noncardiac primary diagnosis, were planned to be discharged to a nursing home or skilled nursing facility, had an irreversible noncardiac medical condition likely to affect

6-month survival, lacked telephone or cellular phone service, or did not regularly fill medications at a VA pharmacy. Patients were randomized to a multifaceted intervention designed to improve medication adherence and outcomes or usual care. The intervention consisted of pharmacist-led medication reconciliation and tailoring, patient education, collaborative care between the study pharmacist and a patient's primary care doctor and/or cardiologist, and two types of voice messaging including educational and medication refill reminder calls.

Exposure variables

Of 254 enrolled patients, 249 (98%) completed the St Louis University Mental Status (SLUMS) exam, a tool assessing for neurocognitive deficits, and the Rapid Estimate of Adult Literacy in Medicine, Revised (REALM-R) survey, an assessment of patients at risk for poor literacy skills, at baseline during their ACS hospitalization.

The SLUMS exam is a 30-point screening questionnaire that tests orientation, memory, attention, and executive function. The sensitivity and specificity of the SLUMS exam has been validated in a VA cohort.¹¹ As compared to the clinical dementia rating, the SLUMS exam has a sensitivity of 74% and specificity of 65% to detect mild neurocognitive disorder (MNCD) and a sensitivity of 93% and specificity of 96% to detect dementia in the VA population.¹² Patient scores are interpreted in the context of the level of patient education. For patients with at least a high school education, a score of 27–30 is normal, 21–26 represents MNCD, and 1–20 represents dementia. For those with less than a high school education, a score of 25–30 is normal, 20–24 represents MCND, and 1–19 represents dementia. In our study, patients with a SLUMS exam score ≤ 20 were categorized as having dementia, those with a score between 21 and 26 were categorized as having MNCD, and those with a score ≥ 27 were categorized as normal.

The REALM-R survey is a commonly used tool for assessing patients at risk for poor health literacy and correlates well with standardized reading tests.¹³ Scores range from 0 to 8, and a score of ≤ 6 corresponds to a sixth grade reading level.¹³ In our analysis, patients with a score ≤ 6 on the REALM-R survey were categorized as at high risk of health illiteracy, while those with a score ≥ 7 were categorized as normal.

Outcomes

The primary outcome was the proportion of patients who were adherent to cardioprotective medication regimens

in the year following ACS. Medications included in the adherence analysis were beta-blockers, statins, clopidogrel, angiotensin converting enzyme inhibitors, and angiotensin receptor blockers. Aspirin was not included in the analysis due to the large proportion of veterans who obtain aspirin over-the-counter.

Of the 249 patients with baseline survey data, 238 had complete follow-up data available for the adherence analysis. Medication adherence was calculated based on the proportion of days covered (PDC) during a follow-up period of 365 days that a patient had a medication available. The calculation was adjusted for inpatient hospitalization, medication cancellation, medication changes within class, medication fills prior to enrollment date, and death. Adherence for each class of medication was scored from 0 to 1.00 (perfect medication coverage) and was then averaged for all prescribed classes of medications to calculate a summary PDC. Patients with a summary PDC >0.80 were categorized as adherent.

Statistical analysis

Demographic variables were compared across SLUMS and REALM-R categories separately using chi-square tests for categorical variables and *t*-tests or *f*-tests for continuous variables. Distributions were summarized as means and standard deviations for continuous variables and as proportions and numbers for categorical variables. Using medication adherence as an outcome, we used three logistic regression models to assess 1) the bivariate association between REALM-R and adherence, 2) the bivariate association between SLUMS and adherence, and 3) a multivariate model with SLUMS and REALM-R as primary predictors, adjusted for potential confounders. We selected confounders from age,

sex, race, smoking history, hypertension, hyperlipidemia, diabetes mellitus, cerebrovascular disease, chronic lung disease, coronary artery disease history, peripheral arterial disease, depression, and chronic kidney disease using backwards stepwise variable selection with a *P*-value cutoff of 0.2. We did not include the primary predictors in the models while selecting the covariates. We fit all three models in the usual care group and separately in the intervention group. An alpha level of 0.05 was used as a cutoff for all *P*-values.

Results

Based on the SLUMS questionnaire, 14% (*n*=36) of patients were categorized as having dementia and 52% (*n*=130) were categorized as having MNCD. Overall, 67% of patients were categorized as having some level of cognitive dysfunction. Both increasing age (*P*=0.01) and non-white race (*P*=0.03) were associated with the presence of dementia (Table 1).

Based on REALM-R survey score, 34% (*n*=85) were categorized as at risk for poor health literacy. Non-white race was the only demographic variable associated with health literacy and was negatively associated with REALM-R (Table 2). Coincident cognitive dysfunction and risk for poor health literacy were present in roughly one-quarter of patients. REALM-R and SLUMS scores were correlated, though the strength of this correlation was low (Kendall correlation =0.18; significance *P*=0.005).

In the usual care arm, 25.6% (30/117) of patients were nonadherent to their cardiovascular medications. Cognitive dysfunction as measured by SLUMS was associated with medication nonadherence in unadjusted analysis (*P*=0.017), where patients with MNCD were more likely to have non-adherence; 35.5% (22/62) of patients with MNCD were

Table 1 SLUMS score and covariates

	Dementia	MNCD	Normal	All	P-value
N subjects	37	130	83	250	
Age	66.54 (9.36)	64.23 (9.28)	61.7 (8.3)	63.8 (9.07)	0.02
Sex – number male	36 (97.3)	128 (98.5)	80 (96.4)	244 (97.6)	0.62
Race – number white	23 (62.2)	102 (78.5)	69 (83.1)	194 (77.6)	0.04
Smoking history	13 (36.1)	42 (32.6)	22 (26.5)	77 (31)	0.50
Hypertension	33 (91.7)	115 (88.5)	79 (95.2)	227 (91.2)	0.24
Hyperlipidemia	31 (86.1)	107 (82.9)	73 (88)	211 (85.1)	0.60
Diabetes mellitus	16 (44.4)	59 (45.4)	39 (47)	114 (45.8)	0.96
Cerebrovascular disease	3 (8.3)	10 (7.7)	6 (7.2)	19 (7.6)	0.98
Chronic lung disease	12 (33.3)	24 (18.5)	15 (18.1)	51 (20.5)	0.11
History of prior CAD	24 (66.7)	83 (63.8)	53 (64.6)	160 (64.5)	0.95
Peripheral arterial disease	5 (13.9)	15 (11.6)	5 (6.1)	25 (10.1)	0.31
Chronic kidney disease	11 (30.6)	28 (21.5)	20 (24.4)	59 (23.8)	0.53
Depression	16 (44.4)	42 (32.3)	29 (34.9)	87 (34.9)	0.40

Notes: The numbers in parentheses are standard deviations for continuous variables and percentages for categorical variables.

Abbreviations: CAD, coronary artery disease; MNCD, mild neurocognitive disorder; SLUMS, St Louis University Mental Status exam.

Table 2 REALM-R score and covariates

	Health illiterate	Normal	All	P-value
N subjects	85	165	250	
Age	64.87 (8.16)	63.24 (9.47)	63.8 (9.07)	0.18
Sex – number male	85 (100)	159 (96.4)	244 (97.6)	0.18
Race – number white	55 (64.7)	139 (84.2)	194 (77.6)	<0.01
Smoking history	28 (32.9)	49 (30.1)	77 (31)	0.75
Hypertension	75 (88.2)	152 (92.7)	227 (91.2)	0.35
Hyperlipidemia	72 (85.7)	139 (84.8)	211 (85.1)	0.99
Diabetes mellitus	39 (45.9)	75 (45.7)	114 (45.8)	1.00
Cerebrovascular disease	7 (8.2)	12 (7.3)	19 (7.6)	0.99
Chronic lung disease	22 (25.9)	29 (17.7)	51 (20.5)	0.18
History of prior CAD	59 (69.4)	101 (62)	160 (64.5)	0.31
Peripheral arterial disease	8 (9.4)	17 (10.5)	25 (10.1)	0.96
Chronic kidney disease	18 (21.2)	41 (25.2)	59 (23.8)	0.59
Depression	36 (42.4)	51 (31.1)	87 (34.9)	0.10


Notes: The numbers in parentheses are standard deviations for continuous variables and percentages for categorical variables.

Abbreviations: CAD, coronary artery disease; REALM-R, Rapid Estimate of Adult Literacy in Medicine, Revised survey.

nonadherent, compared to 17.5% (7/40) of patients with normal cognitive function and 6.7% (1/15) of patients with dementia (Figure 1). In contrast, health illiteracy as measured by REALM-R was not associated with medication nonadherence ($P=0.92$). In multivariable models adjusting for race, age, and history of diabetes, cognitive dysfunction was associated with medication nonadherence ($P=0.007$), primarily due to an association between MNCD and nonadherence (odds ratio [OR] =12.2; 95% confidence interval [CI] =1.9–243), while normal cognition was not associated with medication

nonadherence (OR =4.8; 95% CI =0.6–101.5) (Table 3). Finally, REALM-R was not associated with adherence in the adjusted model ($P=0.31$).

In the intervention group, 10.7% (13/121) of patients were nonadherent to their cardiovascular medications. SLUMS score was not associated with adherence in the intervention group in unadjusted ($P=0.98$) or adjusted ($P=0.91$) analysis. Some 11.3% (7/62) of patients with MNCD were nonadherent, compared to 10% (4/40) of patient with normal cognitive function and 10.5% (2/19) of patients with dementia. In addition, REALM-R score was also not associated with adherence in the intervention group in unadjusted ($P=0.13$) or adjusted ($P=0.17$) analysis.

**Figure 1** Medication adherence across SLUMS score.

Abbreviations: MNCD, mild neurocognitive disorder; SLUMS, St Louis University Mental Status exam.

Discussion

The objectives of this study were to describe the prevalence of cognitive dysfunction and health illiteracy among patients hospitalized for ACS and to evaluate the association with medication nonadherence in the year following ACS hospitalization. We found that cognitive dysfunction was common (67%) and associated with medication nonadherence in the usual care group but not in the intervention group, suggesting that the intervention may have modified this association. In contrast, poor health literacy was common (34%) but not associated with medication adherence. These findings suggest patients with MNCD are at risk for poor medication adherence following ACS, and interventions focusing on adherence may help improve the medication-taking behavior of these patients.

The prevalence of cognitive dysfunction and risk for poor health literacy in our cohort, 67% and 34%, respectively, are higher than prior reports. These studies found a

Table 3 Relationship between SLUMS and REALM-R scores and medication nonadherence

	Unadjusted				Adjusted		
	Probability nonadherent	OR	OR 95% CI	P-value	OR	OR 95% CI	P-value
SLUMS	–	–	–	0.017	–	–	0.01
Dementia	0.07	1.0	–	–	1.0	–	–
MNCD	0.35	7.7	1.4–144.2	–	12.2	1.9–243.0	–
Normal	0.18	3.0	0.5–58.2	–	4.8	0.6–101.5	–
REALM-R	–	–	–	0.92	–	–	0.85
Normal	0.25	1.0	0.4–2.3	–	1.1	0.4–3.0	–
Health illiterate	0.26	1.0	–	–	1.0	–	–
Race							
Non-white	–	–	–	–	3.5	1.2–10.3	0.02
Diabetes							
No history	–	–	–	–	2.7	1.0–8.0	0.06

Note: Table of multivariate model results for usual care.

Abbreviations: CI, confidence interval; MNCD, mild neurocognitive disorder; OR, odds ratio; REALM-R, Rapid Estimate of Adult Literacy in Medicine, Revised survey; SLUMS, St Louis University Mental Status exam.

significantly lower rate of cognitive dysfunction in patients presenting with ACS (2.1% of over 85,000 patients in one series)¹⁴ but relied on claims data, likely underestimating the prevalence. The Pharmacist Intervention for Low Literacy in Cardiovascular Disease (PILL-CVD) study assessed the prevalence of cognitive dysfunction and the level of health literacy in patients enrolled in a randomized control trial assessing the effect of a pharmacist-based intervention on the rate of serious medication errors in the first 30 days following hospital discharge for ACS or decompensated heart failure.⁴ The rate of cognitive dysfunction in that cohort was 11.5%, considerably higher than found in analyses based on claims data, though significantly lower than in our cohort. The rate of poor health literacy in the PILL-CVD study was 18.7%, also lower than that of our cohort. These differences may reflect the different sensitivity of the screening tools used or differences in the patient populations in the PILL-CVD study and in our study.

Medication nonadherence following ACS is common and is associated with an increased risk of recurrent myocardial infarction and death.^{15–18} A number of factors have been associated with medication nonadherence following ACS, including older age, lower level of education, and preexisting cardiovascular disease.¹⁸ While the mechanisms contributing to nonadherence are complex, cognitive dysfunction and low levels of health literacy have been associated with adverse outcomes in those with ACS and heart failure, respectively. An investigation of the prognostic influence of comorbid conditions in a cohort of patients with non-ST elevation ACS found the presence of dementia predicted death (hazard ratio 3.1).¹⁹ In a study of ambulatory heart failure patients with New York Heart Association class II–IV symptoms, 37% had low literacy. Low literacy was associated with a significantly

increased rate of all-cause hospitalization (incidence rate ratio [IRR] 1.31) and heart failure-related hospitalization (IRR 1.46).²⁰ Our findings further support that cognitive dysfunction and poor health literacy are common in those with ACS, and may contribute to medication nonadherence and adverse clinical outcomes in this population.

Our data suggest MNCD may play an important role in medication nonadherence following ACS, and the rate of nonadherence may be improved with multidisciplinary interventions designed to improve adherence. In our cohort, dementia was not associated with adherence. While the mechanisms for the difference in adherence between those with MNCD and dementia are not known, one potential explanation is greater support systems in place for those with recognized significant cognitive dysfunction. Patients with MNCD may lack the social support or assistance from the healthcare system more frequently available to those with dementia. Therefore patients with MNCD are at increased risk for medication nonadherence. More data are needed to better understand whether the combination of cognitive dysfunction and poor adherence leads to worse clinical outcomes after ACS.

Cognitive dysfunction and risk for poor health literacy were poorly predicted by traditional demographic variables (only age and race were significantly associated), and were not associated with preexisting medical comorbidities in our cohort. Non-white race has been associated with medication nonadherence post-ACS by others,²¹ and further efforts to identify factors contributing to this association are needed. Systematic implementation of protocols for screening for MNCD may allow for identification of patients for whom a targeted intervention to improve medication adherence post-ACS is of benefit. Further

studies should assess the impact of screening for MNCD to facilitate implementation of interventions to improve medication adherence and clinical outcomes in patients after ACS hospital discharge. In contrast, we did not find an association between health literacy and adherence. Additional studies should assess whether health numeracy impacts medication adherence.

Several limitations to our study should be acknowledged. First, we analyzed data from a randomized control trial that enrolled patients exclusively from VA medical centers. The majority of these patients were men and white. The generalizability of these findings outside of the VA population is unknown. Second, the VA is an integrated healthcare delivery system. Is it unclear how these findings may be translated to a nonintegrated healthcare delivery system. However, the findings are relevant to veterans who receive care in the VA system as well as the large number of patients who receive care in other large integrated healthcare delivery systems. Third, we do not have data regarding all variables that have been previously associated with medication adherence, including housing status. It is possible this or other unrecognized factors may have contributed to the difference in adherence between those with MNCD and dementia observed in our study. Fourth, we do not have data regarding the level of education achieved for patients in our cohort. We categorized patients' cognitive status based on SLUMS score assuming at least a high school education. It is possible this overestimated the prevalence of significant cognitive dysfunction in our cohort. Lastly, we assessed medication adherence with pharmacy refill data. It is possible that patients refilled medications without ingesting the pills. However, pharmacy refill data as a measure of adherence has been validated in several other medication adherence studies.²²

In conclusion, we found cognitive dysfunction and risk for poor health literacy to be common in a cohort of veterans hospitalized for ACS. Mild cognitive impairment was associated with lower rates of medication adherence in patients randomized to usual care. There was no association between level of cognitive dysfunction and medication adherence in those patients randomized to the intervention, suggesting a role for such a multidisciplinary adherence intervention in patients with cognitive dysfunction and ACS.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Roberts RO, Geda YE, Knopman DS, et al. The Mayo Clinic Study of Aging: design and sampling, participation, baseline measures and sample characteristics. *Neuroepidemiology*. 2008;30(1):58–69.
2. Plassman BL, Langa KM, Fisher GG, et al. Prevalence of cognitive impairment without dementia in the United States. *Ann Intern Med*. 2008;148(6):427–434.
3. Sanf  lix-Gimeno G, Peir   S, Ferreros I, et al. Adherence to evidence-based therapies after acute coronary syndrome: a retrospective population-based cohort study linking hospital, outpatient, and pharmacy health information systems in Valencia, Spain. *J Manag Care Pharm*. 2013;19(3):247–257.
4. Kripalani S, Roumie CL, Dalal, et al; PILL-CVD (Pharmacist Intervention for Low Literacy in Cardiovascular Disease) Study Group. Effect of a pharmacist intervention on clinically important medication errors after hospital discharge: a randomized trial. *Ann Intern Med*. 2012;157(1):1–10.
5. Wolf MS, Curtis LM, Wilson EA, et al. Literacy, cognitive function, and health: results of the LitCog study. *J Gen Intern Med*. 2012; 27(10):1300–1307.
6. Murray MD, Wu J, Tu W, et al. Health literacy predicts medication adherence. *Clin Pharmacol Ther*. 2004;75:P76.
7. Noureldin M, Plake KS, Morrow DG, Tu W, Wu J, Murray MD. Effect of health literacy on drug adherence in patients with heart failure. *Pharmacotherapy*. 2012;32(9):819–826.
8. Davis TC, Wolf MS, Bass PF, et al. Literacy and misunderstanding prescription drug labels. *Ann Intern Med*. 2006;145(12):887–894.
9. Ho PM, Lambert-Kerzner A, Carey EP, et al. Multifaceted intervention to improve medication adherence and secondary prevention measures after acute coronary syndrome hospital discharge: a randomized clinical trial. *JAMA Intern Med*. 2014;174(2):186–193.
10. Lambert-Kerzner A, Del Gaudio EJ, Fahdi IE, et al; Multifaceted Intervention to Improve Cardiac Medication Adherence and Secondary Prevention Measures (Medication) Study Investigators. Patient-centered adherence intervention after acute coronary syndrome hospitalization. *Circ Cardiovasc Qual Outcomes*. 2012;5(4):571–576.
11. Tariq SH, Tumosa N, Chibnall JT, Perry MH, Morley JE. Comparison of the Saint Louis University mental status examination and the mini-mental state examination for detecting dementia and mild neurocognitive disorder – a pilot study. *Am J Geriatr Psychiatry*. 2006; 14(11):900–910.
12. Cummings-Vaughn LA, Chavakula NN, Malmstrom TK, Tumosa N, Morley JE, Cruz-Oliver DM. Veterans Affairs Saint Louis University Mental Status examination compared with the Montreal Cognitive Assessment and the Short Test of Mental Status. *J Am Geriatr Soc*. 2014;62(7):1341–1346.
13. Bass PF, Wilson JF, Griffith CH. A shortened instrument for literacy screening. *J Gen Intern Med*. 2003;18(12):1036–1038.
14. Lin CF, Wu FL, Lin SW, et al. Age, dementia and care patterns after admission for acute coronary syndrome: an analysis from a nationwide cohort under the National Health Insurance coverage. *Drugs Aging*. 2012;29(10):819–828.
15. Ferrari E, Benhamou M, Cerboni P, Marcel B. Coronary syndromes following aspirin withdrawal: a special risk for late stent thrombosis. *J Am Coll Cardiol*. 2005;45(3):456–459.
16. Ho PM, Spertus JA, Masoudi FA, et al. Impact of medication therapy discontinuation on mortality after myocardial infarction. *Arch Intern Med*. 2006;166(17):1842–1847.
17. Rasmussen JN, Chong A, Alter DA. Relationship between adherence to evidence-based pharmacotherapy and long-term mortality after acute myocardial infarction. *JAMA*. 2007;297(2):177–186.
18. Spertus JA, Kettelkamp R, Vance C, et al. Prevalence, predictors, and outcomes of premature discontinuation of thienopyridine therapy after drug-eluting stent placement: results from the PREMIER registry. *Circulation*. 2006;113(24):2803–2809.

19. Sanchis J, Núñez J, Bodí V, et al. Influence of comorbid conditions on one-year outcomes in non-ST-segment elevation acute coronary syndrome. *Mayo Clin Proc.* 2011;86(4):291–296.
20. Wu JR, Holmes GM, DeWalt DA, et al. Low literacy is associated with increased risk of hospitalization and death among individuals with heart failure. *J Gen Intern Med.* 2013;28(9):1174–1180.
21. Crowley MJ, Zullig LL, Shah BR, et al. Medication non-adherence after myocardial infarction: an exploration of modifying factors. *J Gen Intern Med.* 2015;30(1):83–90.
22. Ho PM, Bryson CL, Rumsfeld JS. Medication adherence: its importance in cardiovascular outcomes. *Circulation.* 2009;119(23):3028–3035.

Patient Preference and Adherence

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress