

The value of trabectedin in the treatment of soft tissue sarcoma

Tomoki Nakamura
Akihiko Matsumine
Akihiro Sudo

Department of Orthopaedic Surgery,
Mie University Graduate School of
Medicine, Mie, Japan

Abstract: Soft tissue sarcomas (STSs) are a group of rare tumors accounting for less than 1% of all adult malignant tumors, a heterogeneous group of more than 50 histological subtypes. Five percent to 30% of STS patients experience local recurrence and 10%–38% present with clinically detectable metastases. Doxorubicin either alone or in combination with ifosfamide has been used as first-line chemotherapy for advanced disease. After failure of first-line chemotherapy, high-dose ifosfamide, gemcitabine + docetaxel, and dacarbazine may be applicable, although high-level evidence is lacking. Trabectedin is a synthetic, marine-derived alkylating agent derived from the Caribbean tunicate, *Ecteinascidia turbinata*. Several clinical trials have shown that trabectedin has a favorable toxicity profile and is an alternative therapeutic option in adult patients with advanced STS who have not responded to treatment with doxorubicin and ifosfamide. Several clinical trials also recommend the 24-hour intravenous infusion every 3 weeks regimen. The most frequently reported grade 3/4 adverse events were neutropenia and elevated serum levels of AST/ALT. Steroid pretreatment is an effective way of reducing the extent of hepatotoxicity, and steroids are now given routinely before trabectedin administration. Further studies are ongoing to evaluate the efficacy and safety of combination therapy of trabectedin with other agents.

Keywords: trabectedin, soft tissue sarcoma, efficacy, safety

Introduction

Soft tissue sarcomas (STSs) are a group of rare solid tumors accounting for less than 1% of all adult malignant tumors and 4%–8% of childhood malignancies, a heterogeneous group of more than 50 histological subtypes.^{1–5} Surgical resection with or without adjuvant radiotherapy is the standard treatment of all patients with an adult type, localized STS.^{4–9} Adjuvant chemotherapy may additionally be considered in the localized setting, however, its role in management remains controversial.⁴ Five percent to 30% of STS patients experience local recurrence and 10%–38% present with clinically detectable metastases.^{9–13} Complete surgical resection is reportedly a pivotal therapeutic option to provide patients with prolonged survival.^{4,14} However, even after a seemingly complete resection of metastatic tumors, metastasis will recur in 40%–80% of the patients.¹⁵ Systemic therapy for advanced disease is another therapeutic option in the management of metastases.^{4,16–21} Over the previous decades, doxorubicin either alone or in combination with ifosfamide has been used as first-line chemotherapy, however, it has not greatly improved the patient's prognosis.^{4,16–21} Although standard chemotherapy was performed, the median survival time is 8 to 13 months from initiation of first-line chemotherapy.^{17–21} After failure of standard chemotherapy, high-dose ifosfamide,^{22,23} gemcitabine + docetaxel,²⁴ and dacarbazine²⁵ may be applicable, although high-level evidence is lacking. Recently a randomized

Correspondence: Tomoki Nakamura
Department of Orthopaedic Surgery,
Mie University Graduate School of
Medicine, 2-174 Edobashi, Tsu-city,
Mie, Japan
Tel +81 5 9231 5022
Fax +81 5 9231 5211
Email tomoki66@clin.medic.mie-u.ac.jp

Phase III trial (PALETTE) of pazopanib treatment for advanced STS after failure of conventional chemotherapy demonstrated the potential benefit of pazopanib for a prolonged progression-free survival (PFS).²⁶ Trabectedin is a synthetic, marine-derived alkylating agent derived from the Caribbean tunicate, *Ecteinascidia turbinata*.^{27,28} The success of trabectedin in preliminary clinical trials for STSs has led to the approval of the drug in European countries in 2007 for the treatment of patients with advanced STS after the failure of therapy with doxorubicin either alone or in combination with ifosfamide. With limited systemic therapy options available as a whole, trabectedin has the opportunity to be significantly beneficial for patients with STSs. The purpose of this review is to summarize the efficacy and toxicity of trabectedin in the treatment of STS.

Each author certifies that his institution has approved the human protocol for this investigation and that all investigations were conducted in conformity with the ethical principles of research.

Mechanism of trabectedin

Trabectedin is a tetrahydroisoquinoline alkaloid derived from the Caribbean marine tunicate, *Ecteinascidia turbinata*, and is currently produced synthetically.^{27,28} Trabectedin interacts with the minor groove of DNA double helix and alkylates guanine at the N2 position, which bends toward the major groove,^{29–31} triggering a cascade of events that interferes with several transcription factors, DNA binding proteins, and DNA repair pathways, resulting in G2 – M cell cycle arrest and ultimately apoptosis.²⁹ Furthermore, the pattern of sensitivity observed in cells deficient in DNA repair mechanisms is different. In the case of trabectedin, nucleotide excisional repair and homologous recombination repair are of particular importance.^{29,32–34} In contrast to other DNA-damaging agents such as cisplatin, nucleotide excisional repair-deficient cells are two to ten times less sensitive to trabectedin.³⁵ On the other hand, cells deficient in homologous recombination repair are sensitive to trabectedin. Nevertheless, emerging evidence indicates that trabectedin has pleiotropic mechanisms of action, since, in addition to direct growth inhibition, cell death, and differentiation of malignant cells, trabectedin at therapeutic concentrations has selective immunomodulatory properties as a result of the inhibition of the production of factors that promote tumor growth, progression and inhibition of tumor-promoted angiogenesis. Trabectedin selectively targets monocytes and tumor associated macrophages and downregulates the production of inflammatory mediators such as IL-6 and CCL2, which may

underlie the strong association between chronic inflammation and cancer progression.^{31,36}

Trabectedin also has a specific mechanism against some translocation-related sarcomas. Recent investigations have demonstrated that trabectedin blocks the trans-activating ability of chimaeras by displacing the oncogenic fusion protein FUS-CHOP from its target promoters.^{37,38} This eventually induces adipogenic differentiation of myxoid liposarcoma cells. This effect has been observed in FUS-CHOP expressing experiment models and tumor biopsies taken before and after trabectedin therapy in patients with myxoid liposarcoma.^{37,39}

Efficacy of trabectedin for advanced STSs

Two Phase II trials in 2004 provided the initial analysis of trabectedin in STSs (Table 1).^{40,41} Trabectedin was administered at a dose of 1.5 mg/m², 24-hour intravenous (IV) infusion every 3 weeks (q3ws). The first of these studies was conducted in 54 advanced or metastatic STS patients with disease progression despite prior chemotherapy.⁴⁰ The objective response rate was 4%, however, the disease control rate at 6 months was 24%. The median PFS and overall survival (OS) were 1.9 months and 12.8 months, respectively. The second Phase II trials reported a low response rate of 8% in 36 recurrent or metastatic STS patients with disease progression despite prior chemotherapy.⁴¹ The median PFS and OS were 1.7 months and 12.1 months, respectively. In a small series of chemotherapy naïve patients, a 17.1% response rate with a 16.5-month median PFS was reported. Recently, results from randomized, multicenter, prospective dose-selection Phase IIb trials to evaluate whether trabectedin as first-line chemotherapy for advanced/metastatic STS prolongs the PFS, compared to doxorubicin, were published.⁴² One hundred and thirty-three patients were randomized (1:1:1) to doxorubicin, trabectedin (3-hour [T3h arm] infusion q3ws), or trabectedin (24-hour [T24h arm] infusion q3ws). The median PFS was 2.8 months in the T3h arm, 3.1 months in the T24h arm, and 5.5 months in the doxorubicin arm. No significant improvement in the PFS was observed in the trabectedin arms as compared to the doxorubicin arm (T24h vs doxorubicin: hazard ratio [HR] 1.13; 95% confidence interval [CI] 0.67–1.90, *P*=0.675; T3h vs doxorubicin: HR 1.50, 95% CI 0.91–2.48, *P*=0.944). The authors concluded that doxorubicin continues to be the standard treatment in eligible patients with advanced/metastatic STS as first-line treatment.

The efficacy of trabectedin 1.5 mg/m² 24-hour IV infusion q3ws in patients with pretreated advanced or metastatic

Table I Clinical trials of trabectedin (Phase II and III)

Study design	Patients	Regimen	ORR	Median	
				PFS	OS
Phase II ⁴⁴	LPS or LMS	A; T 1.5 mg/m ² 24-h q3ws	5.6%	3.7 mos	13.9 mos
R (1:1)	n=270	B; T 0.58 mg/m ² 3-h weekly	1.6%	2.3 mos	11.8 mos
Phase II ⁴¹	STS	T 1.5 mg/m ² 24-h q3ws	8%	1.7 mos	12.1 mos
	n=36				
Phase II ⁶⁵	STS	T 1.5 mg/m ² 24-h q3ws	8%	3.4 mos	9.2 mos
	n=104				
Phase II ⁴²	STS	A; T 1.3 mg/m ² 3-h q3ws	14.8%	2.8 mos	
R (1:1:1)	n=133	B; T 1.5 mg/m ² 24-h q3ws	4.7%	3.1 mos	
		C; D 75 mg/m ² q3ws	25.6%	5.5 mos	
Phase II ⁴⁰	STS	T 1.5 mg/m ² 24-h q3ws	4%	1.9 mos	12.8 mos
	n=54				
Phase III ⁴⁶	LPS or LMS	A; T 1.5 mg/m ² 24-h q3ws	9.9%	4.2 mos	12.4 mos
	n=518	B; Dac 1 g/m ² 20- to 120-minute q3ws	6.9%	1.5 mos	12.9 mos
Phase II ⁴⁹	T-STs	A; T 1.2 mg/m ² 24-h q3ws	11%	5.6 mos	Not reached
R (1:1)	n=76	B; Best supportive care	0%	0.9 mos	8 mos
Phase III ⁵⁵	T-STs	A; T 1.5 mg/m ² 24-h q3ws	5.9%	16.1 mos	
R (1:1)	n=121	B; D 75 mg/m ² q3ws or D 60 mg/m ² + I 6–9 g/m ² q3ws	27%	8.8 mos	
Phase II ⁵³	STS	1.5 mg/m ² 24-h q3ws	2.5%		
	n=40				

Abbreviations: R, randomized study; STS, soft tissue sarcoma; A, group A; B, group B; C, group C; T, trabectedin; D, doxorubicin; I, ifosfamide; Dac, dacarbazine; q3ws, every 3 weeks; h, hour; ORR, objective response rate according to Response Evaluation Criteria In Solid Tumors criteria; PFS, progression-free survival; OS, overall survival; mos, months; LPS, liposarcoma; LMS, leiomyosarcoma; T-STs, translocation-related soft tissue sarcoma.

STS was previously evaluated in non-randomized Phase II studies.^{40–42} A weekly trabectedin schedule (0.58 mg/m² 3-hour IV infusion for 3 consecutive weeks in a 4-week cycle) was demonstrated to have substantial anticancer activity in pre-treated ovarian cancer.⁴³ To assess the efficacy and safety of these two schedules in STS, a randomized, open-label, Phase II trial was conducted in patients with advanced and/or metastatic liposarcomas or leiomyosarcomas after the failure of standard therapies.⁴⁴ The time to progression was the primary endpoint. The 24-hour IV q3ws demonstrated a superior time to progression of 3.7 months vs 2.3 months (HR, 0.734; 95% CI, 0.554–0.974; $P=0.0302$). The median PFS was 3.3 months vs 2.3 months (HR, 0.755; 95% CI, 0.574–0.992; $P=0.0418$). The median OS was 13.9 months vs 11.8 months (HR, 0.843; 95% CI, 0.653–1.090; $P=0.1920$). The authors concluded that these data recommend 24-hour IV q3ws regimen. Recently, other Phase II trials showed a similar conclusion that the 1.5 mg/m²/24-hour IV infusion is the optimal approach to delivering trabectedin in the second-line setting.

Due to treatment-related toxicities, most sarcoma experts often propose a drug holiday followed by re-challenge at progression. However, the effect of this approach remains unclear, because no evidence has indicated the absence of deleterious effects of trabectedin discontinuation in non-progressive

patients. Therefore, to elucidate the benefit or harm of trabectedin discontinuation in patients with non-progression, a Phase II trial investigating the clinical benefit of continuation of trabectedin treatment (1.5 mg/m² 24-hour IV infusion q3ws) until progression vs the interruption of therapy after six cycles in patients with advanced STS was performed.⁴⁵ Fifty-three patients were randomly assigned to two groups: 27 to the continuation group and 26 to the interruption group. After randomization, the PFS at 6 months was 51.9% (95% CI, 31.9–68.6) in the continuation group vs 23.1% (95% CI, 9.4–40.3) in the interruption group ($P=0.02$). The authors concluded that trabectedin discontinuation in patients with non-progressive disease is not recommended. Recently, trabectedin has been approved by the US Food and Drug Administration (FDA) based on the result of an open-label, randomized (2:1) Phase III trial of trabectedin (n=345) vs dacarbazine (n=173) in patients with metastatic liposarcoma or leiomyosarcoma.⁴⁶ In the final analysis of PFS, trabectedin administration resulted in a 45% reduction in the risk of disease progression or death compared with dacarbazine. The median PFS was 4.2 months vs 1.5 months (HR, 0.55; 95% CI, 0.44–0.70; $P<0.001$).

It has been reported that translocation-related sarcoma responds well to trabectedin treatment. Trabectedin has shown particular activity in myxoid liposarcoma. In a multicenter,

retrospective analysis of 51 patients, a median overall response rate of 51% was reported, and the median PFS was 14 months.⁴⁷ The PFS at 6 months was also reported to be 88%. In another study assessing the efficacy of trabectedin in specific translocation-related sarcomas, the PFS at 6 months was 64%.⁴⁸ A recent Phase II study in the second-line setting or later has been reported.⁴⁹ This study was a randomized Phase II study of trabectedin monotherapy vs supportive care in patients with translocation-related sarcoma subtypes. The patients were randomized (1:1) to receive trabectedin (1.2 mg/m² 24-hour infusion q3ws) or best supportive care. The trabectedin dose of this trial was 1.2 mg/m² according to the results of a Phase I study in Japanese patients with STSs, in which two of three patients had dose-limiting toxicity at 1.5 mg/m².⁵⁰ The primary endpoint of this trial was the PFS. The median PFS of the trabectedin group was 5.6 months and that of the best supportive care group was 0.9 months (HR, 0.07; 95% CI, 0.03–0.16; $P < 0.0001$). The authors concluded that trabectedin could present a novel treatment option for patients with translocation-related sarcoma who did not respond to doxorubicin-based chemotherapy. The success of trabectedin in this clinical trial for STSs has led to the approval of the drug in Japan. Modulation of transcription by trabectedin has also been shown in Ewing's sarcoma, which is driven by the oncogenic fusion gene *EWS-FLII*.^{51,52} However, a Phase II trial in children with recurrent STS, including Ewing's sarcoma, showed that trabectedin did not demonstrate sufficient activity as a single agent.⁵³ Of eleven patients with Ewing's sarcoma in the Phase II component, only one patient achieved stable disease (SD) after four cycles. The other nine patients had progressive disease.⁵³ Recently, it has been shown that this modulation can be exploited to select a very effective combination treatment of trabectedin followed by irinotecan.⁵⁴ An observational study of combination treatment of trabectedin followed by irinotecan for refractory pediatric sarcoma is currently ongoing ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02509234) Identifier; NCT 02509234). The purpose of this study is to determine if this is a promising treatment option with acceptable toxicity and if the results warrant a prospective study.

One Phase III study in the first-line setting has been reported.⁵⁵ This study was a randomized, Phase III study of first-line trabectedin vs doxorubicin-based chemotherapy in patients with translocation-related sarcoma subtypes.⁵⁵ The primary endpoint of this trial was the PFS. Patients were randomized (1:1) to receive trabectedin (1.5 mg/m² 24-hour IV infusion q3ws), doxorubicin 75 mg/m² IV q3ws, or doxorubicin 60 mg/m² IV plus ifosfamide (range, 6–9 g/m²)

IV q3ws. There was no difference in the median PFS or OS between the groups ($P = 0.9573$ and $P = 0.3659$, respectively). The response rate according to the RECIST (Response Evaluation Criteria In Solid Tumors) criteria was significantly higher in the chemotherapy arm (27%) compared to the trabectedin arm (5.9%). In contrast, the response rate according to the Choi criteria showed fewer differences between the chemotherapy arm (45.9%) and trabectedin arm (37.3%).

Role of trabectedin for neoadjuvant therapy

One Phase II clinical trial in the neoadjuvant setting in patients with advanced localized myxoid liposarcoma has been previously reported.⁵⁶ The treatment consisted of trabectedin 1.5 mg/m² given as a 24-hour IV infusion q3ws. Twenty-nine patients received a minimum of three and a maximum of six cycles before surgery. Of 23 patients who could be evaluated by the pathological response, three patients achieved a pathological complete response. Another 12 of 23 had at least a good regression rate ($>50\%$ regression). Of 29 patients, seven patients (24%) had a partial response and 21 patients had SD according to the RECIST criteria. One patient died prior to the evaluation due to rhabdomyolysis with hepatic and renal failure after the second trabectedin cycle. The authors concluded that trabectedin is a therapeutic option in the neoadjuvant setting of myxoid liposarcoma. A Phase I–II study that combines trabectedin plus radiotherapy for patients with myxoid liposarcoma and metastatic STS is currently recruiting participants ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02275286) Identifier; NCT 02275286). The study is testing the hypothesis that administering trabectedin plus radiotherapy shows synergic activity that turns into tumor shrinkage. A Phase III study currently enrolling patients is comparing the impact of standard neoadjuvant chemotherapy (epirubicin and ifosfamide) with that of neoadjuvant therapy tailored to the specific histology on the disease-specific survival ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01710176) Identifier; NCT 01710176); patients with leiomyosarcoma or myxoid liposarcoma with round cell liposarcoma will receive gemcitabine plus dacarbazine or trabectedin, respectively.

Safety

Trabectedin was well tolerated in a retrospective pooled analysis of five Phase II trials.⁵⁷ The data collected from 350 adult patients were divided into the younger (<60 years; $n = 267$) and the older cohort (≥ 60 years; $n = 83$). The most frequently reported grade 3/4 adverse events were

neutropenia (47.6%) and elevated serum levels of AST/ALT (44.6%/34.4%). Some grade 3/4 adverse events were more common in patients aged ≥ 60 years compared to patients < 60 years, namely anemia 10.1% vs 19.3%, neutropenia 43.6% vs 60.2%, thrombocytopenia 11.3% vs 20.5%, and fatigue 6.4% vs 14.5%, respectively. Deaths associated with drug-related adverse events were infrequent (1.9% and 2.4% of the patients in the younger and older cohorts, respectively). This pooled analysis also showed similar response rates (10.1% and 9.6% of the patients in the younger and older cohorts, respectively), no significant difference in the median PFS (2.5 months vs 3.7 months, respectively) and similar OS rates between the two cohorts (13.0 months vs 14.0 months, respectively). Other trials showed a similar rate of severe neutropenia, while the standard first-line agents in STSs, doxorubicin and ifosfamide, induced substantially more severe hematological toxicities. Randomized trials with doxorubicin showed grade 3/4 neutropenia in 77% of the patients, with 16%–19% febrile neutropenia, and high-dose ifosfamide caused neutropenia and infection in 20% and 4% of the patients, respectively.^{19,44,58} Transaminase increase was the most frequent cause of dose reductions. However, the occurrence of hepatobiliary adverse events was low, further supporting the observation that the liver dysfunction induced by trabectedin is mostly transient, non-cumulative, and without clinical consequences.^{44,57,59} A recurring pattern was observed with increased transaminase levels, typically reaching a peak between days 5 and 7 of each cycle and resolving to grade ≤ 1 by day 15 without implication for the patient.³⁵ Steroid pretreatment is an effective way of reducing the extent of hepatotoxicity, and steroids are now given routinely before trabectedin administration. Premedication with 20 mg of dexamethasone IV 30 minutes prior to trabectedin was shown to provide hepatoprotective effects beyond its antiemetic effect.^{59–61} Although extremely rare, trabectedin-associated rhabdomyolysis is considered to be a potential adverse event. The incidence appears to be approximately 0.5% in patients with sarcoma and ovarian cancer.⁶² Elevations in creatine kinase are observed, however, the incidence of muscle damage is low. No predictive factors have been identified to prospectively identify patients who may be at risk of rhabdomyolysis, and therefore, careful clinical and biochemical monitoring is mandatory at each cycle.

Combination therapy of trabectedin with other agents

Combination therapy of trabectedin with other chemotherapeutic agents has been assessed in Phase I trials. The most

promising results in sarcoma patients have been from the trials administering trabectedin in combination with doxorubicin.^{63,64} One of the trials showed a response rate of 18% and SD in 56% of 29 patients with STS. Another study assessing combination therapy with doxorubicin reported a response rate of 12% with a median PFS of 9.2 months for 41 patients, including 20 liposarcomas and eleven leiomyosarcomas.

Conclusion

Trabectedin has shown a favorable toxicity profile and is an alternative therapeutic option in adult patients with advanced STS who have not responded to treatment with doxorubicin and ifosfamide. Several clinical trials recommend the 24-hour IV q3ws regimen. Furthermore, trabectedin discontinuation in patients with non-progressive disease is not recommended. Steroid pretreatment is an effective way of reducing the extent of hepatotoxicity, and steroids are now given routinely before trabectedin administration. Further study is ongoing to evaluate the efficacy and safety of combination therapy of trabectedin with other agents.

Disclosure

Each author certifies that he has no commercial associations (eg, consultancies, stock ownership, equity interest, patent/licensing arrangements) that might pose a conflict of interest in connection with the submitted article.

References

1. Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin*. 2014;64(1):9–29.
2. Clark MA, Fisher C, Judson I, Thomas JM. Soft-tissue sarcomas in adults. *N Engl J Med*. 2005;353(7):701–711.
3. Le Cense A, Ray-Coquard I, Duffaud F, et al. Trabectedin in patients with advanced soft tissue sarcoma: A retrospective national analysis of the French Sarcoma Group. *Eur J Cancer*. 2015;51(6):742–750.
4. Casali PG, Blay JY, Bertuzzi A, et al. Soft tissue and visceral sarcomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2014;25 Suppl 3:iii102–iii112.
5. Fletcher CD, Bridge JA, Hogendoorn PC, Mertens F, editors. *WHO Classification of Tumours of Soft Tissue and Bone*. Lyon: IARC; 2013.
6. Yang JC, Chang AE, Baker AR, et al. Randomized prospective study of the benefit of adjuvant radiation therapy in the treatment of soft tissue sarcomas of the extremity. *J Clin Oncol*. 1998;16(1):197–203.
7. Beane JD, Yang JC, White D, et al. Efficacy of adjuvant radiation therapy in the treatment of soft tissue sarcoma of the extremity: 20-year follow-up of a randomized prospective trial. *Ann Surg Oncol*. 2014;21(8):2484–2489.
8. Cahlon O, Brennan MF, Jia X, et al. A postoperative nomogram for local recurrence risk in extremity soft tissue sarcomas after limb-sparing surgery without adjuvant radiation. *Ann Surg*. 2012;255(2):343–347.
9. Zager GK, Ballo MT, Pesters PW, et al. Prognostic factors for patients with localized soft-tissue sarcoma treated with conservation surgery and radiation therapy; an analysis of 1225 patients. *Cancer*. 2003;97(10):2530–2543.

10. Nakamura T, Matsumine A, Matsubara T, et al. Restrospective analysis of metastatic sarcoma patients. *Oncol Lett.* 2011;2(2):315–318.
11. Weiss SW, Goldblum JR. Local recurrence. In: Weiss SW, Goldblum JR. *Enzinger and Weiss's Soft Tissue Tumors*. 4th ed. Mosby, St Louis; 2001:28–32.
12. Kane JM, Finley JW, Driscoll D, Kraybill WG, Gibbs JF. The treatment and outcome of patients with soft tissue sarcomas and synchronous metastases. *Sarcoma.* 2002;6(2):69–73.
13. Pollock RE, Karnell LH, Menck HR, Winchester DP. The national cancer data base report on soft tissue sarcoma. *Cancer.* 1996;87(10):2247–2257.
14. Blackmon SH, Shah N, Roth JA, et al. Resection of pulmonary and extrapulmonary sarcomatous metastases is associated with long-term survival. *Ann Thorac Surg.* 2009;88(3):877–884.
15. Weiser MR, Downey RJ, Leung HY, Brennan MF. Repeat resection of pulmonary metastases in patients with soft-tissue sarcoma. *J Am Coll Surg.* 2000;191(2):184–190.
16. Judson I, Verweij J, Gelderblom H, et al. Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft tissue sarcoma: a randomised controlled phase 3 trial. *Lancet Oncol.* 2014;15(4):415–423.
17. Bramwell VH, Anderson D, Charette ML; Sarcoma Disease Site Group. Doxorubicin-based chemotherapy for the palliative treatment of adult patients with locally advanced or metastatic soft tissue sarcoma. *Cochrane Database Syst Rev.* 2003;(3):CD003293.
18. Antman K, Crowley J, Balcerzak SP, et al. An intergroup phase III randomized study of doxorubicin and dacarbazine with or without ifosfamide and mesna in advanced soft tissue and bone sarcomas. *J Clin Oncol.* 1993;11(7):1276–1285.
19. Le Cense, Judson I, Crowther D, et al. Randomized phase III study comparing conventional-dose doxorubicin plus ifosfamide versus high-dose doxorubicin plus ifosfamide plus recombinant human granulocyte-macrophage colony-stimulating factor in advanced soft tissue sarcomas: A trial of the European Organization for Research and Treatment of Cancer/Soft Tissue and Bone Sarcoma Group. *J Clin Oncol.* 2000;18(14):2676–2684.
20. Jelic S, Kovcin V, Milanovic N, et al. Randomized study of high-dose epirubicin versus high-dose epirubicin-cisplatin chemotherapy for advanced soft tissue sarcoma. *Eur J Cancer.* 1997;33(2):220–225.
21. van Glabbeke M, van Oosterom AT, Oosterhuis JW, et al. Prognostic factors for outcome of chemotherapy in advanced soft tissue sarcoma: an analysis of 2,185 patients treated with anthracycline-containing first-line regimens – a European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group Study. *J Clin Oncol.* 1999;17(1):150–157.
22. Le Cesne A, Antoine E, Spielmann M, et al. High-dose ifosfamide: circumvention of resistance to standard-dose ifosfamide in advanced soft tissue sarcomas. *J Clin Oncol.* 1995;13(7):1600–1608.
23. Martin-Liberal J, Alam S, Constantinidou A, et al. Clinical activity and tolerability of a 14-day infusional Ifosfamide schedule in soft-tissue sarcoma. *Sarcoma.* 2013;2013:868973.
24. Maki RG, Wathen JK, Patel SR, et al. Randomized phase II study of gemcitabine and docetaxel compared with gemcitabine alone in patients with metastatic soft tissue sarcomas: results of sarcoma alliance for research through collaboration study 002. *J Clin Oncol.* 2007;25(19):2755–2763.
25. García-Del-Muro X, López-Pousa A, Maurel J, et al. Randomized phase II study comparing gemcitabine plus dacarbazine versus dacarbazine alone in patients with previously treated soft tissue sarcoma: a Spanish Group for Research on Sarcomas study. *J Clin Oncol.* 2011;29(18):2528–2533.
26. Van der Graaf WT, Blay JY, Chawla SP, et al. Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomized, double-blind, placebo-controlled phase 3 trial. *Lancet.* 2012;379(9829):1879–1886.
27. Carter NJ, Keam SJ. Trabectedin: A review of its use in the management of soft tissue sarcoma and ovarian cancer. *Drugs.* 2007;67(15):2257–2276.
28. Cuaves C, Francesch A. Development of Yondelis (trabectedin, ET-743). A semisynthetic process solves the supply problem. *Nat Prod Rep.* 2009;26(3):322–337.
29. D'Incalci M, Galmarini CM. A review of trabectedin (et-743): A unique mechanism of action. *Mol Cancer Ther.* 2010;9(8):2157–2163.
30. Feuerhahn S, Giraudon C, Martinez-Diez M, et al. XPF-dependent DNA breaks and RNA polymerase II arrest induced by antitumor DNA inter-strand crosslinking-mimetic alkaloids. *Chem Biol.* 2011;18(8):988–999.
31. D'Incalci M, Badri N, Galmarini CM, Allavena P. Trabectedin, a drug acting on both cancer cells and the tumor microenvironment. *Br J Cancer.* 2014;111(4):646–650.
32. Scoffski P, Casali M, Taron M, et al. DNA repair functionality modulates the clinical outcome of patients with advanced sarcoma treated with trabectedin (ET-743). *J Clin Oncol.* 2006;24 Suppl 18:9522.
33. Soares DG, Escargueil AE, Poindessous V, et al. Replication and homologous recombination repair regulate DNA double-strand break formation by antitumor alkylator ecteinascidin 743. *Proc Natl Acad Sci U S A.* 2007;104(32):13062–13067.
34. Laroche-Clay A, Chaire V, Le Morvan V, et al. BRCA1 haplotype and clinical benefit of trabectedin in soft-tissue sarcoma patients. *Br J Cancer.* 2015;112(4):688–692.
35. Brodowicz T. Trabectedin in soft tissue sarcomas. *Future Oncol.* 2014;10(8 Suppl):s1–s5.
36. Germano G, Frapolli R, Belgiovine C, et al. Role of macrophage targeting in the antitumor activity of trabectedin. *Cancer Cell.* 2013;23(2):249–262.
37. Di Giandomenico S, Frapolli R, Bello E, et al. Mode of action of trabectedin in myxoid liposarcomas. *Oncogene.* 2014;33(44):5201–5210.
38. Forni C, Minuzzo M, Virdis E, et al. Trabectedin (ET-743) promotes differentiation in myxoid liposarcoma tumors. *Mol Cancer Ther.* 2009;8(2):449–457.
39. Charytonowicz E, Terry M, Coakley K, et al. PPAR γ agonist enhance ET-743-induced adipogenic differentiation in a transgenic mouse model of myxoid round cell liposarcoma. *J Clin Invest.* 2012;122(3): 886–898.
40. Yovine A, Riofrio M, Blay JY, et al. Phase II study of ecteinascidin-743 in advanced pretreated soft tissue sarcoma patients. *J Clin Oncol.* 2004;22(5):890–899.
41. García-Carbonero R, Supko JG, Manola J, et al. Phase II and pharmacokinetic study of ecteinascidin 743 in patients with progressive sarcomas of soft tissue refractory to chemotherapy. *J Clin Oncol.* 2004;22(8):1480–1490.
42. Bui-Nguyen B, Butrynski JE, Penel N, et al. A phase II multicentre study comparing the efficacy of trabectedin to doxorubicin in patients with advanced or metastatic untreated soft tissue sarcoma: the TRUSTS trial. *Eur J Cancer.* 2015;51(10):1312–1320.
43. Krasner CN, McMeekin DS, Chan S, et al. A phase II study of trabectedin single agent in patients with recurrent ovarian cancer previously treated with platinum-based regimens. *Br J Cancer.* 2007;97(12):1618–1624.
44. Demetri GD, Chawla SP, von Mehren M, et al. Efficacy and safety of trabectedin in patients with advanced or metastatic liposarcoma or leiomyosarcoma after failure of prior anthracyclines and ifosfamide: Results of a randomized phase II study of two different schedules. *J Clin Oncol.* 2009;27(25):4188–4196.
45. Le Cense A, Blay JY, Domont J, et al. Interruption versus continuation of trabectedin in patients with soft tissue sarcoma (T-DIS): a randomized phase 2 trial. *Lancet Oncol.* 2015;16(3):312–319.
46. Demetri GD, von Mehren M, Jones RL, et al. Efficacy and safety of trabectedin or dacarbazine for metastatic liposarcoma or leiomyosarcoma after failure of conventional chemotherapy: Results of a phase III randomized multicenter clinical trial. *J Clin Oncol.* Epub 2015 Sep 14.
47. Grosso F, Jones RL, Demetri GD, et al. Efficacy of trabectedin (ecteinascidin-743) in advanced pretreated myxoid liposarcomas: a retrospective study. *Lancet Oncol.* 2007;8(7):595–602.
48. Le Cesne A, Cresta S, Maki RG, et al. A retrospective analysis of antitumor activity with trabectedin in translocation-related sarcomas. *Eur J Cancer.* 2012;48(16):3036–3044.

49. Kawai A, Araki N, Sugiura H, et al. Trabectedin monotherapy after standard chemotherapy versus best supportive care in patients with advanced, translocation-related sarcoma: a randomized, open-label, phase 2 study. *Lancet Oncol*. 2015;16(4):406–416.
50. Ueda T, Kakunaga S, Ando M, et al. Phase I study and pharmacokinetic study of trabectedin, a DNA minor groove binder, administered as a 24-h continuous infusion in Japanese patients with soft tissue sarcoma. *Invest New Drugs*. 2014;32(4):691–699.
51. Grohar PJ, Griffin LB, Yeung C, et al. Ecteinascidin 743 interferes with the activity of EWS-FLI1 in Ewing sarcoma cells. *Neoplasia*. 2011;13(2):145–153.
52. Grohar PJ, Helman LJ. Prospects and challenges for the development of new therapies for Ewing sarcoma. *Pharmacol Ther*. 2013;137(2):216–224.
53. Baruchel S, Pappo A, Krailo M, et al. A phase 2 trial of trabectedin in children with recurrent rhabdomyosarcoma, Ewing sarcoma and non-rhabdomyosarcoma soft tissue sarcomas: A report from the Children's Oncology Group. *Eur J Cancer*. 2012;48(4):579–585.
54. Grohar PJ, Segars LE, Yeung C, et al. Dual targeting of EWS-FLI1 activity and the associated DNA damage response with Trabectedin and SN38 synergistically inhibits Ewing sarcoma cell growth. *Clin Cancer Res*. 2013;20(5):1190–1203.
55. Blay JY, Leahy MG, Nguyen BB, et al. Randomized phase III trial of trabectedin versus doxorubicin-based chemotherapy as first-line therapy in translocation-related sarcomas. *Eur J Cancer*. 2014;50(6):1137–1147.
56. Gronchi A, Bui BN, Bonvalot S, et al. Phase II clinical trial of neoadjuvant trabectedin in patients with advanced localized myxoid liposarcoma. *Ann Oncol*. 2012;23(3):771–776.
57. Le Cense A, Judson I, Maki R, et al. Trabectedin is a feasible treatment for soft tissue sarcoma patients regardless of patient age: a retrospective pooled analysis of five phase II trials. *Br J Cancer*. 2013;109(7):1717–1724.
58. van Oosterom AT, Mouridsen HT, Nielsen OS et al. Results of randomized studies of the EORTC Soft Tissue and Bone Sarcoma Group (STBSG) with two different ifosfamide regimens in first- and second-line chemotherapy in advanced soft tissue sarcoma patients. *Eur J Cancer*. 2002;38(18):2397–2406.
59. Grosso F, Dileo P, Sanfilippo R, et al. Steroid premedication markedly reduces liver and bone marrow toxicity of trabectedin in advanced sarcoma. *Eur J Cancer*. 2006;42(10):1484–1490.
60. Cesne AL, Yovine A, Blay JY, et al. A retrospective pooled analysis of trabectedin safety in 1,132 patients with solid tumors treated in phase II clinical trials. *Invest New Drugs*. 2012;30(3):1193–1202.
61. Amart F, Lorusso D, Mustea A, Duffaud F, Pautier P. Management strategies in advanced uterine leiomyosarcoma: Focus on trabectedin. *Sarcoma*. 2015;2015:704124.
62. Grosso F, D'Incalci M. Problems in dealing with very rare adverse effects of new anticancer drugs: the example of trabectedin. *Tumori*. 2011;97(2):256–257.
63. Sessa C, Perotti A, Noberasco C, et al. Phase I clinical and pharmacokinetic study of trabectedin and doxorubicin in advanced soft tissue sarcoma and breast cancer. *Eur J Cancer*. 2009;45(7):1153–1161.
64. Blay JY, von Mehren M, Samuels BL, et al. Phase I combination study of trabectedin and doxorubicin in patients with soft-tissue sarcoma. *Clin Cancer Res*. 2008;14(20):6656–6662.
65. Le Cesne A, Blay JY, Judson I, et al. Phase II study of ET-743 in advanced soft tissue sarcomas: a European Organization for Research and Treatment of Cancer (EORTC) soft tissue and bone sarcoma group trial. *J Clin Oncol*. 2005;23(3):576–584.

Therapeutics and Clinical Risk Management

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peer-reviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS,

Submit your manuscript here: <http://www.dovepress.com/therapeutics-and-clinical-risk-management-journal>

EMBASE, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress