

Outcomes in adult pectus excavatum patients undergoing Nuss repair

MennatAllah M Ewais
Shivani Chaparala
Rebecca Uhl
Dawn E Jaroszewski

Department of Cardiothoracic
Surgery, Mayo Clinic Hospital,
Phoenix, AZ, USA

Abstract: Pectus excavatum (PEx) is one of the most common congenital chest wall deformities. Depending on the severity, presentation of PEx may range from minor cosmetic issues to disabling cardiopulmonary symptoms. The effect of PEx on adult patients has not been extensively studied. Symptoms may not occur until the patient ages, and they may worsen over the years. More recent publications have implied that PEx may have significant cardiopulmonary implications and repair is of medical benefit. Adults presenting for PEx repair can undergo a successful repair with a minimally invasive “Nuss” approach. Resolution of symptoms, improved quality of life, and satisfying results are reported.

Keywords: complications, minimally invasive surgery, quality of life

Background

Pectus excavatum (PEx) is the most common congenital chest wall anomaly.¹⁻⁴ The deformity is reported to occur more frequently in males than females; however, diagnosis in females may often be missed if obscured with breast tissue.⁵⁻⁷ Depending on the severity, presentation of PEx may range from a minor cosmetic issue to disabling cardiopulmonary symptoms.⁸⁻¹⁰ The internally displaced sternum can cause right-side heart compression and restrictive deficits.^{8,9} As the patient advances in age, the chest wall can become less flexible as a result of increased calcium accumulation in cartilage attachments of the anterior chest wall.^{11,12} Symptoms may occur or show progression as the patient ages.¹¹⁻¹³ Kragten et al¹² reported development of symptoms in the fourth and fifth decade in nearly half of their adult patients with significant improvement after surgical repair. The optimal surgical procedure for adult PEx patients has been controversial, and some surgeons recommend limiting the Nuss procedure to pediatrics and adolescents.^{14,15} We present a review of adult patients with PEx including outcomes after repair with Nuss (“Nuss”) or a minimally invasive repair of pectus excavatum (MIRPEx).

Cardiopulmonary outcomes

The cardiopulmonary effects of PEx have been debated for years,¹⁶⁻¹⁸ and there is a paucity of reports evaluating adult patients.^{19,20} The inward deformity of the anterior chest wall has a negative cardiopulmonary consequence on patients with PEx, as supported by the most recent data.^{9,21} This can cause displacement of the heart into the left chest and varying degrees of heart compression (Figure 1). Decrease in atrial filling and venous return can result in diastolic dysfunction and reduction in cardiac

Correspondence: Dawn E Jaroszewski
Department of Cardiothoracic Surgery,
Mayo Clinic Hospital, 5777 E Mayo Blvd,
Phoenix, AZ 85054, USA
Tel +1 480 342 2270
Fax +1 480 342 2269
Email jaroszewski.dawn@mayo.edu

output with significant compression to the chambers of the right heart.^{9,20–22} Mocchegiani et al²³ reported that the right ventricular outflow tract in PEx patients was significantly narrower and right ventricle (RV) end-diastolic and-systolic areas were significantly smaller. Surgical correction of the PEx has been shown to relieve compression, allowing for a significant increase in right heart chamber size, increased flow velocities, and improved cardiac output^{9,21,24} (Figure 2A and B). Krueger et al²⁵ also noted significant improvement in post-repair cardiac outputs that increased to 66.2% vs 58.4% and the end-diastolic RV volume that increased to a mean of 40.8 mL vs 21.7 mL preoperatively. There may be a greater impact on cardiac function and symptoms in patients over 30 years of age.^{12,24} In post-repair PEx patients ≥ 30 years, a mean increase in right ventricular output of 65% was documented




Figure 1 Computerized tomographic scan of a patient with severe pectus excavatum and Haller index of 24.6. Sternal deformity with compression of the right heart and inflow are seen (arrow).

by intraoperative transesophageal echocardiogram.²⁴ Nevire et al²⁶ found that PEx deformity was associated with reduction in the strength of the inspiratory muscle as evident by reduction in the maximal static respiratory pressure (P_Imax) and sniff nasal inspiratory pressure (SNIP) with significant increase of these values postoperatively which was reflected in enhanced efficacy of the respiratory pump and the cardiovascular function improvement.

Long-term follow-up of corrected PEx patients and correlations between physiologic impact and symptoms are lacking.^{8,27–30} Table 1 reviews some of the major publications reporting cardiopulmonary impairments and postsurgical results. Only six of these represented a mean age of 18 years and older.^{9,19,25,26,31,32} Cardiopulmonary exercise testing has been used to assess exercise capacity and limitations in peak oxygen uptake and O₂ pulse which can result as a consequence of the pectus deformity.^{18,19,33,34} Publications as to the benefits of surgical correction have varied.^{8,19,31,35} Several studies have reported that the cardiopulmonary function has improved significantly with increase in the oxygen consumption (VO₂) and O₂ pulse after surgical repair of PEx.^{8,19} Maagaard et al³⁵ previously reported normalization of the decreased cardiopulmonary function in teenagers with PEx at 3 years following surgical repair; however, in a more recent evaluation of adult patients by Udholm et al³¹ (≥ 21 years), a significant improvement in the maximum oxygen consumption (VO₂ max) was not seen 1 year after PEx repair. These results did show a trend of increased improvement in the VO₂ max which could be more evident with a longer period of follow-up. Adult patients may also differ in their ability to return to normal after PEx repair. During assessment, the patient's baseline exercise history must be considered.




Figure 2 Transesophageal echocardiographic images show preoperative effect (A) of pectus excavatum with compression on the right ventricle due to the inward sternal deformity and relief of the compression following surgical repair (B).

Abbreviations: RV, right ventricle; LV, left ventricle; RA, right atrium; LA, left atrium.

Table 1 Review of major publications reporting cardiopulmonary outcomes and postsurgical results

Study	No of patients	Age (years), mean \pm SD (range)	Haller index mean \pm SD	Surgical method	Variables studied	Cardiopulmonary outcomes
Udholm et al ³¹	19 (15 completed follow-up)	32	NR	MIRPEX	Heart rate at rest, maximum heart rate, maximum VO_2/kg , maximum cardiac output, and maximum cardiac index at baseline and 1 year postoperatively	<ul style="list-style-type: none"> No significant change in any of the variables after 1 year of follow-up. Cardiac output (14.0 ± 0.9 l/min at baseline vs 14.8 ± 1.1 l/min after surgery; $P = 0.2029$), maximum oxygen uptake (30.4 ± 1.9 and 33.3 ± 1.6 mL/kg/min; $P = 0.0940$ after surgery).
Topper et al ³²	38	21 ± 8.3 (12–43)	9.64	Modified Nuss	Heart rate, right and left ventricular ejection fraction, right and left ventricular end diastolic and end systolic volumes, and right and left ventricular stroke volumes before and after repair	<ul style="list-style-type: none"> There was a significant decrease in heart rate ($P < 0.0001$) and significant increase in RVEF ($P = 0.0004$), RVSV ($P = 0.0167$), LVEF ($P = 0.0165$), and in LVSV ($P = 0.0036$) after 1 year of repair. There was no significant increase in RVEDV ($P = 0.7590$), RVESV ($P = 0.0718$), LVEDV ($P = 0.0648$), and in LVESV ($P = 0.8135$).
Chao et al ⁹	168	33 (18–71)	5.7 ± 3.1	Modified Nuss	Right atrium, tricuspid annulus end systolic, right ventricular outflow tract end diastolic and end systolic dimensions, cardiac output, heart rate, systolic blood pressure, and stroke volume	<ul style="list-style-type: none"> There was a significant improvement in right atrium (15.1%), tricuspid annulus end systolic (10.9%), right ventricular outflow tract end diastolic (6.1%) and end systolic dimension (6.1%) size after surgery (all $P < 0.0001$). In a subset of 42 patients, there was a significant improvement in right ventricular cardiac output by 38% ($P < 0.0001$). No change in heart rate, systolic blood pressure, or stroke volume.
O'Keefe et al ³⁷	67	13.9 ± 2.3	4.4 ± 1.3	Nuss	Heart rate (% predicted), maximal oxygen consumption (VO_2 max), O_2 pulse, cardiac index, stroke volume, and cardiac output preoperatively and at 3 months after bar removal	<ul style="list-style-type: none"> O_2 pulse ($P = 0.01$), respiratory quotient ($P < 0.001$), and respiratory reserve ($P = 0.003$) were increased significantly after bar removal, with no significant changes in VO_2 max or heart rate and a decrease in minute volume of breathing.
Neviere et al ²⁶	20	32 ± 11	4.7 ± 1.4	Ravitch	Heart rate, peak VO_2 , peak oxygen pulse, respiratory exchange ratio before and at 1 year post-repair	<ul style="list-style-type: none"> Significant improvement in anaerobic threshold, peak VO_2 ($P < 0.01$), positive correlations were found between postoperative changes for oxygen pulse and PImax ($r^2 = 0.23$; $P = 0.04$) and for oxygen pulse and SNIP ($r^2 = 0.38$; $P < 0.01$). SNIP was the strongest independent predictor factor of oxygen pulse increase after PEx repair, and adding age, weight, Haller index, and pulmonary function test results to the regression equation did not increase the correlation coefficient ($r^2 = 0.378$).
Maagaard et al ³⁵	49 (42 PEx patients) and 24 controls completed follow-up	15.5 ± 1.7	4.9 ± 1.4	MIRPEX	Minimum and maximum heart rate, maximum VO_2/kg , maximum cardiac index, stroke index before repair and at 1 year and 3 years after bar removal between patients and healthy, age-matched controls	<ul style="list-style-type: none"> Preoperatively, patients had lower maximum cardiac index: mean \pm SD (6.6 ± 1.2 l \cdot min⁻¹ \cdot m⁻²) compared with controls (8.1 ± 1.0 l \cdot min⁻¹ \cdot m⁻²) during exercise ($P = 0.0001$). One year and 3 years postoperatively, patients' maximum cardiac index had increased significantly and after 3 years there was no difference between patients and controls (8.1 ± 1.2 l \cdot min⁻¹ \cdot m⁻² vs 8.3 ± 1.6 l \cdot min⁻¹ \cdot m⁻², respectively, $P = 0.572$).

(Continued)

Table 1 (Continued)

Study	No of patients	Age (years), mean \pm SD (range)	Haller index mean \pm SD	Surgical method	Variables studied	Cardiopulmonary outcomes
Tang et al ³⁸	49 patients and 26 controls	15.5 \pm 1.7	5.3 \pm 2.3	Nuss	Minimum and maximum heart rate, VO ₂ max, cardiac index, stroke index, ejection fraction, fractional shortening, left ventricular systolic and diastolic diameter, and right ventricular diastolic diameter before and at 1 year post-repair in patients and in healthy, age-matched controls	<ul style="list-style-type: none"> • There was no difference in the heart rate at any point of the study between patients and controls. • The significantly lower maximum oxygen uptake in patients compared to control group that was found at 1 year ($P = 0.001$) disappeared at 3 years of follow-up ($P = 0.430$). • Maximum cardiac index had increased significantly at 1 ($P = 0.001$) and 3 years ($P = 0.0001$) postoperatively, and it was normalized with no significant difference found between patients and controls at 3-year follow-up ($P = 0.572$). • Stroke index had significantly increased at the 3-year follow-up only in the patients ($P = 0.0002$) with no increase in the controls ($P = 0.299$). • Left ventricular diastolic diameter was significantly increased ($P < 0.05$) at 12 months of follow-up in the PEx patients. • Right ventricular diastolic diameter and VO₂ max were significantly increased in the control group at 12 months of follow-up. • Maximum cardiac index was significantly improved at 1 year postoperatively ($P = 0.0054$) but was still significantly lower than the control group ($P = 0.0008$). • No difference was found in the heart rate, ejection fraction, fractional shortening, and left ventricular systolic diameter between the two groups. • Patients had significantly lower stroke index than controls before operation ($P < 0.01$) but the difference had disappeared at 1 year following repair. • There was a significant increase in peak VO₂ ($P = 0.0005$) and O₂ pulse ($P = 0.001$) at 1 year follow-up, with no significant change in heart rate. • There was a significant increase in all right ventricular dimensions ($P < 0.001$) and in the left ventricular ejection fraction ($P < 0.001$). • VO₂ max, O₂ pulse, respiratory quotient, stroke volume, and cardiac output all showed significant increase ($P = 0.05$) post-bar removal but there was a significant decrease in minute volume breathing ($P = 0.05$) at maximal exercise. • No change in heart rate or cardiac index.
Neviere et al ¹⁹	70 patients	27 \pm 11 (18–62)	4.5 \pm 1.1	Simplified Ravitch-type repair	Heart rate, maximal oxygen uptake (peak VO ₂), O ₂ pulse during rest, and maximal exercise before and at 1 year following surgical repair	
Krueger et al ²⁵	17 patients	28 (17–54)	NR	Ravitch-Shamberger	End-diastolic right ventricular dimension (diameter, area, and volume) and left ventricular ejection fraction	
Sigalet et al ⁸	26 patients	13.2 \pm 2.1	4.5 \pm 1.3	Nuss	Heart rate, VO ₂ max, O ₂ pulse, respiratory quotient, minute volume breathing at maximal exercise, stroke volume, cardiac output, and cardiac index preoperative and after bar removal	

(Continued)

Table 1 (Continued)

Study	No of patients	Age (years), mean \pm SD (range)	Haller index mean \pm SD	Surgical method	Variables studied	Cardiopulmonary outcomes
Bawazir et al ¹⁹	48 (48 patients' data were available at 3 months postoperatively, 22 patients at 21 months, and 11 patients completing the full evaluation after bar removal)	13.5 \pm 1.7	3.9 \pm 0.8	Nuss	VO ₂ max, anaerobic threshold (VO ₂ max % expected), minute volume breathing at maximal exercise, maximal heart rate, stroke volume, cardiac output, and cardiac index before and at 3 and 21 months after repair, and at 3 months after bar removal	<ul style="list-style-type: none"> At preoperative assessment, VO₂ max and anaerobic threshold were significantly below normal for patients of this age and size. There was a significant decrease in VO₂ max and anaerobic threshold at 3 months post-repair ($P < 0.05$) and they had returned to the preoperative values by 21 months post-repair, while maximal heart rate and the minute volume of ventilation showed no significant changes. The cardiac index and cardiac output increased at 3 months post-repair ($P < 0.05$) and cardiac index remained stable. No significant increase in the stroke volume was found throughout the study period.
Haller et al ¹⁸	36 patients and 10 controls	16 \pm 3	NR	Open	Maximal heart rate, exercise duration (min), speed (kph), grade (%), and O ₂ pulse in patients and healthy, age-matched control group before surgery and 6 months post-repair	<ul style="list-style-type: none"> There were significant increases in duration ($P < 0.02$), speed ($P < 0.02$), and grade of exercise ($P < 0.05$) and in O₂ pulse ($P < 0.005$) after surgical repair in the post-repair group, with no changes in maximal heart rate.
Morshuis et al ¹⁷	35 patients	17.9 \pm 5.6 (9.3–29.9)	NR	Open, Daniel technique with modifications	Heart rate, oxygen uptake (VO ₂), and oxygen pulse before and at 1 year post-repair	<ul style="list-style-type: none"> There was a significant increase in VO₂ max, and oxygen pulse ($P = 0.0333$) during exercise with no change in the maximal work performed or in the heart rate.

Abbreviations: MIRPEX, minimally invasive repair of pectus excavatum; RV, right ventricle; LV, left ventricle; EF, ejection fraction; SV, stroke volume; EDV, end-diastolic volume; ESV, end-systolic volume; PImax, maximal static respiratory pressure; SNIP, sniff nasal inspiratory pressure; PEx, pectus excavatum; NR, not reported.

Cardiac output and deconditioning can occur with postoperative inactivity;³⁶ therefore, the patient's exercise history can affect the measurements of VO_2 and may cause more impact on short-term testing results.³⁴

Quality of life and patient satisfaction

Both the exercise limitations and the cosmetic disfigurement with PEx may cause a decrease in quality of life and alteration of social behavior.^{37,40–44} There has been a greater recognition of the physiologic and psychologic impact of these patients.⁴⁵ Lack of self-confidence, poor body image, avoidance of social activities, and emotional difficulties are noted in PEx patients. Feelings of anxiety, depression, sadness, and frustration are also reported.⁴⁰ The importance of corrective surgery for improvement in psychological distress, quality of life, and exercise tolerance has been documented in the literature.^{37,41–44} The majority of these studies report a mixed population of children and adolescents with few adults; therefore, it is difficult to make broad-based assumptions as to their application to the adult population.^{46–50} Table 2 reviews some of the major publications^{44,48–50,53,54} reporting postsurgical quality of life and symptom outcomes.

Kelly et al¹⁰ reported on 264 child patients and 291 parents from multiple centers using a validated Pectus Excavatum Evaluation Questionnaire. Children noted a dramatic improvement in the body image and physical difficulties after surgery. Parents also noticed an improvement in the child's emotional, physical difficulties and social self-consciousness.¹⁰ Patient's satisfaction with the chest appearance was found to be very good, with excellent to good results reported in over 95% of patients at the time of bar removal.⁵¹

In a 2016 study performed by Lomholt et al,⁴⁹ 107 patients and 106 parents completed the generic health-related quality-of-life measure. The Child Health Questionnaire was assessed preoperatively and at 3, 6 months following PEx repair. A control group of 183 school children completed the same measure on one occasion. In the postoperative study, patients and parents reported improved emotional well-being and self-esteem. Additionally, patients at both 3 and 6 months postoperatively reported increased physical and social activities.

There are very few major publications that documented symptoms and quality-of-life improvement after Nuss repair in adult patients. Kragten et al¹² reported on symptomatic seniors with PEx. He found that in 45% of the patients with "serious and sometimes invalidating complaints", symptoms did not start until the fourth or fifth decade of life and were

often labeled as "unexplained cardiovascular complaints". All patients that underwent surgery were repaired by the open Ravitch procedure and reported substantial or complete resolution of the symptoms postoperatively. Tikka et al⁵² used the Brompton's single-step questionnaire (SSQ) to assess the postoperative patient satisfaction and confirmed that Nuss operation had positively impacted the psychological and physical status of their patients along with overall quality-of-life improvement. They reported that their pectus patient information website improved, additionally, their patient's satisfaction and recovery after surgery.

Krasopoulos et al⁴³ proposed the two-step Nuss Questionnaire modified for Adults (NQ-mA) and a SSQ. These questionnaires measured the disease-specific quality-of-life changes after surgery and assessed the effect of surgery on the physical and psychological well-being of postoperative patients. They noted that patients' self-esteem, social functioning, and level of satisfaction were significantly improved following Nuss procedure. Their questionnaire also included the impact of surgical wounds/scars on the overall cosmetic result, consciousness of the presence of metallic bar, the decision to have the operation again, and questions about postoperative pain which may have limited the patient satisfaction after surgery. It was evident from the study that most of the patients were very satisfied with their scars and almost all of them were conscious of the presence of bar, but none of them considered that to be a major inconvenience. Pain was also noted as a concern in the immediate postoperative period; however, it decreased significantly after several weeks. By 4–5 months after surgery, no patient was still requiring analgesics.

Other surgeons have subsequently utilized this modified survey for assessing the patients postoperatively.^{48,50,53} Hokschi et al⁵⁰ performed a prospective study to evaluate the long-term results of Nuss in adults using NQ-mA and SSQ in a shorter and modified format. Initially, a large adult cohort ($n = 129$) was included, but only 19 patients were observed for >10 years after surgery. This has been the only study reporting outcomes for an adult population for more than 10 years after surgery. The results obtained initially after surgery were in the follow-up period of 3, 12 and 36 months showed high levels of satisfaction respectively reported at 97.6%, 97.2%, and 95.7%. Better or much better quality of life was reported at 3, 12, and 36 months in 88%, 89%, and increased to 92.5%, respectively, in the follow-up period. Even after observation for >10 years, continued improvement in quality of life was confirmed in 57.9% of patients. Surgical recommendation for Nuss was given by nearly 95% of patients. Mild pain

Table 2 Review of major publications from 2006–2016 reporting quality of life and patient satisfaction after pectus excavatum repair

Study, years	No of patients/parents in the survey	Age at operation: mean/median (range and/or SD), years	Evaluation	Reported results following surgery
Tikka et al, ⁵⁴ 2016, UK, 2008–2014 "Pectus patient information website"	59 patients, 32 responders	Responders (n = 32), 20 (18–22) years; nonresponders (n = 27), 19 (17.7–20) years	Brompton's SSQ	<ul style="list-style-type: none"> The authors have introduced the website www.pectus.co.uk which provides information about pectus deformities, management options, and advice about surgery to the general population. They have found out that 84% of patients who underwent surgery after visiting the website found it useful. Patients were overall highly satisfied with the surgical outcomes. All of their patients who visited the website said that they would have the operation again in comparison to 76.9% of those who did not ($P = 0.031$). Both patients' and controls' level of health-related quality of life before surgery were comparable except for physical functioning. Boys reported impaired function compared to controls ($P < 0.0001$). There were significant ($P < 0.001$–0.03) improvements in emotional well-being and self-esteem, in addition to an increase in physical and social activities from pre- to postsurgery as reported by the patients and parents.
Lomholt et al, ⁴⁹ 2016, Denmark, 2006–2008 "Prospective study on quality of life in youth after surgical correction"	107 patients, 106 parents completed surveys on the day before surgery, and at 3 and 6 months after surgery. Control group: n = 183, age-matched school children.	Children and adolescents	Child Health Questionnaire (CHQ)	<ul style="list-style-type: none"> Better or much better quality of life after Nuss was observed (n = 88.4% at 3 months, n = 89% at 12 months, and n = 92.5% at 36 months). Pain intensity decreased in the follow-up (pain score visual analog scale [VAS] at 3 months: median 1 (0–7), 12 months: 1 (0–4), 36 months: 0.8 (0–5)). After long-term observation, over 90% described their quality of life after operation as being better or much better. Satisfaction with the cosmetic result was also very high, with >90% being satisfied. Satisfaction with the cosmetic result was still very high (84.2%), and continued improvement in QOL 10 years after surgery was confirmed by 57.9% (n = 19). Recommendation to opt for Nuss was given by 94.7% (n = 18). Results after bar removal were overall satisfactory in 94.4%. About 82% of patients were either satisfied or very satisfied. Overall, 79.5% stated that they would have the operation again. About 1% noted troublesome awareness of the bar for 2 years. Two patients (2.2%) required reoperation for recurrence. About 89.7% of responders reported satisfaction with the chest appearance. Most patients stated that general health and exercise tolerance were improved after operation and 84.6% of responders noted an improvement in social interaction.
Hoksch et al, ⁵⁰ 2016, Switzerland, 2002–2014 "Long-term (> 10 years after surgery) results in adults"	129, bars removed in 72.9%, 19 patients followed for > 10 years	21 (13–56) years	NQ-mA and SSQ in a shorter and modified format	
Sacco Casamassima et al, ⁵³ 2016, Maryland, US, 1998–2011, "Long-term outcomes and overall patient satisfaction after bar removal of adult PEX with Nuss"	98, 89 patients had bar removal, 39 (43.8%) participated in the survey	32.3 (21.8–55.1) years	Modified SSQ to assess patient satisfaction with operative results after bar removal	

(Continued)

Table 2 (Continued)

Study, years	No of patients/parents in the survey	Age at operation: mean/median (range and/or SD), years	Evaluation	Reported results following surgery
Kuru et al, ⁴⁴ 2015, Turkey, "Changes related to quality of life 6 months after Surgery"	88 patients and majority of their parents completed the questionnaires before and 6 months following operation	18.44 ± 3.93 (14–29) years	Turkish version of NQ-mA	<ul style="list-style-type: none"> Patients' median Nuss score increased from 31 (IQR: 31–35) preoperatively to 43 (IQR: 43–46) at 6 months after the operation ($P = 0.000$). The parents' preoperative score of 33 (IQR: 29–36) increased to 38 (IQR: 34–41; $P = 0.000$). Improvements in the physical and psychosocial component scores of Nuss questionnaire were also significant in the patient ($P = 0.000$, $P = 0.000$, respectively) and parent forms ($P = 0.005$, $P = 0.000$, respectively). The authors indicated that the mean self-esteem score was improved from 4.6 of 10 preoperatively to 6.5 postoperatively ($P = 0.002$). The social impact of the pectus deformity became less significant (mean preoperative score 3.6, mean postoperative score 2.8, $P = 0.02$). The severity of initial postoperative pain was much improved on follow-up. Satisfaction with the cosmetic result was achieved in 80%. The decision to have the surgery again was reported by 96%. Over 50% of patients stated that they felt better or much better after the operation. The authors have concluded that the symptoms were reduced markedly or had disappeared completely after surgery (11 patients underwent surgery). All of their patients indicated that their health-related quality of life was significantly improved postoperatively, with an average of 2.7 points on a scale of 1–10. Complete relief of symptoms can be achieved by surgical reconstruction of the chest in seniors with symptomatic PEx.
Hanna et al, ⁴¹ 2013, Canada, 2006–2012, "Midterm results of Nuss in young adults with pectus excavatum"	73; 51 patients participated in the survey (73% response rate)	20 (16–51) years	Validated single-step quality-of-life survey	<ul style="list-style-type: none"> Significant positive postoperative changes were reported by the patients and their parents. Improvements occurred in both physical and psychosocial functioning, including less social self-consciousness and a more favorable body image. For the parent questionnaire, the child's emotional difficulties improved from 1.81 ± 0.70 to 1.24 ± 0.36, social self-consciousness improved from 2.86 ± 1.03 to 1.33 ± 0.68, and physical difficulties improved from 2.14 ± 0.75 to 1.32 ± 0.39 (mean \pm SD). For children, there was an improvement in the body image component from 2.30 ± 0.62 to 1.40 ± 0.42 after surgical repair and also improvement in the physical difficulties component from 2.11 ± 0.82 to 1.37 ± 0.44 (mean \pm SD). About 97% of patients thought that surgery improved how their chest looked. About 90% were either happy or mostly happy after surgery. There was a significant psychosocial improvement independent of Nuss or Ravitch procedures.
Kragten et al, ¹² 2011, the Netherlands, 2006–2009, "Prospective study evaluating the results of surgical correction of PEx"	42 senior patients	Over 50 years of age (seniors)	Symptomatic Pectus Excavatum in Seniors (SPES) score and Pectus Evaluation Index (PEI) score	
Kelly et al, ¹⁰ 2008, Multicenter, 2001–2006, "Before and 1 year after surgery for a large number of patients"	264 patients and 291 parents completed initial questionnaire preoperatively and 247 patients and 274 parents completed postoperative questionnaire	(8–21) years, 63% aged between 13 and 17 years, 11% were > 18 years of age	Pectus Excavatum Evaluation Questionnaire (PEEQ)	

(Continued)

Table 2 (Continued)

Study, years	No of patients/parents in the survey	Age at operation: mean/median (range and/or SD), years	Evaluation	Reported results following surgery
Metzelder et al, ⁴⁸ 2007, Germany, 2002–2006, "Lasting satisfaction after bar removal"	40 patients and 39 parents at 6 months postoperatively	Children and adolescents, 13.5 (6–20) years and 17 (10–24) years when the questionnaire was completed	SSQ with modifications focusing on satisfaction evaluation of the operative result after pectus bar removal	<ul style="list-style-type: none"> • There was a high level of persistent patient satisfaction with MIRPEX after bar removal. Mean total score of the patients was 67 (53–80) and mean score of their parents was 65 (41–79), with a highly significant correlation between self- and parental assessment ($P < 0.001$; Spearman's correlation coefficient = 0.77). • Analysis of specific and total scores revealed a significant improvement of physical and psychosocial well-being after surgical repair, which persisted up to 4 years after bar removal ($P < 0.001$). • There was no significant impact of age and sex on the mean specific and total scores either in patients or in parents. • About 97.5% of patients and 97.4% of parents were satisfied with the overall outcome after removal of pectus bar at the time of assessment by questionnaires. • No interference with overall satisfaction or willingness to have operation again occurred due to postoperative pain. • A highly significant correlation was reported between individual answers of each patient at follow-up 6 months after MIRPEX and after pectus bar removal ($P < 0.001$). • Analysis of the median scores obtained for each question and the total score of the individual patients in the NQ-mA revealed a statistically significant ($P = 0.001$) improvement following surgery. • NQ-mA domains for social function and self-esteem demonstrated a highly significant improvement following surgery ($P = 0.001$–0.008). • NQ-mA domains for physical condition showed a statistically significant improvement only for the degree of dyspnea ($P = 0.005$). • Total scores obtained with SSQ showed significant correlation with the total scores obtained from the postoperative NQ-mA, with a Spearman's correlation coefficient of 0.682 ($P = 0.001$). • Overall, SSQ revealed a high level of satisfaction following Nuss, with the exception of two patients in the low satisfaction zone (mean = 65, range: 49–75). There was a statistically significant improvement with surgery in self-esteem ($P = 0.001$). • About 85% considered that they had made the right decision to undergo operation. • SSQ highlighted that pain was a major concern during hospital stay, even with maximum analgesia. Pain was well controlled after some weeks, and by 4–5 months following surgery no patient was on analgesics.
Krasopoulos et al, ⁴³ 2006, UK, "Short-term results after Nuss"	20 patients	Young male adults, 18 (14–37) years	NQ-mA and SSQ	

Abbreviations: MIRPEX, minimally invasive repair of pectus excavatum; NQ-mA, Nuss questionnaire modified for Adults; SSQ, single-step questionnaire; QOL, quality of life; PE, pectus excavatum; IQR, interquartile range.

Table 3 Review of some of the recent studies and reported results from 2008–2016 after Nuss procedure for pectus excavatum repair in adults

First author, year, study period	Mean/median, age (range and/or SD), years	No of patients	Technique	Operative time, mean/median (range and/or SD), min
Pilegaard, ⁵⁵ 2016 (2001–2016), Retrospective till 2010 and prospective afterword	16 (7–58) >18 y, N = 604 (35%) <18 y, N = 1109 (65%)	1713	MIRPEX Short bar technique Unilateral stabilizer close to the hinge point with sutures to fix the bar on the opposite side In the last 6 years, no additional sutures were used, and the bar has been placed asymmetrically on the chest One bar: 70.6% Two bars: 28.4% Three bars: 1% (in four patients, there was combined correction of PEX with a cardiac operation)	36 (12–270)
Jaroszewski et al, ²⁴ 2016 (2010–2015)	23.7 (18–29), 40.4 (30–72)	266 18–29 y, n = 115 (43.2%) 30–72 y, n = 151 (56.8%)	Thoracoscopic MIRPEX Forced sternal elevation with sternal bone clamp attached to a bedside Rultract retractor. Multiple bars FiberWire multipoint fixation	18–29 y cohort, MIRPEX: 111 (62–178) Hybrid: 247.5 (138–395) 30–72 y cohort, MIRPEX: 121 (60–224) Hybrid: 231.1 (106–390)
Pawlak et al, ⁵⁶ 2016 (2002–2012)	18.2 ± 5.4 (7–49) Groups by age: A: 12.2 ± 2.0 (7–14) y B: 17.2 ± 1.6 (15–20) y C: 25.2 ± 4.8 (21–49) y	680, Groups: A(n = 156), B(n = 328), C(n = 196)	Thoracoscopic MIRPEX Unilateral stabilizers No of bars Groups: A, B, C, respectively One bar: A = 57.6% B = 50.3% C = 36.7% (P = 0.0021) Two bars: A = 41% B = 48.7% C = 61.7% Three bars: A = 1.2% B = 0.9% C = 1.5%	A, B, and C, respectively A: 50.2 ± 19.1 (25–165) B: 50.9 ± 19.8 (15–170) C: 55.4 ± 22 (25–200) (P = 0.0030)

LOS, mean/ median (range and/or SD), days	Complications, %	Redo, %	Patients with bars removed, %	Results
The median LOS decreased over time from 6 (4–29) to 2 days Currently, 1.6% of patients stay more than 4 days, compared with 46% during the early study period	Bar rotation (21 cases [1.2%]) and dislocation (13 cases [0.8%]) Deep infection: 0.9% Fractura sterni (one patient): 0.06% *Sternotomy (two cases): 0.1% Pneumonia: 0.6% Pneumothorax: 1.1% Seroma: 0.4% Pleural effusion: 0.3% Bleeding requiring re-operation: 0.1% Bar end dropped into chest cavity: 0.2% Removal of bar before time: 0.4% Stabilizer was removed because of pain: 1.4%	NR	NR	No mortality. Over the study period, there was increase in the number and decrease in the length of the bars used. The duration of postoperative hospitalization decreased over the study period with the majority of patients (>85%) currently being discharged on the second postoperative day.
2013–2015 18–29 y cohort, MIRPEX: 3.1 (2–6) Hybrid: 6.5 (6–7) 30–72 y cohort, MIRPEX: 3.3 (2–6) Hybrid: 5.6 (3–11)	18–29 y cohort vs 30–72, respectively Bar rotation: 1.7% vs 6.6% Infection: 0.9% vs 1.3% Pleural effusion (and thoracentesis): 2.6% vs 6% Pneumothorax requiring chest tube: 0.9% vs 0.7% Pulmonary embolism: 0% vs 1.3% Bleeding requiring transfusion: 0% vs 0.7% Reoperation for bleeding: 1.7% vs 0.7% Pneumonia 0% vs 4% Urinary tract infection: 0% vs 4% Urinary retention required catheterization: 7.8% vs 8.6% Readmission for pain control: 0% vs 2% One patient with subjective report of regression in > 30 group.	NR	19%, > 30 y	MIRPEX was successfully performed in 88.7% of adults ≥30 years and in 96.5% of patients between 18 and 29 years. Higher percentage of older patients required osteotomy or cartilage resection (11.3% vs 3.5%). Although greater, the frequency of bar rotation requiring reoperation was not significantly increased in the older patients ($P = 0.74$). For complete correction, three bars were required in > 40% of adult patients. Older patients had a 65.2% increase in right ventricular output on transesophageal echo intraoperatively. No recurrences.
NR	A, B, and C respectively. Pneumothorax: A = 14.7% B = 27.1% C = 22.4% ($P = 0.0099$) Requiring drainage: A = 26.1% B = 27% C = 50% Pleural effusion: A = 5.7% B = 6.4% C = 11.2% ($P = 0.1436$) Requiring chest tube or thoracentesis: A = 77.7% B = 90.5% C = 72.7% Pleural hematoma: A = 0.6% B = 0.9% C = 0.5% Fever: A = 1.9% B = 3.6% C = 4.6% ($P = 0.3969$)		NR	Good cosmetic results reported with the use of Nuss irrespective of age of the patients. Good and very good corrective results in 97.7% of the patients. Satisfactory corrective effect: A = 96.8% B = 98.7% C = 96.4% vs unsatisfactory effect: A = 3.2% B = 1.2% C = 3.6% ($P = 0.4563$)

(Continued)

Table 3 (Continued)

First author, year, study period	Mean/median, age (range and/or SD), years	No of patients	Technique	Operative time, mean/median (range and/or SD), min
Ersen et al, ⁵⁷ 2016 (2006–2016)	16.8 (2–45) y Adults: 23.2 (18–45) y	836 Adults: n = 236 (28.2% >18 y)	Thoracoscopic MIRPEX Left-sided stabilizer placed and medial bar secured on right side with polydioxanone (PDS) sutures around ribs. 2/ > bars: 15.8% adults and 7.5% of younger patients (P = 0.068) Shorter bars: median length: 11 inches (9–14) in adults and 10 inches (7–14) in younger patients	44.4 (25–90)
Sacco Casamassima et al, ⁵³ 2016 (1998–2011)	30.9 (21.8–55.1) y	98, 39 patients from 89 who underwent bar removal participated in the survey (43.8% response rate)	MIRPEX without thoracoscopy Bilateral stabilizers One bar: 89.7%; Two bars: 10.2%	62.9 ± 24.9

LOS, mean/ median (range and/or SD), days	Complications, %	Redo, %	Patients with bars removed, %	Results
	Rotation of the bar: A = 0.6% B = 3.6% C = 2% ($P = 0.1232$) Redo corrective surgery due to significant bar displacement in one patient from group A, five patients from group B, and two patients from group C Recurrence: A = 3.2% B = 1.2% C = 1.5% ($P = 0.3251$)			Although the younger patients have the lowest surgical morbidity, the recurrence rate is higher compared to the other groups. Surgical morbidity, reported in most of the patients, was temporary and reversible. Complications did not interfere with a satisfactory outcome of surgical repair.
4.92 ± 2.81 (3–21) in adults and 4.64 ± 1.58 (2–13) in younger patients ($P = 0.637$)	No peri-operative deaths Bar displacement: 5% underwent reoperation Cardiac injury – small ventricular defect: 0.4% Aspiration pneumonia: 0.4% Pleural effusion: 0.8% Wound infection: 1% Thoracic outlet syndrome: 0.4% Prolonged pain: 1% Recurrence: 0%		57%	Overall complications: 26.2%; 11.8%, respectively, for adult and younger patients ($P = 0.007$). Excellent cosmetic results reported in all patients. Patients with > 1 bar had less pain in the adult group. No bar displacement/ rotation in the last 3 y. Same good results can be achieved in adult patients as younger patients with the same operative time and same number of bars. Although adult patients have higher complication rate compared to the younger patients, the length of their hospital stay postoperatively was shorter.
3.6 ± 1.2	Ventricular arrhythmia: 1% Pulmonary embolism: 1% Pleural effusion: 8.2% Pleural effusion with drainage: 1% Hemothorax: 2% Reoperation for bar displacement: 4.1% Reoperation for placement of a second bar due to depression of upper part of sternum: 1% Reoperation for recurrence after bar removal: 2.2% Mild recurrence- 2.2% Residual carinatum: 1.1% Bar infection necessitating removal: 1% Bar removed due to uncontrolled pain: 4.7% Pneumothorax: 11.2% Pneumonia: 2% Wound infection: 10.2% Wound drainage, noninfectious / seroma: 3.1% Allergic reaction: 2 % Prolonged opioid > 8 weeks: 12.2%		90.8% (n = 89) Bar removed <18 months: 6.1% ; persistent chest pain: 4.15%, bar infection: 1%; chronic wound infection: 1%	General health and exercise tolerance were improved after operation in the majority of patients. Satisfactory cosmetic and functional result: 94.4% (n = 84/98). 89.7% reported a subjective improvement in the appearance of their chest wall (n = 35/39). Social interaction improved by 84.6% (n = 33/39); 82% (n = 32/39) were either satisfied or very satisfied. 79.5% stated that they would choose the surgery again (n = 31/39). 1% noted troublesome awareness of bar for 2 years (n = 1/98). Postoperative pain was shown to be the dominant factor in the quality of postoperative course.

(Continued)

Table 3 (Continued)

First author, year, study period	Mean/median, age (range and/or SD), years	No of patients	Technique	Operative time, mean/median (range and/or SD), min
Fibla et al, ⁵⁸ 2016 (2001–2010)	21.2 (10–47)	149, the surgery could not be concluded in two patients due to the inability to elevate the sternum (147 used for calculations in some instances)	Multi center MIRPEX most with thoracoscopy, few Ravitch included Stabilization not reported One bar: 94.6%; Two bars: 5.4%	NR
Park et al, ⁵⁹ 2015 (1999–2012)	10.3 (16 months–53 y) Adults >15 y: 21.4%	1816 Groups according to the bar fixation method: STB: n = 180 MPF: n = 760 CFT: n = 699 CFT + HP : n = 177	MIRPEX thoracoscopy/pectoscopy Series with different bar fixation techniques Stabilizers Multipoint pericostal suture fixation Claw fixator Hinge plate	NR
Zhang et al, ⁶⁰ 2015 (2006–2014)	15.3 ± 5.8 y (2.5–49)	639 2.5–5 y: n = 29; 6–12 y: n = 134; 13–18 y: n = 325; 19–25 y: n = 123; 26–49 y: n = 28	MIRPEX with thoracoscopy Right side secured stabilizer and left sutured to lateral chest wall muscles and rib periosteum Bilateral fixation bars in recurrent patients Limited sternal/coastal cartilage resection performed if necessary One bar: 75.7%; Two bars: 24%; Three bars: 0.3%	64.3 ± 41.7 (40–310)
Park et al, ⁶¹ 2015 (2013–2014)	17.5 (6–38) y	80	Bridge technique connecting two parallel bars using plate-screws at the ends of the bars	NR

LOS, mean/ median (range and/or SD), days	Complications, %	Redo, %	Patients with bars removed, %	Results
NR	Bar displacement: 5.4% (with 3.4% requiring reoperation) Seroma in surgical wounds: 11.6% Breakage of the absorbable stabilizer: 2% Pneumothorax requiring drainage: 2% Hemothorax: 1.4% (one patient required reoperation) Wound infection: 2.7% Pericarditis: 0.7% Pericardial blockage and respiratory failure necessitating emergency bar removal: 0.7% All patients reported postoperative pain with 2% went bar removal due to intense pain		49% Difficult: 7%.	Initial results: Excellent/good: 93.2% Mild: 4.1% Bad: 2.7% After a 1.6-year follow-up period, good results persisted in 98.7%. Result was not satisfactory in 1.4% reoperated using Ravitch. Complications: 30.6% In 32 and 37 year old patients, surgery was converted to Ravitch due to impossibility of raising the sternum.
NR	STB vs MPF vs CFT + HP, respectively Total complication: STB: 20% MPF: 11.2% CFT + HP: 7.3% Early bar displacement: STB: 3.33% MPF: 0.56% CFT + HP: 0% Reoperation: STB: 5% MPF: 1.57% CFT + HP: 3.38% Pneumothorax: STB: 10.3% MPF: 2.6% CFT + HP: 0.5% Pleural effusion: STB: 3.3% MPF: 2.5% CFT + HP: 2.9% Pericardial effusion: STB: 1.6% MPF: 0.3% CFT + HP: 0.2% Wound seroma: STB: 5.4% MPF: 3.7% CFT + HP: 1.5% There were four cases (0.32%) of reoperation because of recurrence after bar removal		Pectus bars were removed from 1231 patients (67.7%).	Total complication rates lower in CFT + HP (14.1%) than STB group (22.7%), ($P < 0.01$). Repair has been durable in more than 99% who had bar removal during follow-up period of 10 y. Pericostal suture fixation is vital for securing the stabilizer to function appropriately. There was a shift from the use of stabilizer to use of pericostal sutures to fix the bar (multipoint fixation). Suture-less claw fixator was equally as effective as MPF.
5.2 ± 2.9 (4–36)	One postoperative death due to right atrial injury: 0.2% Bar displacement: 1.7% Bar displacement requiring reoperation: 0.6% Pericardial perforation: 1.3% Intercostal tear: 0.8% Wound infection: 0.5% Pneumothorax: 2.8% Pneumonia: 0.6% Pleural effusion: 0.8% Hemothorax: 0.2%		47.6%	Outcomes were excellent: 78.9%; good: 16.4%; fair: 4.4%; poor: 0.4%. Good quality rate: 95.3%. Adult patients significantly required more pectus bars compared to the youth. Within the average 31.1 months follow-up, no significant recurrences were reported. Complication rate: 9.7%.
NR	Complication rate: 7.5% Wound hematoma: 1.25% Wound dehiscence: 1.25% Wound infections: 2.5% Pleural effusions: 2.5% Bar displacement: 0%		NR	Over a 4-month follow-up period, there was no reported movement in the upper and lower bars, and there were no cases of bar displacement or reoperation. No suture fixations or invasive devices were required in the bridge technique, making it applicable to use.

(Continued)

Table 3 (Continued)

First author, year, study period	Mean/median, age (range and/or SD), years	No of patients	Technique	Operative time, mean/median (range and/or SD), min
Hanna et al, ⁴¹ 2013 (2006–2012)	20 (16–51) y 16% between 16 and 18 y	73 51 patients agreed to participate in a quality-of-life survey (73% response rate)	MIRPEX with thoracoscopy. Bilateral stabilizers, two wires to adjacent ribs One bar: 81%, Two bars: 19%	NR
Rokitansky and Stanek, ⁶² 2013 (2006–2013)	17.7 ± 7 Majority of patients between 14 and 20 y MMIPR: 15.2±5 MEMIPR: 22.5±8.	262 MMIPR: n = 121 MEMIPR: n = 141	MEMIPR : MMIPR + partial sternotomy (23%) Stabilizer wing (PSI by Hofer Medical, Austria) Slit-rib chondrotomy under thoracoscopic guidance (48%), rib resection (5%), and rib osteotomy MMIPR: ; Symmetrical PEx: 74% Two bars: 13.2% MEMIPR: symmetrical PEx: 57.4%, carinatum /excavatum: 4.9%, two bars: 58.1%	NR
Olbrecht et al, ⁶³ 2008 (1997–2006) Values: median (IQR)	23 (18– 30)	18–30 y: n = 107 (52 bars removed) 6–14 y: n = 137 (80 bars removed)	MIRPEX without routine use of thoracoscopic visualization Lateral stabilizers Sternal wire to anchor bar to crossing ribs bilaterally, One bar: 94.2% Two bars: 5.8% (<i>P</i> = 0.03)	82 (65.5–103.5)

LOS, mean/ median (range and/or SD), days	Complications, %	Redo, %	Patients with bars removed, %	Results
5 (3–9)	Bar displacement: 2.7% (required reoperation) Self-resolving pneumothorax: 4.1% Poor cosmesis: 2.7% (required revision for bar positioning) Bar infection: 1.4% Bruising: 1.4% Ileus: 1.4% Pericarditis: 1.4% Immediate postoperative pain was severe in 51% and very severe in 39%		57%	Mean self-esteem score significantly improved after surgery from 4.6 to 6.5 out of 10 postoperatively ($P = 0.002$). Mean of social impact of pectus deformity became less significant (3.6 to 2.8), ($P = 0.02$) Overall morbidity rate: 15.1%. No incidents of bar displacement were reported after using the stainless steel wires instead of polypropylene. Severity of initial postoperative pain was improved on follow-up with 84% reporting no pain or occasional pain with no need for analgesia. 80% were satisfied with the cosmetic result and 96% would choose to undergo surgical repair again. No documented recurrence of PEX after bar removal.
NR	MMIPR, MEMIPR, respectively No bar displacement, only minimal bar movement: MMIPR: 4.9% MEMIPR: 1.6% Stabilizer dislocation: MMIPR: 2.3% MEMIPR: 0% Subcutaneous infection: MMIPR: 0.7% MEMIPR: 0.8% Pneumothorax: MMIPR: 1.4% MEMIPR: 2.4% Pleural effusion (needs drain): MMIPR: 2.1% MEMIPR: 1.7% Subcutaneous hematoma: MMIPR: 1.7% MEMIPR: 4.1% Tissue necrosis: MMIPR: 2.5% MEMIPR: 0% Recurrence occurred in 0.9% of patients who underwent bar removal, with one patient undergoing early bar removal and required correction of a recurrent funnel chest		103 patients with a mean of 3.4 y (1.4–6.5)	MMIPR and MEMIPR yielded very satisfactory results, especially in older patients with severe deformities and recurrence. Simple MIPR did not yield the desired results in elderly patients with stiff thorax, curved sternum, severe asymmetrical forms, and a mixed pigeon/funnel chest in whom the modification procedures (MMIPR and MEMIPR) resulted in very satisfactory outcomes. About 95% of the patients were satisfied with the cosmetic outcome after bar removal. None of the 262 patients who underwent MMIPR or MEMIPR required re-thoracotomy.
3 (3–4)	Bar displacement requiring operation: 7.7% Pneumothorax: requiring tube: 3.9% Pleural effusion: 3.9% Pneumonia: 3.9% Superficial surgical site infection: 21.6% (requiring surgical revision: 5.9% Requiring early bar removal: 3.9%)		48.6% >18 y 58.4% 6–14 y	No patient required open procedure and two patients required sternal osteotomy. Operative time shorter and postoperative complications were similar to Ravitch. Long-term complication rates were equal between adult and pediatric cohort. Adults often require more than one bar for correction.

(Continued)

Table 3 (Continued)

First author, year, study period	Mean/median, age (range and/or SD), years	No of patients	Technique	Operative time, mean/median (range and/or SD), min
Cheng et al, ³⁰ 2008 (2005–2007), prospective	24.5 (18–42)	96 Young adults: (18–25) y, n = 63 Older adults: (26–42) y n = 33	MIRPEX with bilateral thoracoscopy Two bars: 22.9%, Wire/suture bar fixation at right hinge point and distal bar ends to rib	80 (50–185) Mean time for first repair with one bar: 65 (50–100); redo correction with two bars: 120 (100–185)

Note: *Both sternotomies were done in re-do cases: one was due to inferior vena cava bleeding and one was due to right ventricle tear caused by an adhesion, from the primary Nuss correction.

Abbreviations: NR, not reported; SD, standard deviation; LOS, length of stay; y, year; MIRPEX, minimally invasive repair of pectus excavatum; STB, stabilizer; MPF, multipoint pericostal fixation; CFT, claw fixator; HP, hinge plate; MIPR, minimally invasive pectus repair; MMIPR, modified minimally invasive pectus repair; MEMIPR, modified extended minimally invasive pectus repair; PEX, pectus excavatum; PC, pectus carinatum; QOL, quality of life; IQR, interquartile range; PSI, Pectus Security Implant.

occurring during specific bodily movements was reported in 31.6%, and 63.2% of patients had no pain.

Sacco Casamassima et al⁵³ in 2016 reported long-term results of adults using modified SSQ. Satisfaction with the chest wall appearance was reported in 89% out of 43.8% of responders. Improvement in social interaction was reported by 84% of responders. About 94% of patients obtained overall satisfaction with the results post-bar removal. They also highlighted that the dissatisfaction observed by some patients was due to severe postoperative chest pain (that necessitates more aggressive analgesic regimen) and surgical scars. Willingness to have the operation again was reported by 79% of responders. Generalized conclusions cannot be

drawn from this study as it is limited by small sample size. There is a compelling need for a large number of similar studies commenting on the long-term results in adults to identify the benefits of surgery in this group.

Hanna et al⁴¹ studied the midterm results in young adults who underwent Nuss repair and used the single-step quality-of-life survey for evaluation. With a 73% response rate, they noticed an improvement in both self-esteem and social life. Satisfaction with the cosmetic result was achieved in 80% and recommendation for the surgery was given by 96% of their patients. About 92% reported subjective improvement in the chest wall appearance. As stated by other authors, in-hospital pain despite aggressive analgesic usage was a major concern in the immediate postoperative

LOS, mean/ median (range and/or SD), days	Complications, %	Redo, %	Patients with bars removed, %	Results
7.2 (5–13)	Upper sternal depression requiring reoperation for second bar: 3.9% Development of pectus carinatum requiring repair: 1.9% Prolong pain: 47.1% Only two patients (3.9%) experienced a recurrence after bar removal, and neither had required operative repair Number of young adult vs older adult Bar displacement: 2 vs 0 (both required surgical revision) Pneumothorax: 0 vs 1 Pleural effusion requiring thoracentesis: 1 vs 0 Pericardial effusion with pericardio-centesis: 0 vs 1		7%	Predictors of surgical revision after repair were bar displacement ($P < 0.001$) and early bar removal ($P < 0.001$). Patients did not differ with respect to postoperative complications, including recurrence rates, bar displacement, upper sternal depression, or revision surgery. About 91.6% of patients were satisfied with their surgical correction Satisfaction results were reported as follow in young and old adults, respectively: Excellent: 75.3%, 69.7% Good: 17.4%, 21.2% Fair: 6.3%, 6.1% Failed: 1.6%, 3% Bilateral thoracoscopy facilitated mediastinal dissection, particularly in patients who had previous pectus/thoracic procedure or double-bar insertion. Patients with fair or failed results had more complicated deformities, including PEx combined with PC, long area deformity, severe bony rigidity, residual deformity, or partial recurrence after repair. Longer operation time and higher complication rate were reported in double-bar repair compared to single-bar procedures. Single bar ($n = 74$): 3%; double bar ($n = 22$): 13.6%.

period; however, in the follow-up it was significantly decreased, with almost all patients reporting minimal or no pain.

Most of the data available suggest that patients who had undergone Nuss showed an overall satisfaction with the cosmetic result, had a significant improvement in self-image, and felt that the surgery had a positive impact on their ability to exercise and well-being.

Surgical approaches and outcomes

The Nuss procedure or “MIRPEX” has become the standard of care for PEx repair in children and adolescents.⁶⁴ There is an ongoing discussion in the literature regarding the success of this surgery in adults with PEx. Initial reports of Nuss procedure in adults were criticized due to higher complication rates vs the open Ravitch technique

with most being related to bar migration, postoperative pain, and recurrences.^{65–67} The recommendations of some surgeons were to limit the procedure to pediatrics and adolescents; however, their publications have been replaced with numerous series of successful repairs using a modified MIRPEX approach.^{24,68,69} Table 3 reviews publications reporting on 70 or more adults repaired using an MIRPEX procedure since 2008. The majority of authors considered patients aged 18 years and older as adults.^{28,51,57,63,70,71} Several papers have stratified their results to differentiate younger vs older patients.^{24,30} There is evidence that older patients are more difficult to treat and the risk of complications may be greater.¹⁴ Despite this, excellent results are achieved with an MIRPEX approach even in older adult patients^{15,24,28–30,43,63,68,69,71–73} (Figure 3A–D).

Since the introduction of the original Nuss technique for children in 1998,⁶⁴ several changes have been made in the surgical technique and methods of bar stabilization which have improved the success of the procedure in adult patients.^{24,28,72,74–78} Important modifications include the use of forced sternal elevation,⁷⁸ multiple support bars,^{24,77} and improved fixation methods to secure the bars and prevent rotation.^{24,28,72,74–78} These technical refinements enabled successful MIRPEX repair of older patients and are reviewed in Table 4.

The use of forced sternal elevation may help reduce the force required to insert and rotate bars (Figure 4). This may lessen, but not eliminate, lateral stripping of the intercostal muscles of the more rigid chest wall.^{69,78,79,81,82,84} Several techniques have been proposed for the forceful elevation of sternum. Park et al⁷⁹ reported his Crane technique and discussed the benefits of its use in adult patients with heavier chests and severely asymmetric deformities including prevention of intercostal muscle tear and bar displacement. Similar variations of this technique have been reported by others with similar beneficial results.^{69,78,81,82,84} A more simplified aspect

of handheld retractors can also be utilized, depending on the severity and rigidity of the defect.^{80,83}

Multiple bars may balance the increased pressure of the chest wall and in older patients, the use of two or more bars is frequently reported.^{24,70,77,90} The risk of bar rotation and malposition may also be decreased by distributing the pressure of a more rigid chest wall.^{70,77,90} Pilegaard⁶⁸ reported that 70% of his patients over 30 years of age required two or more bars. In our own practice, two or more bars were utilized in 99% of patients over age 18 years,²⁴ with 40% of patients over 30 years receiving three bars to achieve complete repair. Others have reported decreased risk of bar migration and the need of reoperation when multiple bars were utilized.^{70,77,90} In a study of PEX repair in 44 late adolescent and adult patients, 11.5% of those with single-bar repairs required reoperation for incomplete correction or bar rotation compared with 0% who had a double-bar repair. Double bar also decreases the postoperative pain as described by Nagaso et al.⁸⁹ The risk of bar rotation may be lowered by the use of shorter bars as reported by several surgeons.^{93,94} In a publication reporting




Figure 3 Clinical photographs of a 22-year-old man with severe pectus excavatum are shown before surgery (A, B) and after (C) minimally invasive repair of pectus excavatum, with placement of three Nuss bars as shown in the chest roentgenogram (D).

Table 4 Review of several technical modifications reported for minimally invasive repair of pectus excavatum in adults

Technical modification	Study, years	Reported results
Forced sternal elevation:		
• Crane technique	• Park et al, ⁷⁹ 2008	• Relieves pressure on the hinge points, thus preventing intercostal muscle stripping (type 3 bar displacement).
• Crane technique using Kent retractor	• Yoon et al, ⁶⁹ 2010	• Authors confirmed 0% intraoperative death and 0% 30-day mortality.
• Two Langenbech hand held retractors	• Tedde et al, ⁸⁰ 2012	• They observed that this maneuver reduces the risk of pericardial sac and cardiac injury.
• Horseshoe-shaped sternal elevator	• Takagi et al, ⁸¹ 2012	• No additional skin incision needed for insertion of the elevator, and it widens the retrosternal space for safer passage of thoracoscopically guided introducer.
• Vacuum bell	• Haecker et al, ⁸² 2012	• No cardiac, pericardial, or internal mammary vessel injuries were noted. Facilitates retrosternal dissection and bar insertion.
• Manual sternal lift and anchor	• Johnson et al, ⁸³ 2013	• Utilized even in patients with severe pectus excavatum (Haller index >7); 3 cm subxiphoid incision needed. Improves bar stability and reduces displacement. No intraoperative complications.
• Bone clamp and Rultract retractor	• Jaroszewski et al, ⁷⁸ 2014	• Requires minimal additional incisions, decreases the force required to insert, rotates bars, and reduces the risk of intercostal muscle stripping in adult patients undergoing MIRPEX. The authors reported no intraoperative complications.
• T-fastener suture technique	• Kim et al, ⁸⁴ 2014	• Requires no specialized equipment, no incision in the anterior chest needed, does not cause any fracture or tear to the anterior chest structure. Disadvantage being removal of metal plate after positioning of the bar.
Bar stabilization techniques:		
• Five-point fixation	• Park et al, ⁷⁷ 2004	• All pericostal sutures can be done through the single tiny incision on each side, even in the parallel bar technique. Bar displacement: 3.4%.
• Third point of fixation	• Hebra et al, ²⁸ 2006	• Bar displacement: 6%; stabilizer bar fracture: 3%.
• Bar flipping: Multipoint bar fixation; lateral sliding: insertion of stabilizer on the depressed side; hinge-point disruption: hinge point reinforcement.	• Park et al, ⁷⁹ 2008	• Mechanism-based fixation effective in preventing bar displacement (4.6% vs 1.8% before and after MPF, respectively). Major complications decreased from 6.8% to 2% and reoperation rates decreased to 1.6% from 5.5% after MPF.
• Three-point wire fixation	• Yoon et al, ⁶⁹ 2010	• Narrows the intercostal space, thereby preventing hinge point disruption, bar migration, and rotation. More effective in adults.
• Unilateral stabilizer and multiple polydioxanone (PDS) sutures around ribs	• Kelly et al, ⁵¹ 2010	• Bar displacement requiring surgical repositioning decreased from 12% to 1%. Good to excellent anatomic result was obtained in 95.8%.
• Hinge plate	• Park et al, ⁸⁵ 2011	• Bar displacement rate in patients without the hinge plate: 4% vs 0% in the hinge plate group. Hinge plate is effective in preventing an intercostal strip at hinge points and has a vital role in extending MIRPEX to adults.
• Circumcostal sutures using Deschamps needle under endoscopic survey/lateral stabilizers	• Del Frari and Schwabegger, ⁸⁶ 2014	• Prevents bar displacement. Excellent position of the bar with circumcostal sutures in 96%, incomplete in 1.9%, and poor in 1.9%. With lateral stabilizers, 87.5% showed excellent position, and 12.5% showed poor position.
• FiberWire used to fix the bars circumferentially and bilaterally at multiple points	• McMahon et al, ⁷⁴ 2014	• Effective in preventing bar displacement and rotation. Metal stabilizers are not required.
• Claw fixator (CFT) and hinge plate (HP)	• Park et al, ⁵⁹ 2015	• CFT used for sutureless bar fixation by hooking the rib with blades, whereas HP prevents intercostal muscle stripping at hinge points. Bar dislocation rate with CFT + HP: 0%; reoperation rate: 3.38%; total complication: 14.1%. Authors recommend replacing conventional stabilizer with CFT and HP.
• Bridge technique	• Park et al, ⁶¹ 2015	• Designed to connect two parallel bars using plates and screws to avoid bar displacement, with no use of sutures or invasive devices. During the follow-up, there was no virtual change in bar position, bar dislocation, or reoperation.
• Unilateral stabilizer placed close to the hinge point, fixed to the bar by a steel wire	• Pilegaard, ⁷⁵ 2015	• No death, cardiac perforation, or deep infection occurred, and only 5% of patients experienced a complication.

(Continued)

Table 4 (Continued)

Technical modification	Study, years	Reported results
<ul style="list-style-type: none"> Figure-of-eight FiberWire reinforcement. Bars fixed bilaterally and circumferentially around the rib with FiberWire Stabilizer attached to bar with wire or FiberWire suture on left with multiple pericostal PDS sutures on right 	<ul style="list-style-type: none"> Jaroszewski et al,²⁴ 2016 Nuss et al,⁸⁷ 2016 	<ul style="list-style-type: none"> Prevents lateral-posterior migration when stripping occurs. Bar rotation: 6.6%. Rate of displacement with stabilizers: 5%, and with pericostal sutures: 1%.
Length of the bar:		
<ul style="list-style-type: none"> One inch (2.5 cm) shorter than the measurement from right to left mid-axillary line Eleven inch (7–15) in 2001–2010 and 10 inch (8–14) in 2011–2016 	<ul style="list-style-type: none"> Kelly et al,⁵¹ 2010 Pilegaard,⁵⁵ 2016 	<ul style="list-style-type: none"> Bar displacement requiring reoperation has been reduced from 13% to 1%. Reported lower rate of bar malrotations, and surgery can be done in less than an hour for over 90% of cases. (<2% bars flipped)
Multiple bars:		
<ul style="list-style-type: none"> Two bars Two bars in 32% 	<ul style="list-style-type: none"> Nuss,⁸⁸ 2008 Pilegaard and Licht,⁷⁰ 2008 	<ul style="list-style-type: none"> Bar displacement: 5%; requiring revision: 50%. Use of multiple bars was significantly more common ($P < 0.01$) in adults compared to younger patients, where 86% received one bar and 14% needed two bars.
<ul style="list-style-type: none"> Double-bar application 	<ul style="list-style-type: none"> Nagaso et al,⁸⁹ 2010 	<ul style="list-style-type: none"> Patients in one-bar group required self-injection of intravenous narcotics more frequently than patients in double-bar group (double-bar decreased postoperative pain). Stresses on the thoraces were smaller with double bars than with a single bar.
<ul style="list-style-type: none"> Two bars 	<ul style="list-style-type: none"> Stanfill et al,⁹⁰ 2012 	<ul style="list-style-type: none"> No patient required revision for bar displacement when two bars were used as opposed to 15.5% who required reoperation for bar movement when one bar was initially placed. ($P = 0.05$), with no difference in patient age and Haller index between groups.
<ul style="list-style-type: none"> Three bars 	<ul style="list-style-type: none"> Jaroszewski et al,²⁴ 2016 	<ul style="list-style-type: none"> More than 40% of patients of both adults over 30 year old and patients between 18–29 years groups required three bars
Hybrid approach/osteotomy:		
<ul style="list-style-type: none"> Transverse sternotomy/limited sternal resection/parasternal bar fixation 	<ul style="list-style-type: none"> Dzielicki et al,⁹¹ 2006 	<ul style="list-style-type: none"> Further procedures were essential to achieve and maintain an adequate correction and to decrease sternal rigidity and its pressure on the bar.
<ul style="list-style-type: none"> MOVARPE technique 	<ul style="list-style-type: none"> Del Frari and Schwabegger⁸⁶, 2014 	<ul style="list-style-type: none"> Used in adults with athletic disposition, deformities with deep funnel, and severe asymmetry. Only minor complications (4.4%) were observed.
<ul style="list-style-type: none"> Scoring of deformed cartilages 	<ul style="list-style-type: none"> Nagasao et al,⁹² 2016 	<ul style="list-style-type: none"> Postoperative pain as measured by the frequency of administration of anesthetics for 2 days was reduced: 4.9 vs 2.5.
<ul style="list-style-type: none"> Hybrid approach 	<ul style="list-style-type: none"> Jaroszewski et al,²⁴ 2016 	<ul style="list-style-type: none"> Open-cartilage resection, sternal osteotomy, or both was more commonly performed in patients older than 30 years (mean, 47.8 years vs 39.5 years; $P = 0.0003$) and with defect severity (11.3% vs 3.5% in younger patients).

Abbreviations: MIRPEX, minimally invasive repair of pectus excavatum; MPF, multipoint pericostal bar fixation; MOVARPE, minor open videoendoscopic assisted repair of pectus excavatum.

Nuss revision after procedure failure, too long bars were noted to be a factor related to failure.⁹⁵

The biggest challenge in adult patients continues to be bar fixation. A higher rate of bar displacement is reported in older patients.^{14,24} There are multiple successful ways reported for securing of bars. Medial fixation with a hinge reinforcement plate,⁸⁵ medially placed stabilizers,⁷⁵ multi-point fixation,^{24,69,77} and the Bridge technique, which was more recently published,⁶¹ have all been successful methods for bar fixation in adult patients.^{72,79,87,96}

Chondroplasty or open osteotomy may still be necessary to achieve adequate repair in some adult patients. Patients with complex combined deformities, extensively calcified chest walls, and significant asymmetry may require an open repair for optimal correction. The requirement for osteotomy or cartilage resection is more commonly reported in older patients.^{24,91,97,98} In our experience, over 88% of the patients ≥ 30 years were successfully repaired with MIRPEX; however, some required an osteotomy or open resection for fracture. Postoperative pain may also be reduced by scoring




Figure 4 The Rultract retractor can be utilized to forcefully elevate the sternum when attached by a bone clamp.

of deformed cartilages as illustrated by Nagasao et al.⁹² The use of a hybrid procedure may also be considered and is our procedure of choice for these more difficult deformities.²⁴ Both surgical principles are utilized by incorporating osteotomy cuts and external fixation as well as pectus support bars. Achieving adequate postoperative pain control remains a concern for adults undergoing Nuss.⁹⁹ Various analgesic regimens have been discussed by several authors.¹⁰⁰ Perioperative pain can be well managed by current techniques.¹⁰¹ These include the use of thoracic epidurals, intravenous on-demand patient-administrated narcotics, local paravertebral blocks, and subcutaneous continuous flow catheters.^{102–106} We have had excellent results using a protocol including gabapentin, ibuprofen, acetaminophen, and narcotics along with subcutaneous continuous flow catheters for postoperative pain control.^{24,102,107} Adjuvant medications for postoperative pain management have included the use of ketorolac, diazepam, and gabapentin.^{103–106} Intraoperative use of methadone can also be advantageous.^{24,102}

Discussion

The extension of the Nuss procedure to repair adults with PEx has been controversial in the past.^{14,65–67} There are now multiple publications that report successful repair of adults even beyond 50 years of age.^{9,12,24,55} The difficulty of repair and risk of complications do, however, increase with age.^{14,24,53} Adequate surgical experience with the Nuss procedure in younger patients that are easier to repair is critical prior to attempting the more difficult adult deformity.

Bar rotation and migration can be a significant issue and techniques to minimize intercostal stripping, such as reinforcement of intercostal spaces^{24,59,87} and medially placed stabilizers, may be of benefit in reducing the risks.⁷⁵ The use of forced sternal elevation can also decrease the forces required for bar insertion and positioning.^{69,78–84} The adult chest wall has additional complexities due to the decrease in flexibility and increase in weight. Multiple bars have been noted to decrease the weight supported by an individual bar and decrease the risk of rotation.^{24,70,88–90} Adequate stabilization of bars is also critical due to these factors, and medial and/or multipoint fixation has been shown to reduce bar displacement.^{24,28,51,59,61,69,75,86,87} We did not intend this publication to be an intensive review of surgical techniques in adult patients, and the majority of information presented was based on a larger case series which reported on primary Nuss repair in the adult population. Extension of the Nuss procedure to more complex repairs, such as patients with prior sternotomy or cardiac surgery, is beyond the scope of this paper and can be associated with catastrophic complications.^{55,108}

Conclusion

MIRPEx can be extended to repair the majority of older adult patients. Although adults undergoing Nuss procedure may have a higher rate of complications, continuous technical refinements have significantly reduced the complication rates and contributed to the success of the procedure. As there is increased difficulty in performing this procedure

in adult patients, the experience and expertise of surgeons at specialized centers is critical for successful outcomes. There is enough evidence to validate repair of adults with PEx. Published data support the benefits of repair with good outcomes and improvement of symptoms.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Fokin AA, Steuerwald NM, Ahrens WA, Allen KE. Anatomical, histologic, and genetic characteristics of congenital chest wall deformities. *Semin Thorac Cardiovasc Surg*. 2009;21(1):44–57.
2. Scherer LR, Arn PH, Dressel DA, Pyeritz RM, Haller JA, Jr. Surgical management of children and young adults with Marfan syndrome and pectus excavatum. *J Pediatr Surg*. 1988;23(12):1169–1172.
3. Cobben JM, Oostra RJ, van Dijk FS. Pectus excavatum and carinatum. *Eur J Med Genet*. 2014;57(8):414–417.
4. Chung CS, Myrianthopoulos NC. Factors affecting risks of congenital malformations. I. Analysis of epidemiologic factors in congenital malformations. Report from the Collaborative Perinatal Project. *Birth Defects Orig Artic Ser*. 1975;11(10):1–22.
5. Fonkalsrud EW. Management of pectus chest deformities in female patients. *Am J Surg*. 2004;187(2):192–197.
6. Ma IT, Rebecca AM, Notrica DM, McMahon LE, Jaroszewski DE. Pectus excavatum in adult women: repair and the impact of prior or concurrent breast augmentation. *Plast Reconstr Surg*. 2015;135(2):303e–312e.
7. Park HJ, Gu JH, Jang JC, Dhong ES, Yoon ES. Correction of pectus excavatum with breast hypoplasia using simultaneous pectus bar procedure and augmentation mammoplasty. *Ann Plast Surg*. 2014;73(2):190–195.
8. Sigalet DL, Montgomery M, Harder J, Wong V, Kravarusic D, Alasiri A. Long term cardiopulmonary effects of closed repair of pectus excavatum. *Pediatr Surg Int*. 2007;23(5):493–497.
9. Chao CJ, Jaroszewski DE, Kumar PN, et al. Surgical repair of pectus excavatum relieves right heart chamber compression and improves cardiac output in adult patients – an intraoperative transesophageal echocardiographic study. *Am J Surg*. 2015;210(6):1118–1124.
10. Kelly RE, Jr., Cash TF, Shamberger RC, et al. Surgical repair of pectus excavatum markedly improves body image and perceived ability for physical activity: multicenter study. *Pediatrics*. 2008;122(6):1218–1222.
11. Jaroszewski DE, Fonkalsrud EW. Repair of pectus chest deformities in 320 adult patients: 21 year experience. *Ann Thorac Surg*. 2007;84(2):429–433.
12. Kragten HA, Siebenga J, Hoppener PF, Verburg R, Visker N. Symptomatic pectus excavatum in seniors (SPES): a cardiovascular problem?: a prospective cardiological study of 42 senior patients with a symptomatic pectus excavatum. *Neth Heart J*. 2011;19(2):73–78.
13. Jaroszewski D, Steidley E, Galindo A, Arabia F. Treating heart failure and dyspnea in a 78-year-old man with surgical correction of pectus excavatum. *Ann Thorac Surg*. 2009;88(3):1008–1010.
14. Kim do H, Hwang JJ, Lee MK, Lee DY, Paik HC. Analysis of the Nuss procedure for pectus excavatum in different age groups. *Ann Thorac Surg*. 2005;80(3):1073–1077.
15. Esteves E, Paiva KC, Calcagno-Silva M, Chagas CC, Barbosa-Filho H. Treatment of pectus excavatum in patients over 20 years of age. *J Laparoendosc Adv Surg Tech A*. 2011;21(1):93–96.
16. Kaguraoka H, Ohnuki T, Itaoka T, Kei J, Yokoyama M, Nitta S. Degree of severity of pectus excavatum and pulmonary function in preoperative and postoperative periods. *J Thorac Cardiovasc Surg*. 1992;104(5):1483–1488.
17. Morshuis WJ, Folgering HT, Barentsz JO, Cox AL, van Lier HJ, Lacquet LK. Exercise cardiorespiratory function before and one year after operation for pectus excavatum. *J Thorac Cardiovasc Surg*. 1994;107(6):1403–1409.
18. Haller JA Jr, Loughlin GM. Cardiorespiratory function is significantly improved following corrective surgery for severe pectus excavatum. Proposed treatment guidelines. *J Cardiovasc Surg (Torino)*. 2000;41(1):125–130.
19. Nevieri R, Montaigne D, Benhamed L, et al. Cardiopulmonary response following surgical repair of pectus excavatum in adult patients. *Eur J Cardiothorac Surg*. 2011;40(2):e77–e82.
20. Malek MH, Fonkalsrud EW, Cooper CB. Ventilatory and cardiovascular responses to exercise in patients with pectus excavatum. *Chest*. 2003;124(3):870–882.
21. Gurkan U, Aydemir B, Aksoy S, et al. Echocardiographic assessment of right ventricular function before and after surgery in patients with pectus excavatum and right ventricular compression. *Thorac Cardiovasc Surg*. 2014;62(3):231–235.
22. Coln E, Carrasco J, Coln D. Demonstrating relief of cardiac compression with the Nuss minimally invasive repair for pectus excavatum. *J Pediatr Surg*. 2006;41(4):683–686; discussion 683–686.
23. Mocchegiani R, Badano L, Lestuzzi C, Nicolosi GL, Zanuttini D. Relation of right ventricular morphology and function in pectus excavatum to the severity of the chest wall deformity. *Am J Cardiol*. 1995;76(12):941–946.
24. Jaroszewski DE, Ewais MM, Chao CJ, et al. Success of minimally invasive pectus excavatum procedures (modified Nuss) in adult patients (± 30 years). *Ann Thorac Surg*. 2016;102(3):993–1003.
25. Krueger T, Chassot PG, Christodoulou M, Cheng C, Ris HB, Magnusson L. Cardiac function assessed by transesophageal echocardiography during pectus excavatum repair. *Ann Thorac Surg*. 2010;89(1):240–243.
26. Nevieri R, Benhamed L, Duva Pentiah A, Wurtz A. Pectus excavatum repair improves respiratory pump efficacy and cardiovascular function at exercise. *J Thorac Cardiovasc Surg*. 2013;145(2):605–606.
27. Kelly RE Jr, Mellins RB, Shamberger RC, et al. Multicenter study of pectus excavatum, final report: complications, static/exercise pulmonary function, and anatomic outcomes. *J Am Coll Surg*. 2013;217(6):1080–1089.
28. Hebra A, Jacobs JP, Feliz A, Arenas J, Moore CB, Larson S. Minimally invasive repair of pectus excavatum in adult patients. *Am Surg*. 2006;72(9):837–842.
29. Aronson DC, Bosgraaf RP, van der Horst C, Ekkelkamp S. Nuss procedure: pediatric surgical solution for adults with pectus excavatum. *World J Surg*. 2007;31(1):26–29; discussion 30.
30. Cheng YL, Lee SC, Huang TW, Wu CT. Efficacy and safety of modified bilateral thoracoscopy-assisted Nuss procedure in adult patients with pectus excavatum. *Eur J Cardiothorac Surg*. 2008;34(5):1057–1061.
31. Udholm S, Maagaard M, Pilegaard H, Hjortdal V. Cardiac function in adults following minimally invasive repair of pectus excavatum. *Interact Cardiovasc Thorac Surg*. 2016;22(5):525–529.
32. Topper A, Polleichtner S, Zagrosek A, et al. Impact of surgical correction of pectus excavatum on cardiac function: insights on the right ventricle. A cardiovascular magnetic resonance study dagger. *Interact Cardiovasc Thorac Surg*. 2016;22(1):38–46.
33. Albouaini K, Egred M, Alahmar A, Wright DJ. Cardiopulmonary exercise testing and its application. *Heart*. 2007;93(10):1285–1292.
34. Malek MH, Coburn JW. Strategies for cardiopulmonary exercise testing of pectus excavatum patients. *Clinics (Sao Paulo)*. 2008;63(2):245–254.
35. Maagaard M, Tang M, Ringgaard S, et al. Normalized cardiopulmonary exercise function in patients with pectus excavatum three years after operation. *Ann Thorac Surg*. 2013;96(1):272–278.
36. Perhonen MA, Franco F, Lane LD, et al. Cardiac atrophy after bed rest and spaceflight. *J Appl Physiol (1985)*. 2001;91(2):645–653.

37. O'Keefe J, Byrne R, Montgomery M, Harder J, Roberts D, Sigalet DL. Longer term effects of closed repair of pectus excavatum on cardiopulmonary status. *J Pediatr Surg*. 2013;48(5):1049–1054.
38. Tang M, Nielsen HH, Lesbo M, et al. Improved cardiopulmonary exercise function after modified Nuss operation for pectus excavatum. *Eur J Cardiothorac Surg*. 2012;41(5):1063–1067.
39. Bawazir OA, Montgomery M, Harder J, Sigalet DL. Midterm evaluation of cardiopulmonary effects of closed repair for pectus excavatum. *J Pediatr Surg*. 2005;40(5):863–867.
40. Jaroszewski D, Notrica D, McMahon L, Steidley DE, Deschamps C. Current management of pectus excavatum: a review and update of therapy and treatment recommendations. *J Am Board Fam Med*. 2010;23(2):230–239.
41. Hanna WC, Ko MA, Blitz M, Shargall Y, Comeau CG. Thoracoscopic Nuss procedure for young adults with pectus excavatum: excellent midterm results and patient satisfaction. *Ann Thorac Surg*. 2013;96(3):1033–1036; discussion 1037–1038.
42. Jacobsen EB, Thastum M, Jeppesen JH, Pilegaard HK. Health-related quality of life in children and adolescents undergoing surgery for pectus excavatum. *Eur J Pediatr Surg*. 2010;20(2):85–91.
43. Krasopoulos G, Dusmet M, Ladas G, Goldstraw P. Nuss procedure improves the quality of life in young male adults with pectus excavatum deformity. *Eur J Cardiothorac Surg*. 2006;29(1):1–5.
44. Kuru P, Bostanci K, Ermerak NO, Bahadır AT, Afacan C, Yuksel M. Quality of life improves after minimally invasive repair of pectus excavatum. *Asian Cardiovasc Thorac Ann*. 2015;23(3):302–307.
45. Steinmann C, Krille S, Mueller A, Weber P, Reingruber B, Martin A. Pectus excavatum and pectus carinatum patients suffer from lower quality of life and impaired body image: a control group comparison of psychological characteristics prior to surgical correction. *Eur J Cardiothorac Surg*. 2011;40(5):1138–1145.
46. Kelly RE, Jr. Pectus excavatum: historical background, clinical picture, preoperative evaluation and criteria for operation. *Semin Pediatr Surg*. 2008;17(3):181–193.
47. Krasopoulos G, Goldstraw P. Minimally invasive repair of pectus excavatum deformity. *Eur J Cardiothorac Surg*. 2011;39(2):149–158.
48. Metzelder ML, Kuebler JF, Leonhardt J, Ure BM, Petersen C. Self and parental assessment after minimally invasive repair of pectus excavatum: lasting satisfaction after bar removal. *Ann Thorac Surg*. 2007;83(5):1844–1849.
49. Lomholt JJ, Jacobsen EB, Thastum M, Pilegaard H. A prospective study on quality of life in youths after pectus excavatum correction. *Ann Cardiothorac Surg*. 2016;5(5):456–465.
50. Hokschi B, Kocher G, Vollmar P, Praz F, Schmid RA. Nuss procedure for pectus excavatum in adults: long-term results in a prospective observational study. *Eur J Cardiothorac Surg*. 2016;50(5):934–939.
51. Kelly RE, Goretsky MJ, Obermeyer R, et al. Twenty-one years of experience with minimally invasive repair of pectus excavatum by the Nuss procedure in 1215 patients. *Ann Surg*. 2010;252(6):1072–1081.
52. Tikka T, Kalkat MS, Bishay E, Steyn RS, Rajesh PB, Naidu B. A 20-year review of pectus surgery: an analysis of factors predictive of recurrence and outcomes. *Interact Cardiovasc Thorac Surg*. 2016;23(6):908–913.
53. Sacco Casamassima MG, Gause C, Goldstein SD, et al. Patient satisfaction after minimally invasive repair of pectus excavatum in adults: long-term results of Nuss procedure in adults. *Ann Thorac Surg*. 2016;101(4):1338–1345.
54. Tikka T, Webb J, Agostini P, et al. Pectus patient information website has improved access to care and patient reported outcomes. *J Cardiothorac Surg*. 2016;11(1):69.
55. Pilegaard HK. Single centre experience on short bar technique for pectus excavatum. *Ann Cardiothorac Surg*. 2016;5(5):450–455.
56. Pawlak K, Gasiorowski L, Gabryel P, Galecki B, Zielinski P, Dyszkiewicz W. Early and late results of the Nuss procedure in surgical treatment of pectus excavatum in different age groups. *Ann Thorac Surg*. 2016;102(5):1711–1716.
57. Ersen E, Demirkaya A, Kilic B, et al. Minimally invasive repair of pectus excavatum (MIRPE) in adults: is it a proper choice? *Wideochir Inne Tech Maloinwazyjne*. 2016;11(2):98–104.
58. Fibla JJ, Molins L, Moradiellos J, et al. Experience with the Nuss technique for the treatment of pectus excavatum in Spanish thoracic surgery departments. *Cir Esp*. 2016;94(1):38–43.
59. Park HJ, Kim KS, Lee S, Jeon HW. A next-generation pectus excavatum repair technique: new devices make a difference. *Ann Thorac Surg*. 2015;99(2):455–461.
60. Zhang DK, Tang JM, Ben XS, et al. Surgical correction of 639 pectus excavatum cases via the Nuss procedure. *J Thorac Dis*. 2015;7(9):1595–1605.
61. Park HJ, Kim KS, Moon YK, Lee S. The bridge technique for pectus bar fixation: a method to make the bar un-rotatable. *J Pediatr Surg*. 2015;50(8):1320–1322.
62. Rokitsky AM, Stanek R. Modified minimally invasive pectus repair in children, adolescents and adults: an analysis of 262 patients. *Pneumologia*. 2013;62(4):224–231.
63. Olbrecht VA, Arnold MA, Nabaweesi R, et al. Lorenz bar repair of pectus excavatum in the adult population: should it be done? *Ann Thorac Surg*. 2008;86(2):402–408; discussion 408–409.
64. Nuss D, Kelly RE Jr, Croitoru DP, Katz ME. A 10-year review of a minimally invasive technique for the correction of pectus excavatum. *J Pediatr Surg*. 1998;33(4):545–552.
65. Johnson WR, Fedor D, Singhal S. Systematic review of surgical treatment techniques for adult and pediatric patients with pectus excavatum. *J Cardiothorac Surg*. 2014;9:25.
66. Hebra A, Swoveland B, Egbert M, et al. Outcome analysis of minimally invasive repair of pectus excavatum: review of 251 cases. *J Pediatr Surg*. 2000;35(2):252–257; discussion 257–258.
67. Kanagaratnam A, Phan S, Tchantchaleishvili V, Phan K. Ravitch versus Nuss procedure for pectus excavatum: systematic review and meta-analysis. *Ann Cardiothorac Surg*. 2016;5(5):409–421.
68. Pilegaard HK. Extending the use of Nuss procedure in patients older than 30 years. *Eur J Cardiothorac Surg*. 2011;40(2):334–337.
69. Yoon YS, Kim HK, Choi YS, Kim K, Shim YM, Kim J. A modified Nuss procedure for late adolescent and adult pectus excavatum. *World J Surg*. 2010;34(7):1475–1480.
70. Pilegaard HK, Licht PB. Routine use of minimally invasive surgery for pectus excavatum in adults. *Ann Thorac Surg*. 2008;86(3):952–956.
71. Schalamon J, Pokall S, Windhaber J, Hoellwarth ME. Minimally invasive correction of pectus excavatum in adult patients. *J Thorac Cardiovasc Surg*. 2006;132(3):524–529.
72. Park HJ, Jeong JY, Jo WM, et al. Minimally invasive repair of pectus excavatum: a novel morphology-tailored, patient-specific approach. *J Thorac Cardiovasc Surg*. 2010;139(2):379–386.
73. Mansour KA, Thourani VH, Odessey EA, Durham MM, Miller JJ Jr, Miller DL. Thirty-year experience with repair of pectus deformities in adults. *Ann Thorac Surg*. 2003;76(2):391–395; discussion 395.
74. McMahon LE, Johnson KN, Jaroszewski DE, et al. Experience with FiberWire for pectus bar attachment. *J Pediatr Surg*. 2014;49(8):1259–1263.
75. Pilegaard HK. Nuss technique in pectus excavatum: a mono-institutional experience. *J Thorac Dis*. 2015;7 (Suppl 2):S172–S176.
76. Molik KA, Engum SA, Rescorla FJ, West KW, Scherer LR, Grosfeld JL. Pectus excavatum repair: experience with standard and minimal invasive techniques. *J Pediatr Surg*. 2001;36(2):324–328.
77. Park HJ, Lee SY, Lee CS, Youm W, Lee KR. The Nuss procedure for pectus excavatum: evolution of techniques and early results on 322 patients. *Ann Thorac Surg*. 2004;77(1):289–295.
78. Jaroszewski DE, Johnson K, McMahon L, Notrica D. Sternal elevation before passing bars: a technique for improving visualization and facilitating minimally invasive pectus excavatum repair in adult patients. *J Thorac Cardiovasc Surg*. 2014;147(3):1093–1095.
79. Park HJ, Chung WJ, Lee IS, Kim KT. Mechanism of bar displacement and corresponding bar fixation techniques in minimally invasive repair of pectus excavatum. *J Pediatr Surg*. 2008;43(1):74–78.

80. Tedde ML, de Campos JR, Wihlm JM, Jatene FB. The Nuss procedure made safer: an effective and simple sternal elevation manoeuvre. *Eur J Cardiothorac Surg*. 2012;42(5):890–891.
81. Takagi S, Oyama T, Tomokazu N, Kinoshita K, Makino T, Ohjimi H. A new sternum elevator reduces severe complications during minimally invasive repair of the pectus excavatum. *Pediatr Surg Int*. 2012;28(6):623–626.
82. Haecker FM, Sesia SB. Intraoperative use of the vacuum bell for elevating the sternum during the Nuss procedure. *J Laparoendosc Adv Surg Tech A*. 2012;22(9):934–936.
83. Johnson WR, Fedor D, Singhal S. A novel approach to eliminate cardiac perforation in the nuss procedure. *Ann Thorac Surg*. 2013;95(3):1109–1111.
84. Kim D, Idowu O, Palmer B, Kim S. Anterior chest wall elevation using a T-fastener suture technique during a nuss procedure. *Ann Thorac Surg*. 2014;98(2):734–736.
85. Park HJ, Jeong JY, Kim KT, Choi YH. Hinge reinforcement plate for adult pectus excavatum repair: a novel tool for the prevention of intercostal muscle strip. *Interact Cardiovasc Thorac Surg*. 2011;12(5):687–691.
86. Del Frari B, Schwabegger AH. How to avoid pectus bar dislocation in the MIRPE or MOVARPE technique: results of 12 years' experience. *Ann Plast Surg*. 2014;72(1):75–79.
87. Nuss D, Obermeyer RJ, Kelly RE. Nuss bar procedure: past, present and future. *Ann Cardiothorac Surg*. 2016;5(5):422–433.
88. Nuss D. Minimally invasive surgical repair of pectus excavatum. *Semin Pediatr Surg*. 2008;17(3):209–217.
89. Nagaso T, Miyamoto J, Kokaji K, et al. Double-bar application decreases postoperative pain after the Nuss procedure. *J Thorac Cardiovasc Surg*. 2010;140(1):39–44, 44.e1–2.
90. Stanfill AB, DiSomma N, Henriques SM, Wallace LJ, Vegunta RK, Pearl RH. Nuss procedure: decrease in bar movement requiring reoperation with primary placement of two bars. *J Laparoendosc Adv Surg Tech A*. 2012;22(4):412–415.
91. Dzielicki J, Korlacki W, Janicka I, Dzielicka E. Difficulties and limitations in minimally invasive repair of pectus excavatum – 6 years experiences with Nuss technique. *Eur J Cardiothorac Surg*. 2006;30(5):801–804.
92. Nagasao T, Hamamoto Y, Tamai M, et al. Scoring of deformed costal cartilages reduces postoperative pain after Nuss procedure for pectus excavatum. *Thorac Cardiovasc Surg*. 2016;64(1):62–69.
93. Pilegaard HK. Short Nuss bar procedure. *Ann Cardiothorac Surg*. 2016;5(5):513–518.
94. Ghionzoli M, Ciuti G, Ricotti L, et al. Is a shorter bar an effective solution to avoid bar dislocation in a Nuss procedure? *Ann Thorac Surg*. 2014;97(3):1022–1027.
95. Croitoru DP, Kelly RE Jr, Goretsky MJ, Gustin T, Keever R, Nuss D. The minimally invasive Nuss technique for recurrent or failed pectus excavatum repair in 50 patients. *J Pediatr Surg*. 2005;40(1):181–186; discussion 186–187.
96. Fallon SC, Slater BJ, Nuchtern JG, et al. Complications related to the Nuss procedure: minimizing risk with operative technique. *J Pediatr Surg*. 2013;48(5):1044–1048.
97. Al-Assiri A, Kravarusic D, Wong V, Dicken B, Milbrandt K, Sigalet DL. Operative innovation to the “Nuss” procedure for pectus excavatum: operative and functional effects. *J Pediatr Surg*. 2009;44(5):888–892.
98. Ravenni G, Actis Dato GM, Zingarelli E, Flocco R, Casabona R. Nuss procedure in adult pectus excavatum: a simple artifice to reduce sternal tension. *Interact Cardiovasc Thorac Surg*. 2013;17(1):23–25.
99. Papic JC, Finnell SM, Howenstein AM, Breckler F, Leys CM. Postoperative opioid analgesic use after Nuss versus Ravitch pectus excavatum repair. *J Pediatr Surg*. 2014;49(6):919–923; discussion 923.
100. Muhly WT, Maxwell LG, Cravero JP. Pain management following the Nuss procedure: a survey of practice and review. *Acta Anaesthesiol Scand*. 2014;58(9):1134–1139.
101. Kelly RE, Jr., Shamberger RC, Mellins RB, et al. Prospective multicenter study of surgical correction of pectus excavatum: design, perioperative complications, pain, and baseline pulmonary function facilitated by internet-based data collection. *J Am Coll Surg*. 2007;205(2):205–216.
102. Jaroszewski DE, Temkit M, Ewais MM, et al. Randomized trial of epidural vs. subcutaneous catheters for managing pain after modified Nuss in adults. *J Thorac Dis*. 2016;8(8):2102–2110.
103. Gebhardt R, Mehran RJ, Soliz J, Cata JP, Smallwood AK, Feeley TW. Epidural versus ON-Q local anesthetic-infiltrating catheter for post-thoracotomy pain control. *J Cardiothorac Vasc Anesth*. 2013;27(3):423–426.
104. Ried M, Schilling C, Potzger T, et al. Prospective, comparative study of the On-Q(R) PainBuster(R) postoperative pain relief system and thoracic epidural analgesia after thoracic surgery. *J Cardiothorac Vasc Anesth*. 2014;28(4):985–990.
105. Weber T, Matzl J, Rokitansky A, Klimscha W, Neumann K, Deusch E. Superior postoperative pain relief with thoracic epidural analgesia versus intravenous patient-controlled analgesia after minimally invasive pectus excavatum repair. *J Thorac Cardiovasc Surg*. 2007;134(4):865–870.
106. Futagawa K, Suwa I, Okuda T, et al. Anesthetic management for the minimally invasive Nuss procedure in 21 patients with pectus excavatum. *J Anesth*. 2006;20(1):48–50.
107. Hall Burton DM, Boretzky KR. A comparison of paravertebral nerve block catheters and thoracic epidural catheters for postoperative analgesia following the Nuss procedure for pectus excavatum repair. *Paediatr Anaesth*. 2014;24(5):516–520.
108. Jaroszewski DE, Gustin PJ, Haecker F-M, et al. Pectus excavatum repair after sternotomy: the Chest Wall International Group experience with substernal Nuss bars. *European Journal of Cardio-Thoracic Surgery*. 2017;52(4):710–717.

Patient Related Outcome Measures

Publish your work in this journal

Patient Related Outcome Measures is an international, peer-reviewed, open access journal focusing on treatment outcomes specifically relevant to patients. All aspects of patient care are addressed within the journal and practitioners from all disciplines are invited to submit their work as well as healthcare researchers and patient support groups.

Submit your manuscript here: <http://www.dovepress.com/patient-related-outcome-measures-journal>

Dovepress

The journal is included in PubMed. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.