

Methodical bias for comparison of periodontal ligament injection and local infiltration anesthesia for routine extractions in the maxilla

Peer W Kämmerer
Monika Daubländer

Department of Oral, Maxillofacial
and Facial Plastic Surgery, University
Medical Centre Mainz, Mainz,
Germany

Dear editor

We read the article by Al-Shayyab¹ with great interest, though we think that there is a methodical bias.

Usage of standard dental syringes with 27-gauge needles is not recommended for periodontal ligament (PDL) injections as they are very unlikely to achieve the correct pressure needed for successful single tooth anesthesia. In accordance with this, specialized syringes with short 30-gauge needles are commonly used all over the literature.² The author addresses this in the “Discussion” section and writes that “a standard conventional dental syringe was used in the present study, not a special PDL syringe, since the former is readily available in the clinic and proves equally successful when a standard 27-gauge short needle was used,” citing Malamed from 1982 (a time during which the modern PDL syringes were not developed yet³) and Madan et al who write that “intra-ligamentary injection technique is equally effective when a standard 27-gauge needle is used”.⁴ The second assumption refers to the needle only, not the syringe. In addition, this needle issue is not proven by any reference or study. Therefore, one might come to the conclusion that PDL was not carried out correctly. Also, the authors did not evaluate pulp or tissue anesthesia and started the extraction procedure after a latency period of 5 minutes in all cases. In accordance with this, the success rates of the PDL injection cannot be given, but would be of interest.

Whereas for infiltration anesthesia, 1.8 mL was injected buccally and an additional 0.3 mL on the palatal side, PDL consisted of 0.2 mL on the mesial aspect of each root of the tooth. In accordance with this, one may consider this as infiltration with additional palatal nerve block. Besides, it is difficult to estimate how a mesial injection was possible, for example, for a three-rooted maxillary molar. How did the author approach the mesial side of the distal root? Normally, PDL is recommended at least in the mesiolingual and distolingual aspects. In addition, the newer literature shows that the injection PDL in four sites is significantly more successful than in two sites.⁵

In conclusion, the study of Al-Shayyab rather compares infiltration plus palatal block with a special, unusual and not suitable form of PDL. Therefore, from our point of view, generalization of the results should not be carried out, and further studies on conventional PDL in the maxilla are needed.

Disclosure

The authors report no conflicts of interest in this communication.

Correspondence: Peer W Kämmerer
Department of Oral, Maxillofacial and
Facial Plastic Surgery, University Medical
Centre Mainz, Augustusplatz 2, 55131
Mainz, Germany
Email peer.kaemmerer@unimedizin-mainz.de

References

1. Al-Shayyab MH. Periodontal ligament injection versus routine local infiltration for nonsurgical single posterior maxillary permanent tooth extraction: comparative double-blinded randomized clinical study. *Ther Clin Risk Manag.* 2017;13:1323–1331.
2. Shabazfar N, Daubländer M, Al-Nawas B, Kämmerer PW. Periodontal intraligament injection as alternative to inferior alveolar nerve block—meta-analysis of the literature from 1979 to 2012. *Clin Oral Investig.* 2014;18(2):351–358.
3. Malamed SF. The periodontal ligament (PDL) injection: an alternative to inferior alveolar nerve block. *Oral Surg Oral Med Oral Pathol.* 1982; 53(2):117–121.
4. Madan GA, Madan SG, Madan AD. Failure of inferior alveolar nerve block: exploring the alternatives. *J Am Dent Assoc.* 2002;133(7): 843–846.
5. Lin S, Wigler R, Huber R, Kaufman AY. Anaesthetic efficacy of intraligamentary injection techniques on mandibular molars diagnosed with asymptomatic irreversible pulpitis: a retrospective study. *Aust Endod J.* 2017;43(1):34–37.

Author's reply

Mohammad H Al-Shayyab

Department of Oral and Maxillofacial Surgery, Oral Medicine and Periodontology, School of Dentistry, University of Jordan, Amman, Jordan

Correspondence: Mohammad H Al-Shayyab
Department of Oral and Maxillofacial Surgery, School of Dentistry,
University of Jordan, Amman 11942, Jordan
Tel +962 79 506 1513
Email mshayyab@hotmail.com

Dear editor

I would like to thank you for giving me the opportunity to respond to the concerns raised in Kämmerer and Daubländer's letter. I would also like to thank Kämmerer and Daubländer for giving their time to express their concerns and interest in my article.

In their letter to the editor, Kämmerer and Daubländer expressed potential concerns with the statement written in the "Discussion" section of my article, which should be read in light of the data from the three used references altogether, not separately. Malamed¹ that stated although "special" PDL syringes can be used effectively and safely, a conventional local anesthetic syringe is equally effective in providing PDL anesthesia. The use of a 30-gauge needle for PDL injection is too fragile to withstand the pressure without bending. However, great clinical success, with no increase in patient's discomfort, is achieved using the more readily available 27-gauge short needle.¹ I agree that differences between the two syringes, as well as the needles, may exist, but necessitate further studies rather than recommending one over another. Special dental syringes were dedicated to the PDL technique in the 1970s.² The success of PDL injection is judged when resistance to the deposition of anesthetic solution is felt.^{1,2} Pain during extractions can be evaluated after a 5-minute latency period without evaluating pulp or testing tissue anesthesia.³ This contradicted the recommendation in the letter referenced by a meta-analysis⁴ on PDL injection in the mandible, which is not necessarily applicable to the maxilla, and would indicate that the PDL in the study was carried out in accordance with the literature.^{1,3}

Palatal injection employed in the study cannot be considered as the palatal nerve block, as the anesthetic is not deposited

close to the greater palatine nerve trunk.³ For PDL injection, the local anesthetic was deposited on the mesial aspect of buccal and palatal roots, which provided adequate visibility of the injected site and maintained control of the needle.¹ However, the needle was repositioned and the injection repeated when no backward pressure was felt during injection.^{1,2} Furthermore, I would like to draw the attention of Kämmerer and Daubländer that their argument of two or four sites for PDL injection is based on a retrospective study⁵ related to PDL for molars diagnosed with asymptomatic irreversible pulpitis in the mandible, which is not necessarily applicable to posterior maxillary extractions. Such an argument has to be considered for future PDL anesthesia for maxillary teeth, rather than imposing one methodology over another.

In conclusion, the study still compares routine buccal and palatal infiltration with a special, usual and evidence-based form of PDL. Therefore, within the limitations of the study, generalization of the results should be carried out, and the conclusion of Kämmerer and Daubländer is not supported by specific references. However, I would agree that further studies on conventional PDL in the maxilla are needed.

Disclosure

The author reports no conflicts of interest in this communication.

References

1. Malamed SF. *Handbook of Local Anesthesia*. 6th ed. St Louis, MO: Elsevier; 2015:255.
2. Moore PA, Cuddy MA, Cooke MR, Sokolowski CJ. Periodontal ligament and intraosseous anesthetic injection techniques: alternatives to mandibular nerve blocks. *J Am Dent Assoc*. 2011;142(Suppl 3):13S–18S.
3. Fan S, Chen WL, Pan CB, et al. Anesthetic efficacy of inferior alveolar nerve block plus buccal infiltration or periodontal ligament injections with articaine in patients with irreversible pulpitis in the mandibular first molar. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2009;108(5):e89–e93.
4. Shabazfar N, Daubländer M, Al-Nawas B, Kämmerer PW. Periodontal intraligament injection as alternative to inferior alveolar nerve block: meta-analysis of the literature from 1979 to 2012. *Clin Oral Investig*. 2014;18(2):351–358.
5. Lin S, Wigler R, Huber R, Kaufman AY. Anaesthetic efficacy of intraligamentary injection techniques on mandibular molars diagnosed with asymptomatic irreversible pulpitis: a retrospective study. *Aust Endod J*. 2017;43(1):34–37.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Therapeutics and Clinical Risk Management 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Therapeutics and Clinical Risk Management editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Therapeutics and Clinical Risk Management

Dovepress

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peer-reviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS,

EMBASE, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/therapeutics-and-clinical-risk-management-journal>