

Letter to the editor regarding “To what extent are patients with migraine able to predict attacks?”

Xiang Fang^{1,*}

Weili Kong^{2,*}

Zeping Yu¹

Jianqing Qiu³

Hong Duan¹

¹Department of Orthopedics, West China School of Medicine/West China Hospital, Sichuan University, Chengdu, Sichuan, People's Republic of China;

²Department of Otolaryngology, West China Hospital, Sichuan University, Chengdu, Sichuan, People's Republic of China; ³Department of Epidemiology and Health Statistics, West China School of Public Health, Sichuan University, Chengdu, Sichuan, People's Republic of China

*These authors contributed equally to this work

Dear editor

With interest, we read the article by Gago-Veiga et al published in *Journal of Pain Research* in October 2018.¹ Migraine attacks, encompassing a wide range of symptoms, greatly undermine the quality of life for patients. Premonitory symptoms usually precede and alert the patients of the attack. The objectives of this prospective study¹ were to illuminate if any good predictor or specific combination of premonitory symptoms exists for prediction of migraine attacks. A total of 34 patients recording 229 attacks were analyzed: 67.6% were able to predict at least one attack, while only 35.3% were able to predict >50% of attacks. The positive predictive value was 85.1%. The authors concluded some specific symptoms were predictive, even though only a few were good predictors (predicting >50% of attacks).

However, certain issues regarding the methodology must be addressed. First, the sensitivity, specificity, positive predictive value, negative predictive value, positive likelihood ratio, negative likelihood ratio, and diagnostic odds ratio are the most suitable estimates to assess the validity of a test compared to the gold standard. Nevertheless, also reporting the diagnostic added value using receiver-operating characteristic curves should be considered, since all these validity estimates can be acceptable, whereas the diagnostic added value might be clinically negligible.² Second, correlation, even with statistical significance ($P < 0.01$), cannot guarantee prediction of a certain outcome. Furthermore, for prediction study, we usually need data from two distinct cohorts, or at least from one single cohort but divided into two, first to build a prediction model and then to validate it. Misleading results are generally the major outcome of research without validation of prediction models.^{2,3} Finally, in prediction study, we should evaluate the interactions between important variables.³ Final results could be affected dramatically if qualitative interactions are present, which means without evaluating interaction terms, prediction studies will mainly convey misleading messages.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Gago-Veiga AB, Pagán J, Henares K, et al. To what extent are patients with migraine able to predict attacks? *J Pain Res*. 2018;11:2083–2094.
2. Fang X, Kong W, Yu Z, et al. Letter to the Editor Regarding Can early postoperative FET-PET after the resection of a glioblastoma predict the location of later tumor recurrence. *World Neurosurg*. 2018; In publication.
3. Szklo M, Nieto FJ. *Epidemiology Beyond the Basics*. 2nd ed. Sudbury: Jones and Bartlett Publisher; 2007.

Correspondence: Hong Duan
Department of Orthopedics, West China School of Medicine/West China Hospital, Sichuan University, 37 Guo Xue Lane, Chengdu, Sichuan 610064, People's Republic of China
Tel +86 28 8542 2578
Fax +86 28 8542 2578
Email duanhong1970@126.com

Authors' reply

Ana B Gago-Veiga, Josué Pagán, Kevin Henares, Patricia Heredia, Nuria González-García, María-Irene De Orbe, Jose L Ayala, Mónica Sobrado, Jose Vivancos

Headache Unit, Department of Neurology, Instituto de Investigación Sanitaria Hospital Universitario de la Princesa, Madrid 28006, Spain

Correspondence: Ana B Gago-Veiga
Headache Unit, Department of Neurology, Instituto de Investigación Sanitaria Hospital Universitario de la Princesa, 62 Calle Diego de León, Madrid 28006, Spain
Tel +34 91 520 2416
Fax +34 91 520 2416
Email dra.anagago@gmail.com

Dear editor

We value Fang et al's comments, and thank them for the insights provided on our recent work. We understand some of the comments, but we would like to let you know why we decided to present the results in the most suitable way from the perspective of the analysis performed using machine-learning techniques.

With regard to the issue of, eg, sensitivity and specificity values, we also understand that these are important metrics, and most of Fang et al's requirements were already satisfied in the text. Let us notice that:

- sensitivity = TPR = recall
- PPV = precision
- ROC = ROC

Additionally, all these values, and much more, can be calculated from the confusion matrix. That was the reason we provided this matrix in the text.

Regarding the comment about prediction and the need for two distinct cohorts, let us clarify that our work was not on prediction from the computational point of view (the "time" variable is not present in the model), but on classification (AdaBoost is a classification algorithm). Also, our work uses cross-validation for the validation of the classification model as a computationally valid technique for the estimation of performance of a predictive model. In any case, we have repeated our experiments with two cohorts (70% for building the classification model, 30% for validation), and obtained better results than those presented with our safer cross-validation:

	μ	σ
Accuracy	86.3%	1.2%
Sensitivity = recall = TPR	80.3%	1.7%
Specificity = TNR	89.7%	0.9%
Precision = PPV	80.4%	2.0%
NPV	89.9%	1.0%
LR+	9.6	1.0
LR-	0.2	0
DOR corrected	8.5	1.4

Finally, regarding the comment about the interaction between "important variables", let us say that we already analyzed the effect of sociodemographic variables, and this was found negligible.

Once again, we thank Fang et al for taking the time to read our work and provide comments.

Disclosure

The authors report no conflicts of interest in this communication.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Pain Research 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Pain Research editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Pain Research

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.