

# RETRACTED ARTICLE: MicroRNA-506 regulates apoptosis in retinoblastoma cells by targeting sirtuin 1

This article was published in the following Dove Press journal:  
*Cancer Management and Research*

Zhidu Song<sup>1</sup>  
Hailiang Wang<sup>2</sup>  
Fangwei Zong<sup>1</sup>  
Chao Zhu<sup>1</sup>  
Ying Tao<sup>3</sup>

<sup>1</sup>Department of Ophthalmology, The Second Hospital of Jilin University, Changchun, Jilin 130022, People's Republic of China; <sup>2</sup>Department of Neurosurgery, The Second Hospital of Jilin University, Changchun, Jilin 130031, People's Republic of China; <sup>3</sup>Department of Anesthesiology, The Third Hospital of Jilin University, Changchun, Jilin 130033, People's Republic of China

**Background:** MicroRNAs have been reported to participate in the initiation and progression of retinoblastoma (RB), most common malignancy in children. The refractory mechanisms of chemoresistance and the toxicity of chemotherapy hindered the treatment especially on young children. Novel RB therapies are urgently required. miR-506 is proved to be associated with the tumorigenesis of various cancers while the role of this miRNA in RB remains unclear.

**Methods:** Here, the impact of miR-506 on RB cell survival in vitro and tumorigenesis in vivo was examined. miR-506 expression was examined in human RB samples and cell lines as compared with healthy tissues and non-RB cells. TUNEL staining and colony formation assay were performed to determine the effect of miR-506 on RB cell growth. TdT-mediated dUTP nick end labeling (TUNEL) staining and flow cytometry analysis were applied to detect the apoptotic cell number after miR-506 was downregulated in RB cells. Furthermore, dual-luciferase reporter assay was utilized to confirm the direct interaction between miR-506 and SIRT1 gene.

**Results:** miR-506 expression was upregulated in 20 human RB samples from patients as well as in human RB cell lines WERI-Rb1 and Y79, as compared to that in healthy tissues and non-RB cells. In contrast, the expression of sirtuin 1 (SIRT1), known as NAD-dependent deacetylase, was downregulated in RB samples and cell lines. Aberrant reduced miR-506 expression impaired survival and proliferation of WERI-Rb1 and Y79 cells. The depletion of miR-506 expression promoted apoptosis of the two RB cell lines. The results of bioinformatics analysis and dual-luciferase assay exhibited that miR-506 targeted the 3'-untranslated region of SIRT1 on silencing purpose. The SIRT1 silencing lessened the miR-506 inhibition on RB cell proliferation and undermined apoptosis.

**Conclusion:** The results provided an insight into the role of miR-506 during RB development and offered potential pharmaceutical strategy for RB diagnosis.

**Keywords:** retinoblastoma, miR-506, apoptosis, SIRT1

## Introduction

Retinoblastoma (RB), known as a severe cancer developed in the retina, affects mainly infants and young children under 5 years. The morbidity ratio remains to be ~1:15,000–1:20,000 and responsible for 2–4% of all childhood malignancy.<sup>1</sup> A case report of 2580 based on an epidemiological study was shown in 2015.<sup>2</sup> Various cellular agents and molecules were recognized to involve in proliferation, apoptosis and tumor cell cycle. It is speculated that genetic and epigenetic mutations occur when oncogenes and tumor suppressor genes induced or suppressed RB progression and carcinogenesis.<sup>3,4</sup>

MicroRNAs (miRs) are non-coding RNA molecules (length: <22–25 nucleotides) that exert post-transcriptional effects on the specific gene expression.<sup>5</sup> Increasing

Correspondence: Ying Tao  
Department of Anesthesiology, The Third Hospital of Jilin University, No.2 Xiantai Road, Changchun, Jilin 130033, People's Republic of China  
Tel +86 1 868 668 6675  
Fax +86 4 318 539 5237  
Email taoying.med@yahoo.com

evidences indicated the essential character of miRs on different cellular processes, including apoptosis. MiR-506 can affect cell growth, differentiation, cancer metastasis and Invasion. Consistently, miR-506 dysregulation investigation has been reported in various cancer-associated studies. The variation of upstream factors accompanied with promoter methylation moderated miR-506 expression.<sup>6</sup> In human ductal carcinoma cell lines, miR-506 served as a suppressor to alter Vimentin, CD151 and Snai2 gene expression to regulate epithelial-mesenchymal transition.<sup>7</sup> Similar phenomenon was characterized in HeLa and C33A cells, indicating miR-506 possessed an inhibition property against human cervical cancer.<sup>8</sup> It was also revealed that miR-506 targeted ETS1 to regulate gastric cancer angiogenesis and cell invasion.<sup>9</sup> Although miR-506 has been widely investigated in various tumor studies, researchers rarely probed on its role on eye cancer. Wu et al exhibited that miR-506-3p was significantly downregulated in RB tissues and cell lines. Dual-luciferase reporter assay showed that miR-506-3p directly targeted mitosis Gene A (NIMA)-related kinase 6 (NEK6) in RB cells. Enhanced expression of miR-506-3p remarkably suppressed cell proliferation, induced G0/G1 cell cycle phase arrest and apoptosis in RB cells, which were attenuated by NEK6 overexpression using MTT assay, colony formation and flow cytometry analysis.<sup>10</sup> Recent investigation has demonstrated that knockdown of long-non-coding RNA HOXA11-AS in RB cells suppressed cell proliferation notably. It, furthermore, induced cell cycle arrest at G1/G<sub>0</sub> phase and promoted cell apoptosis. HOXA11-AS served as a competing endogenous RNA that inhibited miR-506-3p expression, which regulated its downstream target NEK6 in RB.<sup>11</sup> Although the work has evaluated the role of miR-506-3p in RB, the detailed information on miR-506 function during the process is still lacking.

As a nicotinamide adenine dinucleotide (NAD<sup>+</sup>)-dependent lysine deacetylase, sirtuin 1 (SIRT1) is vital during metabolism, inflammation and aging. It has been proposed to regulate cell mass.<sup>12</sup> In recent cancer research, it is identified that SIRT1 served as an oncoprotein or a tumor suppressor upon distinctive circumstances. For instance, it enabled to influence on infiltrating immune lymphocytes.<sup>13</sup> Furthermore, an altered expression of SIRT1 led to poor prognosis of gastric cancer,<sup>14</sup> breast cancer<sup>15</sup> and soft tissue sarcoma,<sup>16</sup> respectively. Immunohistochemical analysis unraveled a complicated function of SIRT1 in tumorigenesis: 23 of 82 carcinomas demonstrated a lower level of SIRT1 expression in contrast to another 18 cancer cells among the same test group that exhibited an increased expression

manner relative to normal colonic mucosa.<sup>17</sup> Additionally, SIRT1 mRNA downregulated in gastric cancer.<sup>18</sup> Here, we examined how miR-506 is involved in RB generation. An upregulated MiR-506 expression was identified among RB cells and the specimens from RB patients. MiR-506 inhibition impaired RB cell proliferation as well as suppressed the tumor growth in vitro and in vivo. The depletion of miR-506 expression caused the induction of apoptosis in RB cells (WERI-Rb1 and Y79). Our results demonstrated that miR-506 suppressed apoptosis via suppressing SIRT1 expression in RB cell lines.

## Materials and methods

### RB samples

Retinal tissues were collected from 20 RB patients who did not undergo chemotherapy and radiation therapy prior to enucleation in the Department of Ophthalmology at the second Hospital of Jilin University, China. The patients consisted of 11 male and 9 female ranged from 0.6 to 9.1 years old (mean, 3.4). According to ICRB, the RB patients were divided into Phase I (5 cases), Phase II (9 cases) and Phase III (6 cases). The adjacent healthy tissues were set as the control. Written informed consent was acquired from each patient guardian. The experimental protocol was preapproved by the Medical Ethics Committee of the Second and the Third Hospital of Jilin University, China. Patient tissue samples were conducted in accordance with Declaration of Helsinki.

### Cell culture

WERI-RB1 and Y79 cell lines were obtained from Chinese Academy of Sciences Cell Bank (Shanghai, China), and then cultured to 70% confluence in RPMI-1640 (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (Hyclone, Logan, UT, USA), 100 U/mL penicillin, as well as 100 mg/mL streptomycin at 37°C under normoxic conditions of 100% humidity, 5% CO<sub>2</sub> and 95% air.

### miR-506 and SIRT1 inhibition

The miR-506 inhibitor, NC inhibitor (the scramble control of this inhibitor), SIRT1 siRNA and NC siRNA were obtained from Creative Biogene. Cells were transfected with 2 µg miR-506/NC inhibitor or 4 µg SIRT1/NC siRNA with Lipofectamine 2000 transfection reagent (Thermo Fisher). After cell transfection, the mature endogenous miR-506 was inhibited by the miR-506 inhibitor. Sodium

chloride (NaCl; 0.9%) was used for the preparation of miR-506 and NC inhibitors at 10 mg/mL (final concentration).

## Cell proliferation measurement

The cell proliferation was examined using cell-light 5-ethynyl-20-deoxyuridine (EdU) Apollo Imaging Kit (RiboBio, China). A fluorescence microscopy was applied for the measurement of the rate of EdU-positive cells.

## Western blot analysis

Cell lysates (20 g) was applied to sodium dodecyl sulfate polyacrylamide gel electrophoresis; after protein separation process, polyvinylidene fluoride membranes were utilized for transformation. The membranes were then treated and incubated with primary and secondary antibodies. SIRT1 (ab7343, Abcam, 1:2000) and GAPDH antibodies (ab9485, Abcam, 1:5000) were used for signal detection via Amersham ECL Western Blotting detection system.

## RNA extraction and quantitative polymerase chain reaction

Total RNA was extracted with TRIzol reagent using tissues or treated cells, as per the instructions of the manufacturer. Roche Light-Cycler 480 Real Time PCR system (Roche, Switzerland) and SYBR Green were utilized to examine the levels of miR-506 and SIRT1. U6 and glyceraldehyde-3-phosphate (GAPDH) performed as an internal control, respectively. The primer sequences were shown as follows: miR-506 F: 5'-GGG TAT TGA GCG AGG TGT T-3', R: 5'-CAG TGC GCG TCG TGG ATT-3'; U6 snRNA F: 5'-CTC GCT TCG GCA TCA CAT ATA CT-3', R: 5'-ACG CTT CAC GAA TTT TCG TGT C-3'; SIRT1 F: 5'-TGA TGC TAT TCG TGG AGA TAT TTT T-3', R: 5'-CAT GAT TGG CAG TGG CTC AT-3'; GAPDH F: 5'-AT GGT CAA CCG TTA GGA AA-3', R: 5'-TAA AGC GGT CAT TGA TGG CA-3'. The protocol of quantitative real-time PCR was designed at 95°C in 20 µL reaction volume for 10 mins, followed by 40 cycles at 95°C for 15 s, 60°C for 30 s and 72°C for 30 s. The endogenous reference was used for normalization and the amount of target ( $2^{-\Delta\Delta CT}$ ) was calculated by a calibrator (mean of the controls).

## Colony generation assay

Cells were transfected using various reagents. Cells were re-suspended in 10% FBS-added DMEM after 2 days of transfection and plated on an 8-mm layer of 0.4% top agar,

followed via transferring into 12-well plates with 0.5 mL of 0.5% bottom agar. In fortnight, four regions were randomly chosen from each plate and colonies were quantified.

## TdT-mediated dUTP nick end labeling assay

TUNEL assay was applied to examine the apoptosis-related changes in cells; the nuclei were stained with 4',6-diamidino-2-phenylindole (DAPI, 1:5000; Beyotime, China) and cells were stained using TUNEL fluorescence kit (Roche, Switzerland). The apoptosis of cells was determined via counting the TUNEL-positive cells under SP8 laser scanning confocal microscope (Leica, Germany).

## Analysis of apoptosis

Apoptotic cells were examined via flow cytometry (FC) assay and Annexin V/propidium iodide (PI) staining.<sup>19</sup>

## Dual luciferase reporter assay

Cells that underwent transfection were as follows: miR-506 mimic + SIRT1 WT; NC mimic + SIRT1 WT; miR-506 mimic + SIRT1 MU; and NC mimic + SIRT1 MU, in which SIRT1 WT and SIRT1 MU were synthesized by GeneBio Co. Ltd. (Guangzhou, China). MiR-506 target gene was determined by 3'-untranslated region (UTR) luciferase reporter assay, wherein MU and WT 3'-UTR of SIRT1 were utilized. The reporter luminescence (Rluc) was studied using Renilla luciferase sequence, whereas firefly luciferase sequence was used to examine the calibration luminescence (Luc). Cells were lysed in passive lysing buffer, and Firefly and Renilla luciferase activities were detected via a commercial Dual-Luciferase assay kit (Promega Corp, Madison, WI, USA) according to the manufacturer's instructions. The vectors and miRNA mimic were added into cells and the cells were cultured 24 hrs. Dual-luciferase reporter assay (DLRA) was used for luciferase activity examination.

## Animal tests

BALB/c-nu mice (female, five-week-age) were purchased from Vital River (Beijing, China). The suspension of Y79 cells ( $1 \times 10^6$ ) suspension was loaded in the microliter syringe. Colibri forceps (Storz, SI. Louis, MO) were performed to firmly control the sclera at the limbus area and rotate the globe anteriorly when the animal was appropriately anesthetized. The needle was then manually inserted into mouse sclera and gently rotated until the tip can be

viewed through the retina via a stereomicroscope. The tip was inserted advanced slightly further to elevate the retina when the plastic protective sheath hindered over-penetration of the needle and perforation of the neuroretina. With the bevel of the needle facing the globe, 3- to 4- $\mu$ l injection of cells was injected into the subretinal space. The needle was then quickly withdrawn after the injection and same procedure was repeated at a point 180° opposite to the first injection place on the eye. Typically, two micro-injections were conducted into each eye, in the retinal periphery, anterior to the eye's equator. A topical lubricant was placed on the cornea to prevent drying after the procedure. On day 28 after injection, the mice were sacrificed. All animal experiments were performed under the approval of Animal Operating Principal of the Institutional Animal Care and Use Committee in The Second and the Third Hospital of Jilin University, China (regulation guideline: The Legal Administration Documents for Laboratory Animal in People's Republic of China and Beijing Municipality).

## Statistical analysis

The data obtained in this study are represented as mean  $\pm$  standard deviation (SD). The comparisons between two test groups were examined by one-way analysis of variance (ANOVA), accompanied with the two-tailed Student's *t*-test.  $P < 0.05$  was considered for the detection of the statistically significant difference between groups.


## Results

### Expression of miR-506 was upregulated in RB specimens and RB cell lines

MiR-506 expression among RB specimens was examined using qPCR. According to ICRB guidelines, three groups of patients were selected as follows: good, medium and poor prognosis groups. The poor and medium prognosis groups showed a significantly upregulated miR-506 expression manner compared to the good prognosis group and the normal control group (Figure 1A). In addition, WERI-RB1 (RB1) and Y79 cells displayed increased miR-506 levels compared with the other tumor cell lines (Figure 1B). Therefore, miR-506 expression was upregulated in RB, indicating that the poor prognostic effect is associated with miR-506 expression.

### Inhibition of miR-506 suppressed tumorigenesis of RB in mice


To determine how miR-506 affects xenograft RB tumor formation, the Y79 cells that transfected with Adenoviral-miR-506 inhibitor, Adenoviral-SIRT1 siRNA, Adenoviral-miR-506 inhibitor+SIRT1 siRNA and normal Y79 cells were micro-injected into eyes of BALB/c mice, respectively. The tumor growth was daily monitored. MiR-506 expression and SIRT1 mRNA level were confirmed that occurred in mice tumor from each group and that was depleted in mice from miR-506 inhibition or SIRT1 silencing group (Figure 2A). These mice were executed on day 28 after injection and then the formed tumors were excised and weighed. The miR-506-silenced tumors developed



**Figure 1** miR-506 expression in RB specimens of patients and RB cell lines (RB1 and Y79). (A) miR-506 expression in RB specimens from good prognosis (n=5), medium prognosis (n=9), poor prognosis (n=6) and patients with normal volunteer (n=10), as examined using qPCR. (B) miR-506 expression was compared between RB cell lines and other cancer cell lines, as examined using qPCR. Data were represented in the form of mean  $\pm$  SD. \* $P < 0.05$ .

**Abbreviation:** RB, retinoblastoma.





**Figure 2** miR-506 silencing inhibits xenograft tumor formation. (A) Expression of miR-506 in the RB specimens of each group was examined by qPCR analysis at day 27 post injection. (B) Y79 cells transfected with Adenoviral-miR-506 inhibitor, Adenoviral-siRNA-SIRT1, Adenoviral-miR-506 inhibitor+siRNA-SIRT1 or control cells were injected into eyes of mice (n=10 per group). Mice were sacrificed and tumors were weighed at day 28 post-inoculation. Tumors were weighed after dissection from each group. (C) Tumor growth curve during 28 days post-inoculation. Data represent the mean  $\pm$  SD. \* $P$ <0.05. **Abbreviations:** RB, retinoblastoma.

with a markedly slower manner than control group. It displayed a smaller mean tumor volume in contrast to the control group, whereas the silencing of SIRT1 caused a restoration on tumor volume and weight as compared with inhibitor group (Figure 2B and C).

### Inhibition of miR-506 suppressed RB cell line proliferation and triggered apoptosis

The influence of miR-506 on RB cells was then investigated. In RB1 and Y79 cells, the miR-506 inhibitor

transfection led to a significant reduction in miR-506 level (Figure 3A and B). EdU assay results exhibited a remarkable decrease in the proliferation of these cell types after 24–72 hrs miR-506 inhibitor transfection in contrast to cells from the NC groups (Figure 3C and D). The colony formation assay results demonstrated that the aberrant downregulated expression of miR-506 induced a similar reduction on the colony numbers formed by these two cell types while transfection with NC inhibition presented no effect (Figure 3E and F).


The proliferation of these two RB cells was impaired following with the inhibition of miR-506. A speculative role of miR-506 seems to be that the depletion of the microRNA exerted a stimulatory effect on apoptosis in RB cell lines. Both TUNEL staining (Figure 4A and B) and FC (Figure 4C and D) assay results indicated an enhancement in apoptosis of RB1 and Y79 cell lines with miR-506 inhibitor transfection in comparison to that of cells in the NC groups.

### miR-506 targeted the 3'-UTR of SIRT1

miR-506 was predicted in the Bioinformatics analysis to target the 3'-UTR of SIRT1 (Figure 5A). The correlation of miR-506 to 3'-UTR of SIRT1 was investigated using Dual-Luciferase Assay (Figure 5B). The activity of luciferase declined at 50% in cells transfected with the miR-506 mimic fusing to the 3'-UTR of SIRT1 in comparison to the cells in the control groups. Subsequently, we examined the SIRT1 expression in RB specimens and RB cell lines. As exhibited in Figure 5C and D, SIRT1 level in RB cell lines and specimens decreased in contrast to that observed in normal healthy specimens and other cancer cells. The impacts of miR-506 inhibitor on SIRT1 expression between these two RB cell lines were examined by WB and qPCR. Both the protein and mRNA level of SIRT1 enhanced following with the cells that transfected with miR-506 inhibitor (Figure 5E–H). The results indicated that miR-506 targeted the 3'-UTR of SIRT1 gene.

### SIRT1 silencing counteracted the role of miR-506 on the proliferation and apoptosis of RB cells


To assess whether SIRT1 silencing neutralized the effect of miR-506 on RB cell proliferation and apoptosis, SIRT1 transcription was silenced in the RB1 and Y79 cells. WB and qPCR were used to verify the changes in the expression of SIRT1 (Figure 6A–D). The depletion of SIRT1 expression



**Figure 3** Effect of miR-506 expression on the proliferation of RB1 and Y79 cells. (A, B) Expression of miR-506 in RB1 and Y79 cells transfected with the miR-506 inhibitor or NC inhibitor, respectively, was examined by qPCR. (C, D) Proliferation rate of RB1 and Y79 cells was obtained at 24, 48 and 72 hrs post-transfection using the EdU assay. (E, F) Colony formation assay of RB1 and Y79 cells transfected with the miR-506 inhibitor or NC inhibitor. The lower panel indicated the number of colonies formed in each group. Data were represented as the form of mean  $\pm$  SD. \* $P < 0.05$ , \*\* $P < 0.01$ .

revived the proliferation of the RB1 and Y79 cells repressed by miR-506 inhibition, as shown from the results of EdU assay (Figure 6E and F). Cytometry assay was then performed to examine the apoptosis rate of RB cell lines. Aberrant downregulated expression of SIRT1 caused a visible decreased amount of apoptotic cells corresponding to the transfection with the miR-506 inhibitor (Figure 6G and H). The observable decline of the apoptotic cells subjected to

SIRT1 silencing. Previous studies indicated that Bax and p53 induced apoptosis while NF- $\kappa$ B p65 played an inhibitory role on apoptosis. To further evaluate the mechanism of SIRT1 on apoptosis of RB cell, we examined p53, p65 and Bax expression after transfection. It revealed that inhibition of miR-506 induced the upregulation of both p53 and Bax; however, it led to downregulation of p65. Moreover, the silencing of SIRT1 reversed the effect of miR-506 on the expression of



**Figure 4** Overexpression of miR-506 enhanced the apoptosis of RB cell lines. (A, B) TUNEL staining was carried out to examine the number of apoptotic cells in each group of RB1 and Y79 cells. Magnification,  $\times 200$ . TUNEL staining cells from each group were displayed in the lower panel. (C, D) Annexin V-FITC/PI staining and FACS were performed to determine the number of apoptotic cells. The upper right quadrant of each plot represents early apoptotic cells and apoptotic rate analysis in RB1 and Y79 cells from each group was shown in the lower panel. Data were represented in the form of mean  $\pm$  SD. \* $P < 0.05$ .


**Abbreviations:** RB, retinoblastoma; TUNEL, TdT-mediated dUTP nick end labeling.

these three proteins (Figure 6I and J). Thus, SIRT1 silencing enabled to reverse the inhibition effect of miR-506 on the proliferation and apoptosis of RB cells.

## Discussion

Tumorigenesis is considered as a synergetic intricate process accompanied with several anti-oncogenes and oncogenes. In

the present work, an increased miR-506 expression was observed in RB cells and tumor specimens. It also revealed that miR-506 enables to suppress the expression of SIRT1, the key modulator in malignancies. MiR-506 inhibition caused a suppression of cell viability and proliferation. It promoted apoptosis on both RB1 and Y79 cells. Further investigation demonstrated the interaction between miR-506 and the 3'-UTR



**Figure 5** SIRT1 was a direct target of miR-506. **(A)** Graphical representation of the conserved miR-506 binding motif at the 3'-UTR of SIRT1. **(B)** Luciferase activity displayed by the luciferase reporter constructs carrying either the wild-type (WT) or mutated (MU) human SIRT1 3'-UTR after miR-506 mimic transfection. The luciferase activity was normalized to the activity of pRL-TK-Luc. **(C, D)** qPCR analysis was used to confirm the downregulation during SIRT1 expression in RB specimens and cell lines. **(E, F)** WB and qPCR were performed to examine SIRT1 protein and mRNA expressions, respectively, after transfection of RB1 and Y79 cells with miR-506 inhibitor and NC inhibitor. Data are represented in the form of mean  $\pm$  SD. \* $P < 0.05$ .


**Abbreviation:** RB, retinoblastoma.

of SIRT1 gene, resulting in the abrogation of SIRT1 expression. The knockout of SIRT1 expression neutralized miR-506 on RB cell proliferation. Our findings suggested that miR-506 acted as oncogenic factor by targeting the SIRT1 gene.

The regulating effect of miRNAs on downstream target gene expression has been studied in various cancer types. Multiple target genes downregulated the translation or triggered the cleavage of molecules through the binding of miRs to their 3'-UTR. Therefore, we aimed to examine

the expression of miR-506-regulated downstream target genes. Previous research has demonstrated that miR-506 might target different genes, including YAP, ROCK1, CDK4/6-FOXO, IQGAP1 and SPHK1 which mediated various malignancy progressions.<sup>20-24</sup> It was shown that miR-506 is supposed to target the conservative site of SIRT1 according to bioinformatics analysis. DLRA results further confirmed that 3'-UTR of SIRT1 directly interacted with miR-506 in RB cells. SIRT1 possessed anti-apoptotic





**Figure 6** Silencing of SIRT1 expression counteracted the effect of miR-506 on RB cell lines. (A–D) Expression of SIRT1 was examined in RB1 and Y79 cells at both protein and mRNA levels following transfection with different agents. SIRT1i represents interference of SIRT1 siRNA, NCi represents transfection with NC siRNA. (E, F) SIRT1 silencing recovered the proliferation of RB1 and Y79 cells. (G, H) SIRT1 silencing eliminated the pro-apoptotic effect of the miR-506 inhibitor. Annexin V-FITC/PI staining and FC were carried out to examine the number of early apoptotic RB1 and Y79 cells after 48 hrs of transfection. (I, J) WB was performed to examine the expression of p53, p65 and Bax after 48 hrs of transfection. Results were presented as mean  $\pm$  SD. \* $P$ <0.05, \*\* $P$ <0.01.

**Abbreviation:** RB, retinoblastoma.

property and involved in diverse cellular process including autophagy, proliferation, apoptosis and aging to affect tumorigenesis. It is speculated that the function of miR and SIRT1 interaction varies from the specific tumor-associated oncogenic pathway. For instance, miR-22 acted as an oncogene/tumor-suppressor in prostate cancer and SIRT1 was a target agent to miR-22. Another study showed that SIRT1 overexpression in breast cancer cells induced the downregulated expression of MiR-22.<sup>25</sup> The expression of miR-138 in NSCLC declined due to SIRT1 neutralization function on this tumor suppressor gene. AMP-activated protein kinase (AMPK)-mTOR pathway exerted a vital role in adjusting autophagy. MiR-138 reduced SIRT1 expression to undermine the AMPK pathway activity and subsequently promoted the mTOR phosphorylation.<sup>26</sup> In prostate cancer, downregulation of miR-221 or miR-222 expression suppressed cell proliferation and migration as well as enhanced apoptosis,<sup>27</sup> following with an induced SIRT1 expression. A study implied that downregulation of SIRT1 has the potential to boost migration and reduce cell death,<sup>28</sup> suggesting SIRT1 exhibited an inhibitive feature toward the oncogenic activity of miR-221 and miR-222. The escalated manner of SIRT1 expression succeeding miR-506 inhibition was determined in our study. The knock-down of SIRT1 expression on RB cells recovered the properties of RB cells regulated by miR-506 inhibition. The above evidence indicated that miR-506 contributed essentially to the RB tumorigenesis, modulated by SIRT1.

## Conclusion

Our work proposed the function of miR-506 as an RB oncogene during RB progression. The downregulated miR-506 expression in RB cell line is associated with tumor cell growth impairment, suggestive of its potential role as a biomarker or prognostic agent for RB diagnosis.

## Author contributions

All authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agreed to be accountable for all aspects of the work.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Broadbush E, Topham A, Singh AD. Incidence of retinoblastoma in the USA: 1975–2004. *Br J Ophthalmol*. 2009;93(1):21–23. doi:10.1136/bjo.2008.138750
2. Fernandes AG, Pollock BD, Rabito FA. Retinoblastoma in the United States: a 40-year incidence and survival analysis. *J Pediatr Ophthalmol Strabismus*. 2018;55(3):182–188. doi:10.3928/01913913-20171116-03
3. Beta M, Venkatesan N, Vasudevan M, Vetrivel U, Khetan V, Krishnakumar S. Identification and insilico analysis of retinoblastoma serum microRNA profile and gene targets towards prediction of novel serum biomarkers. *Bioinform Biol Insights*. 2013;7(BBI):S10501. doi:10.4137/BBI.S10501
4. Wang J, Wang X, Wu G, Hou D, Hu Q. MiR-506-3p, down-regulated in retinoblastoma, regulates cell cycle progression and apoptosis of human retinoblastoma cells by targeting PAX6. *FEBS Lett*. 2013;587(12):1779–1786. doi:10.1016/j.febslet.2013.04.029
5. Bartel DP. MicroRNAs: genomic, biogenic, mechanistic, and function. *Cell*. 2004;116(2):281–297. doi:10.1016/j.cell.2002.08.004
6. Li J, Ju J, Ni B, Wang J. The emerging role of miR-506 in cancer. *Oncotarget*. 2016;7(38):6258–6273. doi:10.18632/oncotarget.11294
7. Arora H, Qureshi P, Park W. miR-506 regulates epithelial mesenchymal transition in breast cancer cell lines. *PLoS One*. 2013;8(5):e64273. doi:10.1371/journal.pone.0064273
8. Gong M, Chen C, Zou H, Sun M, Song M. miR-506 suppresses cervical cancer cell proliferation both in vitro and in vivo. *Neoplasma*. 2018;65(3):331–338. doi:10.4149/neo\_2018\_170112N25
9. Vannucci FA, Linhares DC, Barcellos DE, Lam HC, Collins J, Barthaler D. Identification and complete genome of seneca valley virus in vesicular fluid and sera of pigs affected with idiopathic vesicular disease, Brazil. *Transbound Emerg Dis*. 2015;62(6):589–593. doi:10.1111/tbed.12410
10. Chen L, Chen Z, Xing Y. MiR-506-3p inhibits cell proliferation, induces cell cycle arrest and apoptosis in retinoblastoma by directly targeting NEK6. *Cell Biol Int*. 2018. doi:10.1002/cbin.11041
11. Han N, Zuo L, Chen H, Zhang C, He P, Yan H. Long non-coding RNA homeobox A11 antisense RNA (HOXA11-AS) promotes retinoblastoma progression via sponging miR-506-3p. *Oncotargets Ther*. 2019;12:3509–3517. doi:10.2147/OTT.S195404
12. Sebastian C, Satterstrom FK, Haigis MC, Mostoslavsky R. From sirtuin biology to human diseases: an update. *J Biol Chem*. 2012;287(51):42444–42452. doi:10.1074/jbc.R112.402768
13. Yang H, Bi Y, Xue L, et al. Multifaceted modulation of SIRT1 in cancer and inflammation. *Crit Rev Oncog*. 2015;20(1–2):49–64. doi:10.1615/CritRevOncog.2014012374
14. Cha EJ, Noh SJ, Kwon KS, et al. Expression of DBC1 and SIRT1 is associated with poor prognosis of gastric carcinoma. *Clin Cancer Res*. 2009;15(13):4453–4459. doi:10.1158/1078-0432.CCR-08-3329
15. Lee H, Kim KR, Noh SJ, et al. Expression of DBC1 and SIRT1 is associated with poor prognosis for breast carcinoma. *Hum Pathol*. 2011;42(2):204–213. doi:10.1016/j.humpath.2010.05.023
16. Kim JR, Moon YJ, Kwon KS, et al. Expression of SIRT1 and DBC1 is associated with poor prognosis of soft tissue sarcomas. *PLoS One*. 2013;8(9):e74738. doi:10.1371/journal.pone.0074738
17. Kabra N, Li Z, Chen L, et al. Sirt1 is an inhibitor of proliferation and tumor formation in colon cancer. *J Biol Chem*. 2009;284(27):18210–18217. doi:10.1074/jbc.M109.000034
18. Yang Q, Wang B, Gao W, et al. SIRT1 is downregulated in gastric cancer and leads to G1-phase arrest via NF-kappaB/Cyclin D1 signaling. *Mol Cancer Res*. 2013;11(12):1497–1507. doi:10.1158/1541-7786.MCR-13-0214
19. Peng DY, Song H, Liu LB. Resveratrol-downregulated phosphorylated liver kinase B1 is involved in senescence of acute myeloid leukemia stem cells. *J Huazhong Univ Sci Technol Med Sci*. 2015;35(4):485–489. doi:10.1007/s11596-015-1457-7

20. Hua K, Yang W, Song H, et al. Up-regulation of miR-506 inhibits cell growth and disrupt the cell cycle by targeting YAP in breast cancer cells. *Int J Clin Exp Med*. 2015;8(8):12018–12027.
21. Li D, Cao Y, Li J, Xu J, Liu Q, Sun X. miR-506 suppresses neuroblastoma metastasis by targeting ROCK1. *Oncol Lett*. 2017;13(1):417–422. doi:10.3892/ol.2016.5442
22. Liu G, Sun Y, Ji P, et al. MiR-506 suppresses proliferation and induces senescence by directly targeting the CDK4/6-FOXO1 axis in ovarian cancer. *J Pathol*. 2014;233(3):308–318. doi:10.1002/path.4348
23. Sun G, Liu Y, Wang K, Xu Z. miR-506 regulates breast cancer cell metastasis by targeting IQGAP1. *Int J Oncol*. 2015;47(5):1963–1970. doi:10.3892/ijo.2015.3161
24. Lu Z, Zhang W, Gao S, et al. MiR-506 suppresses liver cancer angiogenesis through targeting sphingosine kinase 1 (SPHK1) mRNA. *Biochem Biophys Res Commun*. 2015;468(1–2):8–13. doi:10.1016/j.bbrc.2015.11.008
25. Yi G, He Z, Zhou X, et al. Low concentration of metformin induces a p53-dependent senescence in hepatoma cells via activation of the AMPK pathway. *Int J Oncol*. 2013;43(5):1503–1510. doi:10.3892/ijo.2013.2077
26. Ye Z, Fang B, Pan J, et al. miR-138 suppresses the proliferation, metastasis and autophagy of non-small cell lung cancer by targeting Sirt1. *Oncol Rep*. 2017;37(6):3244–3252. doi:10.3892/or.2017.5619
27. Yang X, Yang Y, Gan R, et al. Down-regulation of mir-221 and mir-222 restrain prostate cancer cell proliferation and migration that is partly mediated by activation of SIRT1. *PLoS One*. 2014;9(6):e98833. doi:10.1371/journal.pone.0098833
28. Fu M, Liu M, Sauve AA, et al. Hormonal control of androgen receptor function through SIRT1. *Mol Cell Biol*. 2006;26(21):8122–8135. doi:10.1128/MCB.00289-06

RETRACTED

## Cancer Management and Research

Dovepress

### Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>