

Transcranial magnetic stimulation for the treatment of anxiety disorder

This article was published in the following Dove Press journal:
Neuropsychiatric Disease and Treatment

Priscila Aparecida

Rodrigues¹

Ana Luiza Zaninotto^{1,2}

Iuri Santana Neville¹

Cintya Yukie Hayashi¹

André R Brunoni³

Manoel Jacobsen

Teixeira¹

Wellingson Silva Paiva¹

¹Department of Neurology, University of São Paulo, São Paulo, Brazil; ²Laboratory of Neuromodulation, Harvard Medical School, Boston, MA, USA; ³Institute of Psychiatry, University of São Paulo, São Paulo, Brazil

Abstract: Anxiety is currently one of the main mood changes and can impair the quality of life of the individual when associated with other neurological or psychiatric disorders. Neuromodulation has been highlighted as a form of treatment of several pathologies, including those involving anxiety symptoms. Among the neuromodulatory options with the potential to improve mood changes, we highlight repetitive transcranial magnetic stimulation (rTMS). rTMS is a viable therapeutical option for neuropsychiatric dysfunctions of high prevalence and is important for the understanding of pathological and neuropsychological adaptation processes. Even with this potential, and high relevance of intervention, we observe the scarcity of literature that covers this subject. The objective of this study was to carry out a survey of the current literature, using scientific databases for the last five years. We found 32 studies reporting the effects of rTMS on anxiety, 7 on anxiety disorders and 25 on anxiety symptoms as comorbidities of neurological or psychiatric disorders. This survey suggests the need for further studies using TMS for anxiety in order to seek strategies that minimize these anxiety effects on the quality of life of the victims of this disorder.

Keywords: transcranial magnetic stimulation, anxiety disorders, review, treatment

Introduction

Individuals may experience anxiety as a warning sign in unknown situations, especially in response to fear and anticipation of danger.¹ However, it is considered pathological when it directly affects the individual's quality of life, affecting social relations, cognitive function, and the wake-sleep cycle.²⁻⁴ Anxiety disorders represent one of the major psychiatric disorders today that can impair the quality of life of adults.²⁻⁴

The Diagnostic and Statistical Manual of Mental Disorders – Fifth Edition (DSM-5) defines the presence of an anxiety disorder when some criteria are met, for example: symptoms occurring more than six months, excessive anxiety and worry, panic attacks, restlessness or feeling nervous, fatigue, irritability, sleep and eating disorder. Based on the symptomatology, anxiety disorders can be classified in: Generalized Anxiety Disorder (GAD), Social Anxiety Disorder, Panic Disorder (PD), Agoraphobia, Separation Anxiety Disorder, Selective Mutism and Specific Phobias. In addition to anxiety disorders, anxiety can be a symptom or comorbidity in several other pathologies, such as Major Depression,⁵ Obsessive Compulsive Disorder,⁶ TBI,⁷ among others.

Neuromodulation minimizes the impact of mood changes⁸⁻¹¹ and repetitive transcranial magnetic stimulation (rTMS) is receiving attention in the last

Correspondence: Priscila Aparecida Rodrigues
Caring - Clinic of Psychology and Neuropsychology, Avenue Vereador Narciso Yague, Guimarães, Number 1145 – Sala 1610, Mogi das Cruzes, São Paulo 08780-500, Brazil
Tel +55 11 199 654 2972
Email pri.ar@outlook.com

decade.^{12,17} TMS is a non-invasive method of stimulating the motor cortex neurons through the scalp and skull based on the principle of electromagnetic induction.^{18,19} TMS was approved by the Food and Drug Administration (FDA) in 2008 as an alternative treatment for Major Depression Disorder,²⁰ and has been shown to decrease the symptoms of Post Traumatic Stress Disorder (PTSD),²¹ Obsessive Compulsive Disorder (OCD),²² and Anxiety Disorders.¹⁴

Non-invasive brain stimulation techniques allow researchers to study in real-time the human brain activity, characterize the balance of excitation and cortical inhibition, and help guide plastic changes.⁸ With TMS, you can apply repetitive induced current pulses that can increase or decrease cortical excitability and stimulate the process of neuronal plasticity.^{9,23,24} A coil of wire is placed on the scalp generating an electric current that flows through the target area, inducing neuronal depolarization. Possible effects are depended on subject's age, pharmacological treatment, number of technical parameters, including the intensity and number of stimulations (ie, frequency), coil orientation, focus, and depth of stimulation.^{9,25,26}

Although TMS pulses (simple and repetitive) are considered safe, it is contraindicated for people who use a pacemaker or other implantable electronic devices.^{21,27} Patients with bone defects and craniotomy pose another concern, because the conductance of the structures would be modified.^{21,24} Some adverse effects might occur, such as headache and minor muscle spasms at the stimulation site.^{21,25,28,29}

Recent literature reviews discussed the use of TMS as an intervention strategy for anxiety. Vicario et al's³⁰ study addressed the use of TMS and transcranial direct current stimulation (tDCS) and other studies portray its efficacy for other psychiatric disorders, without major insights.^{14,30–32} However, there is a gap in the literature regarding rTMS efficacy in anxiety symptoms (primary and secondary outcomes) in clinical trials. Our study focused on the effectiveness of TMS for Anxiety Disorders and as an intervention for Anxiety Symptoms arising from other pathologies. We make the division between anxiety arising from Neurological Disorders and Psychiatric Disorders. Our hypothesis was that high-frequency rTMS on the dorsolateral prefrontal cortex (DLPFC) will decrease anxiety symptoms in patients with anxiety disorders, considering anxiety as the primary outcome.

Methods

We conducted a literature online search including Web of Science Medline/PubMed MEDLINE databases. We have included publications from any time until March 2019. We included clinical trial and open-label using the keywords: “TMS”, “transcranial magnetic stimulation”, “noninvasive brain stimulation”, and “anxiety”. The first exclusion was by title and followed by abstracts and full texts. Abstracts and full text, and studies were included if fulfilled, the inclusion criteria: (a) use of rTMS as intervention; (b) anxiety was assessed as primary or secondary outcome; (c) sample was adults; (d) published in peer-review journals; (e) full text written in English. The exclusion criteria: (a) animal studies; (b) case report; (c) systematic review or meta-analysis; (d) paper not written in English; (e) study with healthy; and (f) studies that did not report the use of rTMS in anxiety symptoms. Searching and data analysis were performed by Rodrigues PA and Zaninotto AL. This method follows PRISMA guidelines.

Results

Figure 1 shows the systematic review that initially 358 papers were found (Pubmed =172; Web of Science =186). We found 32 studies that fulfilled our eligibility criteria.

For a better understanding of the results, we divided the target population of the trials in two categories: Category 1) patients with Anxiety Disorders, according to DSM-5 (Table 1); Category 2) Anxiety symptoms, from neurological and psychiatric disorders that also evaluated anxiety symptoms as comorbidity (Table 2).

Treatment for anxiety disorders

In our search, five studies were randomized double-blind clinical trial (RCT) and two studies were open-label. Seven papers described Anxiety Disorders as the main outcome for rTMS,^{5,33–38} one was related to Anxiety Disorders, two were Panic Disorder (PD), and the other four were GAD (Table 1). The sample size of these studies ranged from 13 to 25 patients.

The right dorsolateral prefrontal cortex (DLPFC) was the most frequent stimulation region and in one study they stimulated the left region of the DLPFC. The predominant frequency used was 1 Hz, although one study compared the use of 1 hz and 10 hz and another used only 20 Hz.

The number of sessions ranged between 10 and 30 and the number of pulses per stimulation ranged from 750 to 3600. Most papers used an intensity of 110% of the rest




Figure 1 Flow diagram of the papers selected in our study, following PRISMA statement guidelines.

motor threshold (RMT), one used 80% and other used 90% of RMT and one study did not specify.

Treatment for other psychiatric neurological diseases

We found 23 studies reporting rTMS as an alternative for the treatment of neurological or psychiatric disorders and as secondary intervention to anxiety symptoms (Table 2).^{6,21,39–61}

Papers that cited symptoms of anxiety as comorbidity were grouped in Neurological Disorders (Table 3)^{40,45,47,51,54,55,59} and Psychiatric Disorders (Table 4).^{6,21,39,41–44,46,48–50,52,53,56–58,60,61} We found that MDD is the psychiatric disorder with the highest occurrence appearing in seven papers while pain is the highest Neurological Disorders occurrence appearing in three papers.

Fourteen papers reported the stimulation in the left DLPFC, three papers both left and right DLPFC and in one study the left auditory cortex was stimulated. Two papers focused the stimulation in the left temporoparietal

region and left primary motor cortex. The stimulation target of right DLPFC appeared in three papers, in addition to those already mentioned. These regions were stimulated in one paper each: primary motor cortex contralateral to the amputated leg, motor cortex area of the frontal lobe, and the epileptogenic focus. Two studies described stimulation in the left temporoparietal region and left primary motor cortex.

High pulse frequency (>5 Hz) used in rTMS protocols can have an excitatory effect while a low frequency (1 Hz) has an inhibitory effect. These effects are not limited to the target of stimulation, favoring the improvement of mood symptoms since there is complex connectivity of the cerebral cortex with other deep brain regions, favoring, ultimately, the improvement of mood symptoms. Most studies analyzed in this review used 10 stimulation sessions and half used excitatory stimulation for all subjects with rTMS frequency range between 5 Hz and 20 Hz. In the treatment of anxious symptoms, studies reported using between 4 and 31 sessions with 200–5000 pulses per stimulation and an intensity of 20–130% of the RMT.

Table 1 Patients with anxiety disorders, according to DSM 5

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	NP tests	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Zhang, 2019 ³⁷	117 rTMS	Mood or Anxiety Disorders	RCT	Left DLPFC	10	10 Hz	2400	120% of RMT	HAMA	All the age groups showed significant improvements in clinical symptoms	No safety or tolerability concerns were identified	Yes for adolescents, but not in adults
Lu R, 2018 ³⁸	28 rTMS	GAD	Open- label	Right DLPFC + left DLPFC	10	1 Hz	750	80% of RMT	HRSA	In this study, low- frequency rTMS which was successively delivered to right and left DLPFC effectively alleviated anxiety symptoms in GAD patients	All patients tolerated rTMS well without any adverse effects	Yes
Dilkov D, 2017 ³³	9 rTMS/ 12 sham	GAD	RCT	Right DLPFC	25	20 Hz	3600	110% of RMT	HRSA	Participants receiving rTMS showed clinically significant decrease in reported anxiety symptoms.	One participant in the active group had seizure; transient dizziness was reported in three patients	Yes
Diefenbach GJ, 2016 ³⁵	13 rTMS/12 sham	GAD	RCT	Right DLPFC	30	1 Hz	900	90% of RMT	HRSA; DASS	ANOVA demonstrated significant group x time interaction for the primary and secondary outcomes with the gains maintained only in the rTMS group at follow-up.	No seizures occurred; mild adverse effects were reported and was similar in rTMS and sham groups	Yes

(Continued)

Table 1 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	NP tests	Relevant results following treatment	Tolerability and safety	Sustained effect of response
White D and Tavakoli S, 2015 ⁵	13 rTMS	Comorbid MDD and GAD	Open- label, pilot study	Right DLPFC + left DLPFC	2436	1Hz (right side)+10Hz (left side)	1000	NR	GAD-7	Paired T-test comparing baseline to final scores showed a markedly significant difference for both GAD-7 and the HDRS-21, suggesting a decrease of 65% and 75% in depression and anxiety, respectively.	NR	Not applicable
Mantovani A, 2013 ³⁴	12 rTMS/13 sham	PD and comorbid MDD	RCT. Two phases: Phase I 4-week double- blind followed by Phase II (optional) 4-week open-label	Right DLPFC	Phase I: 20; Phase II: 20	1Hz	1800	110% of RMT	PDSS; HRSA- 14	Repeated-measures ANOVA revealed a significant time by group interaction with PDSS, with greater reduction in active group.	No seizures were reported. None of the patients reported significant side effects	Yes
Prasko J, 2007 ³⁶	7 rTMS/ 8 sham	PD or PD with agoraphobia	RCT	Right DLPFC	10	1 Hz	1800	110% of RMT	HRSA; BAI; PDSS	No statistical difference between groups were observed after treatment.	No seizures, headaches, neurological and cognitive decline occurred	Not applicable

Abbreviations: BAI, Beck Anxiety Inventory; DASS, Depression Anxiety Stress Scales; DLPFC, Dorsolateral Prefrontal Cortex; GAD, Generalized Anxiety Disorder; GAD-7, Generalized Anxiety Disorder scale; HAMA, Hamilton Rating Scale for Anxiety; HRSA, Hamilton Rating Scale for Anxiety; MDD, major depressive disorder; NP, neuropsychological; NR, Not Reported; PD, Panic Disorder; PDSS, Panic Disorder Severity Scale; RCT, Randomized Controlled Trial; RMT, Resting Motor Threshold; rTMS, repetitive Transcranial Magnetic Stimulation.

Table 2 Anxiety symptoms, from neurological and psychiatric disorders

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Kaur M, 2019 ³⁹	16 rTMS	Mood disorders	Open- label	Left DLPFC	20	10 Hz	NR	110% of RMT	Anxiety subscale of the HDRS	To determine whether sleep-wake disturbance, cognition or depression chronicity are associated with rTMS outcome in young depressed adults	Show reduced depressive and anxiety symptoms and no cognitive deterioration	NR	Yes
Lin H, 2018 ⁴⁰	7 rTMS	Thalamic pain	Open- label	Motor cortex	10	10 Hz	1000	90% of RMT	HAMA	To evaluate the analgesic lasting effect to treat thalamic pain	High-frequency rTMS can provide long-term pain relief in patients with thalamic pain	All patients tolerated rTMS well with out any adverse effects	No
Reyes- Lopez J, 2018 ⁴¹	29 rTMS	Borderline	RCT	Right DLPFC + left DLPFC	15	5 Hz or 1 Hz	900 or 1500	100% of RMT	HRSA	To evaluate clinical improvement in patients with BPD after application of rTMS	HAM-A scores reduced after rTMS treatment. Both protocols produced global improvement in severity and symptoms of BPD, particularly in impulsiveness, affective instability, and anger	NR	Yes
Durmaz O, 2017 ⁴²	36 rTMS	MDD	Open- label	Left DLPFC	15	20 Hz	1000	110% of RMT	HRSA	To evaluate the efficacy of rTMS in patients with treatment-resistant major depression	The findings suggested that comorbid anxiety symptoms, particularly somatic anxiety, could predict the response to rTMS in treatment-resistant major depressive disorder	Headache in eight patients, dizziness in four patients, and lacrimation in three patients. Only one patient dropped out of the study due to side effects or intolerance	Yes
Noh TS, 2017 ⁴³	17 rTMS	Tinnitus	RCT	Left auditory cortex (AC) and left DLPFC or only the left DLPFC	4	1 Hz	2000 or 3000	110% of RMT	STAI	We evaluated treatment outcomes following single-site rTMS in the DLPFC and dual-site rTMS in the AC and DLPFC	Group 1 exhibited significant improvements in scores on the STAI for both state anxiety and trait anxiety at 12 weeks posttreatment. Group 2 showed an improvement in only the STAI- X2 score at 12 weeks posttreatment.	NR	No

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Tovar- Perdomo S, 2017 ⁴⁴	24 rTMS	MDD	Open- label, pilot study	Left DLPFC	20	10 Hz	3000	120% of RMT	BAI	To explore the effects of a course of accelerated high- frequency rTMS on two neurocognitive domains in patients with MDD	Depression and anxiety scores significantly improved from pre-post HF-rTMS treatment.	The absence of practice effects in our longitudinal design raises the possibility that rTMS may also have cognitive side effects which, like antidepressant effects, may recede and reveal cognitive improvements after treatment cessation and sustained recovery	No
Elbeh KAM, 2016 ⁵	30 rTMS/15 sham	OCD	RCT	Right DLPFC + left DLPFC	10	1 Hz or 10 Hz	200 or 500	100% of RMT	HRSA	To evaluate the impact of different frequencies of rTMS in OCD	1 Hz rTMS over the right DLPFC has medium-term effect on obsessive-compulsive symptoms and anxiety	All patients tolerated rTMS well without any adverse effects except for two patients in the active stimulation group who experienced transient headache that disappeared with in a few hours	Yes
Malavera A, 2016 ⁴⁵	27 rTMS/27 sham	Pain	RCT	Primary Motor Cortex contralateral to the amputated leg	10	10 Hz	1200	90% of RMT	Zung Self- Rating Anxiety Scale	The score change in the Visual Analogue Scale for pain	No statistically significant between-group difference was found when comparing the absolute scores of the depression and anxiety scales at day 15 or day 30	Some patients experienced minor adverse effects such as headache, neck pain, and sleepiness without significant differences between groups	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Billci S, 2015 ⁴⁶	45 rTMS/30 sham	Tinnitus	RCT	Left temporoparietal region	10	1 Hz or 10 Hz	600 or 900	110% of RMT	BAS	To compare the effects of rTMS and paroxetine on tinnitus in terms of effectiveness and medium-term results	The positive effects that were observed might reflect a relationship between tinnitus and auditory cortex areas related to emotions	In the 10 Hz TMS treatment group, two patients complained of neck and shoulder stiffness, which disappeared in 2 days without any medical treatment; two patients had jaw fasciculation, which continued for 1 hr; and one patient had a headache for 1 day. One patient in the 1 Hz rTMS group had mild jaw pain for 2 days. All patients in the paroxetine-treated group were asked to report possible side effects of the drug. Among those, three had cephalgia, two had gastrointestinal disorders, and one had sexual dysfunction; all these side effects were mild-to-moderate in intensity and did not delay the therapy	Yes
Lin YC, 2015 ⁴⁷	14 rTMS	RLS	Open- label	Motor cortex area of the frontal lobe	14	15 Hz	600	100% of RMT	HRSA	To investigate whether rTMS could have any beneficial effects in restless legs syndrome	rTMS can markedly alleviate the motor system symptoms, sleep disturbances, and anxiety in RLS patients	No adverse effects were observed during stimulation or after treatment, and all patients showed good compliance	Yes
Oznur T, 2014 ⁴⁸	20 rTMS	PTSD	Open- label	Right DLPFC	20	1 Hz	NR	80% of RMT	BAI	To examine the effectiveness of rTMS in patients with treatment-resistant posttraumatic stress disorder	The effectiveness of rTMS on the anxiety and depression scores in the patients was not determined in this study	NR	No
Diefenbach GJ, 2013 ⁴⁹	32 rTMS	MDD	Open- label	Left DLPFC	31	10 Hz	3000– 5000	80 at 130% of RMT	Anxiety/ somatization subscale of the HDRS	To determine if anxious depression is associated with attenuated response to rTMS	Both depression and anxiety symptoms improved from pre- to post-treatment with moderate to large treatment effects	NR	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Pretalli JB, 2012 ⁵⁰	75 rTMS	MDD	RCT	Left DLPFC	10	10 Hz	NR	95% of RMT	HAMD anxiety subscale (items 9, 10, 11, and 15); Tyrer scale for anxiety; STAI	To investigate whether or not RMT changes during the treatment of resistant depression	These RMT changes influenced the outcome of the 10 sessions concerning the severity of depressive and anxiety symptoms	Only mild side-effects were reported (pain at the site of the coil placement or headache)	Yes
Sun W, 2012 ⁵¹	60 rTMS	Focal epilepsy	RCT	Epileptogenic focus	10	0.5 Hz	1500	90% or 20% of RMT	SCL-90	To evaluate the therapeutic effect of low-frequency rTMS on patients with refractory partial epilepsy	All the subscale scores of somatization, obsession- compulsion, interpersonal sensitivity, depression, anxiety, paranoid ideation and psychoticism in group 1 was lower than that of group 2 when evaluated at the end of follow-up period	The most common adverse events were mild or moderate headache and tinnitus. Adverse events occurred more often in group one patients who received "90% rMT" rTMS	Yes
Watts BV, 2012 ⁵¹	10 rTMS/10 sham	PTSD	RCT	Right DLPFC	10	1 Hz	NR	90% of RMT	STAI	This study seeks to examine the efficacy of rTMS for PTSD	Anxiety symptoms showed improvement with rTMS, but those improvements were not statistically significant compared with sham	NR	Yes
Berlim MT, 2011 ⁵²	15 rTMS	MDD	Open- label	Left DLPFC	20	10 Hz	3000	120% of RMT	HRSA; BAI	To address an augmenting strategy in subjects with chronic, severely treatment-resistant MDD	Clinically meaningful improvements in anxious and depressive symptoms	Only one of the 15 participants withdrew from the study at week 1 because of lack of tolerability (ie, severe scalp pain)	Yes
Boggio PS, 2010 ⁵³	20 rTMS/10 sham	PTSD	RCT	Right DLPFC or left DLPFC	10	20 Hz	1600	80% of RMT	HRSA	To investigate the efficacy of rTMS for the relief of posttraumatic stress disorder (PTSD)- associated symptoms	Right rTMS generated a significant improvement in the measure of anxiety at days 5 and 10 while left rTMS did not	There were no seizures and only mild adverse effects such as mild headache, neck pain, sleepiness, and dizziness were reported similarly in the 3 groups of treatment	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Epstein CM, 2007 ⁵⁴	14 rTMS	Parkinson disease	Open- label	Left DLPFC	10	10 Hz	1000	110% of RMT	HRSA	To investigate about rTMS as a potential treatment for depression in PD and for the movement disorder of PD	Open rTMS treatment of PD patients with treatment- resistant depression was followed by highly significant improvement in mood scores and anxiety ratings	There were no seizures and no complaints of headache or neurological deterioration	Yes
Passard A, 2007 ⁵⁵	15 rTMS/15 sham	Fibromyalgia	RCT	Left primary motor cortex	10	10 Hz	2000	80% of RMT	Hospital Anxiety Scale	To assess the effects of unilateral rTMS of the motor cortex on chronic widespread pain in patients with fibromyalgia	The analgesic effects were observed from the fifth stimulation onwards and were not related to changes in mood or anxiety.	Minor and transient side effects were reported during the stimulation period only. Nine patients reported headaches four in the active- stimulation group and five in the sham stimulation group. These headaches, reported after only 1 of the 10 daily sessions, were mild and transient in all cases. Other side effects included nausea after the fifth session in one patient in the active-treatment group. Two patients reported transient tinnitus and one patient reported mild dizziness after one sham-stimulation session	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Rossi S, 2007 ⁵⁶	14 rTMS	Tinnitus	Open- label	Left temporoparietal region	5	1 Hz	1200	120% of RMT	HRSA	To assess about rTMS on chronic tinnitus in which also eventual mood changes are monitored	In patients with chronic tinnitus, psychiatric comorbidity as mood or anxiety disorders are relevant and may partly found their functional counterpart in the activation of higher-order processing	The majority of patients did not complain of side effects due to rTMS, a part a slight transient headache on the stimulation site, which however did not require pharmacological treatment. About 30% of patients complained of tongue paraesthesia occurring during the active rTMS. Most of patients reported a transient worsening of their tinnitus in the first two-three days of active rTMS. Two male patients, one receiving active rTMS and one sham rTMS as first intervention, dropped out from the study for this reason	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Cohen H, 2004 ⁵⁷	18 rTMS/6 sham	PTSD	RCT	Right DLPFC	10	1 Hz or 10 Hz	NR	80% of RMT	HRSA	To evaluate the therapeutic effects of two different frequencies of active rTMS of PTSD patients	Active 10-Hz rTMS, relative to 1-Hz treatment and sham, significantly reduced Hamilton anxiety scale scores but not Hamilton depression scale scores	Headache was the main side effect reported, regardless of stimulation group. It was reported by 14 patients: eight patients reported headache after one rTMS treatment, five patients reported it after two sessions, and one (receiving sham treatment) reported it after three sessions. In most cases, this side effect was reported several hours after the stimulation or on the following morning. Only four patients reported headache immediately after the stimulation. In three cases, headache was a symptom before the study. However, the total number of headaches after treatment was 21. Two patients receiving high-frequency rTMS reported neck pain and muscular contraction in the area. Another patient receiving high-frequency treatment reported an exacerbation of previously existing dizziness. One patient in the group receiving slow-frequency rTMS and one patient from the high- frequency group developed a manic episode; in both cases, this occurred after the third session of rTMS. One patient reported a mild rage attack, probably related to the stimulation. Although we did not use earplugs, only two patients reported ear discomfort, which lasted less than 1 min	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Loo CK, 2003 ⁵⁸	18 rTMS	MDD	RCT	Left DLPFC	NR	15 Hz or 1 Hz	NR	90% of RMT	7-item scale	To understand further the mechanisms of action of high- and low- frequency rTMS by examining their acute effects on regional cerebral blood flow (rCBF) in depressed patients	Anxiety scores did not differ between the 15 Hz and 1 Hz groups. rTMS can produce functional changes in areas of the brain involved in mood control, including changes tending toward correction of deficits associated with depression	NR	Yes
Münchau A, 2002 ⁵⁹	12 rTMS	GTS	Open- label	Left premotor	6	1 Hz	1200	80% of RMT	Hospital Anxiety Scale	To study whether rTMS targeted to left motor and premotor cortex can improve tics in GTS	There was no significant improvement of symptoms after any of the rTMS conditions as assessed	One patient reported mild headache after premotor rTMS. Two patients reported excessive tiredness after both premotor and motor rTMS lasting for about 1 day	Yes
George MS, 2000 ⁶⁰	20 rTMS/10 sham	MDD	RCT	Left prefrontal cortex	10	5 Hz or 20 Hz	NR	100% of RMT	HRSA	To conduct a study to address whether 2 weeks of daily TMS over the left prefrontal cortex has antidepressant activity greater than sham	Expressed as a percent change from baseline, active TMS subjects had significantly greater improvement on the BDI as well as the HARS than did those who received sham	Two subjects with average MTs (60% and 70% of machine output) elected to stop the study because of the pain of stimulation. One subject tried for 2 consecutive days, whereas the other decided to stop after only 2 mins of the first session. Ten subjects reported mild headaches following at least one session (beginning immediately after to 3 hrs after rTMS), which were relieved by acetaminophen	Yes

(Continued)

Table 2 (Continued).

Author/ Year	Patient/ Sham group	Diagnosis	Study design	Target	# of rTMS sessions	Frequency	Pulses per session	Stimulation Intensity	Anxiety questionnaire	Primary outcome	Relevant results following treatment	Tolerability and safety	Sustained effect of response
Rollnik JD, 2000 ⁶¹	6 rTMS/ 6 sham	Schizophrenia	RCT	Left DLPFC	10	20 Hz	800	80% of RMT	STAI	To investigate the therapeutic efficacy of rTMS in schizophrenic patients with acute exacerbation of their psychosis	STAI, BDI, and NCT scores tended to improve during active rTMS and to worsen during sham stimulation, but the observed changes were not significant	NR	Yes

Abbreviations: STAI, State-Trait Anxiety Inventory; HRSA, Hamilton Rating Scale for Anxiety; BAI, Beck Anxiety Inventory; HAMA, Hamilton Rating Scale for Anxiety; RMT, Resting Motor Threshold; rTMS, repetitive Transcranial Magnetic Stimulation; DLPFC, dorsolateral prefrontal cortex; MDD, major depressive disorder; PTSD, Post-Traumatic Stress Disorder; RLS, Restless Legs Syndrome; OCD, Obsessive Compulsive Disorder.

Table 3 Symptoms of anxiety present in neurological disorders

Primary pathology	Number of studies
Pain	3 (Lin et al, 2018 ⁴⁰ Malavera et al, 2016 ⁴⁵ Passard et al, 2007 ⁵⁵)
Restless Legs Syndrome	1 (Lin et al, 2015 ⁴⁷)
Focal epilepsy	1 (Sun W, 2012 ⁵¹)
Parkinson's disease	1 (Epstein et al, 2007 ⁵⁴)
Gilles de la Tourette Syndrome	1 (Münchau et al, 2002 ⁵⁹)

Table 4 Symptoms of anxiety present in psychiatric disorders

Primary pathology	Number of studies
Borderline MDD	1 (Reyes-Lopez. J et al, 2018 ⁴¹) 8 (Kaur et al, 2019 ³⁹ Durmaz et al, 2017 ⁴² Tovar-Perdomo et al, 2017 ⁴⁴ Diefenbach et al, 2013 ⁴⁹ Pretalli et al, 2012 ⁵⁰ Berlim et al, 2011 ⁵² Loo et al, 2003 ⁵⁸ George et al, 2000 ⁶⁰)
Tinnitus	3 (Noh et al, 2017 ⁴³ Bilici et al, 2015 ⁴⁶ Rossi et al, 2007 ⁵⁶)
Obsessive Compulsive Disorder	1 (Elbeh et al, 2016 ⁶)
Post Traumatic Stress Disorder	4 (Oznur et al, 2014 ⁴⁸ Watts et al, 2012 ²¹ Boggio et al, 2010 ⁵³ Cohen et al, 2004 ⁵⁷)
Schizophrenia	1 (Rollnik et al, 2000 ⁶¹)

Abbreviation: MDD, major depressive disorder.

Measures of anxiety symptoms

One of the criteria used to include the paper in our systematic review was to contain measures to assess the symptoms of anxiety pre and post rTMS sessions. The Hamilton Rating Scale for Anxiety (HRSA) was the main scale found in this review; overall, this scale appeared in 14 papers; secondly the Beck Anxiety Inventory (BAI) and the State-Trait Anxiety Inventory (STAI), both appearing in four papers. In Table 5, we observe the other scales of assessment of the symptoms of anxiety found and their frequency in the papers.

Clinical findings

We observed, in general, that most of the studies found satisfactory results with the use of rTMS in Anxiety Disorders and Anxiety as comorbidity.

In the studies on Anxiety Disorders, it was observed that three papers reported a sustained effect of response on

Table 5 Anxiety scales found in papers

Anxiety questionnaire papers	Number of citations
HRSA	15
BAI	4
STAI	4
Hospital Anxiety Scale	2
PDSS	2
Anxiety/somatization subscale of the HDRS	2
HAMA	2
7-item scale	1
BAS	1
DASS	1
GAD-7	1
HAMD anxiety subscale (items 9, 10, 11 and 15)	1
SCL-90	1
Tyrer scale for anxiety	1
Zung Self-Rating Anxiety Scale	1

Abbreviations: HRSA, Hamilton Rating Scale for Anxiety; BAI, Beck Anxiety Inventory; STAI, State-Trait Anxiety Inventory; PDSS, Panic Disorder Severity Scale; HAMA, Hamilton Rating Scale for Anxiety; DASS, Depression Anxiety Stress Scales; GAD, Generalized Anxiety Disorder.

the improvement of anxious symptoms, one paper reported improvement of symptoms, although this response did not sustain in the long term and one paper reported that there was no significant improvement.

The papers on Anxious Symptoms showed that most of the studies obtained the sustained effect of response on the anxiety symptoms observed in 21 studies. Two studies showed improvement but not sustained over time and one of them did not find a positive response to the improvement in anxiety symptoms.

Tolerability and safety

Adverse effects were minimal, showing that there is safety in the application of rTMS in anxious symptoms independent of the primary outcome.

In the papers on Anxiety Disorders, only one study reported that one participant in the active group had seizures and three participants reported transient dizziness; in the other papers, no adverse effects were reported.

In the papers on Anxious Symptoms, mild side effects were observed. The most frequent effect was a headache that appeared in 12 studies. In two studies, some patients were withdrawn because of side effects or treatment intolerance. Table 6 has the description and frequency of other side effects found in the papers on Anxious Symptoms.

Table 6 Side effects of rTMS found in papers

Side effects	Frequency in papers
Dizziness	4
Pain and stiffness in the neck and shoulder	2
Ringing in the ear	3
Drowsiness	2
Tearing	1
Regression in cognitive improvement during treatment	1
Neck pain	1
Fasciculation and mandibular pain	1
Gastrointestinal disorders	1
Sexual dysfunction	1
Muscle contraction	1
Rage attack	1
Ear discomfort	1
Excessive fatigue	1

Discussion

The rTMS is already established as a non-invasive valid alternative to measure plastic alterations of the cerebral cortex for the treatment of depression,² borderlines,⁴¹ OCD,⁶ among other pathologies. We observed in this review that anxiety is one of the main symptoms related to current mood disorders, with a direct impact on individuals' quality of life.⁷

TMS have a direct correlation between the parameters associated with cortical excitability and plasticity, suggesting the existence of mechanisms that partially overlap and probably act in the same neurophysiological framework.⁶²

Pennisi et al suggest that lesion in the ischemic sub-cortical and prefrontal region might have implications in cognition and mood and may result in functional changes of the intracortical system, in addition to associating an increase in global cortical excitability, together with a significant worsening of frontal lobe abilities, but without substantial functional impairment.⁶²

Cortical plasticity plays a clear and fundamental role in patients with depression, while this still seems to be firmly established in anxiety disorders.⁶³

Studies observed that impaired brain plasticity may be one of the pathophysiological mechanisms underlying cognitive decline and major depression.⁶³ Some degree of cognitive impairment is often observed across the clinical spectrum of mood disorders, and between depression and cognition often bidirectional.⁶³ Depressed patients have

significant differences between the brain hemispheres, the intracortical neurochemical circuit (inhibitory or excitatory) might be unbalanced, the excitability of the motor cortex and TMS may indicate a disruption of plasticity.⁶³ The data suggest that the motor cortex is more refractory to modulatory inputs from other non-motor areas within the CNS in depressed individuals.⁶³ In addition, several studies have shown that functional abnormalities in cortical connections may play crucial roles in patients with depression and other mood disorders.⁶³

Studies with depressive disorder usually predict anxiety symptoms as a secondary outcome. It is possible to observe how this mechanism occurs in cases of anxiety disorders, since it is directly related to the functioning of neurotransmitters and to the cerebral circuits. TMS may interfere directly in this functioning and may influence the regulation of anxiety as a symptom arising from another psychopathology or as an anxiety disorder.⁶³

Other studies^{14,30,32} have found that TMS have promising results as a treatment for anxiety disorders; we observed comparable results, showing TMS intervention having positive effects on anxiety symptoms. However, there is still no standard intervention protocol for the use of TMS, neither for anxiety or depression disorders, as a comorbidity of psychiatric and neurological disorders.^{14,30,32}

A few studies describe protocols for TMS application on anxiety disorders, Dilkov et al³³ evidenced the effects of rTMS on participants with GAD. Most participants who received treatment from 25 rTMS sessions had a clinically significant decrease in anxiety symptoms as identified by the efficacy symptom scale.³³

Another study also observed positive results in subjects with GAD after 30 rTMS sessions according to pre- and post-treatment comparative assessment scales.³⁵ However, the Sham coil group also had reduced anxiety symptoms; they had a limited sample size and further assessment is necessary.³⁵

The authors used 2 sessions of rTMS in the prefrontal ventromedial cortex region for the treatment of Acrophobia and fear of irrational stature.⁶⁴ Their study used virtual reality technology to assist in the therapy and they observed a better response for fear and symptoms.⁶⁴

There is no clear pattern for using high or low frequencies of TMS stimulation in which regions of the brain. One review study cited the use of both high- and low-frequency TMS stimulation under the right and left stimulation target DLPFC.³⁰ Iannone et al¹⁴ cited papers that used both high and low frequencies of TMS on anxiety disorders. Another review study found that the right and left DLPFC were

targets for TMS stimulation in most papers.³¹ However, when searching for anxiety as a secondary outcome, the stimulation region was dependable on the primary outcome of the TMS intervention. TMS stimulation for pathologies involving motor aspects poses another difficulty.⁶⁵ In these cases, the stimulation target was primarily in specific regions of the motor cortex.⁶⁵

Patients with Restless Legs Syndrome (RLS) who receive TMS intervention had improvements on anxiety symptoms with the improvement of the RLS symptoms.⁴⁷ A similar study found that using TMS to stimulate the motor cortex area alleviated motor sensory complaints of RLS patients. TMS excitation and inhibition rates indicate intracortical injury and corticospinal imbalance, involving GABAergic and glutamatergic circuits, as well as impairment of the short-term mechanisms of plasticity.⁶⁵ The activation induced by rTMS with the consequent increase in dopamine release may have contributed to the clinical and neurophysiological outcome in those patients.⁶⁵ The occurrence of anxiety in patients with RLS is related to an abnormal sensorimotor integration, suggesting an interrupted connectivity in the RLS.⁶⁶ Using TMS in the sensory cortex motor region may promote improvements of physical symptoms and therefore reduction in anxiety.⁶⁶

TMS has also been shown to be effective as a sustained response effect compared to drug intervention. It is observed that drug-resistant patients on MDD treatment achieved improvement of symptoms with the use of TMS after a period of six months post-intervention, while this period is reduced when only medication is used.⁶⁷

Although there is no consensus in the TMS intervention parameters for anxiety disorders and symptoms due to neurological or psychiatric disorders, we observed that the DLPFC region is preferred among researcher's stimulation target. TMS have promising results in high frequency, promoting excitatory stimulation, and low frequency, promoting inhibitory stimulation. The number of sessions should range from 10 to 20 sessions so that the sustained response effect is possible. The stimulation of the motor cortex, especially in the sensorimotor region, also shows promising results in the stimulation of anxiety symptoms due to neurological disorders, mainly focusing on motor aspects.

Conclusion

We observed in this review that TMS might be a satisfactory intervention measure to improve anxiety, although there are a limited number of reports on the use of this intervention. In addition, satisfactory results have also

noted that TMS is a safe treatment strategy with low side effect. Further studies using TMS could direct the treatment with psychological or psychiatric intervention.

Disclosure

Miss Cintya Yukie Hayashi reports grant from Coordination for the Improvement of Higher Education Personnel (CAPES) – Brazil, outside the submitted work. The authors report no other conflicts of interest in this work.

References

- Cabrera CC, Sponholz Júnior A. Ansiedade e insônia. In: *Prática Psiquiátrica No Hospital Geral: Interconsulta E Emergência*. Porto Alegre, Brazil: Artmed Editora; 2005:283–304.
- Young AS, Klap R, Sherbourne CD, Wells KB. The quality of care for depressive and anxiety disorders in the United States. *Arch Gen Psychiatry*. 2001;58(1):55–61. doi:10.1001/archpsyc.58.1.55
- Borkovec TD, Newman MG, Pincus AL, Lytle R. A component analysis of cognitive-behavioral therapy for generalized anxiety disorder and the role of interpersonal problems. *J Consult Clin Psychol*. 2002;70(2):288. doi:10.1037/0022-006X.70.2.288
- Balconi M, Ferrari C. Left DLPFC rTMS stimulation reduced the anxiety bias effect or how to restore the positive memory processing in high-anxiety subjects. *Psychiatry Res*. 2013;209:554–559. doi:10.1016/j.psychres.2013.03.032
- White D, Tavakoli S. Repetitive transcranial magnetic stimulation for treatment of major depressive disorder with comorbid generalized anxiety disorder. *Ann Clin Psychiatry*. 2015;27(3):192–196.
- Elbeh KAM, Elserogy YMB, Khalifa HE, Ahmed MA, Hafez MH, Khedr EM. Repetitive transcranial magnetic stimulation in the treatment of obsessive-compulsive disorders: double blind randomized clinical trial. *Psychiatry Res*. 2016;238:264–269. doi:10.1016/j.psychres.2016.02.031
- Gould KR, Ponsford JL, Spitz G. Association between cognitive impairments and anxiety disorders following traumatic brain injury. *J Clin Exp Neuropsychol*. 2014;36(1):1–14. doi:10.1080/13803395.2013.863832
- Bashir S, Mizrahi I, Weaver K, Fregni F, Pascual-Leone A. Assessment and modulation of neural plasticity in rehabilitation with transcranial magnetic stimulation. *PM&R*. 2010;2(12):S253–S268. doi:10.1016/j.pmrj.2010.10.015
- Kobayashi M, Pascual-Leone A. Transcranial magnetic stimulation in neurology. *Lancet Neurol*. 2003;2(3):145–156.
- Barker AT. An introduction to the basic principles of magnetic nerve stimulation. *J Clin Neurophysiol*. 1991;8(1):26–37.
- Hallett M. Transcranial magnetic stimulation and the human brain. *Nature*. 2000;406(6792):147–150. doi:10.1038/35018000
- Kim YI, Kim SM, Kim H, Han DH. The effect of high-frequency repetitive transcranial magnetic stimulation on occupational stress among health care workers: a pilot study. *Psychiatry Investig*. 2016;13(6):622–629. doi:10.4306/pi.2016.13.6.622
- Sagliano L, D'Olimpio F, Panico F, Gagliardi S, Trojano L. The role of the dorsolateral prefrontal cortex in early threat processing: a TMS study. *Soc Cogn Affect Neurosci*. 2016;11(12):1992–1998. doi:10.1093/scan/nsw105
- Iannone A, Cruz APM, Brasil-Neto JP, Boechat-Barros R. Transcranial magnetic stimulation and transcranial direct current stimulation appear to be safe neuromodulatory techniques useful in the treatment of anxiety disorders and other neuropsychiatric disorders. *Arg Neuropsychiatr*. 2016;74(10):829–835. doi:10.1590/0004-282X20160115
- Luber BM, Davis S, Bernhardt E, et al. Using neuroimaging to individualize TMS treatment for depression: toward a new paradigm for imaging-guided intervention. *NeuroImage*. 2017;148:1–7. doi:10.1016/j.neuroimage.2016.12.083
- Vidor LP, Torres ILS, Medeiros LF, et al. Association of anxiety with intracortical inhibition and descending pain modulation in chronic myofascial pain syndrome. *BMC Neurosci*. 2014;15:42. doi:10.1186/1471-2202-15-42
- Mak ADP, Chan SSM, Lam LCW, et al. Preliminary results from a randomized sham-controlled trial of augmentative neuro-navigated right-dorsolateral prefrontal cortex low-frequency repetitive transcranial magnetic stimulation for antidepressant non-responding bipolar depression. *Brain Stimul*. 2017;10(2):378. doi:10.1016/j.brs.2017.01.121
- Bashir S, Vernet M, Yoo WK, Mizrahi I, Theoret H, Pascual-Leone A. Changes in cortical plasticity after mild traumatic brain injury. *Restor Neurol Neurosci*. 2012;30(4):277–282. doi:10.3233/RNN-2012-110207
- Li SS, Zaninotto AL, Neville IS, Paiva WS, Nunn D, Fregni F. Clinical utility of brain stimulation modalities following traumatic brain injury: current evidence. *Neuropsychiatr Dis Treat*. 2015;11:1573–1586. doi:10.2147/NDT.S65816
- Weir K, Gray S. Can magnets cure depression. Transcranial magnetic stimulation is gaining ground as a therapy for treatment-resistant depression. *APA*. 2015;46(2):50.
- Watts BV, Landon B, Groft A, Young-Xu Y. A sham controlled study of repetitive transcranial magnetic stimulation for posttraumatic stress disorder. *Brain Stimul*. 2012;5(1):38–43. doi:10.1016/j.brs.2011.02.002
- Ma X, Huang Y, Liao L, Jin Y. A randomized double-blinded sham-controlled trial of a electroencephalogram-guided transcranial magnetic stimulation for obsessive-compulsive disorder. *Chin Med J*. 2014;127(4):601–606.
- Hegde S. Music-based cognitive remediation therapy for patients with traumatic brain injury. *Front Neurol*. 2014;5. doi:10.3389/fneur.2014.00034
- Nielson DM, McKnight CA, Patel RN, Kalnin AJ, Mysiw WJ. Preliminary guidelines for safe and effective use of repetitive transcranial magnetic stimulation in moderate to severe traumatic brain injury. *Arch Phys Med Rehabil*. 2015;96(4):S138–S144. doi:10.1016/j.apmr.2014.09.010
- Araújo HA, Iglesias RF, de Camargo Correia GS, et al. Estimulação magnética transcraniana e aplicabilidade clínica: perspectivas na conduta terapêutica neuropsiquiátrica. *Rev Med*. 2011;90(1):3–14. doi:10.11606/issn.1679-9836.v90i1p3-14
- Barker A, Freeston I, Jalinous R, Jarratt J. Magnetic stimulation of the human brain and peripheral nervous system: an introduction and the results of an initial clinical evaluation. *Neurosurgery*. 1987;20(1):100–109. doi:10.1097/00006123-198701000-00024
- Lefaucheur JP. Stroke recovery can be enhanced by using repetitive transcranial magnetic stimulation (rTMS). *Clin Neurophysiol*. 2006;36(3):105–115. doi:10.1016/j.neucli.2006.08.011
- Miniussi C, Rossini PM. Transcranial magnetic stimulation in cognitive rehabilitation. *Neuropsychol Rehabil*. 2011;21(5):579–601. doi:10.1080/09602011.2011.562689
- Ricci R, Ramsey D, Johnson K, et al. A pilot feasibility study of daily rTMS to modify corticospinal excitability during lower limb immobilization. *Ther Clin Risk Manag*. 2008;4(5):1127–1134. doi:10.2147/term.s2719
- Vicario CM, Salehinejad MA, Felmingham K, Martino G, Nitsche MA. A systematic review on the therapeutic effectiveness of non-invasive brain stimulation for the treatment of anxiety disorders. *Neurosci Biobehav Rev*. 2018;96:219–231.
- Paes F, Machado S, Arias-Carrion O, et al. The value of repetitive transcranial magnetic stimulation (rTMS) for the treatment of anxiety disorders: an integrative review. *CNS Neurol Disord Drug Targets*. 2011;10(5):610–620.

32. Lefaucheur JP, André-Obadia N, Antal A, et al. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). *Clin Neurophysiol.* 2014;125(11):2150–2206. doi:10.1016/j.clinph.2014.05.021
33. Dilkov D, Hawken ER, Kaludiev E, Milev R. Repetitive transcranial magnetic stimulation of the right dorsal lateral prefrontal cortex in the treatment of generalized anxiety disorder: a randomized, double-blind sham controlled clinical trial. *Prog Neuro-Psychopharmacol Biol Psychiatry.* 2017;78:61–65. doi:10.1016/j.pnpbp.2017.05.018
34. Mantovani A, Aly M, Dagan Y, Allart A, Lisanby SH. Randomized sham controlled trial of repetitive transcranial magnetic stimulation to the dorsolateral prefrontal cortex for the treatment of panic disorder with comorbid major depression. *J Affect Disord.* 2013;144(1–2):153–159. doi:10.1016/j.jad.2012.05.038
35. Diefenbach GJ, Bragdon LB, Zertuche L, et al. Repetitive transcranial magnetic stimulation for generalised anxiety disorder: a pilot randomised, double-blind, sham-controlled trial. *Br J Psychiatry.* 2016;209(3):222–228. doi:10.1192/bjp.bp.115.168203
36. Prasko J, Zalesky R, Bares M, et al. The effect of repetitive transcranial magnetic stimulation (rTMS) add on serotonin reuptake inhibitors in patients with panic disorder: a randomized, double blind sham controlled study. *Neuroendocrinol Lett.* 2007;28(1):33–38.
37. Zhang T, Zhu J, Xu L, et al. Add-on rTMS for the acute treatment of depressive symptoms is probably more effective in adolescents than in adults: evidence from real-world clinical practice. *Brain Stimul.* 2019;12(1):103–109. doi:10.1016/j.brs.2018.09.007
38. Lu R, Zhang C, Liu Y, Wang L, Chen X, Zhou X. The effect of bilateral low-frequency rTMS over dorsolateral prefrontal cortex on serum brain-derived neurotrophic factor and serotonin in patients with generalized anxiety disorder. *Neurosci Lett.* 2018;684:67–71. doi:10.1016/j.neulet.2018.07.008
39. Kaur M, Naismith SL, Lagopoulos J, et al. Sleep-wake, cognitive and clinical correlates of treatment outcome with repetitive transcranial magnetic stimulation for young adults with depression. *Psychiatry Res.* 2019;271:335–342. doi:10.1016/j.psychres.2018.12.002
40. Lin H, Li W, Ni J, Wang Y. Clinical study of repetitive transcranial magnetic stimulation of the motor cortex for thalamic pain. *Medicine.* 2018;97(27):e11235.
41. Reyes-López J, Ricardo-Garcell J, Armas-Castañeda G, et al. Clinical improvement in patients with borderline personality disorder after treatment with repetitive transcranial magnetic stimulation: preliminary results. *Rev Bras Psiquiatr.* 2018;40(1):97–104. doi:10.1590/1516-4446-2016-2112
42. Durmaz O, Ebrinc S, Ates MA, Algul A. Evaluation of repetitive transcranial magnetic stimulation for treatment resistant major depression and the impact of anxiety symptoms on outcome. *Psychiatry Clin Psychopharmacol.* 2017;27(1):14–18. doi:10.1080/24750573.2017.1293239
43. Noh TS, Kyong JS, Chang MY, et al. Comparison of treatment outcomes following either prefrontal cortical-only or dual-site repetitive transcranial magnetic stimulation in chronic tinnitus patients: a double-blind randomized study. *Otol Neurotol.* 2017;38(2):296–303. doi:10.1097/MAO.0000000000001266
44. Tovar-Perdomo S, McGirr A, Van Den Eynde F, Dos Santos NR, Berlim MT. High frequency repetitive transcranial magnetic stimulation treatment for major depression: dissociated effects on psychopathology and neurocognition. *J Affect Disord.* 2017;217:112–117. doi:10.1016/j.jad.2017.03.075
45. Malavera A, Silva FA, Fregni F, Carrillo S, Garcia RG. Repetitive transcranial magnetic stimulation for phantom limb pain in land mine victims: a double-blinded, randomized, sham-controlled trial. *J Pain.* 2016;17(8):911–918. doi:10.1016/j.jpain.2016.05.003
46. Bilici S, Yigit O, Taskin U, Gor AP, Yilmaz ED. Medium-term results of combined treatment with transcranial magnetic stimulation and antidepressant drug for chronic tinnitus. *Eur Archiv Oto-Rhino-Laryngol.* 2015;272(2):337–343. doi:10.1007/s00405-013-2851-z
47. Lin YC, Feng Y, Zhan SQ, et al. Repetitive transcranial magnetic stimulation for the treatment of restless legs syndrome. *Chin Med J.* 2015;128(13):1728. doi:10.4103/0366-6999.159344
48. Ozgur T, Akarsu S, Celik C, et al. Is transcranial magnetic stimulation effective in treatment-resistant combat related posttraumatic stress disorder? *Neurosciences (Riyadh).* 2014;19(1):29–32.
49. Diefenbach GJ, Bragdon L, Goethe JW. Treating anxious depression using repetitive transcranial magnetic stimulation. *J Affect Disord.* 2013;151(1):365–368. doi:10.1016/j.jad.2013.05.094
50. Pretalli JB, Nicolier M, Chopard G, et al. Resting motor threshold changes and clinical response to prefrontal repetitive transcranial magnetic stimulation in depressed patients. *Psychiatry Clin Neurosci.* 2012;66(4):344–352. doi:10.1111/j.1440-1819.2012.02341.x
51. Sun W, Mao W, Meng X, et al. Low-frequency repetitive transcranial magnetic stimulation for the treatment of refractory partial epilepsy: a controlled clinical study. *Epilepsia.* 2012;53(10):1782–1789. doi:10.1111/j.1528-1167.2012.03626.x
52. Berlim MT, McGirr A, Beaulieu MM, Turecki G. High frequency repetitive transcranial magnetic stimulation as an augmenting strategy in severe treatment-resistant major depression: a prospective 4-week naturalistic trial. *J Affect Disord.* 2011;130(1–2):312–317. doi:10.1016/j.jad.2010.10.011
53. Boggio PS, Rocha M, Oliveira MO, et al. Noninvasive brain stimulation with high-frequency and low-intensity repetitive transcranial magnetic stimulation treatment for posttraumatic stress disorder. *J Clin Psychiatry.* 2010;71(8):992. doi:10.4088/JCP.08m04638blu
54. Epstein CM, Evatt ML, Funk A, et al. An open study of repetitive transcranial magnetic stimulation in treatment-resistant depression with Parkinson's disease. *Clin Neurophysiol.* 2007;118(10):2189–2194. doi:10.1016/j.clinph.2007.07.010
55. Passard A, Attal N, Benadhira R, et al. Effects of unilateral repetitive transcranial magnetic stimulation of the motor cortex on chronic widespread pain in fibromyalgia. *Brain.* 2007;130(10):2661–2670. doi:10.1093/brain/awm189
56. Rossi S, De Capua A, Olivelli M, et al. Effects of repetitive transcranial magnetic stimulation on chronic tinnitus. A randomised, cross over, double blind, placebo-controlled study. *Journal of Neurology, Neurosurgery & Psychiatry.* 2007. doi:10.1136/jnnp.2006.105007
57. Cohen H, Kaplan Z, Kotler M, Kouperman I, Moisa R, Grisaru N. Repetitive transcranial magnetic stimulation of the right dorsolateral prefrontal cortex in posttraumatic stress disorder: a double-blind, placebo-controlled study. *Am J Psychiatry.* 2004;161(3):515–524. doi:10.1176/appi.ajp.161.3.515
58. Loo CK, Sachdev PS, Haindl W, et al. High (15 Hz) and low (1 Hz) frequency transcranial magnetic stimulation have different acute effects on regional cerebral blood flow in depressed patients. *Psychol Med.* 2003;33(6):997–1006.
59. Münchau A, Bloem BR, Thilo KV, Trimble MR, Rothwell JC, Robertson MM. Repetitive transcranial magnetic stimulation for Tourette syndrome. *Neurology.* 2002;59(11):1789–1791. doi:10.1212/01.wnl.0000036615.25044.50
60. George MS, Nahas Z, Molloy M, et al. A controlled trial of daily left prefrontal cortex TMS for treating depression. *Biol Psychiatry.* 2000;48(10):962–970. doi:10.1016/s0006-3223(00)01048-9
61. Rollnik JD, Huber TJ, Mogk H, et al. High frequency repetitive transcranial magnetic stimulation (rTMS) of the dorsolateral prefrontal cortex in schizophrenic patients. *Neuroreport.* 2000;11(18):4013–4015. doi:10.1097/00001756-200012180-00022
62. Pennisi G, Bella R, Lanza G. Motor cortex plasticity in subcortical ischemic vascular dementia: what can TMS say? *Clin Neurophysiol.* 2015;126(5):851–852. doi:10.1016/j.clinph.2014.09.001
63. Cantone M, Bramanti A, Lanza G, et al. Cortical plasticity in depression: a neurochemical perspective from transcranial magnetic stimulation. *ASN Neuro.* 2017;9(3):1759091417711512. doi:10.1177/1759091417711512

64. Herrman MJ, Katzorke A, Busch Y, et al. Medial prefrontal cortex stimulation accelerates therapy response of exposure therapy in acrophobia. *Brain Stimul.* 2017;10(2):291. doi:10.1016/j.brs.2016.11.007
65. Lanza G, Cantone M, Aricò D, et al. Clinical and electrophysiological impact of repetitive low-frequency transcranial magnetic stimulation on the sensory-motor network in patients with restless legs syndrome. *Ther Adv Neurol Disord.* 2018;11:1756286418759973. doi:10.1177/1756286418759973
66. Lanza G, Bachmann CG, Ghorayeb I, Wang Y, Ferri R, Paulus W. Central and peripheral nervous system excitability in restless legs syndrome. *Sleep Med.* 2017;31:49–60. doi:10.1016/j.sleep.2016.05.010
67. Concerto C, Lanza G, Cantone M, et al. Repetitive transcranial magnetic stimulation in patients with drug-resistant major depression: a six-month clinical follow-up study. *Int J Psychiatry Clin Pract.* 2015;19(4):252–258. doi:10.3109/13651501.2015.1084329

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and

is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>