

The Impact Of Succinate Dehydrogenase Gene (SDH) Mutations In Renal Cell Carcinoma (RCC): A Systematic Review

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Seyed Mohammad Kazem Aghamir¹
Ramin Heshmat²
Mehdi Ebrahimi³
Seyed Ebrahim Ketabchi⁴
Somayeh Parichehreh Dizaji⁵
Fatemeh Khatami¹

¹Urology Research Center, Tehran University of Medical Sciences, Tehran, Iran; ²Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran; ³Department of Internal Medicine, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran; ⁴Department of Neurosurgery, Sina Hospital, Tehran, Iran; ⁵Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran

Introduction: Renal cell cancer (RCC) syndrome is linked to Krebs cycle compartments and their coding genes' alterations like *succinate dehydrogenase* genes (*SDHx*). Here we present a systematic review of the SDH genes' mutations and their impact on both RCC diagnosis and prognosis.

Methods: This systematic review includes any study in which tissue samples of RCC are considered in correlation with the SDHx mutations, microsatellite instability (MSI), and protein expression. For this purpose, a systematic search of MEDLINE (PubMed), Scopus, Embase, and Web of Science databases was conducted and finally 5384 articles were recruited. All studies' content was checked to find the related ones which were 145 articles, which with data extraction were limited to nineteen.

Results: The final selected nineteen studies investigating the *SDHx* role in RCC tumor genesis were included, among which fifteen were mutation analysis, three were just *SDHx* protein expression, and two were MSI and mutation analysis studies. A total of 432 RCC patients were reported by *SDH* mutations, and 64 patients with MSI and *SDH* expression change were reported in 514 surgically resected renal epithelial tumors. The most common mutation was the single nucleotide variant rs772551056 (c.137G>A) of *SDHB*. For *SDHC*, c.380A>G presented in 48 RCC patients, and for *SDHA* a novel germline mutation c.2T>C: p.M1T in an occasional case of gastrointestinal stromal tumor intricate with RCC.

Conclusion: RCC as an aggressive type of kidney cancer needs some biomarkers to be diagnosed exactly. It was shown recently that the succinate dehydrogenase gene variations can provide this diagnostic and prognostic biomarker. For this purpose, *SDHB* rs772551056 associated with its protein expression alterations can be taken into account. It is possible that a novel mutation of *SDHA* (c.2T>C: p.M1T) can provide evidence of GIST associated with RCC as well.

Keywords: renal cell cancer, RCC, succinate dehydrogenases, mutation, expression, MSI

Introduction

Kidney cancers with different genetic changes have different histology and clinical significance. The genes relating to kidney cancer are usually involved in metabolic stress or nutrient stimulation pathways including *Von Hippel-Lindau(VHL)*, *mesenchymal-epithelial transition factor(MET)*, *Folliculin(FLCN)*, *Tuberous sclerosis(TSC1 and TSC2)*, *Microphthalmia-associated transcription factor(MITF)*, *Phosphatase and tensin homolog(PTEN)*, *fumarate hydratase(FH)*, and *succinate dehydrogenase (SDH)*.^{1,2} One of the usual forms of kidney cancers is RCC that initiates in the lining

Correspondence: Fatemeh Khatami
Urology Research Center, Sina Hospital,
Hassan Abad Square, Tehran, Iran
Tel +98 21 88220086
Email f-khatami@alumnus.tums.ac.ir

of the proximal convoluted tubule responsible for about 90–95% of kidney cancer cases.^{3–6} It commonly is described by an absence of early-warning signs (which outcomes in an extraordinary proportion of patients with metastases), various medical manifestations, and fighting radiotherapy and chemotherapy with a possible impact on immunomodulation in the tumor growth suppression.^{7–9} RCC has the highest mortality rate of the genitourinary cancers and its prevalence has ascended steadily over the past few decades.¹⁰ Some recent research has proved the benefit of genomic data, principally gene expression hallmarks, as medical predictive factors in personalized cancer managements.^{11–13} Succinate dehydrogenase (*SDH* or *SQR*) is an enzyme complex of the inner mitochondrial membrane that contributes to the citric acid cycle as well as the electron transport chain.¹⁴ Succinate dehydrogenase is made of four protein subunits (*SDHA*, *SDHB*, *SDHC*, and *SDHD*) having a role in the TCA cycle and electron transport chain in mitochondria.^{15,16} Patients with mutations in the *SDH* genes are under the risk of autonomic nervous system tumors like pheochromocytomas and paragangliomas (PPGLs), both head and neck, and in the thorax and abdomen, gastrointestinal stromal tumors (GISTs), and renal cell carcinoma (RCC). The evaluation of protein expression by immunohistochemistry routinely is used to find a discriminative biomarker of benign and malignant tumors including RCC.^{17–19} Succinate dehydrogenase mutation in PPGL and GIST tumor cells had a microscopic result of predominantly epithelioid morphology, and epithelioid morphology, but not in RCC.²⁰ In fact, renal tumors are usually known with typical morphology of identical cells with eosinophilic or oncocytic cytoplasm that have cytoplasmic vacuoles or flocculent inclusions.^{21–23} RCCs with additional histologic presence have been described in patients with germline mutations of succinate dehydrogenase genes, contrary to some limited histological types of RCCs with no succinate dehydrogenase mutation.^{24,25}

In recent years the genetic basis of RCC has been known and both genetic and epigenetic alterations of succinate dehydrogenase have been discussed. Interestingly, germline mutations of the genes coding for the succinate dehydrogenase subunits (*SDHB*, *SDHC*, and *SDHD*) have been identified in patients with a combination of gastrointestinal stromal tumors (GISTs) and paraganglioma (PGL).^{26,27} The co-occurrence of RCC with paraganglioma or pheochromocytoma suggested a succinate dehydrogenase gene mutation presence.^{28,29} A newly characterized specific subtype of RCC is in the new World Health Organization (WHO) classification and was published in 2016.^{30–32}

In spite of the fact that recently the focus of scientists is on RCC genetic and epigenetic modifications, knowledge of the clinical features and management of patients associated with the *SDH* mutations is limited. The current systematic review study is run with the purpose of delivering the first meticulous summary of all the available primary research over the *SDH* mutations, expression, and microsatellite instability (MSI) in RCC management and screening recommendations.

Research Design And Methods

Search Strategy

The study was based on an international prospective register of systematic reviews with PROSPERO 2018 code CRD42018087806 available from http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42018087806. All linked works were searched from four targeted databases including: MEDLINE (PubMed), Scopus, Embase, and Web of Science. Publication dates of relevant articles were limited from 1st January 1990 to 30th March 2018. Our search syntax were “Succinic Dehydrogenase”, “*SDH* mutations”, “Succinic Oxidase”, “succinate-coenzyme *Q* reductase”, “(*SQR*)”, “respiratory Complex II” combined with “Collecting Duct Carcinoma”, “RCC”, “Renal Cell Adenocarcinoma”, “Kidney adenocarcinoma”, and “kidney Cancer” (Supplementary materials). In order to decrease the selection bias, two separate investigators (FK and RH) individually reviewed titles, abstracts, and available full-text articles to choose the related ones with *SDH* and RCC. Additional related papers were documented above searching the reference lists of selected studies. Disagreements were solved by the third independent investigator (SMKA).

Eligibility Criteria

All chosen studies were studied by two authors individually and based on their English full text were included or excluded. The considering inclusion criteria were: 1) participants included kidney cancer patients pathologically classified as RCC, renal epithelial tumors, or renal tumors; 2) all *SDHx* alteration detection techniques such as immunohistochemistry (IHC), DNA sequencing, western blotting, PCR-based methods, SNaPshot Assay, and PCR-RFLP were included; and 3) *SDHx* alterations were composed mutations, MSI, and protein expression. Studies were excluded if they: 1) analyzed *SDHx* mutations or expression in animals (in vivo studies); 2) studied them in cell culture

(in vitro studies); or (3) did not have an appropriate explanation of selected case groups.

Data Extraction And Quality Assessment

All data on population characteristics and *SDHx* deficiency results were extracted in the Excel file. One investigator completed the data extraction step (SPD), which was established via another investigator (FK). Due to the insufficient quantity of studies relevant to *SDHx* deficiencies and RCC, together with the significant variations in targeted genetic variations and study population characteristics, a meta-analysis for the current data was not applicable. The study team used the Newcastle–Ottawa Scale (NOS) scale assessment tools to assess the quality of all selected final studies.^{33,34} The methodologies of studies scoring 6/8 or more on the NOS were categorized as “high quality”, whereas those that scored 5/8 or lower were reported as “low quality”.

Results

Study Selection And Characteristics

The selection flow chart and results of the study selection procedure are presented in Figure 1. A total of 5384 articles was retrieved and after duplication deletion 3964 remained, in which there were 2905 from PubMed, 559

from Scopus, 325 from Web of Science, and 130 from Embase. When the review articles, in vivo/in vitro studies, and book or conference papers were deleted the total number of nineteen articles was chosen for advance considerations. A total of sixteen studies were mainly focused on *SDHx* mutations, three studied the *SDHx* protein expression without mutation analysis, and two targeted *SDHx* MSI with or without mutation (Table 1).

In sixteen studies, *SDHB* was checked for mutation, and protein expression. In these studies, *SDHB* was evaluated alone or simultaneously with other candidate genes like *FH*, *TFE3 gene rearrangement*, *TMEM127*, *MAX*, *HIF1a*, *Cathepsin*, *PAX8*, *Cathepsin*, *CK20*, and *EMA*. In these studies, several mutations of *SDHB* were checked including exonic mutations (c.137G>A), splice site acceptor or donor mutations (c.72+1G→T), exon 1 splice acceptor site or c.268C>T (p. Arg90X) in exon 3 splice site mutation, and (c.136C>T Stop) mutation of stop codons which are resulting in truncated inactive forms of the protein. Five studies were mainly focused on *SDHC* in which loss of heterozygosity (LOH) in two telomeric regions (D3S369, D3S1597) and five centromeric regions (D3SVHL3, D3S1337, D3SVHL7, D3SVHL8, D3S3611) more than c.380A>G mutations were tested. Especially, in an aggressive example of the Warburg Effect in succinate




Figure 1 Flow chart of study selection.

Table 1 Characterization Of Nineteen Related Studies To SDHx Genetic Modifications And Protein Expression In RCC Patients

Target SDH	Title	First Author	Publication Year	Country	Sample Size	Age	Type Of Study	Detected Mutation
SDHB	Renal carcinoma with giant mitochondria associated with germline mutation and somatic loss of the succinate dehydrogenase B gene ³⁹	Sarah L Housley	2010	UK	1 RCC	58	Case report	c.72+1G→T exon 1 splice acceptor site
	Germline SDHB mutations and familial renal cell carcinoma ²¹	Christopher Ricketts	2008	UK	68 RCC	50	Case series	c.137G>A, c.32G>A, c.136C>T Stop
	Early-onset renal cell carcinoma as a novel extraparaganglial component of SDHB-associated heritable paraganglioma ³⁶	Sakari Vanharanta	2004	Finland	60 sporadic RCCs	15–34	Case series	e R27X mutation (N8168)
	Succinate dehydrogenase (SDH)-deficient renal carcinoma: a morphologically distinct entity: a clinicopathologic series of 36 tumors from 27 patients ⁴⁰	Anthony J Gill	2014	Australia	36 tumors from 27 patients	37	Cohort	c.137G>A c.725G>A c.423+1G>A SDHB exon 3 deletion c.423+1G>A c.338G>A c.423+1G>A c.749C>A SDHC c.380A>G
	Renal cell carcinoma with TFE3 translocation and succinate dehydrogenase B mutation ⁴¹	Anna Calió	2017	USA	4 RCCs	From 19 to 65	Case series	c.423+1G 4A p.V140F c.72+1G 4T
	Immunohistochemical characterization of fumarate hydratase (FH) and succinate dehydrogenase (SDH) in cutaneous leiomyomas for detection of familial cancer syndromes ⁴²	Cody S Carter	2017	USA	96 consecutive specimens of cutaneous leiomyomas from 87 RCC patients		Case/control	SDHB expression

(Continued)

Table 1 (Continued).

Target SDH	Title	First Author	Publication Year	Country	Sample Size	Age	Type Of Study	Detected Mutation
	Clinical and molecular features of renal and pheochromocytoma/paraganglioma tumour association syndrome (RAPTAS): case series and literature review ⁴³	Ruth T Casey	2017	UK	22 probands with non-VHL RAPTAS	30	Case series	N/A
	Succinate dehydrogenase B: a new prognostic biomarker in clear cell renal cell carcinoma ³⁷	Kristine M Cornejo	2015	USA	420 surgically resected renal epithelial tumors	62.6 (33–92)	Case series	SDHB expression
	Utility of the succinate: fumarate ratio for assessing SDH dysfunction in different tumor types ⁴⁴	Edward Kim	2017	Australia	11 RCCs 18 PPGLs, 10 GISTs	–	Case/control	c.380T>G, p.Ile127Ser
	Increased HIF1 α in SDH and FH deficient tumors does not cause microsatellite instability ⁴⁵	Heli J Lehtonen	2007	Finland	11 RCCs, 12 ULMs 1 ULMs		Case series	N/A
	Renal cell carcinoma occurring in patients with prior neuroblastoma ⁴⁶	Sara M Falzarano	2016	USA	7 RCCs	40–64	Case series	SDHB expression
	Renal tumors associated with germline SDHBmutation show distinctive morphology ²⁰	Anthony J Gill	2011	Australia	4 renal tumors	32	Case reports	c.166-170delCCTCA in exon 2 c.72+1G>T c.268C>T (p.Arg90X) in exon 3 splice site mutation (c.423+1G>A)
	SDHB-associated renal oncocytoma suggests a broadening of the renal phenotype in hereditary paragangliomatosis ⁴⁷	Alex Henderson	2009	UK	11 Renal tumors	16–73	Case series	c.136C>T c.137G>A c.79C>T c.715_718 delTCTC c.32G>A c.141G>A c.3G>A c.600G>T

(Continued)

Table 1 (Continued).

Target SDH	Title	First Author	Publication Year	Country	Sample Size	Age	Type Of Study	Detected Mutation
SDHC	Molecular genetic analysis of <i>FIH-1</i> , <i>FH</i> , and <i>SDHB</i> candidate tumour suppressor genes in renal cell carcinoma ³⁵	M R Morris	2004	UK	29 RCCs	–	Case series	IVS2+33 A R G IVS2+35 G R A c.18 A R C; A6A DIS2697
	Renal carcinoma associated with <i>succinate dehydrogenase B</i> mutation: a new and unique subtype of renal carcinoma ⁴⁸	Julie Y Paik	2014	Australia	A unique case RCC	27	Case report	c.88delC (p.Gln30ArgfsX47) in exon 2
	Papillary renal cell carcinoma with cytologic and molecular genetic features overlapping with renal oncocytoma: analysis of 10 cases. ⁴⁹	Kvetoslava Michalova	2018	Czech Republic	147 PRCCs	67	Case series	N/A
	Clinical and molecular features of renal and pheochromocytoma/paraganglioma tumour association syndrome (RAPTAS): case series and literature review ⁴³	Ruth T Casey	2017	UK	22 probands with non-VHL RAPTAS	30	Case series	N/A
	Increased HIF1 α in SDH and FH deficient tumors does not cause microsatellite instability ⁴⁵	Heli J Lehtonen	2007	Finland	11 RCCs, 12 ULMs 1 ULMs		Case series	N/A
	Biallelic inactivation of the <i>SDHC</i> gene in renal carcinoma associated with paraganglioma syndrome type 3 ²⁴	Angelica Malinoc	2012	Germany	35 head and neck paragangliomas	–	Case series	LOH at <i>SDHC</i> : 2 telomeric D3S3691 D3S1597 5 centromeric D3SVHL3, D3S1337, D3SVHL7, D3SVHL8, D3S3611;
	Succinate dehydrogenase (<i>SDH</i>)-deficient renal carcinoma: a morphologically distinct entity a clinicopathologic series of 36 tumors from 27 patients ⁴⁰	Anthony J Gill	2014	Australia	28 tumor samples	37	Cohort	<i>SDHC</i> c.380A>G

(Continued)

Table 1 (Continued).

Target SDH	Title	First Author	Publication Year	Country	Sample Size	Age	Type Of Study	Detected Mutation
SDHA	Succinate dehydrogenase kidney cancer (SDH-RCC): an aggressive example of the Warburg Effect in cancer ⁵⁰	Christopher J Ricketts	2012	USA	14 patients from SDHB mutation families	47	Case series	R133X
	A novel germline mutation in SDHA identified in a rare case of gastrointestinal stromal tumor complicated with renal cell carcinoma ³⁸	Quan Jiang	2015	China	A case of GIST RCC	23	Case report	c.2T>C; p.M1T
	Early-onset renal cell carcinoma as a novel extraparaganglial component of SDHB-associated heritable paraganglioma ³⁶	Sakari Vanharanta	2004	Finland	60 sporadic RCCs	15–34	Case series	N/A
SDHD	Germline SDHB mutations and familial renal cell carcinoma ²¹	Christopher Ricketts	2008	UK	68 RCC	50	Case series	N/A
	Early-onset renal cell carcinoma as a novel extraparaganglial component of SDHB-associated heritable paraganglioma ³⁶	Sakari Vanharanta	2004	Finland	60 sporadic RCCs	15–34	Case series	N/A
	Clinical and molecular features of renal and pheochromocytoma/paraganglioma tumour association syndrome (RAPTAS); case series and literature review ⁴³	Ruth T Casey	2017	UK	22 probands with non-VHL RAPTAS	30	Case series	N/A
	Biallelic inactivation of the SDHC gene in renal carcinoma associated with paraganglioma syndrome type 3 ²⁴	Angelica Malinoc	2012	Germany	35 head and neck paragangliomas	–	Case series	N/A
	Increased HIF1 α in SDH and FH deficient tumors does not cause microsatellite instability ⁴⁵	Hei J Lehtonen	2007	Finland	11 RCCs, 12 ULMs 1 ULMs		Case series	N/A

Abbreviations: RAPTAS, pheochromocytoma/paraganglioma tumor association syndrome; SDH, succinate dehydrogenase; PPGLs, pheochromocytoma and paraganglioma; GISTs, gastrointestinal stromal tumors; RCCs, renal cell carcinomas; HLRCC, hereditary leiomyomatosis and renal cell cancer; ULM, benign uterine leiomyomas; ULMs, uterine leiomyosarcoma; HPGL, hereditary paragangliomatosis; NB, neuroblastoma; GIST RCC, gastrointestinal stromal tumor complicated with renal cell carcinoma; PRCC, papillary RCC; cRCC, clear cell RCC; LOH, loss of heterozygosity; OPRCC, oncocytic variant of papillary renal cell carcinoma; HIF, hypoxia inducible factor; FH, fumarate hydratase; N/A, not available.

dehydrogenase kidney cancer (SDH-RCC), the germline SDHC mutation (R133X) (NM_003001.3-c.397C>T (p. Arg133Ter)) was identified. Two studies were related to SDHA (c.2T>C: p.M1T/rs864622194) resulting in substitution of Methionine with Threonine in a NM_006493.2: c.2T>C missense variant that change the amino acid sequence in protein and resulted in non-functional protein (<https://www.ncbi.nlm.nih.gov/clinvar/variation/219649/>).

Five studies concerned *SDHD* with no exact determined targeted mutations. Except for one study in sporadic RCC indicating no mutations in the three *FH*, *FIH-1*, and *SDHB*, others mainly suggested early-onset RCC with unusual histology (e.g., solid) must be observed to take an extended family history and *SDHB* mutant-related RCC in the way of better diagnosis and prognosis.^{35,36} In fact the weak staining, particularly in clear neoplasms, usually can be interpreted as negative by mistake.³⁷ A total of 432 RCC patients were reported by *SDH* mutations, and 64 patients with MSI and *SDH* expression change were reported in 514 surgically resected renal epithelial tumors. The most common mutation was the single nucleotide variant rs772551056 (c.137G>A) of *SDHB* with genomic location Chr1: 17044824 resulting in protein change *R46Q* which was reported in 106 RCC patients. After that, two mutations of c.32G>A and c.136C>T Stop were the most with 79 reported RCC patients. The mutation of c.423+1G>A was in 44 RCC patients and c.72+1G→T in 20 patients. The *SDHC* c.380A>G (p. His127Arg) was in 48 RCC, 18 PPGLs, and 11 GIST. The rs786201095 (c.380T>G (p. Ile127Ser)) with chromosome location Chr1: 17028643 is the common mutation found in RCC in addition to GIST and PPGL. *SDHB* expression in ccRCCs with high nucleolar grade (G3–G4) was considerably linked to patient's low survival so it can be an excellent candidate biomarker for RCC diagnosis and prognosis.³⁷ The novel germline mutation of chromosome X rs864622194 (c.2T>C: p.M1T) with genomic locus ChrX: 103776997 in *SDHA* was recognized in a rare case of gastrointestinal stromal tumor complicated with RCC.³⁸

Discussion

RCC arises from the cells of the proximal renal tubular epithelium and has two subtypes: sporadic (non-hereditary) and hereditary.⁵¹ Inherited predisposition to RCC is dependent on cellular metabolism responsible genes.^{52,53} RCC is basically a metabolic disease so metabolism involving genes like SDH can trigger the cellular transformation of renal cells in reaction to sensing oxygen, iron, nutrients, and

energy.^{1,54} The SDH enzyme is a very conserved heterotrimeric protein made up of four subunits (*SDHA*, *SDHB*, *SDHC*, and *SDHD*) in which *SDHA* and *SDHB* are catalytic subunits and *SDHC* and *SDHD* are anchored to the inner membrane.⁵⁵ Recently the importance of SDH subunit mutations has been highlighted in different malignancies including RCC.^{56,57} There is a rare and aggressive type of RCC which is called SDH-deficient RCC (abbreviated SDH-RCC) that lately has been added to the WHO classification of renal neoplasia.⁵⁸ The incidence of SDH-deficient RCC is assessed among 0.05–0.2% of all renal carcinomas and among SDH-deficient RCCs, *SDHB*-mutated RCC is the most frequent,⁵⁹ followed by *SDHC* and *SDHD*-deficient RCC.^{24,40} Moreover, a patient case with *SDHA*-deficient RCC has been reported currently.⁶⁰ Early age of onset for RCC has been observed in patients with *SDHB* germline mutations.^{20,61,62}

Common genetic alterations of *SDHB* consist of a nonsense mutation (c.268C>T p.Arg90X), four missense mutations (c.137G>A p.Arg46Gln, c.286G>A p.Gly96Ser, c.379A>C p.Ile127Leu, c.689G>A p.Arg230His), two splice site altering mutations (c.286+2T>A, c.541-2A>G), and three complete deletions of the first exon of *SDHB*.⁵⁰ The mutation c.72+1G→T which is in the exon 1 splice acceptor site resulting in production of a truncated inactive form of the protein was in RCC with giant mitochondria.³⁹ *SDHB* mutation results in distinctive morphology of RCC and these RCCs have a respectable prognosis subsequently of whole excision unless there is sarcomatoid dedifferentiation.²⁰ The involvement of *SDHB* mutations in RCC recommends that *SDHB* mutations must be checked when renal tumors are presented in families with other tumors consistent with hereditary paraganglioma syndrome.⁴⁷ The two mutations c.541-2A>G (Splice) and c.689G>A (p.Arg230His) have a role in impairing iron–sulfur cluster delivery and are highlighted in several cancers.^{63,64} According to our result the most common *SDHB* mutation was the single nucleotide variant rs772551056 (c.137G>A). *SDHB* is the Fe–S subunit of mitochondrial complex II including extremely consensus L (I)YR motifs essential for gaining of Fe–S clusters by recruiting the Fe–S transfer machinery.^{65,66} Importantly, the c.137G>A (p.Arg46Gln or *R46Q*) mutation occurs in the first L(I)YR motif of *SDHB* and can be detected in familial paraganglioma/pheochromocytoma/GIST/renal cell carcinoma tumor syndromes.^{67,68} KDM4/JMJD2 proteins are demethylases that target histone H3 on lysine 9 and 36, and histone H1.4 on lysine 26, and are key

epigenetic regulators of several cancer cells.^{69–71} However, succinate-mediated competitive suppression of 2-oxoglutarate-dependent dioxygenases, histone demethylases (JMJD), and the ten–eleven translocation (TET) family of 5-methyl cytosine (5-mC) family of hydroxylases are important in SDH-deficient tumor phenotypes.⁷² Possible succinate-induced modifications contain stabilization of HIF- α propyl hydroxylase and hypermethylation of histones and DNA.^{73–75} The morphological study of SDH-deficient renal carcinoma represents a distinct and rare renal neoplasm, which is defined by loss of IHC staining for *SDHB*.⁴⁰ It was in 2000 that the first germline mutation of *SDHD* was reported in HOGL families and then in link with familial PCC.^{76,77} Moreover, the *SDHD* mutations together with *SDHB* mutations were reported in familial PCC and HPGL.^{78,79} SDH-deficient PPGLs, GISTs, and RCCs keep a considerably advanced succinate:fumarate ratio versus their SDH-sufficient equivalents so metabolomic analysis is essential to directly measure SDH dysfunction linking to the numerous types of malignancies.⁴⁴

Germline mutation c.380A>G (p.His127Arg) of *SDHC* is the important mutation of this gene which is reported as a recurrent biomarker of SDH-deficient GIST and renal carcinoma.^{80,81} This variant was described in two brothers with head and neck paragangliomas and in a patient with papillary thyroid cancer, renal cell cancer, and GIST; both the renal tumor and GIST demonstrated absence of *SDHB* by IHC.⁴⁰ This mutation was not detected in about 6,500 persons from Europeans and African Americans in the National Heart, Lung, and Blood Institute (NHLBI) Exome Sequencing Project (ESP), showing that it is not a public benign alternative in these people.⁸² Subsequently, the amino acids of Histidine and Arginine share analogous characteristics; this is measured as a conservative amino acid substitution mutation. *SDHC* c.380A>G happens at a conserved protein region through different species and is located at the metal attachment site for iron and in the helical transmembrane topological domain.⁸³ In silico analyses predict that this polymorphism can perhaps change the protein structure and function, and suggested novel mutations in sporadic head and neck paraganglioma and familial paraganglioma and/or pheochromocytoma.^{84–86} Based on the currently available evidence, *SDHC* c.380A>G can be considered the RCC pathogenic variant. These are conflicting because germline *SDHC* mutations are comparatively uncommon, so it was supposed that carriers of mutations of the *SDHC* gene have a special risk for head and neck paragangliomas more than a risk for adrenal pheochromocytoma.⁸⁷ Therefore, not only

the *SDHC* mutations but also the LOH and MSI were taken into consideration in RCC patients.²⁴ Two telomeric (D3S3691, D3S1597) and five centromeric (D3SVHL3, D3S1337, D3SVHL7, D3SVHL8, D3S3611) were evaluated and renal carcinoma. It was reported by Malinoc et al that for the clear cell renal carcinoma LOH was established in *D3S3691* and, *D3S1597* as the telomeric ones and in two centromeric signs (*D3SVHL3* and *D3S3611*) in comparison with undesirable controls so it might be a new molecular indicator for the pathogenesis of RCC and ought to be checked in both heritable and sporadic forms.²⁴

A report about *SDHA* by Jiang et al in 2015 diagnosed a rare case as a wild-type gastrointestinal stromal tumor (WT GIST) intricated with renal chromophobe cancer cells and distinguished an innovative germline mutation chromosome X rs864622194 (c.2T>C: p.M1T) in the position of a transcription initial codon within the *SDHA* gene sequence.³⁸ There was another described patient of an RCC linked with an *SDHA* mutation.⁸⁸ As the main catalytic subunit of *SDH* complex, the (c.2T>C: p.M1T) mutation certainly deactivates the entire *SDH* complex. It has been presented that *SDHA* homozygous deletion mutation of *SDHA* and *SDHB* following altered protein expression evident by IHC and decreased gene expression of *SDHA* noticeable by IHC faultlessly can be coordinated with *SDHA* mutation.⁶⁰ Nevertheless, *SDHA* mutation does not result in loss of *SDHA* protein expression, which directs that the role of the other allele is normal, so *SDHA* in addition to *SDHB* have been recommended as diagnostic biomarkers for screening for potential *SDH* mutations in RCC cases.^{89,90}

More than genetic change there are some epigenetic modifications that change the gene expression with no change in the DNA sequences. In fact, epigenetics acts as an interface between environmental/exogenous factors, cellular responses, and pathological processes.⁹¹ Epigenetic signatures (DNA methylation, mRNA and microRNA expression, etc) can be biomarkers for risk stratification, early detection, and disease classification, as well as targets for therapy and chemoprevention.⁹² To better understand the interplay between etiological factors, cellular molecular characteristics, and disease evolution, the field of “molecular pathological epidemiology (MPE)” has emerged as an interdisciplinary integration of “molecular pathology” and “epidemiology”. The widespread application of epigenome (e.g., methylome) analyses will increase our understanding of disease heterogeneity, epigenotypes (CpG island methylator phenotype, LINE-1 (long interspersed nucleotide element-1; also called long

interspersed nuclear element-1; long interspersed element-1; L1) hypomethylation, etc), and host–disease interactions.⁹³

Conclusion

Succinate dehydrogenase is an important metabolic enzyme in the TCA cycle and electron transport chain. Germline mutations in *SDHB*, *SDHC*, *SDHA*, and *SDHD* are associated with RCC. To our knowledge, as the first systematic review on the succinate dehydrogenase genetic alterations, we can say that the most frequently detected mutation is *SDHB* rs772551056 and its protein expression. Moreover, the *c.380A>G* mutation with four MSI markers (D3S3691, D3S1597, D3SVHL3, D3S3611) of *SDHC* can bring a morphologically distinct entity of RCC and be a predictor of its recurrence and aggressive behavior. The newly suggested mutation of *SDHA* (*c.2T>C: p.MIT*) can provide evidence of GIST associated with RCC as well.

Acknowledgment

Special thanks to the Urology Research Center, Tehran University of Medical Sciences.

Disclosure

The authors report no conflicts of interest in this work.

References

- Linehan WM, Srinivasan R, Schmidt LS. The genetic basis of kidney cancer: a metabolic disease. *Nat Rev Urol*. 2010 May;7(5):277–85.
- Khatami F, Noorinayer B, Ghiasi S, Mohebi R, Hashemi M, Zali MR. Single nucleotide polymorphisms of DNA methyltransferase 1 gene and gastric cancer in Iranian patients: a case control study. *Iran J Cancer Prev*. 2008;1(3):111–118.
- Rini BI, Campbell SC, Escudier B. Renal cell carcinoma. *Lancet*. 2009;373(9669):1119–1132. doi:10.1016/S0140-6736(09)60229-4
- Abdollahi A, Khatami F, Tavangar SM. Clear cell sarcoma: a case report and review of literature. *Int J Hematol Oncol Stem Cell Res*. 2018;12(1):65–68.
- Talab SS, Elmi A, Emami H, et al. Protective effects of acute lithium preconditioning against renal ischemia/reperfusion injury in rat: role of nitric oxide and cyclooxygenase systems. *Eur J Pharmacol*. 2012;681(1–3):94–99. doi:10.1016/j.ejphar.2012.01.042
- Kajbafzadeh A-M, Sabetkish N, Sabetkish S, Javan-Farazmand N, Harsini S, Tavangar S-M. The ameliorative effect of various antioxidants on Adriamycin-induced fetal renal abnormalities. *J Pediatr Urol*. 2013;9(6):1084–1092. doi:10.1016/j.jpuro.2013.03.010
- Motzer RJ, Bander NH, Nanus DM. Renal-cell carcinoma. *N Engl J Med*. 1996;335(12):865–875. doi:10.1056/NEJM199609193351207
- Talab SS, Emami H, Elmi A, et al. Chronic lithium treatment protects the rat kidney against ischemia/reperfusion injury: the role of nitric oxide and cyclooxygenase pathways. *Eur J Pharmacol*. 2010;647(1–3):171–177. doi:10.1016/j.ejphar.2010.08.036
- Kajbafzadeh A-M, Payabvash S, Salmasi AH, Monajemzadeh M, Tavangar SM. Smooth muscle cell apoptosis and defective neural development in congenital ureteropelvic junction obstruction. *J Urol*. 2006;176(2):718–723. doi:10.1016/j.juro.2006.03.041
- Cairns P. Renal cell carcinoma. *Cancer Biomark*. 2011;9(1–6):461–473. doi:10.3233/CBM-2011-0176
- Khatami F, Larijani B, Tavangar SM. Circulating tumor BRAF mutation and personalized thyroid cancer treatment. *Asian Pac J Cancer Prev*. 2017;18(2):293. doi:10.22034/APJCP.2017.18.1.23
- Khatami F, Aghayan HR, Sanaei M, Heshmat R, Tavangar SM, Larijani B. The potential of circulating tumor cells in personalized management of breast cancer: a systematic review. *Acta Med Iran*. 2017;55(3):175–193.
- Khatami F, Tavangar SM. Liquid biopsy in thyroid cancer: new insight. *Int J Hematol Oncol Stem Cell Res*. 2018;12(3):234–247.
- Cecchini G, Schröder I, Gunsalus RP, Maklashina E. Succinate dehydrogenase and fumarate reductase from *Escherichia coli*. *Biochim Biophys Acta Bioenerg*. 2002;1553(1–2):140–157. doi:10.1016/S0005-2728(01)00238-9
- Gill AJ. Succinate dehydrogenase (SDH) and mitochondrial driven neoplasia. *Pathology*. 2012;44(4):285–292. doi:10.1097/PAT.0b013e3283539932
- Parvizi MR, Parviz M, Tavangar SM, et al. Protective effect of magnesium on renal function in STZ-induced diabetic rats. *J Diabetes Metab Disord*. 2014;13(1):84. doi:10.1186/s40200-014-0084-3
- Tavangar SM, Shojaei A, Tabriz HM, et al. Immunohistochemical expression of Ki67, c-erbB-2, and c-kit antigens in benign and malignant pheochromocytoma. *Pathol Res Pract*. 2010;206(5):305–309. doi:10.1016/j.prp.2010.01.007
- Sarmadi S, Izadi-Mood N, Sotoudeh K, Tavangar SM. Altered PTEN expression; a diagnostic marker for differentiating normal, hyperplastic and neoplastic endometrium. *Diagn Pathol*. 2009;4(1):41. doi:10.1186/1746-1596-4-41
- Haghighpanah V, Shooshtarizadeh P, Heshmat R, Larijani B, Tavangar SM. Immunohistochemical analysis of survivin expression in thyroid follicular adenoma and carcinoma. *Appl Immunohistochem Mol Morphol*. 2006;14(4):422–425. doi:10.1097/01.pai.0000213100.88074.b8
- Gill AJ, Pachter NS, Chou A, et al. Renal tumors associated with germline *SDHB* mutation show distinctive morphology. *Am J Surg Pathol*. 2011;35(10):1578–1585. doi:10.1097/PAS.0b013e318227e7f4
- Ricketts C, Woodward ER, Killick P, et al. Germline *SDHB* mutations and familial renal cell carcinoma. *J Natl Cancer Inst*. 2008;100(17):1260–1262. doi:10.1093/jnci/djn254
- Miettinen M, Sarlomo-Rikala M, Cue PM, et al. Mapping of succinate dehydrogenase losses in 2258 epithelial neoplasms. *Appl Immunohistochem Mol Morphol*. 2014;22(1). doi:10.1097/PAI.0b013e31828bfdd3.
- Alimoghaddam K, Sharifabrizi A, Tavangar M, et al. Anti-leukemic and anti-angiogenesis efficacy of arsenic trioxide in new cases of acute promyelocytic leukemia. *Leuk Lymphoma*. 2006;47(1):81–88. doi:10.1080/10428190500300373
- Malinoc A, Sullivan M, Wiech T, et al. Biallelic inactivation of the *SDHC* gene in renal carcinoma associated with paraganglioma syndrome type 3. *Endocr Relat Cancer*. 2012;19(3):283–90.
- Papathomas TG, Gaal J, Corssmit EP, et al. Non-pheochromocytoma (PCC)/paraganglioma (PGL) tumors in patients with succinate dehydrogenase-related PCC–PGL syndromes: a clinicopathological and molecular analysis. *Eur J Endocrinol*. 2014;170(1):1–12. doi:10.1530/EJE-13-0623
- Pasini B, McWhinney SR, Bei T, et al. Clinical and molecular genetics of patients with the Carney–stratakis syndrome and germline mutations of the genes coding for the succinate dehydrogenase subunits *SDHB*, *SDHC*, and *SDHD*. *Eur J Hum Genet*. 2008;16(1):79. doi:10.1038/ejhg.2008.52
- Janeyway KA, Kim SY, Lodish M, et al. Defects in succinate dehydrogenase in gastrointestinal stromal tumors lacking KIT and PDGFR mutations. *Proc Natl Acad Sci*. 2011;108(1):314–318. doi:10.1073/pnas.1009199108

28. Srirangalingam U, Walker L, Khoo B, et al. Clinical manifestations of familial paraganglioma and pheochromocytomas in succinate dehydrogenase B (SDH-B) gene mutation carriers. *Clin Endocrinol (Oxf)*. 2008;69(4):587–596. doi:10.1111/j.1365-2265.2008.03274.x
29. Schimke RN, Collins DL, Stolle CA. Paraganglioma, neuroblastoma, and a SDHB mutation: resolution of a 30-year-old mystery. *Am J Med Genet Part A*. 2010;152(6):1531–1535.
30. Agaimy A. [Succinate dehydrogenase (SDH)-deficient renal cell carcinoma]. *Pathologie*. 2016;37(2):144–152. doi:10.1007/s00292-016-0158-8
31. Khatami F, Larijani B, Tavangar SM. The presence of tumor extrachromosomal circular DNA (ecDNA) as a component of liquid biopsy in blood. *Med Hypotheses*. 2018;114:5–7. doi:10.1016/j.mehy.2018.02.018
32. Khatami F, Tavangar SM. Current diagnostic status of pheochromocytoma and future perspective: a mini review. *Iran J Pathol*. 2017;12(3):313–322.
33. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–605. doi:10.1007/s10654-010-9491-z
34. Wells G, Shea B, O'connell D, et al. *The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomised Studies in Meta-analyses*. Ottawa Hospital Research Institute, 2014. oxford: ASP; 2015.
35. Morris M, Maina E, Morgan N, et al. Molecular genetic analysis of FH1, FH, and SDHB candidate tumour suppressor genes in renal cell carcinoma. *J Clin Pathol*. 2004;57(7):706–711. doi:10.1136/jcp.2003.011767
36. Vanharanta S, Buchta M, McWhinney SR, et al. Early-onset renal cell carcinoma as a novel extraparaganglial component of SDHB-associated heritable paraganglioma. *Am J Hum Genet*. 2004;74(1):153–159. doi:10.1086/381054
37. Cornejo KM, Lu M, Yang P, et al. Succinate dehydrogenase B: a new prognostic biomarker in clear cell renal cell carcinoma. *Hum Pathol*. 2015;46(6):820–826. doi:10.1016/j.humpath.2015.02.013
38. Jiang Q, Zhang Y, Zhou Y-H, et al. A novel germline mutation in SDHA identified in a rare case of gastrointestinal stromal tumor complicated with renal cell carcinoma. *Int J Clin Exp Pathol*. 2015;8(10):12188.
39. Housley SL, Lindsay RS, Young B, et al. Renal carcinoma with giant mitochondria associated with germ-line mutation and somatic loss of the succinate dehydrogenase B gene. *Histopathology*. 2010;56(3):405–408. doi:10.1111/j.1365-2559.2010.03482.x
40. Gill AJ, Hes O, Papathomas T, et al. Succinate dehydrogenase (SDH)-deficient renal carcinoma: a morphologically distinct entity: a clinicopathologic series of 36 tumors from 27 patients. *Am J Surg Pathol*. 2014;38(12):1588. doi:10.1097/PAS.0000000000000115
41. Calió A, Grignon DJ, Stohr BA, Williamson SR, Eble JN, Cheng L. Renal cell carcinoma with TFE3 translocation and succinate dehydrogenase B mutation. *Mod Pathol*. 2017;30(3):407. doi:10.1038/modpathol.2016.200
42. Carter CS, Skala SL, Chinnaiyan AM, et al. Immunohistochemical characterization of fumarate hydratase (FH) and succinate dehydrogenase (SDH) in cutaneous leiomyomas for detection of familial cancer syndromes. *Am J Surg Pathol*. 2017;41(6):801. doi:10.1097/PAS.0000000000000840
43. Casey RT, Warren AY, Martin JE, et al. Clinical and molecular features of renal and pheochromocytoma/paraganglioma tumor association syndrome (RAPTAS): case series and literature review. *J Clin Endocrinol Metab*. 2017;102(11):4013–4022. doi:10.1210/jc.2017-00562
44. Kim E, Wright MJ, Sioson L, et al. Utility of the succinate: fumarate ratio for assessing SDH dysfunction in different tumor types. *Mol Genet Metab Rep*. 2017;10:45–49. doi:10.1016/j.ymgmr.2016.12.006
45. Lehtonen HJ, Mäkinen MJ, Kiuru M, et al. Increased HIF1 α in SDH and FH deficient tumors does not cause microsatellite instability. *Int J Cancer*. 2007;121(6):1386–1389. doi:10.1002/ijc.22819
46. Falzarano SM, McKenney JK, Montironi R, et al. Renal cell carcinoma occurring in patients with prior neuroblastoma. *Am J Surg Pathol*. 2016;40(7):989–997. doi:10.1097/PAS.0000000000000632
47. Henderson A, Douglas F, Perros P, Morgan C, Maher E. SDHB-associated renal oncocytoma suggests a broadening of the renal phenotype in hereditary paragangliomatosis. *Fam Cancer*. 2009;8(3):257–260. doi:10.1007/s10689-009-9234-z
48. Paik JY, Toon CW, Benn DE, et al. Renal carcinoma associated with succinate dehydrogenase B mutation: a new and unique subtype of renal carcinoma. *J Clin Oncol*. 2014;32(6):e10–e3. doi:10.1200/JCO.2012.47.2647
49. Michalova K, Steiner P, Alaghebandan R, et al. Papillary renal cell carcinoma with cytologic and molecular genetic features overlapping with renal oncocytoma: analysis of 10 cases. *Ann Diagn Pathol*. 2018;35:1–6. doi:10.1016/j.anndiagpath.2018.01.010
50. Ricketts CJ, Shuch B, Vocke CD, et al. Succinate dehydrogenase kidney cancer (SDH-RCC): an aggressive example of the warburg effect in cancer. *J Urol*. 2012;188(6). doi:10.1016/j.juro.2012.08.030
51. Sonker P, Tewari AK, Chaube SK, et al. World Journal of Pharmaceutical Sciences. 2018.
52. Network CGAR. Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature*. 2013;499(7456):43. doi:10.1038/nature12222
53. DeBerardinis RJ, Thompson CB. Cellular metabolism and disease: what do metabolic outliers teach us? *Cell*. 2012;148(6):1132–1144. doi:10.1016/j.cell.2012.02.032
54. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *cell*. 2011;144(5):646–674. doi:10.1016/j.cell.2011.02.013
55. Scheffler IE. Molecular genetics of succinate: quinone oxidoreductase in eukaryotes. In: *Progress in Nucleic Acid Research and Molecular Biology*. Vol. 60. Elsevier; 1998:267–315.
56. Bardella C, Pollard PJ, Tomlinson I. SDH mutations in cancer. *Biochim Biophys Acta Bioenerg*. 2011;1807(11):1432–1443. doi:10.1016/j.bbabi.2011.07.003
57. Hao X, Al-Ahmadie HA, Gopalan A, et al. editors. Succinate dehydrogenase (SDH)-deficient renal cell carcinoma, a report of 7 cases with an emphasis on high grade morphologic features. In: *Laboratory Investigation*. New York, NY: Nature Publishing Group; 2016
58. Srigley JR, Delahunt B, Eble JN, et al. The International Society of Urological Pathology (ISUP) vancouver classification of renal neoplasia. *Am J Surg Pathol*. 2013;37(10):1469–1489. doi:10.1097/PAS.0b013e318299f2d1
59. Kuroda N, Yorita K, Nagasaki M, et al. Review of succinate dehydrogenase-deficient renal cell carcinoma with focus on clinical and pathobiological aspects. *Pol J Pathol*. 2016;67(1):3–7.
60. Yakirevich E, Ali SM, Mega A, et al. A novel SDHA-deficient renal cell carcinoma revealed by comprehensive genomic profiling. *Am J Surg Pathol*. 2015;39(6):858–863. doi:10.1097/PAS.0000000000000403
61. Ricketts CJ, Forman JR, Rattenberry E, et al. Tumor risks and genotype-phenotype-proteotype analysis in 358 patients with germline mutations in SDHB and SDHD. *Hum Mutat*. 2010;31(1):41–51. doi:10.1002/humu.21136
62. Gill AJ, Pachter NS, Clarkson A, et al. Renal tumors and hereditary pheochromocytoma-paraganglioma syndrome type 4. *N Engl J Med*. 2011;364(9):885–886. doi:10.1056/NEJMc1012357
63. Saxena N, Maio N, Crooks DR, et al. SDHB-deficient cancers: the role of mutations that impair iron sulfur cluster delivery. *J Natl Cancer Inst*. 2016;108(1):djv287. doi:10.1093/jnci/djv287
64. Evenepoel L, Papathomas TG, Krol N, et al. Toward an improved definition of the genetic and tumor spectrum associated with SDH germ-line mutations. *Genet Med*. 2015;17(8):610. doi:10.1038/gim.2014.162
65. Maio N, Singh A, Uhrigshardt H, Saxena N, Tong W-H, Rouault TA. Co-chaperone binding to LYR motifs confers specificity of iron sulfur cluster delivery. *Cell Metab*. 2014;19(3):445–457. doi:10.1016/j.cmet.2014.01.015

66. Gimenez-Roqueplo A-P, Favier J, Rustin P, et al. Functional consequences of a SDHB gene mutation in an apparently sporadic pheochromocytoma. *J Clin Endocrinol Metab.* 2002;87(10):4771–4774. doi:10.1210/jc.2002-020525
67. Andrews KA, Ascher DB, Pires DEV, et al. Tumour risks and genotype-phenotype correlations associated with germline variants in succinate dehydrogenase subunit genes SDHB, SDHC and SDHD. *J Med Genet.* 2018;55(6):384–394. doi:10.1136/jmedgenet-2017-105127
68. Niemeijer ND, Rijken JA, Eijkelenkamp K, et al. The phenotype of SDHB germline mutation carriers; a nationwide study. *Eur J Endocrinol.* 2017;177(2):115–125.
69. Berry WL, Janknecht R. KDM4/JMJD2 histone demethylases: epigenetic regulators in cancer cells. *Cancer Res.* 2013;73:2936–2942. doi:10.1158/0008-5472.CAN-12-4300
70. Rose NR, Woon EC, Kingham GL, et al. Selective inhibitors of the JMJD2 histone demethylases: combined nondenaturing mass spectrometric screening and crystallographic approaches. *J Med Chem.* 2010;53(4):1810–1818. doi:10.1021/jm901680b
71. Liang Y, Vogel JL, Arbuckle JH, et al. Targeting the JMJD2 histone demethylases to epigenetically control herpesvirus infection and reactivation from latency. *Sci Transl Med.* 2013;5(167):167ra5–ra5. doi:10.1126/scitranslmed.3005145
72. Piruat JJ, Millán-Uclés Á. Genetically modeled mice with mutations in mitochondrial metabolic enzymes for the study of cancer. *Front Oncol.* 2014;4:200. doi:10.3389/fonc.2014.00200
73. Selak MA, Armour SM, MacKenzie ED, et al. Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF- α prolyl hydroxylase. *Cancer Cell.* 2005;7(1):77–85. doi:10.1016/j.ccr.2004.11.022
74. Killian JK, Kim SY, Miettinen M, et al. Succinate dehydrogenase mutation underlies global epigenomic divergence in gastrointestinal stromal tumor. *Cancer Discov.* 2013;3:648–657. doi:10.1158/2159-8290.CD-13-0092
75. Letouze E, Martinelli C, Liorot C, et al. SDH mutations establish a hypermethylator phenotype in paraganglioma. *Cancer Cell.* 2013;23(6):739–752. doi:10.1016/j.ccr.2013.04.018
76. Baysal BE, Ferrell RE, Willett-Brozick JE, et al. Mutations in SDHD, a mitochondrial complex II gene, in hereditary paraganglioma. *Science.* 2000;287(5454):848–851. doi:10.1126/science.287.5454.848
77. Hao H-X, Khalimonchuk O, Schraders M, et al. SDH5, a gene required for flavination of succinate dehydrogenase, is mutated in paraganglioma. *Science.* 2009;325(5944):1139–1142. doi:10.1126/science.1175689
78. Andrici J, Gill AJ, Hornick JL, editors. Next generation immunohistochemistry: emerging substitutes to genetic testing? Seminars in diagnostic pathology; 2017; Elsevier.
79. Jackson CB, Nuoffer JM, Hahn D, et al. Mutations in SDHD lead to autosomal recessive encephalomyopathy and isolated mitochondrial complex II deficiency. *J Med Genet.* 2013. doi:10.1136/jmedgenet-2013-101932
80. Gill AJ, Lipton L, Taylor J, et al. Germline SDHC mutation presenting as recurrent SDH deficient GIST and renal carcinoma. *Pathol J RCPA.* 2013;45(7):689–691.
81. Rattenberry E, Vialard L, Yeung A, et al. A comprehensive next generation sequencing-based genetic testing strategy to improve diagnosis of inherited pheochromocytoma and paraganglioma. *J Clin Endocrinol Metab.* 2013;98(7):E1248–E56. doi:10.1210/jc.2013-1319
82. Dorschner MO, Amendola LM, Turner EH, et al. Actionable, pathogenic incidental findings in 1,000 participants' exomes. *Am J Hum Genet.* 2013;93(4):631–640. doi:10.1016/j.ajhg.2013.08.006
83. Betts MJ, Russell RB. Amino-acid properties and consequences of substitutions. *Bioinf Geneticists.* 2007;2:311–339.
84. Bayley J-P, van Minderhout I, Weiss MM, et al. Mutation analysis of SDHB and SDHC: novel germline mutations in sporadic head and neck paraganglioma and familial paraganglioma and/or pheochromocytoma. *BMC Med Genet.* 2006;7(1):1. doi:10.1186/1471-2350-7-1
85. Bayley J-P, Devilee P, Taschner PE. The SDH mutation database: an online resource for succinate dehydrogenase sequence variants involved in pheochromocytoma, paraganglioma and mitochondrial complex II deficiency. *BMC Med Genet.* 2005;6(1):39. doi:10.1186/1471-2350-6-39
86. Astuti D, Hart-Holden N, Latif F, et al. Genetic analysis of mitochondrial complex II subunits SDHD, SDHB and SDHC in paraganglioma and pheochromocytoma susceptibility. *Clin Endocrinol (Oxf).* 2003;59(6):728–733. doi:10.1046/j.1365-2265.2003.01914.x
87. Schiavi F, Boedeker CC, Bausch B, et al. Predictors and prevalence of paraganglioma syndrome associated with mutations of the SDHC gene. *Jama.* 2005;294(16):2057–2063. doi:10.1001/jama.294.16.2057
88. Tufton N, Ghelani R, Srirangalingam U, et al. A novel SDHA-deficient renal cell carcinoma revealed by comprehensive genomic profiling. *Endocr Relat Cancer.* 2017;24(7):T91–T103.
89. Miettinen M, Lasota J. Succinate dehydrogenase deficient gastrointestinal stromal tumors (GISTs)—A review. *Int J Biochem Cell Biol.* 2014;53:514–519. doi:10.1016/j.biocel.2014.05.033
90. Miettinen M, Killian JK, Wang Z-F, et al. Immunohistochemical loss of succinate dehydrogenase subunit A (SDHA) in gastrointestinal stromal tumors (GISTs) signals SDHA germline mutation. *Am J Surg Pathol.* 2013;37(2):234. doi:10.1097/PAS.0b013e3182671178
91. Ogino S, Lochhead P, Chan AT, et al. Molecular pathological epidemiology of epigenetics: emerging integrative science to analyze environment, host, and disease. *Mod Pathol.* 2013;26(4):465.
92. Ogino S, Nishihara R, VanderWeele TJ, et al. The role of molecular pathological epidemiology in the study of neoplastic and non-neoplastic diseases in the era of precision medicine. *Epidemiology (Cambridge, Mass).* 2016;27(4):602. doi:10.1097/EDE.0000000000000471
93. Hamada T, Keum N, Nishihara R, Ogino S. Molecular pathological epidemiology: new developing frontiers of big data science to study etiologies and pathogenesis. *J Gastroenterol.* 2017;52(3):265–275. doi:10.1007/s00535-016-1272-3

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress