

Hemitruncus Arteriosus in a 10-Day-Old Neonate with Patent Ductus Arteriosus and Thrombocytopenia

This article was published in the following Dove Press journal:
Vascular Health and Risk Management

Ali Zolfi-Gol¹
Mohammad Radvar¹
Amin Sedokani² 

¹Department of Pediatrics, Pediatric Cardiology, Urmia University of Medical Sciences, Urmia, Iran; ²Student Research Committee, Urmia University of Medical Sciences, Urmia, Iran

Abstract: Hemitruncus arteriosus refers to an uncommon congenital cardiovascular abnormality. It usually presents in infancy and leads to the development of pulmonary hypertension, heart failure, and in severe cases maybe death. Herein, we reported hemitruncus arteriosus in a 10-day-old neonate with respiratory distress, patent ductus arteriosus, and thrombocytopenia.

Keywords: pulmonary artery, cardiac anomaly, Hemitruncus arteriosus, patent ductus arteriosus, thrombocytopenia

Introduction

In 1868, Fraentzel described an abnormal origin of one pulmonary artery in a 25-year-old woman as hemitruncus arteriosus for the first time. Hemitruncus arteriosus (HA) is an uncommon congenital cardiovascular abnormality which refers to the abnormal originating of one branch of the pulmonary artery from the aorta with the other branch arising normally from the right ventricle. Mostly the condition is diagnosed in infancy but rarely it may be undiagnosed until adulthood. Commonly, in most HA cases, the right pulmonary artery (RPA) originates from the ascending aorta next to the aortic valve. In most patients, the right pulmonary artery originates from the posterior side of the ascending aorta next to the aortic valve. In some patients, the left pulmonary artery originates from the ascending aorta, which is usually associated with the right aortic arch.¹⁻³ The persistent flow of systemic high-pressure blood causes progressive diseases like pulmonary hypertension, and death in severe cases.^{3,4} The 3-month and first-year mortality rates of HA are 30% and 70% respectively which the condition can be corrected by early cardiac surgery.⁵⁻⁷

Case Presentation

A 10-day-old male neonate with birth bodyweight of 3700 g, firstborn of non-relative parents in which the pregnancy was induced by intrauterine insemination (IUI) and the baby was born by cesarian section (C/S) with a gestational age of 40 weeks and good Apgar score was referred to the pediatric emergency department of Motahari hospital of Urmia. According to the mother's statements, the chief complaints were shortness of breath, rapid breathing, and vomiting. Also, the patient had a history of NICU admission on 2nd day after birth because of

Correspondence: Amin Sedokani
Tel +98 914 324 1375
Fax +98 443 222 2090
Email A.sedokani@gmail.com

dehydration and poor feeding for 1st day. The patient could not take enough oral feeding for the last 2 days and vomiting was progressive.

On physical examinations, the anthropometric measures were normal, the patient was non-dysmorphic and chest X-ray showed mild to moderate cardiomegaly with bilateral perihilar infiltrates. Vital signs were as follows; PR: 162/min, RR: 62/min, BT: 36.5°C, and oxygen saturation of 94% (room air). Mucous membranes were dry but skin turgor and fontanels were normal. The chest shape was normal and symmetric with subcostal retraction and tachypnea. Cardiac auscultation revealed a grade II/VI continuous murmur. Peripheral pulses were palpable. On abdominal examination, the liver span was increased on palpitation and generalized edema was obvious. Primitive neonatal reflexes such as Moro, sucking, and rooting were normal.

The urgent ABG evaluation revealed PH of 7.14, PaCO₂ of 29 mmHg, HCO₃ of 9 mEq/L, and PaO₂ of 43 mmHg. Serum electrolytes, BUN, creatinine, serum albumin, and blood sugar were in normal ranges. Primary complete blood count (CBC) and coagulation tests were normal except for the platelet count and INR. The patient had thrombocytopenia (platelet count of 95×10³/mm³) and an INR of 1.19.

Due to severe pulmonary distress, the patient was intubated and admitted to NICU. After stabilizing the vital signs and hydration, a pediatric cardiology consultation was done. Echocardiography demonstrated no visible RPA, PDA with right to left flow, mild supravalvular aortic stenosis, mild aortic insufficiency, a severe reversal flow in aortic arch, secundum type ASD, dilated right atrium, right ventricle, and main pulmonary artery, severe tricuspid regurgitation and severe pulmonary hypertension, moderate mitral regurgitation and pulmonary insufficiency, dilated inferior vena cava, mild bilateral pleural effusion, and severe ascites. Due to the findings, close cardiological follow up and CT-angiography was performed.

Multi-slice computed tomography of the heart and major vasculature with special reconstructed views (dynamic and delayed images with contrast medium) revealed; anomalous arising of RPA from ascending aorta (hemitruncus), PDA 3.5 mm in size, stretched PFO (6 mm), atelectasis and congestion in both lungs with pleural effusion (Figure 1).

Serious efforts to perform cardiac surgery in addition to medical therapy such as inotropes and diuretics began. However, pediatric cardiac surgery was not available in




Figure 1 CT angiogram image showing abnormal origin of RPA from ascending aorta (AAO).

Abbreviations: SVC, superior vena cava; MPA, main pulmonary artery; DAO, descending aorta; LPA, left pulmonary artery.

the health care center or nearby cities. So, the patient was maintained on medical therapy and supportive care until cardiac surgery was a possibility.

Gradually, the patient's respiratory distress increased and liver function tests were more impaired and eventually cardiopulmonary arrest occurred on the 24th day of admission in the NICU. Immediate CPR was begun by cardiac massage, intravenous epinephrine, and positive pressure ventilation. After 30 mins the patient had no vital response and CPR was stopped. The last blood tests revealed metabolic and respiratory acidosis. The last blood test values were in Table 1.

Discussion

Hemitruncus arteriosus is an unusual cardiac structural abnormality that, as Agati et al described, "results from failure of development of the left sixth arch and persistence of the left fifth arch". In the disease, the pulmonary pressure increases with time and the mortality rate varies from 30% to 70% depending on case severity and receiving cardiac surgery as soon as possible.¹⁻⁸ Early surgical repair can prevent pulmonary vascular disease progression.⁴ The surgical repair should be done within the first six months to prevent severe pulmonary vasculature disease. In our case, the severity of the disease, refractory acidosis, and unavailability of immediate surgery and coagulopathy and thrombocytopenia led to death.

Table I The Last Blood Tests of the Patient on 24th Day of NICU Admission

Test	Results	Unit	Reference Value
PH	7.18	–	7.35–7.45
PaCO ₂	51	mmHg	35–45
HCO ₃	18.6	mEq/L	22–26
PaO ₂	85	mmHg	80–100
BUN	16	mg/dL	7–20
Creatinine	1	mg/dL	0.5–1.3
AST	41	U/L	5–40
ALT	7	U/L	5–40
Blood sugar	<20	mg/dL	70–100
Serum Na	137	mEq/L	135–148
Serum K	3.6	mEq/L	3.5–5.5
Serum Ca	10.1	mg/dL	8.6–10.3
WBC	7.2	10 ³ /mm ³	4–12
RBC	3.08	10 ⁶ /mm ³	4.5–6.3
Hb	9.8	gr/dL	14–18
Hct	29.9	%	39–52
Platelet	63	10 ³ /mm ³	140–440
MCV	97.08	fL	77–97
MCH	31.82	Pgm	26–32
MCHC	32.78	%	32–36
PT	25	Sec	10.1–12.9
INR	3.2	–	0.9–1
PTT	39	Sec	25.4–38.4

Conclusion

The anomalous origin of the left pulmonary artery results from the failure of development of the left sixth arch and persistence of the left fifth arch. If left untreated, the pulmonary bed is vulnerable to early onset of pulmonary vascular obstructive disease owing to the large blood supply to both lungs. It is important to recognize this rare congenital cardiac anomaly early and contemplate surgical repair for a better prognosis.

Ethics and Consent Statement

The patient's parents had the opportunity not to accept the study. The patient's personal information remained confidential. There was no cost or harm to the patient through the study. All of the stages of the study were conducted under supervision and confirmation of the ethical committee of Taleghani hospital and Urmia medical sciences university.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Fraentzel O. Ein Fall von abnormer Communication der Aorta mit der Arteria pulmonalis. *Virchows Arch.* 1868;43(3):420–426. doi:10.1007/BF01915623
2. Haddadin RB. Left hemitruncus: a rare congenital heart condition. *East Mediterr Health J.* 2014;19 Suppl 3:S224–S226.
3. Cheng W, Xiao Y, Zhong Q, Wen R. Anomalous origin of left pulmonary artery branch from the aorta with Fallot's tetralogy. *Thorac Cardiovasc Surg.* 2008;56(7):432–434. doi:10.1055/s-2008-1038466
4. Penkoske P, Castaneda A, Fyler D, Van RP. Origin of pulmonary artery branch from ascending aorta. Primary surgical repair in infancy. *J Thorac Cardiovasc Surg.* 1983;85(4):537–545. doi:10.1016/S0022-5223(19)37538-5
5. Fontana GP, Spach MS, Effmann EL, Sabiston DC Jr. Origin of the right pulmonary artery from the ascending aorta. *Ann Surg.* 1987;206(1):102–113. doi:10.1097/00000658-198707000-00016
6. Amir G, Frenkel G, Bruckheimer E, et al. Anomalous origin of the pulmonary artery from the aorta: early diagnosis and repair leading to immediate physiological correction. *Cardiol Young.* 2010;20(6):654–659. doi:10.1017/S1047951110000892
7. Peng EW, Shanmugam G, Macarthur KJ, Pollock JC. Ascending aortic origin of a branch pulmonary artery—surgical management and long-term outcome. *Eur J Cardiothorac Surg.* 2004;26(4):762–766. doi:10.1016/j.ejcts.2004.07.007
8. Agati S, Sousa CG, Calvaruso FD, et al. Anomalous aortic origin of the pulmonary arteries: case series and literature review. *Ann Pediatr Cardiol.* 2019;12(3):248–253. doi:10.4103/apc.APC_89_18

Vascular Health and Risk Management

Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement

of metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/vascular-health-and-risk-management-journal>

Dovepress