

Treating COPD with PDE 4 inhibitors

William M Brown

VaxDesign Corp, Orlando, FL, USA

Abstract: While the pathogenesis of chronic obstructive pulmonary disease (COPD) is incompletely understood, chronic inflammation is a major factor. In fact, the inflammatory response is abnormal, with CD8⁺ T-cells, CD68⁺ macrophages, and neutrophils predominating in the conducting airways, lung parenchyma, and pulmonary vasculature. Elevated levels of the second messenger cAMP can inhibit some inflammatory processes. Theophylline has long been used in treating asthma; it causes bronchodilation by inhibiting cyclic nucleotide phosphodiesterase (PDE), which inactivates cAMP. By inhibiting PDE, theophylline increases cAMP, inhibiting inflammation and relaxing airway smooth muscle. Rather than one PDE, there are now known to be more than 50, with differing activities, substrate preferences, and tissue distributions. Thus, the possibility exists of selectively inhibiting only the enzyme(s) in the tissue(s) of interest. PDE 4 is the primary cAMP-hydrolyzing enzyme in inflammatory and immune cells (macrophages, eosinophils, neutrophils). Inhibiting PDE 4 in these cells leads to increased cAMP levels, down-regulating the inflammatory response. Because PDE 4 is also expressed in airway smooth muscle and, in vitro, PDE 4 inhibitors relax lung smooth muscle, selective PDE 4 inhibitors are being developed for treating COPD. Clinical studies have been conducted with PDE 4 inhibitors; this review concerns those reported to date.

Keywords: COPD, asthma, phosphodiesterase IV inhibitor

Introduction

Chronic obstructive pulmonary disease (COPD) is a serious and increasing global public health problem; physiologically, it is characterized by progressive, irreversible airflow obstruction and pathologically, by an abnormal airway inflammatory response to noxious particles or gases (MacNee 2005a). The COPD patient suffers a reduction in forced expiratory volume in 1 second (FEV₁), a reduction in the ratio of FEV₁ to forced vital capacity (FVC), compared with reference values, absolute reductions in expiratory airflow, and little improvement after treatment with an inhaled bronchodilator. Airflow limitation in COPD patients results from mucosal inflammation and edema, bronchoconstriction, increased secretions in the airways, and loss of elastic recoil. Patients with COPD can experience 'exacerbations,' involving rapid and prolonged worsening of symptoms (Seneff et al 1995; Connors et al 1996; Dewan et al 2000; Rodriguez-Roisin 2006; Mohan et al 2006). Many are idiopathic, though they often involve bacteria; airway inflammation in exacerbations can be caused or triggered by bacterial antigens (Murphy et al 2000; Blanchard 2002; Murphy 2006; Veeramachaneni and Sethi 2006). Increased IL-6, IL-1 β , TNF- α , GRO- α , MCP-1, and IL-8 levels are found in COPD patient sputum; their levels increase further during exacerbations. COPD has many causes and significant differences in prognosis exist, depending on the cause (Barnes 1998; Madison and Irwin 1998).

COPD is already the fourth leading cause of death worldwide, according to the World Health Organization (WHO); the WHO estimates that by the year 2020, COPD will be the third-leading cause of death and the fifth-leading cause of disability worldwide (Murray and Lopez 1997). COPD is the fastest-growing cause of death

Correspondence: William M Brown
VaxDesign Corp, 2721 Discovery Drive,
Suite 400, Orlando, FL 32826, USA
Tel +1 407 249 3650
Fax +1 407 249 3649
Email wbrown@vaxdesign.com

in developed nations and is responsible for over 2.7 million deaths per year worldwide. In the US, there are currently estimated to be 16 million people with COPD. There are estimated to be up to 20 million sufferers in Japan, which has the world's highest per capita cigarette consumption and a further 8–12 million in Europe. In 2000, COPD accounted for over 20 million outpatient visits, 3.4 million emergency room visits, 6 million hospitalizations, and 116,500 deaths in the US (National Center for Health Statistics 2002). Factors associated with COPD, including immobility, often lead to secondary health consequences (Polkey and Moxham 2006).

Risk factors for the development of COPD include cigarette smoking, and occupational exposure to dust and chemicals (Senior and Anthonisen 1998; Anthonisen et al 2002; Fabbri and Hurd 2003; Zaher et al 2004). Smoking is the most common cause of COPD and the underlying inflammation typically persists in ex-smokers. Oxidative stress from cigarette smoke is also an issue in COPD (Domej et al 2006). Despite this, relatively few smokers ever develop COPD (Siafakas and Tzortzaki 2002).

While many details of the pathogenesis of COPD remain unclear, chronic inflammation is now recognized as a major factor, predominantly in small airways and lung parenchyma, characterized by increased numbers of macrophages, neutrophils, and T-cells (Barnes 2000; Stockley 2002). As recently as 1995, the American Thoracic Society issued a statement defining COPD without mentioning the underlying inflammation (American Thoracic Society 1995). Since then, the Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines have made it clear that chronic inflammation throughout the airways, parenchyma, and pulmonary vasculature plays a central role (Pauwels et al 2001; GOLD 2003). The comparatively recent realization of the role of airway inflammation in COPD has altered thinking with regard to potential therapies (Rogers and Giembycz 1998; Vignola 2004).

Most pharmacological therapies available for COPD, including bronchodilator and anti-inflammatory agents, were first developed for treating asthma. The mainstays of COPD treatment are inhaled corticosteroids (McEvoy and Niewoehner 1998; Borron and deBoisblanc 1998; Pauwels 2002; Gartlehner et al 2006; D'Souza 2006), supplemental oxygen (Petty 1998; Austin and Wood-Baker 2006), inhaled bronchodilators (Costello 1998; Doherty and Briggs 2004), and antibiotics (Taylor 1998), especially in severely affected patients (Anthonisen et al 1987; Saint et al 1995; Adams et al 2001; Miravittles et al 2002; Donnelly

and Rogers 2003; Sin et al 2003; Rabe 2006), though the use of antibiotics remains controversial (Ram et al 2006). Long-acting β_2 -agonists (LABAs) improve the mucociliary component of COPD. Combination therapy with LABAs and anticholinergic bronchodilators resulted in modest benefits and improved health-related quality of life (Buhl and Farmer 2005; Appleton et al 2006). Treatment with mucolytics reduced exacerbations and the number of days of disability (Poole and Black 2006). The combined use of inhaled corticosteroids and LABAs has been demonstrated to produce sustained improvements in FEV₁ and positive effects on quality of life, number of hospitalizations, distance walked, and exacerbations (Mahler et al 2002; Szafranski et al 2003; Sin et al 2004; Miller-Larsson and Selroos 2006; van Schayck and Reid 2006). However, all of these treatments are essentially palliative and do not impact COPD progression (Hay 2000; Gamble et al 2003; Antoniu 2006a).

A further complication in drug development and therapy is that it can be difficult to determine the efficacy of therapy, because COPD has a long preclinical stage, is progressive, and patients generally do not present for treatment until their lung function is already seriously impaired. Moreover, because COPD involves irreversible loss of elasticity, destruction of the alveolar wall, and peribronchial fibrosis, there is often little room for clinical improvement.

Smoking cessation remains the most effective intervention for COPD. Indeed, to date, it is the only intervention shown to stop the decline in lung function, but it does not resolve the underlying inflammation, which persists even in ex-smokers. Smoking cessation is typically best achieved by a multifactor approach, including the use of bupropion, a nicotine replacement product, and behavior modification (Richmond and Zwar 2003).

In COPD, there is an abnormal inflammatory response, characterized by a predominance of CD8⁺ T-cells, CD68⁺ macrophages, and neutrophils in the conducting airways, lung parenchyma, and pulmonary vasculature (Soto and Hanania 2005; O'Donnell et al 2006; Wright and Churg 2006). Inflammatory mediators involved in COPD include lipids, inflammatory peptides, reactive oxygen and nitrogen species, chemokines, cytokines, and growth factors. COPD pathology also includes airway remodeling and mucociliary dysfunction (mucus hypersecretion and decreased mucus transport). Corticosteroids reduce the number of mast cells, but CD8⁺ and CD68⁺ cells, and neutrophils, are little affected (Jeffery 2005). Inflammation in COPD is not suppressed by corticosteroids, consistent with it

being neutrophil-, not eosinophil-mediated. Corticosteroids also do not inhibit the increased concentrations of IL-8 and TNF- α (both neutrophil chemoattractants) found in induced sputum from COPD patients. Neutrophil-derived proteases, including neutrophil elastase and matrix metalloproteinases (MMPs), are involved in the inflammatory process and are responsible for the destruction of elastin fibers in the lung parenchyma (Mercer et al 2005; Gueders et al 2006). MMPs play important roles in the proteolytic degradation of extracellular matrix (ECM), in physiological and pathological processes (Corbel, Belleguic et al 2002). PDE 4 inhibitors can reduce MMP activity and the production of MMPs in human lung fibroblasts stimulated with pro-inflammatory cytokines (Lagente et al 2005). In COPD, abnormal remodeling results in increased deposition of ECM and collagen in lungs, because of an imbalance of MMPs and TIMPs (Jeffery 2001). Fibroblast/myofibroblast proliferation and activation also occur, increasing production of ECM-degrading enzymes (Crouch 1990; Segura-Valdez et al 2000). Additionally, over-expression of cytokines and growth factors stimulates lung fibroblasts to synthesize increased amounts of collagen and MMPs, including MMP-1 (collagenase-1) and MMP-2 and MMP-9 (gelatinases A and B) (Sasaki et al 2000; Zhu et al 2001).

It is now generally accepted that bronchial asthma is also a chronic inflammatory disease (Barnes et al 1988; Barnes 1995). The central role of inflammation of the airways in asthma's pathogenesis is consistent with the efficacy of corticosteroids in controlling clinical symptoms. Eosinophils are important in initiating and continuing the inflammatory state (Holgate et al 1987; Bruijnzeel 1989; Underwood et al 1994; Teixeira et al 1997), while other inflammatory cells, including lymphocytes, also infiltrate the airways (Holgate et al 1987; Teixeira et al 1997). The familiar acute symptoms of asthma are the result of airway smooth muscle contraction. While recognition of the key role of inflammation has led to an emphasis on anti-inflammatory therapy in asthma, a significant minority of patients remains poorly controlled and some exhibit accelerated declines in lung function, consistent with airway remodeling (Martin and Reid 2006). Reversal or prevention of structural changes in remodeling may require additional therapy (Burgess et al 2006).

There is currently no cure for asthma; treatment depends primarily on inhaled glucocorticoids to reduce inflammation (Taylor 1998; Petty 1998), and inhaled bronchodilators to reduce symptoms (Torphy 1994; Costello 1998; Georgitis 1999; DeKorte 2003). Such treatments, however, do not address disease progression.

COPD and asthma are both characterized by airflow obstruction, but they are distinct in terms of risk factors and clinical presentation. While both involve chronic inflammation and cellular infiltration and activation, different cell types are implicated and there are differences in the inflammatory states (Giembycz 2000; Fabbri and Hurd 2003; Barnes 2006). In COPD, neutrophil infiltration into the airways and their activation appear to be key (Stockley 2002); in asthma, the inflammatory response involves airway infiltration by activated eosinophils and lymphocytes, and T-cell activation of the allergic response (Holgate et al 1987; Sætta et al 1998; Barnes 2006). While macrophages are present in both conditions, the major controller cells are CD8⁺ T-cells in COPD (O'Shaughnessy et al 1997; Sætta et al 1998) and CD4⁺ T-cells in asthma. IL-1, IL-8, and TNF- α are the key cytokines in COPD, while in asthma, IL-4, IL-5, and IL-13 are more important. There are differences in histopathological features of lung biopsies between COPD patients and asthmatics; COPD patients have many fewer eosinophils in lung tissue than asthmatics.

While the early phases of COPD and asthma are distinguishable, there are common features, including airway hyper-responsiveness and mucus hypersecretion. MUC5AC is a major mucin gene expressed in the airways; its expression is increased in COPD and asthmatic patients. At least in vitro, epidermal growth factor stimulates MUC5AC mRNA and protein expression; this can be reversed by PDE 4 inhibitors, which may contribute to their clinical efficacy in COPD and asthma (Mata et al 2005). Similar structural and fibrotic changes make COPD and asthma much less distinguishable in extreme cases; the chronic phases of both involve inflammatory responses, alveolar detachment, mucus hypersecretion, and subepithelial fibrosis. The two conditions have been linked epidemiologically; adults with asthma are up to 12 times more likely to develop COPD over time than those without (Guerra 2005).

Theophylline

Theophylline and related xanthine compounds have been used for decades to treat asthma (Weinberger 1988; Torphy and Undem 1991; Manganiello et al 1995; Dent and Giembycz 1996; Weinberger and Hendeles 1996; Torphy 1998; Ram et al 2002; Barnes 2003; Barr et al 2003). The use of these drugs has been limited, however, by their side effects and modest efficacy (Persson 1986; Rabe et al 1995). Additionally, theophylline is a difficult drug to use, requiring titration and plasma monitoring, because of the risk of cardiovascular and CNS side effects, even at therapeutic doses (Boswell-Smith, Cazzola et al 2006).

The second messenger cyclic 3',5'-adenosine monophosphate (cAMP) controls many cellular functions and it is well established that an elevated cAMP level can inhibit some inflammatory processes. Thus, inhibitors of enzymes that catalyze cAMP hydrolysis would seem to be good candidates to treat inflammatory conditions.

Phosphodiesterase (PDE)

Theophylline is believed to cause bronchodilation by inhibiting cyclic nucleotide phosphodiesterase (PDE), an enzyme that catalyzes the hydrolysis of cAMP and cyclic 3',5'-guanosine monophosphate (cGMP) to inactive 5'-nucleotide products (Muller et al 1996). cAMP and cGMP exhibit many intracellular effects, mediated largely through their stimulatory effect on multisubstrate protein kinases (Torphy and Undem 1991; Montminy 1997; Daniel et al 1998; Spina 2003). By inhibiting PDE, theophylline increases the level of cAMP and cGMP, resulting in relaxation of airway smooth muscle and inhibition of inflammatory cell activation (Holgate et al 1987; Bryson and Rodger 1987; Schramm and Grunstein 1992; Cortijo et al 1993; Kotlikoff and Kamm 1996; Spina 2003).

Theophylline is now known to have many properties in addition to that of a bronchodilator (Persson 1986; Sullivan et al 1994; Rabe et al 1995; D'Alonzo 1996; Weinberger and Hendeles 1996; Vassallo and Lipsky 1998). Theophylline also causes pulmonary arterial vasodilatation, enhances diaphragmatic contractility, and increases CNS respiratory drive. Theophylline is a cardiac ionotrope and chromotrope. It is also a weak diuretic and increases mucociliary sweep. Theophylline has anti-inflammatory effects in COPD, reducing neutrophil counts, IL-8, and the total number inflammatory cells in sputum. Theophylline is also subject to many drug interactions and has adenosine receptor antagonist activity (Barnes 2003).

Far from there being a single PDE, it is now clear that there are many with differing activities, substrate preferences, and tissue distributions (Nicholson et al 1991; Thompson 1991; Lowe and Cheng 1992; Beavo et al 1994; Manganiello et al 1995; Torphy 1998; Silver et al 1988; Matsumoto et al 2003); theophylline is actually a non-specific PDE inhibitor (Persson 1986; Rabe et al 1995; D'Alonzo 1996; Weinberger and Hendeles 1996; Vassallo and Lipsky 1998). Some of its various properties and side-effects have been attributed to non-selective inhibition of PDEs (Barnes 2003).

Indeed, there are believed to be at least 11 gene families of PDE enzymes in mammals, encoding more than 50 enzymes, because of alternative splicing and alternative transcriptional

start sites (Bolger 1994; Lobban et al 1994; Horton et al 1995; Bushnik and Conti 1996; Jin et al 1998; Houslay et al 1998; Conti and Jin 1999; Giembycz 2000; Giembycz 2001; Scapin et al 2004; Giembycz 2005a; Bender and Beavo 2006). PDE enzymes share approximately 25% sequence identity over the conserved catalytic domain of about 300 amino acids (Ke 2004). While all PDEs catalyze the hydrolysis of cAMP and/or cGMP, the enzymes differ in their biochemical and pharmacological properties and exhibit different affinities for various inhibitors (Silver et al 1988; Torphy and Undem 1991; Manganiello et al 1995; Muller et al 1996; Torphy 1998). PDE 4, PDE 7, and PDE 8 are specific for cAMP (Conti and Yin 1999; Soderling and Beavo 2000). This diversity of enzyme type and tissue-specific expression raises the possibility of selectively inhibiting only the enzyme(s) in the tissue(s) of interest, if sufficiently specific inhibitors can be found (Giembycz and Dent 1992; Card et al 2004).

PDE 4 in COPD

With regard to COPD, PDE 4 is the primary cAMP-hydrolyzing enzyme in inflammatory and immune cells, especially macrophages, eosinophils, and neutrophils, all of which are found in the lungs of COPD and asthma patients (Torphy et al 1992; Karlsson and Aldous 1997; De Brito et al 1997; Wang et al 1999; Torphy and Page 2000). Inhibition of PDE 4 leads to elevated cAMP levels in these cells, down-regulating the inflammatory response (Dyke and Montana 2002).

PDE 4 has also attracted much attention because it is expressed in airway smooth muscle (Ashton et al 1994; Undem et al 1994; Nicholson et al 1995; Kerstjens and Timens 2003; Mehats et al 2003; Lipworth 2005; Fan Chung 2006). In vitro, PDE 4 inhibitors relax lung smooth muscle (Undem et al 1994; Dent and Giembycz 1995). In COPD and asthma, a selective PDE 4 inhibitor with combined bronchodilatory and anti-inflammatory properties would seem desirable (Nicholson and Shahid 1994; Lombardo 1995; Palfreyman 1995; Cavalia and Frith 1995; Palfreyman and Souness 1996; Karlsson and Aldous 1997; Compton et al 2001; Giembycz 2002; Jacob et al 2002; Soto and Hanania 2005).

PDE 4 inhibitors in COPD

So, because PDE 4 inhibitors suppress inflammatory functions in several cell types involved in COPD and asthma (Huang and Mancini 2006) and because, at least in vitro, PDE 4 inhibitors relax lung smooth muscle, selective PDE 4 inhibitors, originally intended for use in treating depression (Renau 2004), have been developed for the treatment of

COPD and asthma (Torphy et al 1999; Spina 2000; Huang et al 2001; Spina 2004; Gienbycz 2005a, 2005b; Lagente et al 2005; Boswell-Smith, Spina et al 2006). PDE 4 enzymes are strongly inhibited by the antidepressant drug rolipram (Pinto et al 1993), which decreases the influx of inflammatory cells at sites of inflammation (Lagente et al 1994; Lagente et al 1995; Alves et al 1996). PDE 4 inhibitors down-regulate cytokine production in inflammatory cells, *in vivo* and *in vitro* (Undem et al 1994; Dent and Gienbycz 1995). TNF- α is an important inflammatory cytokine in COPD; its release is reduced by PDE 4 inhibitors (Souness et al 1996; Chambers et al 1997; Griswold et al 1998; Gonçalves de Moraes et al 1998; Corbel, Belleguic et al 2002). Some PDE 4 inhibitors, including cilomilast and AWD 12-281, can inhibit neutrophil degranulation, a property not shared by theophylline (Ezeamuzie 2001; Jones et al 2005). PDE 4 inhibitors reduce overproduction of other pro-inflammatory mediators, including arachidonic acid and leukotrienes (Torphy 1998). PDE 4 inhibitors also inhibit cellular trafficking and microvascular leakage, production of reactive oxygen species, and cell adhesion molecule expression *in vitro* and *in vivo* (Sanz et al 2005). PDE 4 inhibitors, including cilomilast and CI-1044, inhibit LPS-stimulated TNF- α production in whole blood from COPD patients (Burnouf et al 2000; Ouagued et al 2005).

There are now thought to be at least four PDE 4s, A, B, C, and D, derived from four genes (Lobbam et al 1994; Muller et al 1996; Torphy 1998; Conti and Jin 1999; Matsumoto et al 2003). Alternative splicing and alternative promoters add further complexity (Manganiello et al 1995; Horton et al 1995; Torphy 1998). Indeed, the four genes encode more than 16 PDE 4 isoforms, which can be divided into short (~65–75 kDa) and long forms (~80–130 kDa); the difference between the short and long forms lies in the N-terminal region (Bolger et al 1997; Huston et al 2006). PDE 4 isoforms are regulated by extracellular signal-related protein kinase (ERK), which can phosphorylate PDE 4 (Houslay and Adams 2003).

The four PDE 4 genes are differentially expressed in various tissues (Silver et al 1988; Lobbam et al 1994; Manganiello et al 1995; Horton et al 1995; Muller et al 1996; Torphy 1998). PDE 4A is expressed in many tissues, but not in neutrophils (Wang et al 1999). PDE 4B is also widely expressed and is the predominant PDE 4 subtype in monocytes and neutrophils (Wang et al 1999), but is not found in cortex or epithelial cells (Jin et al 1998). Upregulation of the PDE 4B enzyme in response to pro-inflammatory agents suggest that it has a role in inflammatory processes (Manning et al 1999). PDE 4C is expressed in lung and testis, but not in circulating inflammatory cells, cortex, or hippocampus

(Oberholte et al 1997; Manning et al 1999; Martin-Chouly et al 2004). PDE 4D is highly expressed in lung, cortex, cerebellum, and T-cells (Erdogan and Houslay 1997; Jin et al 1998). PDE 4D also plays an important role in airway smooth muscle contraction (Mehats et al 2003).

A major issue with early PDE 4 inhibitors was their side effect profile; the signature side effects are largely gastrointestinal (nausea, vomiting, increased gastric acid secretion) and limited the therapeutic use of PDE 4 inhibitors (Dyke and Montana 2002). The second generation of more selective inhibitors, such as cilomilast and roflumilast, have improved side effect profiles and have shown clinical efficacy in COPD and asthma (Barnette 1999; Spina 2000; Lagente et al 2005). However, even cilomilast and roflumilast, the most advanced clinical candidates, discussed below, cause some degree of emesis (Spina 2003).

It is now thought that the desirable anti-inflammatory properties and unwanted side effects of nausea and emesis are associated with distinct biochemical activities (Torphy et al 1992; Jacobitz et al 1996; Barnette et al 1996; Souness et al 1997; Souness and Rao 1997). Specifically, the side effects are believed to be associated with the so-called 'high-affinity rolipram binding site' (HARBS) (Barnette et al 1995; Muller et al 1996; Jacobitz et al 1996; Kelly et al 1996; Torphy 1998) and/or inhibition of the form of PDE 4 found in the CNS (Barnette et al 1996). The exact nature of HARBS remains unclear, although it has been described as a conformer of PDE 4 (Souness and Rao 1997; Barnette et al 1998). Using mice deficient in PDE 4B or PDE 4D, it appears that emesis is the result of selective inhibition of PDE 4D (Robichaud et al 2002; Lipworth 2005), which is unfortunate, because the most clinically advanced PDE 4 inhibitors are selective for PDE 4D. Also, from animal studies, it appears that the nausea and vomiting are produced via the CNS, though there may also be direct effects on the gastrointestinal system (Barnette 1999).

While beyond the scope of this review, it has been proposed that PDE 4 inhibitors may be useful in treating inflammatory bowel disease (Banner and Trevethick 2004), cystic fibrosis (Liu et al 2005), pulmonary arterial hypertension (Growcott et al 2006), myeloid and lymphoid malignancies (Lerner and Epstein 2006), Alzheimer's disease (Ghavami et al 2006), rheumatoid arthritis and multiple sclerosis (Dyke and Montana 2002), infection-induced preterm labor (Oger et al 2004), depression (Wong et al 2006), and allergic disease (Crocker and Townley 1999). Varying degrees of *in vitro*, *in vivo*, and clinical data exist to support these claims.

So, after that theoretical buildup, we reach the proof of the pudding; clinical studies have been conducted with PDE 4 inhibitors. A potent, but not-very-selective, PDE 4 inhibitor is approved in Japan and is used clinically, including for treating asthma. Another is awaiting approval in the US. One is in advanced clinical development and others are at earlier stages.

Ibutilast

The drug ibutilast (3-isobutyl-2-isopropylpyrazolo[1,5-a]pyridine) is a nonselective PDE inhibitor. It is approved in Japan and has been widely used to treat bronchial asthma and ischemic stroke. Ibutilast preferentially inhibits PDE 3A, PDE 4, PDE 10, and PDE 11. Ibutilast potently inhibits purified human PDE 4A, 4B, 4C and 4D with IC_{50} values of 54, 65, 239 and 166 nM, respectively (Huang et al 2006). It may be useful in treating a range of neurological conditions, linked to its ability to elevate cellular cyclic nucleotide concentrations (Gibson et al 2006).

Cilomilast

Cilomilast is a second-generation PDE 4 inhibitor that was developed to separate activity at the HARBS and PDE 4 (Christensen et al 1998; Barnette et al 1998; Griswold et al 1998; Underwood et al 1998; Torphy et al 1999). Cilomilast is as strong an anti-inflammatory as rolipram, but causes much less nausea and gastric acid secretion. Cilomilast is also negatively charged at physiological pH, limiting its penetration into the CNS.

Cilomilast is being developed as a treatment for COPD; the drug has been assessed in phase III trials (Norman 1999; Norman 2000; Barnette and Underwood 2000; Torphy and Page 2000; Martina et al 2006). The compound had previously been in development for asthma, and phase II trials were conducted in the US and Japan in 2001; however, development for asthma was apparently discontinued.

Cilomilast is a potent ($K_i = 92$ nM), selective PDE 4 inhibitor (Christensen et al 1998; Barnette et al 1998; Griswold et al 1998; Underwood et al 1998). Cilomilast is considerably more selective for PDE 4D ($IC_{50} = 12$ nM) than 4A ($IC_{50} = 115$ nM), 4B ($IC_{50} = 86$ nM), or 4C ($IC_{50} = 308$ nM). The drug is essentially inactive against PDEs 1, 2, 3, 5, and 7 (Christensen et al 1998). Cilomilast inhibited human TNF- α production and PDE 4, and increased intracellular cAMP levels in both neutrophils and PBMCs (Chambers et al 1997). Cilomilast (10 μ M) inhibited the degradation of three-dimensional collagen gel by fibroblasts (Kohyama et al 2002).

The anti-inflammatory effects of cilomilast have been assessed in bronchial epithelial cells and sputum cells from smokers, COPD patients, and normal controls (Profita et al 2003). TNF- α and IL-8 were released at a significantly higher level in bronchial epithelial and sputum cells from patients with COPD than in controls or smokers. Cilomilast significantly reduced TNF- α release by bronchial epithelial and sputum cells, and GM-CSF release by sputum cells; IL-8 release was not significantly changed. Thus, cilomilast inhibited the production of some neutrophil chemoattractants by airway cells (Profita et al 2003). In bronchial biopsies from COPD patients, cilomilast treatment was associated with reductions in CD8⁺ and CD68⁺ cells; both cell types are increased in COPD and correlate with disease severity (Gamble et al 2003).

The cilomilast COPD clinical program included over 4000 patients in phase II and III trials. Evidence of safety and efficacy was based on four pivotal trials, involving 2883 patients. In addition, two phase-III open-label extension studies followed 1069 cilomilast patients for as long as three years. Inclusion criteria for the pivotal studies were patients between the ages of 40 and 80-years, with a diagnosis of COPD. Two primary endpoints were used: FEV₁ and total score on the St George's respiratory questionnaire (SGRQ), a self-administered questionnaire intended to determine the impact of chronic respiratory disease on health-related quality of life and well-being.

To date, four clinical trials have evaluated the efficacy of cilomilast and demonstrated improvement in lung function (FEV₁) and quality of life and reduction in the occurrence of COPD exacerbations, compared with placebo. Cilomilast was generally well tolerated, with adverse effects being overall mild and self-limiting.

The phase I and phase II studies demonstrated that cilomilast significantly improved lung function and quality of life to a clinically meaningful extent. A phase III program followed, to evaluate efficacy, safety, and mechanism of action.

By late 2003, GSK had performed four pivotal studies of cilomilast in a total of 2883 patients (n = 647 (study 039), 700 (study 042), 711 (study 091) and 825 (study 156)), comparing cilomilast (15 mg bid) with placebo over 24-weeks. Data from these trials, relating to various combinations of patients, have now been reported in several publications.

Phase III results in 2058 stable COPD patients were reported comparing cilomilast (15 mg bid for 6-months) with placebo. Cilomilast caused a sustained improvement in lung function and a reduction in the risk of exacerbation. Using

the SGRQ to assess quality of life, there was an improvement in health status in the cilomilast-treated group.

In a 6-month study, involving 647 patients with stable COPD (431 received 15 mg cilomilast bid and 216 received placebo), the cilomilast-treated patients exhibited an improved health status (assessed by SGRQ). These patients also demonstrated improved lung function (FEV_1 : 40 mL improvement over placebo), reduced healthcare resources utilization (physician visits, emergency room visits, hospitalization), and a lower rate of COPD exacerbation (39% lower than placebo).

After a 4-week, single-blind, run-in phase, 1411 patients with stable COPD received placebo or cilomilast (15 mg bid) for 24-weeks. FEV_1 was maintained in patients receiving cilomilast versus placebo, with a treatment difference of 300 mL. Cilomilast achieved a clinically significant reduction (26%) in the risk of moderate-to-severe COPD exacerbations, compared with placebo.

Thus, two of the four pivotal studies (studies 039 and 156) reached clinical significance and two (042 and 091) did not. There was a mean change in FEV_1 of 10 mL from baseline following cilomilast treatment in the two positive trials, compared with 20 and 30 mL reductions, for placebo in these trials (for studies 156 and 039, respectively).

Side effects and contraindications

GI-related side effects, including nausea, diarrhea, dyspepsia, vomiting, and abdominal pain, have been observed; they are believed to be dose-related and were monitored specifically because of preclinical studies finding vasculitis in cilomilast-treated mice and rats. Ischemic colitis (a consequence of mesenteric arteritis) was a monitored adverse event in the clinical program; it was observed in three patients receiving cilomilast and in two receiving placebo, a low rate consistent with the normal incidence in the general population. The frequency of GI symptoms that concerned the patients or interfered with daily activities was 3-fold higher in patients receiving cilomilast than placebo.

Roflumilast

Roflumilast (3-cyclopropylmethoxy-4-difluoromethoxy-N-[3,5-dichloropyrid-4-yl]-benzamide) is a potent and selective PDE 4 inhibitor. It is being developed as an oral therapy for COPD and asthma (Reid 2002; Christie 2005; Cowan 2005; Antoniu 2006b; Boswell-Smith and Page 2006). It is an effective anti-inflammatory agent in COPD and asthma. Animal data and clinical trials to date have demonstrated

favorable efficacy and safety, and no documented drug interactions (Hatzelmann and Schudt 2001).

Roflumilast inhibits PDE 4 activity in human neutrophils ($IC_{50} = 0.8$ nM), without affecting PDE 1, 2, 3, or 5, even at 10,000-fold higher concentrations. Roflumilast has good bioavailability following oral administration, a long half-life (~10 h), and an active metabolite (roflumilast-N-oxide, with a half-life of ~20 h). Roflumilast is approximately equipotent with its major in vivo metabolite (roflumilast-N-oxide).

It can be given once a day; it has been studied as an oral tablet at doses of 250 or 500 μ g/day. Roflumilast is thus convenient to administer and has a favorable side effect profile in clinical studies reported to date (Karish and Gagnon 2006).

Roflumilast has a range of anti-inflammatory properties and has potential for treating inflammatory diseases. Roflumilast increases levels of cellular cAMP and inhibits microvascular leakage, trafficking, and the release of cytokines and chemokines from inflammatory cells (Christie et al 2005). Roflumilast apparently mediates some of its anti-inflammatory effects by inducing heme oxygenase-1 expression in macrophages (Kwak et al 2005). The anti-inflammatory and immunomodulatory potential of roflumilast has been assessed in human leukocytes. Regardless of cell type and the response investigated, the IC_{50} values were in a narrow range (2–21 nM), similar to that of roflumilast N-oxide (3–40 nM) (Hatzelmann and Schudt 2001).

Roflumilast has shown encouraging efficacy in patients with COPD, with significant improvements observed in FEV_1 and PEF versus baseline (Cowan 2005). COPD patients receiving roflumilast experienced fewer exacerbations. The most common adverse effects reported in clinical trials were diarrhea, nausea, headache, and abdominal pain (Cowan 2005). In a biopsy study of COPD patients, roflumilast significantly reduced the numbers of CD8⁺ T-cells and caused lesser reductions in the numbers of CD4⁺ T-cells and neutrophils, and no changes in the expression of IL-8 or TNF- α .

A 6-month dose ranging study of roflumilast in COPD patients has been reported. Patients receiving roflumilast exhibited a significant, although modest, improvement in FEV_1 . Patients in the roflumilast group had a 48% reduction in the number of exacerbations, as compared with an 8% reduction in the placebo group.

In a phase III, multicenter, double-blind, randomized, placebo-controlled study undertaken in an outpatient setting, 1411 patients with moderate-to-severe COPD were randomly assigned to receive roflumilast 250 μ g (n = 576), roflumilast

500 µg (n = 555), or placebo (n = 280) given orally once daily for 24-weeks. Primary outcomes were post-bronchodilator FEV₁ and health-related quality of life. Secondary outcomes included exacerbations. 1157 (82%) patients completed the study. Post-bronchodilator FEV₁ at the end of treatment significantly improved with roflumilast 250 µg and 500 µg, compared with placebo. Most adverse events were mild-to-moderate in intensity and resolved during the study. Roflumilast improved lung function and reduced exacerbations compared with placebo (Rabe et al 2005).

Roflumilast has also been assessed in several clinical studies of asthma. In a double-blind, randomized study, roflumilast was compared with inhaled beclomethasone dipropionate. 499 patients (FEV₁ = 50%–85% predicted) received roflumilast 500 µg once daily or beclomethasone dipropionate 200 µg twice daily for 12 weeks. Roflumilast and beclomethasone dipropionate significantly improved FEV₁ and FVC. Once daily roflumilast (500 µg oral) was comparable with inhaled twice-daily beclomethasone dipropionate (400 µg/d) in improving pulmonary function and asthma symptoms and reducing rescue medication use (Bousquet et al 2006). In a dose-ranging study of roflumilast in patients with mild-to-moderate asthma, patients (n = 693) were randomized in a double-blind, parallel-group manner. After a 1–3-week placebo run-in period, patients (mean FEV₁ 73% of predicted) were randomized to receive roflumilast 100, 250, or 500 µg once daily for 12-week. The primary end point was change in FEV₁ from baseline. Secondary end points included change in morning and evening peak expiratory flow from baseline. Roflumilast significantly increased FEV₁ (improvements from baseline at the last visit were 260, 320, and 400 mL for the 100, 250, and 500 µg doses, respectively). Roflumilast was well tolerated at all doses tested; most adverse events were mild to moderate in intensity and transient (Bateman et al 2006). Roflumilast was assessed in a placebo-controlled, randomized, double-blind, crossover study in 16 patients with exercise-induced asthma. Patients received placebo or roflumilast (500 µg/d) for 28 d. Exercise challenge was performed 1 h after dosing on days 1, 14, and 28. FEV₁ was measured before exercise challenge, immediately after the end of exercise, and then 1, 3, 5, 7, 9, and 12 min later. The mean percentage fall in FEV₁ after exercise was reduced by 41%, compared with placebo. The median TNF-α level decreased by 21% during roflumilast treatment, but remained constant with the placebo. Roflumilast was effective in treating exercise-induced asthma and there was a significant reduction of TNF-α levels ex vivo (Timmer et al 2002).

Roflumilast's anti-allergy properties have been assessed in several clinical studies. The efficacy of oral roflumilast (500 µg/day) has been investigated in allergic rhinitis, in a randomized, placebo-controlled, double-blind, crossover study. Twenty five people (16 male, 9 female; median age, 28 years) with histories of allergic rhinitis, but who were asymptomatic at screening, received roflumilast (500 µg once daily) and placebo for 9 d each with a washout period of at least 14 d between treatments. Controlled intranasal allergen provocation with pollen extracts was performed daily beginning on the third day of treatment, ~2 h after drug administration. After allergen provocation, rhinal airflow was measured and subjective symptoms (obstruction, itching, rhinorrhea) were assessed. Rhinal airflow improved during roflumilast treatment and was significantly higher at study day 9 than with placebo. Thus, roflumilast, effectively controlled symptoms of allergic rhinitis (Schmidt et al 2001).

The effects of repeated doses of roflumilast (250 or 500 µg oral) on asthmatic airway responses to allergen were examined in a randomized, double-blind, placebo-controlled, crossover study. Patients (n = 23) with mild asthma (FEV₁ ≥ 70% of predicted value) participated in 3 treatment periods (7–10 d), separated by washout periods (2–5-week). Patients received roflumilast (250 µg or 500 µg oral) or placebo once daily. Allergen challenge was performed at the end of each treatment period, followed by FEV₁ measurements over the next 24 h. Once-daily oral roflumilast attenuated late asthmatic reactions and, to a lesser degree, early asthmatic reactions to allergen in patients with mild allergic asthma (van Schalkwyk et al 2005).

Several studies on roflumilast's pharmacokinetic properties and metabolism have been reported. In an open, randomized, single-dose crossover study, the effects of a high-fat meal on the pharmacokinetics of roflumilast and its N-oxide metabolite were investigated. Twelve healthy subjects received roflumilast (2 × 250 µg orally) after an overnight fast and after breakfast. Blood was sampled up to 54 h for pharmacokinetic profiling of roflumilast and its N-oxide. After the meal, roflumilast C_{max} was modestly reduced and the N-oxide C_{max} was unchanged. Roflumilast t_{max} was delayed in the fed (2 h) versus the fasted state (1 h), while the N-oxide t_{max} was unchanged. No significant food effect was seen on roflumilast or the N-oxide AUC_{0–last} or AUC_{0–8}. Thus, roflumilast can be taken with or without food (Hauns et al 2006).

Roflumilast is partly metabolized by cytochrome P450 (CYP) 3A4, and may inhibit its activity. Therapeutic steady-state concentrations of roflumilast and its active

metabolite roflumilast N-oxide did not alter the metabolism of the CYP3A substrate midazolam in healthy adult male subjects, suggesting that roflumilast is not likely to affect the clearance of drugs that are metabolized by CYP3A4 (Nassr et al 2006).

BAY 19-8004

The effects of a 1-week treatment with BAY 19-8004 (5 mg once per day) have been examined on trough FEV₁ and markers of inflammation in induced sputum in patients with COPD or asthma. Eleven patients with COPD (FEV₁ ~60% predicted, all smokers) and 7 patients with asthma (FEV₁ ~70% predicted, all non-smokers) took part in a randomized, double-blind, placebo-controlled trial. FEV₁ was measured before and after 1 week of treatment; sputum was induced by 4.5% saline inhalation on the last day of treatment. FEV₁ did not improve during either treatment in either patient group. Sputum cell counts were not different following placebo and BAY 19-8004 treatment in COPD and asthma patients. In patients with COPD, small but significant reductions in sputum levels of albumin and eosinophil cationic protein were observed. Thus, a 1-week treatment with BAY 19-8004 did not affect FEV₁ or sputum cell numbers in patients with COPD or asthma. However, such treatment did reduce levels of albumin and eosinophil cationic protein in sputum samples obtained from patients with COPD (Grootendorst, Gauw, Benschop et al 2003).

Other PDE 4 inhibitors for COPD in development

CC3

CC3 is another PDE 4 inhibitor with low affinity for the HARBS. Its airway-relaxing properties were analyzed using rat precision-cut lung slices (PCLS) in which airways were contracted by methacholine or in passively sensitized PCLS exposed to ovalbumin. The anti-inflammatory properties were investigated by measuring the release of TNF from endotoxin-treated human monocytes. CC3 in combination with motapizone, attenuated methacholine-induced bronchoconstriction in a concentration-dependent manner. CC3 has bronchospasmolytic and anti-inflammatory properties (Martin et al 2002).

AWD 12-281

AWD 12-281 (N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide), is a potent (IC₅₀ = 9.7 nM) and selective inhibitor of PDE 4, with a low affinity for the HARBS (Kuss et al 2003; Gutke et al

2005). The compound was optimized for topical treatment of COPD, asthma, and allergic rhinitis. The compound has a low oral bioavailability and a low solubility. It exerts long-lasting pharmacological effects after intratracheal administration, indicating persistence in lung tissue in various animal models. It has high plasma-protein binding and hepatic metabolism (primarily glucuronidation); both contribute to low systemic exposure after intratracheal dosing. The drug has a large difference between emetic and anti-inflammatory dose levels (a factor of more than 100 in ferrets) (Kuss et al 2003).

SCH 351591

SCH 351591 (N-(3,5-dichloro-1-oxido-4-pyridinyl)-8-methoxy-2-(trifluoromethyl)-5-quinoline carboxamide) has been identified as a potent (IC₅₀ = 58 nM) and highly selective PDE 4 inhibitor with oral bioactivity in several animal models of lung inflammation and is being investigated as a potential therapeutic for COPD and asthma.

Ciclamilast

Ciclamilast is a piclamilast (RP 73-401) analog, though is a more potent inhibitor of PDE 4 and airway inflammation and has a more favorable side-effect profile than piclamilast (Deng et al 2006). In a murine model, oral administration of ciclamilast dose-dependently inhibited changes in lung resistance and lung dynamic compliance, up-regulated cAMP-PDE activity, and increased PDE 4D, but not PDE 4B, mRNA expression in lung tissue. Ciclamilast also dose-dependently reduced mRNA expression of eotaxin, TNF- α and IL-4, but increased mRNA expression of IFN- γ in lung tissue. There was a correlation between increases in PDE 4D mRNA expression and airway hyper-responsiveness (Deng et al 2006).

Piclamilast

Piclamilast (RP 73-401) reduced antigen challenge induced-cell recruitment in airways of sensitized mice, and reduced gelatinase B (MMP-9) (Belleguic et al 2000). Piclamilast also reduced MMP-9 activity and TGF- β 1 release during acute lung injury in mice, suggesting that PDE 4 inhibitors might modulate tissue remodeling in lung injury (Corbel, Germain et al 2002). Fibroblasts cultured with PMA or TNF- α released increased amounts of pro-MMP-1, whereas TGF- β 1 had no effect (Martin-Chouly et al 2004). Incubation with CI-1044 or cilomilast significantly prevented the TNF- α increase in pro-MMP-1. These results suggest that PDE 4 inhibitors are effective in inhibiting the pro-MMP-2 and pro-MMP-1 secretion induced by TNF- α and might

indicate the potential therapeutic benefit of selective PDE 4 inhibitors in lung diseases associated with abnormal tissue remodeling (Martin-Chouly et al 2004).

CGH2466

CGH2466 resulted from a study to identify a theophylline-like compound with improved effectiveness. CGH2466 antagonized the adenosine A1, A2b and A3 receptors and inhibited the p38 mitogen-activated protein (MAP) kinases α and β and PDE 4D. CGH2466 inhibited the production of cytokines and oxygen radicals by human peripheral blood leucocytes in vitro. When given orally or locally into the lungs, CGH2466 potently inhibited the ovalbumin- or LPS-induced airway inflammation in mice (Trifileff et al 2005).

The in vitro activity of CI-1044 has been compared with that of rolipram and cilomilast and to the glucocorticoid dexamethasone in reducing LPS-induced TNF- α release in whole blood from COPD patients. In whole blood from COPD patients, pre-incubation with PDE 4 inhibitors or dexamethasone resulted in a dose-dependent inhibition of LPS-induced TNF- α release. There was a similar inhibition using whole blood from healthy volunteers, however, at higher IC₅₀ values. Thus, CI-1044 inhibited in vitro LPS-induced TNF- α release in whole blood from COPD patients (Ouagued et al 2005).

Other treatments for COPD in development

Many other treatments for COPD and asthma are in various stages of development (Donnelly and Rogers 2003; Buhl and Farmer 2004; Buhl and Farmer 2005; Malhotra et al 2006). They include aids to smoking cessation (Richmond and Zwar 2003), antiproteases, including inhibitors of neutrophil elastase (Ohbayashi 2002), matrix metalloproteinase inhibitors (Owen 2005; Gueders et al 2006), cathepsin inhibitors (de Garavilla et al 2005), selectin antagonists (Romano 2005), inhibitors of TNF- α (Spond et al 2003), adenosine A2a receptor agonists (Bonneau et al 2005), ω -3 polyunsaturated fatty acids (Matsuyama et al 2005), inhibitors of mucus hypersecretion (Knight 2004; Rogers and Barnes 2006), purinoceptor P2Y₂ receptor agonists to increase mucus clearance (Kellerman 2002), inhibitors of p38 mitogen-activated protein (MAP) kinase (Adcock et al 2006), inhibitors of NF- κ B kinase-2 (IKK2) (Caramori et al 2004), leukotriene (LT) blockers (Riccioni et al 2004), anti-chemokine therapy (Panina et al 2006; Hardaker et al 2004), anti-cytokine therapy (Chung 2006), statins, angiotensin-converting enzyme inhibitors, and angiotensin receptor

blockers (Mancini et al 2006), antioxidant therapy (Bowler et al 2004; Owen 2005; MacNee 2005b; Rahman and Kilty 2006; Hanta et al 2006; Sadowska et al 2006; Rahman and Adcock 2006), and activators of histone deacetylase (Adcock et al 2005; Antoniu 2006; Kirkham and Rahman 2006; Barnes 2006; Ito et al 2006).

Conclusion

The concept of PDE 4 inhibitors as treatments for COPD, asthma, and other inflammatory airway conditions has been widely discussed in the literature in recent years and may soon come to fruition. Taking one step back, PDE inhibitors have proven to be successful drugs. Today, the first-line oral pharmacotherapy for most patients with erectile dysfunction is a PDE 5 inhibitor: sildenafil (Viagra), tadalafil (Cialis), or vardenafil (Levitra) (Briganti et al 2005; Boswell-Smith, Spina, et al 2006). Furthermore, ibudilast, a drug that does inhibit PDE 4, is marketed in Japan and is used to treat asthma.

There is much preclinical data supporting the use of PDE 4 inhibitors in treating COPD. In vitro, PDE 4 inhibitors relax lung smooth muscle and decrease the production of cytokines from inflammatory cells (Torphy and Undem 1991; Undem et al 1994; Dent and Giembycz 1995; Teixeira et al 1997). PDE 4 inhibitors also reduce TNF- α release (Profita et al 2003). Furthermore, some PDE 4 inhibitors inhibit neutrophil degranulation (Jones et al 2005). These inhibitors also suppress the activity of many pro-inflammatory and immune cells (Lipworth 2005).

To date, only limited clinical data is available to assess PDE 4 inhibitors. Results from large, phase III COPD studies of cilomilast have been reported; cilomilast was well-tolerated, improved health status, and lung function, and reduced the utilization of healthcare resources and incidence of COPD exacerbations. However, the results of these phase III trials were somewhat unremarkable and disappointing (Giembycz 2006).

Questions remain about both the efficacy and safety of cilomilast, the one PDE 4 inhibitor to have undergone a full clinical program to date. Its efficacy has been somewhat limited and, furthermore, somewhat inconsistent results have been reported; indeed, the FDA has yet to approve the drug, apparently because of its limited efficacy. In two of the four pivotal phase III trials, cilomilast did not reach statistical significance over the placebo. In a 6-week phase II study in 424 patients with moderate COPD, significant improvements in lung function were seen in patients receiving cilomilast. Administered at 15 mg bid, it resulted in

significant improvements in FEV₁ compared with placebo (130 mL versus -30 mL at week 6); FVC and peak expiratory flow rate also improved. However, no improvement in quality of life (SGRQ) was found. The observed difference in FEV₁ compared with placebo after 12-weeks was 70 mL (10 mL versus -60 mL at week 12; statistically insignificant). This was compared with 160 mL (130 mL versus -30 mL at week 6) in the larger study, despite patients having similar levels of baseline function (Compton et al 1991). In a smaller (59-patient) phase II study, no significant change in FEV₁ was found (Gamble et al 2003).

The FDA expressed concern about cilomilast's toxicity and side effects. Vasculitis was seen in rats at doses lower than those used in the phase III studies and GI-related side effects were seen in patients receiving cilomilast at three times the frequency seen in those taking placebo.

It remains unclear whether the effects of cilomilast on lung function are the result of bronchodilator activity or of an anti-inflammatory effect; the relatively slow improvement in FEV₁ suggests an anti-inflammatory action, not bronchodilation. The acute bronchodilating effects of a single dose of cilomilast have been assessed in COPD patients. FEV₁ was measured before and at up to 8 h intervals after patients received placebo, cilomilast, or cilomilast and inhaled salbutamol (400 µg) and/or ipratropium bromide (80 µg). A single dose of cilomilast did not cause acute bronchodilation in COPD patients who were responsive to inhaled bronchodilators (Grootendorst, Gauw, Baan et al 2003). Anti-inflammatory properties of cilomilast have been assessed in several studies. In one, patients with COPD received cilomilast (15 mg bid) or placebo for 12-weeks. In bronchial biopsies, cilomilast treatment was associated with reductions in CD8⁺ and CD68⁺ cells; this was the first report of a reduction in airway tissue inflammatory cells characteristic of COPD by any agent (Gamble et al 2003).

While the FDA issued an 'approvable' letter for cilomilast to treat COPD, significant safety and efficacy issues remain. In two of four pivotal phase III studies, the drug failed to reach statistical significance in FEV₁ change, the co-primary endpoint. Indeed, in the two studies that did, this was largely the result of decreases in FEV₁ in the placebo group, not increases in those receiving cilomilast (FEV₁ remained close to baseline, even after 6-months of treatment, in all four studies). The FEV₁ changes seen were small and there is a question as to whether even statistically significant results would be of much clinical significance. For comparison, the changes were smaller than those reported in a meta-analysis of theophylline studies (Ram et al 2002). Three of the four

phase III studies failed to reach statistical significance in SGRQ, the other primary endpoint. However, cilomilast did reduce the incidence of exacerbations.

In rat studies, cilomilast was associated with vasculitis and death at doses lower than the human dose, although there is some reason to believe that rats may be more sensitive to PDE inhibitor toxicity (Bian et al 2004). Vasculitis has been seen with other PDE inhibitors (Larson et al 1996; Slim et al 2003). The FDA was apparently not satisfied with the investigation by GSK of patients with GI-related side effects.

Cilomilast seems unlikely to be a replacement for existing COPD therapies (supplemental oxygen, inhaled bronchodilators, corticosteroids and antibiotics (Sin et al 2003)). Cilomilast, however, may be a useful additional drug (for example, in combination with corticosteroids), especially if it can be shown in longer-term studies that the increases in FEV₁ are more substantial. Cilomilast has other anti-inflammatory properties, which may also be of clinical significance (Gamble et al 2003).

The concept of using PDE 4 inhibitors to treat COPD may well be sound, but the first drug in the class may be roflumilast (Antoniou 2006b), not cilomilast. The completion and publication of its clinical development is awaited with interest.

References

- Adams SG, Melo J, Luther M, et al. 2001. Antibiotics are associated with lower relapse rates in outpatients with acute exacerbations of COPD. *Chest*, 117:1345-52.
- Adcock IM, Chung KF, Caramori G, et al. 2006. Kinase inhibitors and airway inflammation. *Eur J Pharmacol*, 533:118-32.
- Adcock IM, Ito K, Barnes PJ. 2005. Histone deacetylation: an important mechanism in inflammatory lung diseases. *COPD*, 2:445-55.
- Alves AC, Pires ALA, Cruz HN, et al. 1996. Selective inhibition of phosphodiesterase type IV suppresses the chemotactic responsiveness of rat eosinophils in vitro. *Eur J Pharmacol*, 312:89-96.
- American Thoracic Society. 1995. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. *Am J Resp Crit Care Med*, 152:S77-120.
- Anthonisen NR, Connett JE, Murray RP. 2002. Smoking and lung function of Lung Health Study participants after 11 years. *Am J Respir Crit Care Med*, 166:675-9.
- Anthonisen NR, Manfreda J, Warren CP, et al. 1987. Antibiotic therapy in exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med*, 106:196-204.
- Antoniou SA. 2006a. Histone deacetylase pathway: an evolving therapeutic target in chronic obstructive pulmonary disease. *Expert Opin Ther Targets*, 10:329-32.
- Antoniou SA. 2006b. Roflumilast for the treatment of chronic obstructive pulmonary disease. *Curr Opin Investig Drugs*, 7:412-17.
- Appleton S, Jones T, Poole P, et al. 2006. Ipratropium bromide versus long-acting beta-2 agonists for stable chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, 19:CD006101.
- Ashton MJ, Cook DC, Fenton G, et al. 1994. Selective type IV phosphodiesterase inhibitors as antiasthmatic agents. The syntheses and biological activities of 3-(cyclopentyloxy)-4-methoxybenzamides and analogues. *J Med Chem*, 37:1696-703.

- Austin M, Wood-Baker R. 2006. Oxygen therapy in the pre-hospital setting for acute exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, 19:CD005534.
- Banner KH, Trevelthick MA. 2004. PDE4 inhibition: a novel approach for the treatment of inflammatory bowel disease. *Trends Pharmacol Sci*, 25:430–6.
- Barnes CJ, Chung KF, Page CP. 1988. Inflammatory mediators and asthma. *Pharmacol Rev*, 40:49–84.
- Barnes PJ. 1995. Molecular mechanisms of antiasthma therapy. *Ann Med*, 27:531–5.
- Barnes PJ. 1998. New therapies for chronic obstructive pulmonary disease. *Thorax*, 53:137–47.
- Barnes PJ. 2000. Chronic obstructive pulmonary disease. *N Engl J Med*, 343:269–80.
- Barnes PJ. 2003. Theophylline: new perspectives for an old drug. *Am J Respir Crit Care Med*, 67:813–18.
- Barnes PJ. 2006a. Novel signal transduction modulators for the treatment of airway diseases. *Pharmacol Ther*, 109:238–45.
- Barnes PJ. 2006b. Reduced histone deacetylase in COPD: clinical implications. *Chest*, 129:151–5.
- Barnette MS, Bartus JO, Burman M, et al. 1996. Association of the anti-inflammatory activity of phosphodiesterase 4 (PDE4) inhibitors with either inhibition of PDE4 catalytic activity or competition for [3H]rolipram binding. *Biochem Pharmacol*, 51:949–56.
- Barnette MS, Christensen SB, Essayan DM, et al. 1998. SB 207499 (Ariflo), a potent and selective second-generation phosphodiesterase 4 inhibitor: in vitro anti-inflammatory actions. *J Pharmacol Exp Ther*, 284:420–6.
- Barnette MS, Christensen SB, Underwood DC. 1996. Phosphodiesterase 4: Biological underpinnings for the design improved inhibitors. *Pharmacol Rev Commun*, 8:65–73.
- Barnette MS, Grous M, Cieslinski LB, et al. 1995. Inhibitors of phosphodiesterase IV (PDE IV) increase acid secretion in rabbit isolated gastric glands: correlation between function and interaction with a high-affinity rolipram binding site. *J Pharmacol Exp Ther*, 273:1396–402.
- Barnette MS, Underwood DC. 2000. New phosphodiesterase inhibitors as therapeutics for the treatment of chronic lung disease. *Curr Opin Pulmon Med*, 6:164–9.
- Barnette MS. 1999. Phosphodiesterase 4 (PDE4) inhibitors in asthma and chronic obstructive pulmonary disease (COPD). *Prog Drug Res*, 53:193–229.
- Barr RG, Rowe BH, Camargo CA Jr. 2003. Methylxanthines for exacerbations of chronic obstructive pulmonary disease: meta-analysis of randomised trials. *Brit Med J*, 327:643–8.
- Bateman ED, Izquierdo JL, Harnest U, et al. 2006. Efficacy and safety of roflumilast in the treatment of asthma. *Ann Allergy Asthma Immunol*, 96:679–86.
- Beavo JA, Conti M, Heaslip RJ. 1994. Multiple cyclic nucleotide phosphodiesterases. *Mol Pharmacol*, 46:399–405.
- Belleguic C, Corbel M, Germain N, et al. 2000. Reduction of matrix metalloproteinase-9 activity by the selective phosphodiesterase 4 inhibitor, RP 73–401 in sensitized mice. *Eur J Pharmacol*, 404:369–73.
- Bender AT, Beavo JA. 2006. Cyclic nucleotide phosphodiesterases: molecular regulation to clinical use. *Pharmacol Rev*, 58:488–520.
- Bian H, Zhang J, Wu P, et al. 2004. Differential type 4 cAMP-specific phosphodiesterase (PDE4) expression and functional sensitivity to PDE4 inhibitors among rats, monkeys and humans. *Biochem Pharmacol*, 68:2229–36.
- Blanchard AR. 2002. Treatment of COPD exacerbations: pharmacologic options and modification of risk factors. *Postgrad Med*, 111:65–75.
- Bolger GB, Erdogan S, Jones RE, et al. 1997. Characterization of five different proteins produced by alternatively spliced mRNAs from the human cAMP-specific phosphodiesterase PDE4D gene. *Biochem J*, 328:539–48.
- Bolger GB. 1994. Molecular biology of the cyclic AMP-specific cyclic nucleotide phosphodiesterases: a diverse family of regulatory enzymes. *Cell Signalling*, 6:851–9.
- Bonneau O, Wyss D, Ferretti S, et al. 2006. Effect of adenosine A2A receptor activation in murine models of respiratory disorders. *Am J Physiol Lung Cell Mol Physiol*, 290:L1036–43.
- Borron W, deBoisblanc BP. 1998. Steroid therapy for chronic obstructive pulmonary disease. *Curr Opin Pulmon Med*, 4:61–5.
- Boswell-Smith V, Cazzola M, Page CP. 2006. Are phosphodiesterase 4 inhibitors just more theophylline? *J Allergy Clin Immunol*, 117:1237–43.
- Boswell-Smith V, Page CP. 2006. Roflumilast: a phosphodiesterase-4 inhibitor for the treatment of respiratory disease. *Expert Opin Investig Drugs*, 15:1105–13.
- Boswell-Smith V, Spina D, Page CP. 2006. Phosphodiesterase inhibitors. *Br J Pharmacol*, 147 (Suppl 1):S252–7.
- Bousquet J, Aubier M, Sastre J, et al. 2006. Comparison of roflumilast, an oral anti-inflammatory, with beclomethasone dipropionate in the treatment of persistent asthma. *Allergy*, 61:72–8.
- Bowler RP, Barnes PJ, Crapo JD. 2004. The role of oxidative stress in chronic obstructive pulmonary disease. *COPD*, 1:255–77.
- Briganti A, Salonia A, Gallina A, et al. 2005. Oral phosphodiesterase type 5 inhibitors for erectile dysfunction. *Nat Clin Pract Urol*, 2:239–47.
- Bruijnzeel PL. 1989. Contribution of eosinophil derived mediators in asthma. *Intl Arch Allergy Appl Immunol*, 90 (Suppl 1): 57–63.
- Bryson SE, Rodger IW. 1987. Effects of phosphodiesterase inhibitors on normal and chemically-skinned isolated airway smooth muscle. *Br J Pharmacol*, 92:673–81.
- Buhl R, Farmer SG. 2004. Current and future pharmacologic therapy of exacerbations in chronic obstructive pulmonary disease and asthma. *Proc Am Thorac Soc*, 1:136–42.
- Buhl R, Farmer SG. 2005. Future directions in the pharmacologic therapy of chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 2:83–93.
- Burgess JK, Oliver BG, Poniris MH, et al. 2006. A phosphodiesterase 4 inhibitor inhibits matrix protein deposition in airways in vitro. *J Allergy Clin Immunol*, 118:649–57.
- Burnouf C, Auclair E, Avenel N, et al. 2000. Synthesis, structure-activity relationships, and pharmacological profile of 9-amino-4-oxo-1-phenyl-3,4,6,7-tetrahydro[1,4]diazepino[6,7,1-h]indoles: discovery of potent, selective phosphodiesterase type 4 inhibitors. *J Med Chem*, 43:4850–67.
- Bushnik T, Conti M. 1996. Role of multiple cAMP-specific phosphodiesterase variants. *Biochem Soc Trans*, 24:1014–19.
- Caramori G, Adcock IM, Ito K. 2004. Anti-inflammatory inhibitors of IkappaB kinase in asthma and COPD. *Curr Opin Investig Drugs*, 5:1141–7.
- Card GL, England BP, Suzuki Y, et al. 2004. Structural basis for the activity of drugs that inhibit phosphodiesterases. *Structure*, 12:2233–47.
- Cavalla D, Frith R. 1995. Phosphodiesterase IV inhibitors: structural diversity and therapeutic potential in asthma. *Curr Med Chem*, 2:561–72.
- Chambers RJ, Marfat A, Cheng JB, et al. 1997. Biarylcarboxamide inhibitors of phosphodiesterase IV and tumor necrosis factor-alpha. *Bioorg Med Chem Lett*, 7:739–44.
- Christensen SB, Guider A, Forster CJ, et al. 1998. 1,4-cyclohexanecarboxylates: potent and selective inhibitors of phosphodiesterase 4 for the treatment of asthma. *J Med Chem*, 41:821–35.
- Christie P. 2005. Roflumilast: a selective phosphodiesterase 4 inhibitor. *Drugs Today*, 41:667–75.
- Chung KF. 2006. Cytokines as targets in chronic obstructive pulmonary disease. *Curr Drug Targets*, 7:675–81.
- Compton CH, Gubb J, Nieman R, et al. 2001. Cilomilast, a selective phosphodiesterase-4 inhibitor for treatment of patients with chronic obstructive pulmonary disease: a randomised, dose-ranging study. *Lancet*, 358:265–70.
- Connors AF Jr, Dawson NV, Thomas C, et al. 1996. Outcomes following acute exacerbation of severe chronic obstructive lung disease. The SUPPORT investigators (Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments). *Am J Respir Crit Care Med*, 154:959–67.

- Conti M, Jin SL. 1999. The molecular biology of cyclic nucleotide phosphodiesterases. *Prog Nucleic Acid Res Mol Biol*, 63:1–38.
- Corbel M, Belleguic C, Boichot E, et al. 2002. Involvement of gelatinases (MMP-2 and MMP-9) in the development of airway inflammation and pulmonary fibrosis. *Cell Biol Tox*, 18:51–61.
- Corbel M, Germain N, Lanchou J, et al. 2002. The selective phosphodiesterase 4 inhibitor RP 73-401 reduced matrix metalloproteinase 9 activity and transforming growth factor-beta release during acute lung injury in mice: the role of the balance between tumor necrosis factor-alpha and interleukin-10. *J Pharmacol Exp Ther*, 301:258–65.
- Cortijo J, Bou J, Beleta J, et al. 1993. Investigation into the role of phosphodiesterase IV in bronchorelaxation, including studies with human bronchus. *Br J Pharmacol*, 108:562–8.
- Costello J. 1998. Cardiac effects of beta-agonists in patients with COPD (chronic obstructive pulmonary disease). *Chest*, 114:353–4.
- Cowan C. 2005. Roflumilast for asthma and chronic obstructive pulmonary disease. *Issues Emerg Health Technol*, 74:1–4.
- Crocker IC, Townley RG. 1999. Therapeutic potential of phosphodiesterase 4 inhibitors in allergic diseases. *Drugs Today*, 35:519–35.
- Crouch E. 1990. Pathobiology of pulmonary fibrosis. *Am J Lung Cell Mol Physiol*, 259:L159–84.
- D'Alonzo GE. 1996. Theophylline revisited. *Allergy Asthma Proc*, 17:335–9.
- Daniel PB, Walker WH, Habener JF. 1998. Cyclic AMP signaling and gene regulation. *Ann Rev Nutr*, 18:353–83.
- De Brito FB, Souness JE, Warne PJ. 1997. Type 4 phosphodiesterase inhibitors and their potential in the treatment of inflammatory disease. *Emerging Drugs*, 2:249–68.
- de Garavilla L, Greco MN, Sukumar N, et al. 2005. A novel, potent dual inhibitor of the leukocyte proteases cathepsin G and chymase: molecular mechanisms and anti-inflammatory activity in vivo. *J Biol Chem*, 280:18001–7.
- DeKorte CJ. 2003. Current and emerging therapies for the management of chronic inflammation in asthma. *Am J Health Syst Pharm*, 60:1949–59.
- Deng YM, Xie QM, Tang HF, et al. 2006. Effects of cilomilast, a new PDE 4 PDE4 inhibitor, on airway hyperresponsiveness, PDE4D expression and airway inflammation in a murine model of asthma. *Eur J Pharmacol*, 547:125–35.
- Dent G, Giembycz MA. 1995. Selective phosphodiesterase inhibitors in the therapy of asthma. *Clin Immunother*, 3:423–37.
- Dent G, Giembycz MA. 1996. Phosphodiesterase inhibitors: Lily the Pink's medicinal compound for asthma? *Thorax*, 51:647–9.
- Dewan NA, Rafique S, Kanwar B, et al. 2000. Acute exacerbation of COPD: factors associated with poor treatment outcome. *Chest*, 117:662–71.
- Doherty DE, Briggs DD Jr. 2004. Long-term pharmacologic management of patients with chronic obstructive pulmonary disease. *Clin Cornerstone*, Suppl 2:S29–34.
- Domej W, Foldes-Papp Z, Fogel E, et al. 2006. Chronic obstructive pulmonary disease and oxidative stress. *Curr Pharm Biotechnol*, 7:117–23.
- Donnelly LE, Rogers DF. 2003. Therapy for chronic obstructive pulmonary disease in the 21st century. *Drugs*, 63:1973–98.
- D'Souza AO, Smith MJ, Miller LA, et al. 2006. An appraisal of pharmacoeconomic evidence of maintenance therapy for COPD. *Chest*, 129:1693–708.
- Dyke HJ, Montana JG. 2002. Update on the therapeutic potential of PDE4 inhibitors. *Expert Opin Investig Drug*, 11:1–13.
- Erdogan S, Houslay MD. 1997. Challenge of human Jurkat T-cells with the adenylate cyclase activator forskolin elicits major changes in cAMP phosphodiesterase (PDE) expression by up-regulating PDE3 and inducing PDE4D1 and PDE4D2 splice variants as well as down-regulating a novel PDE4A splice variant. *Biochem J*, 321:165–75.
- Ezeamuzie CI. 2001. Requirement of additional adenylate cyclase activation for the inhibition of human eosinophil degranulation by phosphodiesterase IV inhibitors. *Eur J Pharmacol*, 417:11–18.
- Fabbri LM, Hurd SS. 2003. Global strategy for the diagnosis, management and prevention of COPD: 2003 update. *Eur Respir J*, 22:1–2.
- Fan Chung K. 2006. Phosphodiesterase inhibitors in airways disease. *Eur J Pharmacol*, 533:110–17.
- Gamble E, Grootendorst DC, Brightling CE, et al. 2003. Antiinflammatory effects of the phosphodiesterase-4 inhibitor cilomilast (Ariflo) in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 168:976–82.
- Gartlehner G, Hansen RA, Carson SS, et al. 2006. Efficacy and safety of inhaled corticosteroids in patients with COPD: a systematic review and meta-analysis of health outcomes. *Ann Fam Med*, 4:253–62.
- Georgitis JW. 1999. The 1997 asthma management guidelines and therapeutic issues relating to the treatment of asthma. *Chest*, 115:210–17.
- Ghavam A, Hirst WD, Novak TJ. 2006. Selective phosphodiesterase (PDE)-4 inhibitors: a novel approach to treating memory deficit? *Drugs R D*, 7:63–71.
- Gibson LC, Hastings SF, McPhee I, et al. 2006. The inhibitory profile of Ibudilast against the human phosphodiesterase enzyme family. *Eur J Pharmacol*, 538:39–42.
- Giembycz MA, Dent G. 1992. Prospects for selective cyclic nucleotide phosphodiesterase inhibitors in the treatment of bronchial asthma. *Clin Exp Allergy*, 22:337–44.
- Giembycz MA. 2000. Phosphodiesterase 4 inhibitors and the treatment of asthma: where are we now and where do we go from here? *Drugs*, 59:193–212.
- Giembycz MA. 2001. Cilomilast: a second generation phosphodiesterase 4 inhibitor for asthma and chronic obstructive pulmonary disease. *Expert Opin Invest Drugs*, 10:1361–79.
- Giembycz MA. 2002. Development status of second generation PDE4 inhibitors for asthma and COPD: the story so far. *Monaldi Arch Chest Dis*, 57:48–64.
- Giembycz MA. 2005a. Phosphodiesterase-4: selective and dual-specificity inhibitors for the therapy of chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 2:326–33.
- Giembycz MA. 2005b. Life after PDE4: overcoming adverse events with dual-specificity phosphodiesterase inhibitors. *Curr Opin Pharmacol*, 5:238–44.
- Giembycz MA. 2006. An update and appraisal of the cilomilast Phase III clinical development programme for chronic obstructive pulmonary disease. *Br J Clin Pharmacol*, 62:138–52.
- [GOLD] Global Initiative for Chronic Obstructive Lung Disease. 2003. Executive summary (updated 2003). Accessed May 28, 2007. URL: <http://www.goldcopd.com>
- Gonçalves de Moraes VL, Singer M, Vargaftig BB, et al. 1998. Effects of rolipram on cyclic AMP levels in alveolar macrophages and lipopolysaccharide-induced inflammation in mouse lung. *Br J Pharmacol*, 123:631–6.
- Griswold DE, Webb EF, Badger AM, et al. 1998. SB 207499 (Ariflo), a second generation phosphodiesterase 4 inhibitor, reduced tumor necrosis factor alpha and interleukin-4 production in vivo. *J Pharmacol Exp Ther*, 287:705–11.
- Grootendorst DC, Gauw SA, Baan R, et al. 2003. Does a single dose of the phosphodiesterase 4 inhibitor, cilomilast (15 mg), induce bronchodilation in patients with chronic obstructive pulmonary disease. *Pulmon Pharmacol Ther*, 16:115–20.
- Grootendorst DC, Gauw SA, Benschop N, et al. 2003. Efficacy of the novel phosphodiesterase-4 inhibitor BAY 19-8004 on lung function and airway inflammation in asthma and chronic obstructive pulmonary disease (COPD). *Pulm Pharmacol Ther*, 16:341–7.
- Growcott EJ, Spink KG, Ren X, et al. 2006. Phosphodiesterase type 4 expression and anti-proliferative effects in human pulmonary artery smooth muscle cells. *Respir Res*, 7:9.
- Gueders MM, Foidart JM, Noel A, et al. 2006. Matrix metalloproteinases (MMPs) and tissue inhibitors of MMPs in the respiratory tract: potential implications in asthma and other lung diseases. *Eur J Pharmacol*, 533:133–44.
- Guerra S. 2005. Overlap of asthma and chronic obstructive pulmonary disease. *Curr Opin Pulmon Med*, 11:7–13.

- Gutke HJ, Guse JH, Khobzaoui M, et al. 2005. AWD-12-281 (inhaled) (elbion/ GlaxoSmithKline). *Curr Opin Investig Drugs*, 6:1149–58.
- Hanta I, Kocabas A, Canacankatan N, et al. 2006. Oxidant-antioxidant balance in patients with COPD. *Lung*, 184:51–5.
- Hardaker EL, Bacon AM, Carlson K, et al. 2004. Regulation of TNF- α - and IFN- γ -induced CXCL10 expression: participation of the airway smooth muscle in the pulmonary inflammatory response in chronic obstructive pulmonary disease. *FASEB J*, 18:191–3.
- Hatzelmann A, Schudt C. 2001. Anti-inflammatory and immunomodulatory potential of the novel PDE4 inhibitor roflumilast in vitro. *J Pharmacol Exp Ther*, 297:267–79.
- Hauns B, Hermann R, Hunnemeyer A, et al. 2006. Investigation of a potential food effect on the pharmacokinetics of roflumilast, an oral, once-daily phosphodiesterase 4 inhibitor, in healthy subjects. *J Clin Pharmacol*, 46:1146–53.
- Hay DWP. 2000. Chronic obstructive pulmonary disease: emerging therapies. *Curr Opin Chem Biol*, 4:412–19.
- Holgate ST, Twentyman OP, Rafferty P, et al. 1987. Primary and secondary effector cells in the pathogenesis of bronchial asthma. *Intl Arch Allergy Appl Immunol*, 82:498–506.
- Horton YM, Sullivan M, Houslay MD. 1995. Molecular cloning of a novel splice variant of human type IVa (PDE-IVa) cyclic AMP phosphodiesterase and localization of the gene to the p13.1-q12 region of human chromosome 19. *Biochem J*, 308:683–91.
- Houslay MD, Adams DR. 2003. PDE4 cAMP phosphodiesterases: modular enzymes that orchestrate signalling cross-talk, desensitization and compartmentalization. *Biochem J*, 370:1–18.
- Houslay MD, Sullivan M, Bolger GB. 1998. The multienzyme PDE4 cyclic adenosine monophosphate-specific phosphodiesterase family: intracellular targeting, regulation, and selective inhibition by compounds exerting anti-inflammatory and antidepressant actions. *Adv Pharmacol*, 44:225–342.
- Huang Z, Ducharme Y, Macdonald D, et al. 2001. The next generation of PDE4 inhibitors. *Curr Opin Chem Biol*, 5:432–8.
- Huang Z, Liu S, Zhang L, et al. 2006. Preferential inhibition of human phosphodiesterase 4 by ibudilast. *Life Sci*, 78:2663–8.
- Huang Z, Mancini JA. 2006. Phosphodiesterase 4 inhibitors for the treatment of asthma and COPD. *Curr Med Chem*, 13:3253–562.
- Huston E, Houslay TM, Baillie GS, et al. 2006. cAMP phosphodiesterase-4A1 (PDE4A1) has provided the paradigm for the intracellular targeting of phosphodiesterases, a process that underpins compartmentalized cAMP signalling. *Biochem Soc Trans*, 34:504–9.
- Ito K, Yamamura S, Essilfie-Quaye S, et al. 2006. Histone deacetylase 2-mediated deacetylation of the glucocorticoid receptor enables NF- κ B suppression. *J Exp Med*, 203:7–13.
- Jacob C, Lepout M, Szilagy C, et al. 2002. DMSO-treated HL60 cells: a model of neutrophil-like cells mainly expressing PDE4B subtype. *Intl Immunopharmacol*, 2:1647–56.
- Jacobitz S, McLaughlin MM, Livi GP, et al. 1996. Mapping the functional domains of human recombinant phosphodiesterase 4A: structural requirements for catalytic activity and rolipram binding. *Mol Pharmacol*, 50:891–9.
- Jeffery P. 2005. Phosphodiesterase 4-selective inhibition: novel therapy for the inflammation of COPD. *Pulmon Pharmacol Therapeut*, 18:9–17.
- Jeffery PK. 2001. Remodeling in asthma and chronic obstructive lung disease. *Am J Respir Care Med*, 164:S28–38.
- Jin SL, Bushnik T, Lan L, et al. 1998. Subcellular localization of rolipram-sensitive, cAMP-specific phosphodiesterases. Differential targeting and activation of the splicing variants derived from the PDE4D gene. *J Biol Chem*, 273:19672–8.
- Jones NA, Boswell-Smith V, Lever R, et al. 2005. The effect of selective phosphodiesterase isoenzyme inhibition on neutrophil function in vitro. *Pulm Pharmacol Ther*, 18:93–101.
- Karish SB, Gagnon JM. 2006. The potential role of roflumilast: the new phosphodiesterase-4 inhibitor. *Ann Pharmacother*, 40:1096–104.
- Karlsson JA, Aldous D. 1997. Phosphodiesterase 4 inhibitors for the treatment of asthma. *Curr Opin Ther Patents*, 7:989–1003.
- Ke H. 2004. Implications of PDE4 structure on inhibitor selectivity across PDE families. *Int J Impot Res*, 16 Suppl 1:S24–7.
- Kellerman DJ. 2002. P2Y(2) receptor agonists: a new class of medication targeted at improved mucociliary clearance. *Chest*, 121(5 Suppl):201S–205S.
- Kelly JJ, Barnes PJ, Giembycz MA. 1996. Phosphodiesterase 4 in macrophages: relationship between cAMP accumulation, suppression of cAMP hydrolysis and inhibition of [3H]R-(-)-rolipram binding by selective inhibitors. *Biochem J*, 318:425–36.
- Kerstjens HA, Timens W. 2003. Phosphodiesterase 4 inhibitors: antiinflammatory therapy for chronic obstructive pulmonary disease at last? *Am J Respir Crit Care Med*, 168:914–15.
- Kirkham P, Rahman I. 2006. Oxidative stress in asthma and COPD: antioxidants as a therapeutic strategy. *Pharmacol Ther*, 111:476–94.
- Knight D. 2004. Talniflumate (Genaera). *Curr Opin Investig Drugs*, 5:557–62.
- Kohyama T, Liu X, Zhu YK, et al. 2002. Phosphodiesterase 4 inhibitor cilomilast inhibits fibroblast-mediated collagen gel degradation induced by tumor necrosis factor- α and neutrophil elastase. *Am J Resp Cell Mol Biol*, 27:487–94.
- Kotlikoff MI, Kamm KE. 1996. Molecular mechanisms of beta-adrenergic relaxation of airway smooth muscle. *Ann Rev Physiol*, 58:115–41.
- Kuss H, Hoefgen N, Johanssen S, et al. 2003. In vivo efficacy in airway disease models of N-(3,5-dichloropyrid-4-yl)-[1-(4-fluorobenzyl)-5-hydroxy-indole-3-yl]-glyoxylic acid amide (AWD 12-281), a selective phosphodiesterase 4 inhibitor for inhaled administration. *J Pharmacol Exp Ther*, 307:373–85.
- Kwak HJ, Song JS, No ZS, et al. 2005. The inhibitory effects of roflumilast on lipopolysaccharide-induced nitric oxide production in RAW264.7 cells are mediated by heme oxygenase-1 and its product carbon monoxide. *Inflamm Res*, 54:508–13.
- Lagente V, Martin-Chouly C, Boichot E, et al. 2005. Selective PDE4 inhibitors as potent anti-inflammatory drugs for the treatment of airway diseases. *Mem Inst Oswaldo Cruz*, 100 (Suppl 1):131–6.
- Lagente V, Moodley I, Perrin S, et al. 1994. Effects of isozyme-selective phosphodiesterase inhibitors on eosinophil infiltration in the guinea-pig lung. *Eur J Pharmacol*, 255:253–6.
- Lagente V, Pruniaux MP, Junien JL, et al. 1995. Modulation of cytokine-induced eosinophil infiltration by phosphodiesterase inhibitors. *Am J Respir Crit Care Med*, 151:1720–4.
- Larson JL, Pino MV, Geiger LE, et al. 1996. The toxicity of repeated exposures to rolipram, a type IV phosphodiesterase inhibitor, in rats. *Pharmacol Toxicol*, 78:44–9.
- Lerner A, Epstein PM. 2006. Cyclic nucleotide phosphodiesterases as targets for treatment of haematological malignancies. *Biochem J*, 393:21–41.
- Lipworth BJ. 2005. Phosphodiesterase-4 inhibitors for asthma and chronic obstructive pulmonary disease. *Lancet*, 365:167–75.
- Liu S, Veilleux A, Zhang L, et al. 2005. Dynamic activation of cystic fibrosis transmembrane conductance regulator by type 3 and type 4D phosphodiesterase inhibitors. *J Pharmacol Exp Ther*, 314:846–54.
- Lobban M, Shakur Y, Beattie J, et al. 1994. Identification of two splice variant forms of type IVB cyclic AMP phosphodiesterase, DPD (rPDE-IVB1) and PDE-4 (rPDE-IVB2) in brains; selective localization in membrane and cytosolic compartments and differential expression in various brain regions. *Biochem J*, 304:399–406.
- Lombardo LJ. 1995. Phosphodiesterase-IV inhibitors: novel therapeutics for the treatment of inflammatory diseases. *Curr Pharmaceut. Design*, 1:255–68.
- Lowe JA, III, Cheng JB. 1992. The PDE IV family of calcium-independent phosphodiesterase enzymes. *Drugs Fut*, 17:799–807.
- MacNee W. 2005a. Pathogenesis of chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 2:258–66.
- MacNee W. 2005b. Pulmonary and systemic oxidant/antioxidant imbalance in chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 2:50–60.
- Madison JM, Irwin RS. 1998. Chronic obstructive pulmonary disease. *Lancet*, 352:467–73.

- Mahler DA, Wire P, Horstman D, et al. 2002. Effectiveness of fluticasone propionate and salmeterol combination delivered via the Diskus device in the treatment of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 166:1084–91.
- Malhotra S, Man SF, Sin DD. 2006. Emerging drugs for the treatment of chronic obstructive pulmonary disease. *Expert Opin Emerg Drugs*, 11:275–91.
- Mancini GB, Etminan M, Zhang B, et al. 2006. Reduction of morbidity and mortality by statins, angiotensin-converting enzyme inhibitors, and angiotensin receptor blockers in patients with chronic obstructive pulmonary disease. *J Am Coll Cardiol*, 47:2554–60.
- Manganiello VC, Murata T, Taira M, et al. 1995. Diversity in cyclic nucleotide phosphodiesterase isoenzyme families. *Arch Biochem Biophys*, 322:1–13.
- Manning CD, Burman M, Christensen SB, et al. 1999. Suppression of human inflammatory cell function by subtype-selective PDE4 inhibitors correlates with inhibition of PDE4A and PDE4B. *Br J Pharmacol*, 128:1393–8.
- Martin C, Goggel R, Dal Piaz V, et al. 2002. Airway relaxant and anti-inflammatory properties of a PDE4 inhibitor with low affinity for the high-affinity rolipram binding site. *Naunyn Schmiedeberg's Arch Pharmacol*, 365:284–9.
- Martin N, Reid PT. 2006. The potential role of phosphodiesterase inhibitors in the management of asthma. *Treat Respir Med*, 5:207–17.
- Martina SD, Ismail MS, Vesta KS. 2006. Cilomilast: orally active selective phosphodiesterase-4 inhibitor for treatment of chronic obstructive pulmonary disease. *Ann Pharmacother*, 40:1822–8.
- Martin-Chouly CA, Astier A, Jacob C, et al. 2004. Modulation of matrix metalloproteinase production from human lung fibroblasts by type 4 phosphodiesterase inhibitors. *Life Sci*, 75:823–40.
- Mata M, Sarria B, Buenestado A, et al. 2005. Phosphodiesterase 4 inhibition decreases MUC5AC expression induced by epidermal growth factor in human airway epithelial cells. *Thorax*, 60:144–52.
- Matsumoto T, Kobayashi T, Kamata K. 2003. Phosphodiesterases in the vascular system. *J Smooth Mus Res*, 39:67–86.
- Matsuyama W, Mitsuyama H, Watanabe M, et al. 2005. Effects of omega-3 polyunsaturated fatty acids on inflammatory markers in COPD. *Chest*, 128:3817–27.
- McEvoy CE, Niewoehner DE. 1998. Adverse effects of corticosteroid therapy for COPD: a critical review. *Chest*, 111:732–43.
- Mehats C, Jin SL, Wahlstrom J. 2003. PDE4D plays a critical role in the control of airway smooth muscle contraction. *FASEB J*, 17:1831–41.
- Mercer PF, Shute JK, Bhowmik A, et al. 2005. MMP-9, TIMP-1 and inflammatory cells in sputum from COPD patients during exacerbation. *Respir Res*, 22:151.
- Miller-Larsson A, Selroos O. 2006. Advances in asthma and COPD treatment: combination therapy with inhaled corticosteroids and long-acting beta 2-agonists. *Curr Pharm Des*, 12:3261–79.
- Miravittles M, Alvarez-Sala JL, Lamarca R, et al. 2002. Treatment and quality of life in patients with chronic obstructive pulmonary disease. *Qual Life Res*, 11:329–38.
- Mohan A, Premanand R, Reddy LN, et al. 2006. Clinical presentation and predictors of outcome in patients with severe acute exacerbation of chronic obstructive pulmonary disease requiring admission to intensive care unit. *BMC Pulm Med*, 6:27.
- Montminy M. 1997. Transcriptional regulation by cyclic AMP. *Ann Rev Biochem*, 66:807–22.
- Muller T, Engels P, Fozard JR. 1996. Subtypes of the type 4 cAMP phosphodiesterases: structure, regulation and selective inhibition. *Trends Pharmacol Sci*, 17:294–8.
- Murphy TF, Sethi S, Niederman MS. 2000. The role of bacteria in exacerbations of COPD: a constructive view. *Chest*, 118:204–9.
- Murphy TF. 2006. The role of bacteria in airway inflammation in exacerbations of chronic obstructive pulmonary disease. *Curr Opin Infect Dis*, 19:225–30.
- Murray CJ, Lopez AD. 1997. Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *Lancet*, 349:1498–504.
- Nassar N, Lahu G, von Richter O, et al. 2006. Lack of a pharmacokinetic interaction between steady-state roflumilast and single-dose midazolam in healthy subjects. *Br J Clin Pharmacol*, Sep 29.
- National Center for Health Statistics. 2002. NCHS data on chronic obstructive pulmonary disorder [online]. Accessed May 28, 2007. URL: <http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5106a1.htm>
- Nicholson CD, Challiss RA, Shahid M. 1991. Differential modulation of tissue function and therapeutic potential of selective inhibitors of cyclic nucleotide phosphodiesterase isoenzymes. *Trends Pharmacol Sci*, 12:19–27.
- Nicholson CD, Shahid M, Bruin J, et al. 1995. Characterization of ORG 20241, a combined phosphodiesterase IV/III cyclic nucleotide phosphodiesterase inhibitor for asthma. *J Pharmacol Exp Ther*, 274:678–87.
- Nicholson CD, Shahid M. 1994. Inhibitors of cyclic nucleotide phosphodiesterase isoenzymes: their potential utility in the therapy of asthma. *Pulmon Pharmacol*, 7:1–17.
- Norman P. 1999. PDE4 inhibitors. *Curr Opin Ther Patents*, 9:1101–18.
- Norman P. 2000. PDE4 inhibitors: sustained patenting activity as leading drugs near the market. *Curr Opin Ther Patents*, 10:1415–27.
- Oberholte R, Ratzliff J, Baecker PA, et al. 1997. Multiple splice variants of phosphodiesterase PDE4C cloned from human lung and testis. *Biochim Biophys Acta*, 1353:287–97.
- O'Donnell R, Breen D, Wilson S, et al. 2006. Inflammatory cells in the airways in COPD. *Thorax*, 61:448–54.
- Oger S, Mehats C, Barnette MS, et al. 2004. Anti-inflammatory and utero-relaxant effects in human myometrium of new generation phosphodiesterase 4 inhibitors. *Biol Reprod*, 70:458–64.
- Ohbayashi H. 2002. Neutrophil elastase inhibitors as treatment for COPD. *Expert Opin Investig Drugs*, 11:965–80.
- O'Shaughnessy TC, Ansari TW, Barnes NC, et al. 1997. Inflammation in bronchial biopsies of subjects with chronic bronchitis: inverse relationship of CD8⁺ T lymphocytes with FEV₁. *Am J Respir Crit Care Med*, 155:852–7.
- Ouagued M, Martin-Chouly CA, Brinchault G, et al. 2005. The novel phosphodiesterase 4 inhibitor, CI-1044, inhibits LPS-induced TNF-alpha production in whole blood from COPD patients. *Pulm Pharmacol Ther*, 18:49–54.
- Owen CA. 2005. Proteinases and oxidants as targets in the treatment of chronic obstructive pulmonary disease. *Proc Am Thorac Soc*, 2:373–85.
- Palfreyman MN, Souness JE. 1996. Phosphodiesterase type IV inhibitors. *Prog Med Chem*, 33:1–52.
- Palfreyman MN. 1995. Phosphodiesterase type IV inhibitors as antiinflammatory agents. *Drugs Fut*, 20:793–804.
- Panina P, Mariani M, D'Ambrosio D. 2006. Chemokine receptors in chronic obstructive pulmonary disease (COPD). *Curr Drug Targets*, 7:669–74.
- Pauwels R. 2002. Inhaled glucocorticosteroids and chronic obstructive pulmonary disease: how full is the glass? *Am J Respir Crit Care Med*, 165:1579–80.
- Pauwels RA, Buist AS, Calverley PM, et al. 2001. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Resp Crit Care Med*, 163:1256–76.
- Persson CG, Erjefält I, Gustafsson B. 1988. Xanthines-symptomatic or prophylactic in asthma? *Agents Actions*, (Suppl 23):137–55.
- Persson CG. 1986. Overview of effects of theophylline. *J Allergy Clin Immunol*, 78:780–7.
- Petty TL. 1998. Supportive therapy in COPD. *Chest*, 113: Suppl 256S–262S.
- Pinto IL, Buckle DR, Readshaw SA, et al. 1993. The selective inhibition of phosphate phosphodiesterase from vascular smooth muscle by rolipram analogues. *Bioorg Med Chem Lett*, 3:1743–6.
- Polkey MI, Moxham J. 2006. Attacking the disease spiral in chronic obstructive pulmonary disease. *Clin Med*, 6:190–6.

- Poole PJ, Black PN. 2006. Mucolytic agents for chronic bronchitis or chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, 19:CD001287.
- Profita M, Chiappara G, Mirabella F, et al. 2003. Effect of cilomilast (Ariflo) on TNF-alpha, IL-8, and GM-CSF release by airway cells of patients with COPD. *Thorax*, 58:573–9.
- Rabe KF, Bateman ED, O'Donnell D, et al. 2005. Roflumilast – an oral anti-inflammatory treatment for chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet*, 366:563–71.
- Rabe KF, Magnussen H, Dent G. 1995. Theophylline and selective PDE inhibitors as bronchodilators and smooth muscle relaxants. *Eur Resp J*, 8:637–42.
- Rabe KF. 2006. Improving dyspnea in chronic obstructive pulmonary disease: optimal treatment strategies. *Proc Am Thorac Soc*, 3:270–5.
- Rahman I, Adcock IM. 2006. Oxidative stress and redox regulation of lung inflammation in COPD. *Eur Respir J*, 28:219–42.
- Rahman I, Kilty I. 2006. Antioxidant therapeutic targets in COPD. *Curr Drug Targets*, 7:707–20.
- Ram FS, Jones PW, Castro AA, et al. 2002. Oral theophylline for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, 4:CD003902.
- Ram FS, Rodriguez-Roisin R, Granados-Navarrete A, et al. 2006. Antibiotics for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*, 19:CD004403.
- Reid P. 2002. Roflumilast Altana Pharma. *Curr Opin Investig Drugs*, 3:1165–70.
- Renau TE. 2004. The potential of phosphodiesterase 4 inhibitors for the treatment of depression: opportunities and challenges. *Curr Opin Investig Drugs*, 5:34–9.
- Riccioni G, Di Ilio C, Conti P, et al. 2004. Advances in therapy with anti-leukotriene drugs. *Ann Clin Lab Sci*, 34:379–87.
- Richmond R, Zwar N. 2003. Review of bupropion for smoking cessation. *Drug Alcohol Rev*, 22:203–20.
- Robichaud A, Stamatiou PB, Jin C, et al. 2002. Deletion of phosphodiesterase 4D in mice shortens alpha2-adrenoceptor-mediated anesthesia, a behavioural correlate of emesis. *J Clin Invest*, 110:1045–52.
- Rodriguez-Roisin R. 2006. COPD exacerbations. 5: management. *Thorax*, 61:535–44.
- Rogers DF, Barnes PJ. 2006. Treatment of airway mucus hypersecretion. *Ann Med*, 38:116–25.
- Rogers DF, Giembycz MA. 1998. Conquering airway inflammation in the 21st century. *Drug Disc Today*, 3:532–5.
- Romano SJ. 2005. Selectin antagonists: therapeutic potential in asthma and COPD. *Treat Respir Med*, 4:85–94.
- Sadowska AM, Klebe B, Germonpre P, et al. 2006. Glucocorticosteroids as antioxidants in treatment of asthma and COPD. New application for an old medication? *Steroids*, 72:1–6.
- Saetta M, Di Stefano A, Turato G, et al. 1998. CD8⁺ T-lymphocytes in peripheral airways of smokers with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 157:822–6.
- Saint S, Bent S, Vittinghoff E, et al. 1995. Antibiotics in chronic obstructive pulmonary disease exacerbations: a meta-analysis. *JAMA*, 273:957–60.
- Sanz MJ, Cortijo J, Morcillo EJ. 2005. PDE4 inhibitors as new anti-inflammatory drugs: effects on cell trafficking and cell adhesion molecules expression. *Pharmacol Ther*, 106:269–97.
- Sasaki M, Kashima M, Ito T, et al. 2000. Differential regulation of metalloproteinases production proliferation and chemotaxis of human lung fibroblasts by PDGF, interleukin-1b and TNFa. *Mediators Inflamm*, 9:155–60.
- Scapin G, Patel SB, Chung C, et al. 2004. Crystal structure of human phosphodiesterase 3B: atomic basis for substrate and inhibitor specificity. *Biochem*, 43:6091–100.
- Schmidt BM, Kusma M, Feuring M, et al. 2001. The phosphodiesterase 4 inhibitor roflumilast is effective in the treatment of allergic rhinitis. *J Allergy Clin Immunol*, 108:530–6.
- Schramm CM, Grunstein MM. 1992. Assessment of signal transduction mechanisms regulating airway smooth muscle contractility. *Am J Physiol-Lung Cell Molec Physiol*, 262:L119–39.
- Segura-Valdez L, Pardo A, Gaxiola M, et al. 2000. Upregulation of gelatinases A and B, collagenases 1 and 2, and increased parenchymal cell death in COPD. *Chest*, 117:684–94.
- Seneff MG, Wagner DP, Wagner RP, et al. 1995. Hospital and 1-year survival of patients admitted to intensive care units with acute exacerbation of chronic obstructive pulmonary disease. *JAMA*, 274:1852–7.
- Senior RM, Anthonisen NR. 1998. Chronic obstructive pulmonary disease (COPD). *Am J Resp Crit Care Med*, 157:139–47.
- Siafakas NM, Tzortzaki EG. 2002. Few smokers develop COPD. Why? *Respir Med*, 96:615–24.
- Silver PJ, Hamel LT, Perrone MH, et al. 1988. Differential pharmacologic sensitivity of cyclic nucleotide phosphodiesterase isozymes isolated from cardiac muscle, arterial and airway smooth muscle. *Eur J Pharmacol*, 150:85–94.
- Sin DD, Johnson M, Gan WQ, et al. 2004. Combination therapy of inhaled corticosteroids and long-acting beta2-adrenergics in management of patients with chronic obstructive pulmonary disease. *Curr Pharm Des*, 10:3547–60.
- Sin DD, McAlister FA, Man SF, et al. 2003. Contemporary management of chronic obstructive pulmonary disease: scientific review. *JAMA*, 290:2301–12.
- Slim RM, Song Y, Albassam M, et al. 2003. Apoptosis and nitrate stress associated with phosphodiesterase inhibitor-induced mesenteric vasculitis in rats. *Toxicol Pathol*, 31:638–45.
- Soderling SH, Beavo JA. 2000. Regulation of cAMP and cGMP signaling: new phosphodiesterases and new functions. *Curr Opin Cell Biol*, 12:174–9.
- Soto FJ, Hanania NA. 2005. Selective phosphodiesterase-4 inhibitors in chronic obstructive lung disease. *Curr Opin Pulmon Med*, 11:129–34.
- Souness JE, Griffin M, Maslen C, et al. 1996. Evidence that cyclic AMP phosphodiesterase inhibitors suppress TNF alpha generation from human monocytes by interacting with a 'low-affinity' phosphodiesterase 4 conformer. *Br J Pharmacol*, 118:649–58.
- Souness JE, Houghton C, Sardar N, et al. 1997. Evidence that cyclic AMP phosphodiesterase inhibitors suppress interleukin-2 release from murine splenocytes by interacting with a 'low-affinity' phosphodiesterase 4 conformer. *Br J Pharmacol*, 121:743–50.
- Souness JE, Rao S. 1997. Proposal for pharmacologically distinct conformers of PDE4 cyclic AMP phosphodiesterases. *Cell Signalling*, 9:227–36.
- Spina D. 2000. The potential of PDE4 inhibitors in asthma or COPD. *Curr Opin Investig Drugs*, 1:204–13.
- Spina D. 2003. Phosphodiesterase-4 inhibitors in the treatment of inflammatory lung disease. *Drugs*, 63:2575–94.
- Spina D. 2004. The potential of PDE4 inhibitors in respiratory disease. *Curr Drug Targets Inflamm Allergy*, 3:231–6.
- Spond J, Case N, Chapman RW, et al. 2003. Inhibition of experimental acute pulmonary inflammation by pirfenidone. *Pulm Pharmacol Ther*, 16:207–14.
- Stockley RA. 2002. Neutrophils and the pathogenesis of COPD. *Chest*, 121 5 (Suppl 1):S1S–155S.
- Sullivan P, Bekir S, Jaffar Z, et al. 1994. Anti-inflammatory effects of low-dose oral theophylline in atopic asthma. *Lancet*, 343:1006–8.
- Szafranski W, Cukier A, Ramirez A, et al. 2003. Efficacy and safety of budesonide/formoterol in the management of chronic obstructive pulmonary disease. *Eur Respir J*, 21:74–81.
- Taylor DR. 1998. Chronic obstructive pulmonary disease; no longer any justification for therapeutic nihilism. *Br Med J*, 316:7143.
- Teixeira MM, Gristwood RW, Cooper N, et al. 1997. Phosphodiesterase (PDE)4 inhibitors: Anti-inflammatory drugs of the future? *Trends Pharmacol Sci*, 18:164–70.
- Thompson WJ. 1991. Cyclic nucleotide phosphodiesterases: pharmacology, biochemistry and function. *Pharmacol Ther*, 51:13–33.
- Timmer W, Leclerc V, Birraux G, et al. 2002. The new phosphodiesterase 4 inhibitor roflumilast is efficacious in exercise-induced asthma and leads to suppression of LPS-stimulated TNF-alpha ex vivo. *J Clin Pharmacol*, 42:297–303.

- Torphy TJ, Barnette MS, Underwood DC, et al. 1999. Ariflo (SB 207499), a second generation phosphodiesterase 4 inhibitor for the treatment of asthma and COPD: From concept to clinic. *Pulmon Pharmacol Ther*, 12:131–5.
- Torphy TJ, Page C. 2000. Phosphodiesterases: the journey towards therapeutics. *Trends Pharmacol Sci*, 21:157–9.
- Torphy TJ, Stadel JM, Burman M, et al. 1992. Coexpression of human cAMP-specific phosphodiesterase activity and high affinity rolipram binding in yeast. *J Biol Chem*, 267:1798–804.
- Torphy TJ, Undem BJ. 1991. Phosphodiesterase inhibitors: new opportunities for the treatment of asthma. *Thorax*, 46:512–23.
- Torphy TJ. 1994. Beta-Adrenoceptors, cAMP and airway smooth muscle: challenges to the dogma. *Trends Pharmacol Sci*, 15:370–4.
- Torphy TJ. 1998. Phosphodiesterase isozymes: molecular targets for novel antiasthma agents. *Am J Resp Crit Care Med*, 157:351–70.
- Trifilieff A, Keller TH, Press NJ, et al. 2005. CGH2466, a combined adenosine receptor antagonist, p38 mitogen-activated protein kinase and phosphodiesterase type 4 inhibitor with potent in vitro and in vivo anti-inflammatory activities. *Br J Pharmacol*, 144:1002–10.
- Undem BJ, Meeker SN, Chen J. 1994. Inhibition of neurally mediated nonadrenergic, noncholinergic contractions of guinea pig bronchus by isozyme-selective phosphodiesterase inhibitors. *J Pharmacol Exp Ther*, 271:811–17.
- Underwood DC, Bochnowiz S, Osborn RR, et al. 1998. Antiasthmatic activity of the second generation phosphodiesterase 4 (PDE4) inhibitor SB 207499 (Ariflo) in the guinea pig. *J Pharmacol Exp Ther*, 287:988–95.
- Underwood DC, Kotzer CJ, Bochnowicz S, et al. 1994. Comparison of phosphodiesterase III, IV, and dual III/IV inhibitors on bronchospasm and pulmonary eosinophil influx in guinea pigs. *J Pharmacol Exp Ther*, 270:250–9.
- van Schalkwyk E, Strydom K, Williams Z, et al. 2005. Roflumilast, an oral, once-daily phosphodiesterase 4 inhibitor, attenuates allergen-induced asthmatic reactions. *J Allergy Clin Immunol*, 116:292–8.
- van Schayck CP, Reid J. 2006. Effective management of COPD in primary care – the role of long-acting beta agonist/inhaled corticosteroid combination therapy. *Prim Care Respir J*, 15:143–51.
- Vassallo R, Lipsky JJ. 1998. Theophylline: recent advances in the understanding of its mode of action and uses in clinical practice. *Mayo Clinic Proc*, 73:346–54.
- Veeramachaneni SB, Sethi S. 2006. Pathogenesis of bacterial exacerbations of COPD. *COPD*, 3:109–15.
- Vignola AM. 2004. PDE4 inhibitors in COPD—a more selective approach to treatment. *Respir Med*, 98:495–503.
- Wang P, Wu P, Ohleth KM, et al. 1999. Phosphodiesterase 4B2 is the predominant phosphodiesterase species and undergoes differential regulation of gene expression in human monocytes and neutrophils. *Mol Pharmacol*, 56:170–4.
- Weinberger M, Hendeles L. 1996. Theophylline in asthma. *N Engl J Med*, 334:1380–8.
- Weinberger M. 1988. Treatment of chronic asthma with theophylline. *ISI Atlas Sci Pharmacol*, 2:53–61.
- Wong ML, Whelan F, Deloukas P, et al. 2006. Phosphodiesterase genes are associated with susceptibility to major depression and antidepressant treatment response. *Proc Natl Acad Sci USA*, 103:15124–29.
- Wright JL, Churg A. 2006. Advances in the pathology of COPD. *Histopathol*, 49:1–9.
- Zaher C, Halbert R, Dubois R, et al. 2004. Smoking-related diseases: the importance of COPD. *Intl J Tuberculosis Lung Dis*, 8:1423–8.
- Zhu YK, Liu XD, Skold CM, et al. 2001. Synergistic neutrophil elastase-cytokine interaction degrades collagen in three-dimensional culture. *Am J Physiol Lung Cell Mol Physiol*, 281:L868–78.

