ORIGINAL RESEARCH **RETRACTED ARTICLE: Efficient miRNA Inhibitor** with GO-PEI Nanosheets for Osteosarcoma Suppression by Targeting PTEN

This article was published in the following Dove Press journal: International Journal of Nanomedicine

Lingling Ou¹ Haiyingjie Lin² Yuwei Song¹ Guogiang Tan¹ Xiujuan Gui¹ Jinyuan Li¹ Xiaoting Chen³ Zhendong Deng³ Shaogiang Lin¹

¹Department of Stomatology, The First Affiliated Hospital of Jinan University, Guangzhou 510632, People's Republic of China; ²Department of Orthopedics, The Third Affiliated Hospital, Southern Medical University, Guangzhou 510630, People's Republic of China; ³Integrated Traditional and Western Medicine Research Center of the First Affiliated Hospital of Guangdong Pharmaceut University, Guangzhou 510000, Pople's Republic of China



The First Affiliated spital of Jinan University, No. 613 West Huangpu Avenue, Guangzhou 510632, People's Republic of China Email oulingling 123@126.com

Shaogiang Lin

Integrated Traditional and Western Medicine Research Center of the First Affiliated Hospital of Guangdong Pharmaceutical University, No. 19 Nonglinxia Road, Guangzhou 510000, People's Republic of China Email tlshq@jnu.edu.cn



vel way to steosarcoma, and Background: Gene therapy is considered a microRNAs are potential therapeutic targets for steosar ma. miR-244 has been found to nene oxid (GO) is widely used for promote osteosarcoma aggression and met asis. gene delivery for the distinct physiocher cal properties and mimal cytotoxicity. Methods: Polyethyleneimine (PEI) zed GO convex was well-prepared and loaded

with miR-214 inhibitor at different concentration. The load efficacy was tested by gel retardation assay and the cy3-labeler fluorescence of cearlar uptake. The experiments of wound healing, immunofluorescence staining, Weetern blot, qRT-PCR and immunohistochemical staining were performed to meaning the inhibitor ry effect of the miR-214 inhibitor systematically released from the complexes a just MG/ U2OS cells and xenograft tumors.

Results: The sy mechanistic elucidation of the efficient delivery of the miR-214 indi the inhibition of cellular miR-214 caused a decrease in inhibitor by GO-F na cell i on and migration and an increase in apoptosis by targeting phososteosar nsin he olog (PTEN). The synergistic combination of the GO-PEI-miR-214 e and phat bitor a CDDP comotherapy showed significant cell death. In a xenograft mouse GO-PEI-miR-214 inhibitor significantly inhibited tumor volume growth. mo

Conclusion: This study indicates the potential of functionalized GO-PEI as a vehicle for miRNA in pitor delivery to treat osteosarcoma with low toxicity and miR-214 can be rood target for osteosarcoma therapy.

Ke, ords: osteosarcoma, GO-PEI-miR-214 inhibitor, tumor suppression

Introduction

Osteosarcoma is one of the most common types of aggressive malignant bone tumors.¹ Since the high invasiveness and rapid metastasis, the conventional therapy of osteosarcoma was subject to serious restriction. microRNAs (miRNA) have acted as important prognostic factors in patients with osteosarcoma,^{2,3} and miR-214 was proved to be linked to osteosarcoma progression and adverse prognosis. $^{4-6}$ The depletion of miR-214 or miR-214 inhibition increased radiosensitivity or chemosensitivity in both osteosarcoma cells and mouse xenograft models.4,7 However, naked miRNAs or miRNA inhibitors can be rapidly degraded in culture medium or in vivo. Therefore, vectors that protect and deliver miRNAs or miRNA inhibitors into cells are imperative for miRNA therapy.

Many gene delivery tools have been developed, including liposomes, viruses, polymer particles, carbon nanomaterials, etc.^{8,9} Viral vectors are efficient gene delivery

CO 0 S C2020 Ou et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

but the biosafety concerns limit its extensive application.¹⁰ However, the transfection efficiency of nonviral vectors is low and needs to be improved.¹¹ The major obstacle is to develop nonviral-based safe and efficient gene delivery vehicles.

Graphene oxide (GO) nanosheets have been demonstrated to utilize in drug delivery for cancer therapy for their abundant oxidized functional groups and large surface area to adsorb nucleobases or aromatic compounds.^{12–14} In this study, we designed a GO-PEI complex to deliver the miRNA into cells in vitro and in vivo. Negatively charged GO is able to bind cationic polyethyleneimine (PEI) polymers to form stable GO-PEI complexes by electrostatic interactions. GO-PEI complexes were highly enriched in positive charges that could bind to negative miRNA and wrap miRNA. Therefore, it was hypothesized that the highly enriched positive charges of GO-PEI complexes allow effective loading of miRNA inhibitors and delivery inhibitors into cells for tumor suppression. To test this hypothesis, we delivered miR-214 inhibitor into two different osteosarcoma cell lines by means of GO-PEI complexes. And the progression of osteosarcoma was inhibited by GO-PEI-miR -214-inhibitor (GP-inhibitor) complexes through PTEN/ PI3K/Akt and ERK1/2 signaling pathways. A schematic diagram was shown that GP-inhibitor delivered to osteosarcoma cells for therapy in Figure 1.

Experiments and Methods Preparation of GO-PEI

The GO-PEI complexes were obtained by GO (Aladdin, Shanghai, China) conjugated with PEI (Sigma-Aldrich, Missouri, USA) through electrostatic adsorption. Briefly, PEI solutions were mixed with diluted GC solution at a GO:PEI weight ratio of 1:3. A 25-kDa PK solution (1 h (mL) was added to a GO solution (apprecimate. 1 mg/mL) in 10 min. The mixture was ultrasonic ded for 15 m and stirred overnight. The mixed sole on was washed with ddH2O and ns, 4°C d filtered. The size centrifuged at 3000 pm, 5 distributions of and GO-LL we measured by a DLS spectrophoto leter tsuka Electronics, Japan). The FTIR used to the presence of functional groups spectrum surface of GO, PLI, and GO-PEI complexes. The on um was musured from 4000 to 400 cm⁻¹ using an spe instrument Vertex 70, Bruker, Germany) at room FTI



Figure I A schematic diagram depicting the efficiency of miR-214 inhibitor delivery into cells by means of graphene oxide (GO)-polyethyleneimine (PEI). GO was conjugated with PEI polymers to form positively charged GO-PEI complexes. Negatively charged miR-214 inhibitor was wrapped into the GO-PEI complexes by electrostatic interactions. GP-inhibitor complexes were delivered into the cytoplasm through endocytosis and inhibited the expression of miR-214 in osteosarcoma cells. Due to miR-214 inhibitor by GP-inhibitor, the expression of PTEN could be increased and demonstrated a therapeutic effect in vitro. The GP-inhibitor also showed good therapeutic effect on the subcutaneous xenograft tumor mouse model through intratumoral injection.

temperature. The samples suspended in deionized water were analyzed the surface charge using a Zeta-sizer (Malvern Nano ZS, Malvern, UK). GO and GO-PEI were also examined by TEM (HT7700, Hitachi, Japan) analysis. The ultravioletvisible (UV-vis) absorption spectrum (190 to 800 nm) for GO-PEI and GO-PEI-inhibitor (10 µg/mL) samples was obtained by a UV-vis spectrophotometer (Shimadzu, Japan). To determine the gel retardation assay, the GO-PEI-miRNA inhibitor solution (9 µL) was added with 3 µL of 4× loading buffer, loaded on a 1% (w/v) agarose gel and then electrophoresed at 100 V for 25 min.

Cell Viability Assay

Human U2OS and MG-63 cell lines (Geneseed, Guangzhou, China) were used in this study. The CCK-8 assay was applied to measure the cytotoxicity of GO and GO-PEI complexes. Briefly, U2OS and MG-63 cells were cultured with GO-PEI at a series of concentrations (0, 5, 10, 20, 30, 40 and 50 μ g/mL) for 24 hrs. Then, the CCK-8 test solution was added to each well and incubated at 37°C for 2 hrs. The absorbance of each well was measured at 450 nm using a microplate reader (Molecular Devices, CA, USA). The experiment was repeated 3 times.

Measurement of Cellular Uptake of GO-EL

To measure the cellular uptake of GO-PF PEI v stained with FITC dye.¹⁵ GO-PEI and TC-BS solution (Bioss Inc., Beijing, China) (1 mg/mL) ver mixe a mass ratio of 1:1 at 37°C and in soated for thrs, the supernatant containing any unbound IN BSA was depended after centrifuged at 10,000 g. For the cellular ptake test, U2OS and MG-63 cells were see if a in 24-well plan $(5 \times 10^4 / \text{well})$ and then incubated with afferent oses of FITC-labeled GO-PEI in serum-free medius for 24 h. GO PEI-treated U2OS and weight fixed th 49 (w/v) paraformaldehyde at MG-63 cell 4°C for 5 mins. The cell ytoskeleton was stained with phelonum 10, Invitrogen, CA, USA) for rhodam 30 mins at 4 and then incubated with 4.6-diamidino-2-phenylindole (DAN Vector Laboratories, USA) (1:2000) for 15 mins. The stained cells were examined under a laser scanning confocal microscope (Carl Zeiss, Inc, Jena Germany).

RNA Delivery and RNase Protection Assay of GP-Inhibitor

To analyze the miRNA inhibitor delivery efficiency of GO-PEI, U2OS and MG-63 cells were seeded in 24-well plates $(5\times10^4/\text{well})$ overnight before transfection. GO-PEI was

mixed with a cy3-conjugated miRNA inhibitor (N/P ratio of 30) for 1 h and then added into U2OS and MG-63 culture media (5 μ g/mL) for 1, 4, 8, 16, 24, 36, 48 and 72 hrs. The miR-214 inhibitor was delivered by Lipofectamine 2000 (Life Technologies, CA, USA) taken as one of the controls according to the manufacturer's instructions. Cy3-miRNA inhibitor delivery was analyzed by fluorescence microscopy (Olympus Co., Ltd., Japan). To evaluate the stability of GPinhibitor complexes in RNase A solution (30 μ g/mL, Sigma), miR-214 inhibitor complexes were added simultaneously or GP-inhibitor complexes were added into the solution. The cells were observed using a fluorescence microscope, and the fluorescence intensity was analyzed.

Invasion Assay and Yound realing

To measure whether Conhibitor inhibited the biological function of _____eosarcon___ells, ____nswell assays and wound healing a systere perform. For the invasion assay, matrigel (BD Bioscien s, CA, USA, 25 µg/50 µL) was added σ the upper welk with 8 μm pore size polycarbonate hembrane filters. DMEM culture medium (containing 10%) BS) was added into the lower layer of the transwell. MG63 an V2OS cons were added into the upper chamber with FBS free cuture medium (5 μ g/mL lipo-inhibitor or GP-inhibitor) . A 24 h of incubation, the cells in the upper well were fixed with 4% paraformaldehyde and stained with 0.4% crystal violet. Cells in the upper part of the filter were removed, and the number of stained migrated cells was counted under an inverted microscope (Olympus Co., Ltd., Japan). For the wound healing, the confluent MG63 cells cultured in six-well plates were wounded with a yellow tip (tip of 100 μ L), and the cells were observed to migrate over a cell-free gap after treatment with lipo-inhibitor or GP inhibitor (5 µg/mL) for 0, 24 and 48 hrs. The cells were then examined under a microscope (Olympus Co., Ltd., Japan).

Immunofluorescence and Western Blotting Analysis

MG63 cells were seeded onto coverslips in 24-well plates $(5 \times 10^4$ /well) and co-treated with lipo-inhibitor or GP-inhibitor for 24 h. The cells were fixed and permeabilized as described, ¹⁶ blocked in 3% horse serum, and incubated with the primary antibody (1:250 dilution) overnight at 4°C. The primary antibody was specific for p-PTEN (CST, MA, USA) and cleaved caspase-3 (ab2302, Abcam, MA, USA), and the secondary antibody (1:300, Jackson ImmunoResearch, PA,

USA) was incubated for 1 h at room temperature and counterstained with DAPI. The images were visualized with a fluorescence microscope and analyzed by ImageJ. For Western blot analysis, cells were collected and lysed for protein, and the same amount of proteins was loaded onto a 12% SDS-PAGE gel (Beyotime, Nanjing, China) and transferred onto PVDF membranes. The membranes were blocked and incubated with monoclonal antibodies directed against p-PTEN, PTEN, p-Akt, Akt, p-ERK1/2, ERK1/2, cleaved caspase-3 and cleaved caspase-8 (CST, MA, USA), followed by HRP-conjugated secondary antibodies (CST, MA, USA). GAPDH (CST, MA, USA) was used as an internal control. The transferred membranes were reacted with an ECL Western blot substrate kit (Beyotime, Nanjing, China) for exposure and analyzed using a FluorChem™ M System (Protein Simple, CA, USA).

qRT-PCR Analysis

Total RNA or miRNA were isolated by RNeasy Mini or miRNeasy Mini kits (Qiagen, Valencia, CA, USA). The mRNA levels for genes were determined using SYBR Green qRT-PCR mix (Promega, Madison, USA) on a 7500 Fast Real-Time PCR System (Applied Biosystems, MA, USA). The expression of miR-214 was measured using Bulge-LoopTM miRNA qRT-PCR kit (Ribobio, Guangzhou China). Primers used for qRT-PCR are listed below 🕻 twist. forward, TTCTCGGTCTGGAGGATGGA; erse, CAT TG TTTCTCCTTCTCTGGAAACA; N-cadhe forw GGAAATGGAAACTTGATGGC; rey se, 0 GCTCA CTGCTCTCATA; E-cadherin, for 1, AGTCA TTCAG ACTCCAGCC; reverse, TGT, GCTC CGGCGTCAAA; GAPDH, forward, GTC/ GGCTGAO, ACGGGAAG, reverse, GAGGGGGGCAC AGATCATGACC.

In vivo Antitumor Micacy

Six-week-old camale ALB/c up a mice (20–22 g) were obtained from Southore Medical University Animal Center. The animal experiments were performed with the approval of Jinan University Institutional Animal Care and Use Committee and followed the guidelines of the National Act on the Use of Experimental Animals (China). Five mice were housed per cage in a temperature and humidity-controlled room with a 12 hr light/dark cycle. All mice had free access to sterilized water and food.

MG63 cells (1×10^6 cells) suspended in 100 µL of Matrigel were subcutaneously injected into the right flank of female BALB/c nude mice. Once the tumor volume reached approximately 100 mm³, the mice got the treatment

of PBS, miR-214 inhibitor, GO-PEI, and GP inhibitor by peritumoral injection. The dosage of miR-214 inhibitor was 1 mg/kg per mouse, and PEI-miR-214 inhibitor was 5 mg/kg per mouse. The beginning of the treatment was defined as day 0, and the formulations were given to the mice on days 0, 4, 8, and 12. The mice were weighed, and tumor size was measured with a Vernier caliper at different time points. The tumor volume was calculated as follows: tumor volume = length × width²/2. Each tumor volume was independently measured, and fold changes in volume were calculated according to the tumor volume on day 0

Histology and Immunoh, tochemiliry

The mice were anesthetize and sach used at any 20, and the xenograft tumors are isolated anotated with 4% paraformaldehyde. For immunohistochemical staining, sections were blocked with 2% goat erum in PBS for 30 min, follower by the primary antibody (1:200 dilute) incubation at 4°C exernight. The primary antibodies were second for p-PTEs caspase-3 and caspase-8 (CST, MA USA) and followed by the secondary antibody incubation for 2 h at nom temperature. Three samples for each group were measured by light microscopy (Olympus, X71, Japany.

Statistical Analysis

he data are presented as the mean \pm SD (standard deviation). Statistical comparisons between different groups or two groups were evaluated by one-way ANOVA test or *t*-test comparison. Statistical significance was determined with a P-value <0.05.

Results

Characterization of GO-PEI Complexes

It has been reported that GO needs to be modified for loading nucleic acids, and PEI is known as the gold standard of cationic polymers for gene transfection.^{17,18} GO-PEI complexes were synthesized at a GO: PEI weight ratio of 1:5. Pristine GO and GO-PEI were measured by transmission electron microscopy (TEM). It revealed that GO sheets might be multiple layers and highly agglomerated, GO-PEI slightly increased the surface area (Figure 2A). Dynamic light scattering (DLS) results showed that the size distributions of GO and GO-PEI were 158.6 ± 46.8 nm and 205.5 ± 65.9 nm, respectively (Figure 2B). The zeta potential of GO was negative (-32.06 ± 3.26 mV), and that of GO-PEI was positive (28.17 ± 5.09 mV), indicating that the negatively



Figure 2 Characterization of GO-PEI complexes. (A) SEM images of GO and GO-PEI complexes. Scale bars: 200 nm. (B) The size distributions of GO and GO-PEI were detected by DLS analysis. (C) Measurement of the zeta potential of GO and GO-PEI complexes. (D) FTIR spectra of PEI (top) and GO-PEI complexes (bottom). (E) UV-vis spectra normalized by their extinction coefficients at 260 nm. Blue line for GO-PEI-miR-214 inhibitor complex, red line for free miR-214 inhibitor sense strand, and dark line for GO-PEI; experiments were carried out in phosphate buffer (pH 7.4). (F) A gel retardation assay of the GP-inhibitor at different N/P ratios (0, 15, 30, 45).

charged GO is able to bind cationic PEI polymers by electrostatic interactions and the conjugated PEI changed the negative charge on the surface of GO.¹² Moreover, the potential of the GO-PEI inhibitor was slightly lower than that of GO-PEI, which meant that the negatively charged miR-214 inhibitor might be coupled with GO-PEI (Figure 2C). It is suggested that the GO-PEI complex can bond with negatively charged miRNA inhibitors to deliver

miRNA inhibitors into cells. Fourier transform infrared red (FTIR) analysis confirmed that GO-PEI and PEI showed four peaks at 665-910, 720-725, 1020-1250, and 2850–3000 cm⁻¹ denoting C-H, N-H, C-N and C-H groups, respectively. GO-PEI possessed all peaks of GO and PEI, with peaks in the 1000-1320, 1665-1760, and 3200-3500 cm⁻¹ ranges corresponding to the C-O, C=O, and O-H groups, respectively. These peaks indicated that PEI was successfully grafted onto GO by electrostatic interactions rather than by covalent bonding (Figure 2D). The UV-vis spectrum demonstrated an absorption peak at 280 nm for miRNA and GP-inhibitor, which meant that the miRNA inhibitor was loaded on GO-PEI (Figure 2E). To further examine the binding capacity of GO-PEI complexes to miR-inhibitor, we used agarose gel electrophoresis. GO-PEI was complexed with the miR-inhibitor at various N/P ratios (molar ratio of nitrogen of PEI to phosphate groups of miRNA). The GPM complex demonstrated significant electrophoretic retardation at the N/P ratio of 30, which meant that GO-PEI would wrap miR-inhibitor inside and prevent miR-inhibitor being degraded at the N/P ratio of 30 (Figure 2F).

Cellular Uptake of GO-PEI Analysis

To ensure that the uptake of GO-PEI complexes by cells was the first important step for GO-PEI-inhibitor de very, TEM images were used to confirm cell ar uptal . As shown in Figure 3A, GO-PEI sheets ar Identified enclosed in a membrane in the cytosol. In addition, FITC-labeled GO-PEI was also used track the Hular uptake and localization of O-PEI in N63 and U2OS anning confocal microscopy cells through laser (LSCM). FITC-labele GO-P/ (10 µg/mL, green fluorescence) was observed to unside the cells or bind on the plasma membrine and 8 h in poation, and more GO-PEI completes were aken up into the cells as the incuba-. An enlarged image of GO-PEI uptake tion time inclusion was shown in the right panel of Figure 3B. Next, the cytotoxicity of GO-EI to MG63 and U2OS cells was measured by a cell counting kit (CCK-8) assay. The cytotoxicity of MG63 or U2OS cells had no significant difference in different GO-PEI-treated groups (5, 10, 20, and 30 μ g/mL). The cells showed mild toxicity only when the concentration of GO-PEI reached 40 or 50 µg/mL. Moreover, GO-PEI-treated MG63 and U2OS cells were viable after 24 or 48 h (Figure 3C). These GO-PEI complexes showed low cytotoxicity, even the high-molecular-

5136

weight PEI (25 kDa) was used, suggesting that GO-PEI complexes were much safer than the single PEI polymers.

GO-PEI-Inhibitor Synthesis and Transfection Evaluation

To determine whether miRNA inhibitors were successfully transfected into cells by GO-PEI complexes, a cy3-labeled miRNA inhibitor was used to study the gene delivery. MG63 cells were incubated with free miR-214 inhibitor, GO-PEI, and different N/P ratios of GO-PEI-miR-214 inhibitor. After 24 h of incubation, the 15 W detected by fluorescence microscopy. The transfection efficiency was highest while the N/P ratio (GQ: PL inhibitor) v s 30, and the relative fluorescence of 3-labeled, iRN/ nhibitor in cells was weaker when the pration as below 30 (Figure 4A). The CO-N mplexer could release the inhibitors slowly in the cell and se fluorescence in the cells was very strong at 24 h a weakened as the time increased: however, the cells maintained cy3 fluorescence for much than 72 h post-tic sfection (Figure 4B). To further asse s the protective ability of GO-PEI complexes, we incu ted naked niRNA inhibitors or GO-PEI-inhibitors (GP-in. bitors) oth RNase A at 37°C. GO-PEI complexes h an N/P ratio of 30 were successfully loaded with ative charged miRNAs and delivered miRNAs into the cells. When RNase A was added in after GP-inhibitor pmplex formation, the cy3-labeled miRNAs were successfully delivered into the cells without degradation, and the fluorescence of cy3 in the cells was strong. However, when RNase A was added during the formation of the GP inhibitor complex resulted in loss of the effectiveness of cy3miRNA delivery (Figure 4C). These results demonstrated that GO-PEI could successfully load miRNA inhibitors into cells without the hydrolysis of RNA enzyme.

GP-Inhibitor Inhibits Migration and Invasion in MG63 and U2OS Cells

miR-214 was upregulated in osteosarcoma tissues and cells, miR-214 overexpression enhanced osteosarcoma cell proliferation and invasion.¹⁹ To investigate the effect of GP-inhibitors on cell migration and invasion, the MG63 and U2OS cell lines were examined by wound healing migration and transwell invasion assays after incubation with GP-inhibitor complexes. The transwell assays showed that GP inhibitor significantly reduced the invasion ability of MG63 and U2OS cells, and the relative migrated cell number dramatically decreased compared



Figure 3 Uptake of GO-PEI into osteosarcoma cells. (A) TEM images of GO-PEI-incubated and non-incubated cells (3 μ g/mL GO-PEI in the cell culture medium). The right panel is enlarged images of white squares in the image of GO-PEI-incubated cells. Scale bars: 1 μ m. (B) FITC-BSA-labeled GO-PEI (green) is visualized in cells through confocal laser scanning microscopy. Fluorescence images of FITC-labeled GO-PEI (green) within MG63 and U2OS cells are shown. The cell cytoskeleton was stained with α -tubulin (red) for MG63 cells and rhodamine-phalloidin (red) for U2OS cells, and the nuclei were stained with DAPI (blue). The right panel is enlarged images of white squares in the image of GO-PEI-incubated cells. Scale bars: 2 μ m. (C) Cell viability of MG63 and U2OS cells exposed to different concentrations of GO-PEI was measured by CCK8 assays. *p < 0.05.



with that of the neative control-transfected cells and naked inhibitor-transfected cells (Figure 5A and B). Similarly, the capacity of wound healing in MG63 cells was significantly attenuated by GP-inhibitor complexes, and the relative wound area was much larger than those of the control and naked inhibitor groups after 24 h or 48 h of treatment (Figure 5C and D). Moreover, it was reported that twist was highly expressed, accompanied with N-cadherin high expressed and E-cadherin low expressed, in osteosarcoma with metastasis.^{20,21} qRT-PCR results showed that the GP-inhibitor decreased the expression of twist and N-cadherin and increased the expression of E-cadherin compared to that of the control or the lipo-inhibitor group (Figure 5E). These results demonstrated that miR-214 has an important role in the regulation of osteosarcoma cellular motility and that the GP-inhibitor has good transfection and inhibits the invasive and metastatic capacity of osteosarcoma cells.



Figure 5 GP inhibitor inhibits in usion and migration in 563 and U2OS cells. (**A**) The invasion of MG63 and U2OS cells were measured for 24 h using a Matrigel-coated chamber with polycarbonate ters. (**B**) Statistical analysis of the invasion cell ratio is presented for each group. *p < 0.05. (**C**) The wounds of confluent MG63 cells were created with 100 µL yellow by and interacted with a lipo-inhibitor or GP-inhibitor (5 µg/mL). Images of the wound at different time points are shown, and the borders of the wound are shown by a pair to lack use. (**D**) Statistical analysis of cell migration by calculating the ratio of the final cell-free gap to the initial wound area. *p < 0.05. (**E**) The expression levels of the initial of the interacted with lipo-inhibitor or GP-inhibitor were evaluated by qRT-PCR. *p < 0.05.

GP-Intibiter And ted miR-214-Regulated Pathways in Osteosarcoma Cells

To assess the enct of GP-inhibitor complexes on osteosarcoma cells, the level of miR-214 was analyzed by qRT-PCR. The level of miR-214 was significantly decreased by the GP-inhibitor (Figure 6A). Bioinformatic analysis showed that miR-214 binding site is highly conserved and PTEN was suggested to be a target of miR-214 in osteosarcoma cells (Figure 6B), which was in accordance with previous reports.^{19,22} miR-214 has been reported to target the 3'-UTR of PTEN and induced the activation of the Akt and ERK signaling pathways to enhance cell growth.^{22,32,33} Thus, to analyze the underlying mechanism by which GP-inhibitors prevent osteosarcoma cell proliferation and metastasis, Western blot analysis was used to examine the effect of GP-inhibitors on the PTEN expression in MG63 cells. PTEN protein expression level was increased after GP inhibitor treatment in MG63 cells. Furthermore, GP inhibitors were found to significantly decrease the protein levels of p-Akt, and total Akt protein level was slightly decreased. In addition, the protein expression of p-ERK1/2 was significantly decreased by



Figure 6 GP-inhibitors affected miR-214-regulated pathways in osteosarcoma cells. (A) Measurements of miR-214 levels in cells after treatment with lipo-inhibitor or GP-inhibitor. *p < 0.05. (B) Bioinformatic analysis revealed that miR-214 potentially targeted PTEN by the 3' untranslated region (3'-UTR), and the miR-214 binding site was conserved among different species. (C) The expression levels of p-PTEN, PTEN, p-Akt, Akt, p-ERK1/2, ERK1/2, cleaved caspase-3 and cleaved caspase-8 after incubation with lipo-inhibitor or GP inhibitor or GP inhibitor. *D Representative immunofluorescence microscopy images of p-PTEN and caspase-3 after incubation with lipo-inhibitor or GP inhibitor. Scale bars: 20 μ m.

the GP inhibitor, and total ERK1/2 protein level remained identical (Figure 6C). PTEN acts as a tumor suppressor by inhibiting PI3K and ERK1/2 signaling, activating diverse

signaling pathways involved in cell growth and cell survival.^{23–26} Furthermore, the inhibition of miR-214 by the GP-inhibitor increased the expression of caspase-3 and

caspase-8 (Figure 6C). Caspase-3 and caspase-8 are major apoptosis effectors.^{27,28} In addition, immunofluorescence staining was performed to further confirm the effect of GPinhibitors in osteosarcoma cells. Consistent with the WB results, the expression level of PTEN was notably increased by the GP-inhibitor compared to those of the control and lipo-inhibitor groups. Accordantly, the expression of caspase-3 was also increased in GP-inhibitortreated cells (Figure 6D). Together, these data indicated that GP inhibitors contributed to promoting apoptosis and suppressing tumor growth by targeting PTEN via PI3K/ Akt- and ERK1/2-dependent signaling.

GP-Inhibitor Improved the Cytotoxicity of Cisplatin on Osteosarcoma Cells

One of the abilities of miR-214 is to inhibit apoptotic cell death and decrease chemotherapy sensitivity, which is the main cause of chemotherapy failure and disease progression.²⁹ Based on the WB and immunofluorescence results in Figure 6, GP-inhibitor complexes could efficiently increase the expression of apoptotic proteins. It was suggested that GP-inhibitors might have synergistic anticancer effects with chemotherapy. To explore whether GP-inh sensitize chemotherapy, we incubated MG63 and U2OS ells for 24 h and 48 h with GP-inhibitors and cisting (CDL As shown in Figure 7A, CDDP (5 µc nL)-treated ce. showed a slightly lower cell density the the c a significantly ever, the CDDP and GP-inhibitor -treat. decreased the cell density computed to the content The CCK-8 assay results showed that DDP (sug/mL) imparted mild cytotoxicity compared the untreated trol. On the other hand, obvious cell rowth inhibition and cell death were observed in cells treat with GP-inhibitor (5 µg/mL) and CDDP (50 mL), adding to a precipitous decrease in mpared to that of the control cell viability to 255 ± 1.8 $(82.7 \pm 5\%)$ at the inhibitor (56.33 ± 2.2%) after 48 h of gure 7B). These results indicated that the GPincubation inhibitor and DP synergistically enhanced the sensitivity of MG63 and U20S cells.

GP-Inhibitor Showed Superior Antitumor Activities in an MG63 Xenograft Mouse Model

To assess whether GP-inhibitors could inhibit tumor growth in vivo, GP-inhibitors were used to treat tumor-bearing mice. Saline, miR-214 inhibitor, GO-PEI or GP-inhibitor complexes (30 µL) were injected into mice 4 times by peritumoral injection, the mice were weighed, and the tumors were measured at different time points. Representative tumor images are shown in Figure 8A. The tumors with PBS treatment increased rapidly, and the tumor volume in the naked inhibitor and GO-PEI treatment groups showed no significant difference, which suggests that the naked inhibitor and GO-PEI had a negligible effect on the osteosarcoma tumors, showing 26.9% and 20% reduction in tumor volume, respectively, compared to the PBS-treated mice. On the other hand, GP-inhibitor showed a strong effection suppressing tumor growth (the inhibition rate was ver 73.0), with minimal change in mouse body weight using the observation period (Figure 8B and C). The proce with ferent transmission of the second seco show serious body yinght loss, when singested that there was no significant vicity from the GO-PEI and inhibitor treatment with the examental criod.

, we remain and sectioned the tumors Subsec at day 21 for his logy assessment. As shown in Figure 8D, HE ing results lowed that the tumor cells were large id dense, with hyperchromatism and pleomorphism. The esults illustrated that the saline and naked inhibitor treatment Lno effect on cell growth in the tumor tissues, and the GO not cause cell death in tissues. Notably, the GPtream bitor treatment-induced obvious cell death in tumor tissues. Moreover, the expression of PTEN, caspase-3 and caspase-8 proteins in tumor tissues largely increased with GP-inhibitor treatment compared to that in saline-treated tumors, which was consistent with the results in vitro. In addition, there were no noticeable signs of organ damage in the liver, spleen, lung or kidneys of mice treated with GPinhibitor, GO-PEI or miR-214-inhibitor compared with those of the control group (Figure 9). Thus, these results indicated that GO-PEI has a latent capacity to be used as a drug delivery platform and that GP-inhibitors have a good inhibitory effect on osteosarcoma tumor progression.

Discussion

Gene therapy has been found to hold tremendous potential for the treatment of cancers beyond conventional methods.^{30–32} Nonviral vector provides safer methods without the potential side effects of viral transfection, such as immunogenicity and carcinogenesis.³³ GO has been considered an ideal vector for gene delivery.^{34,35} In this study, we demonstrated that GO-PEI was an effective nonviral carrier for miRNA delivery to suppress osteosarcoma cancer progression. GO bonded to PEI by electrostatic interactions and formed GO-PEI complexes with



Figure 7 Combination of GP inhibitor and CDDP for improved using of osteosarcoma cells. (A) Image of MG63 and U2OS cell morphology changes after co-treatment with GP-inhibitor and CDDP for 24 h. Scale bars: μ (μ). (B) MG6s, μ viability measured by CCK-8 assay after co-treatment with GP-inhibitor and CDDP for 12, 24 and 48 hrs. *p < 0.05.

a positive charge (Figure 2). PEL with a may of cationic charge could wrap not sic acid to protect it from RNase releas of nucleic acid.³⁶ and facilitate the endost pale of endocytosis and FITC-labeled JO-PL was aggregated the cy plasm in osteosarcoma cancer cells ticular, GO-PEI exhibited high cell (Figure 3). a concentration of 50 µg/mL, likely viabilities even because GO with high biocompatibility in the complexes reduced the cell cytotoxicity of PEI.

As miR-214 has been revealed to upregulate and function as oncogenic miRNA in osteosarcoma cancer,^{19,22,37} the synthetic miR-214 inhibitor might be a novel therapeutic approach for osteosarcoma cancer. In this study, we revealed that GO-PEI was capable of intracellular delivery of cy3-labeled miR-214 inhibitors and that the sequence specificity of miR-214 inhibitors was toward endogenous miR-214; however, we found no observable fluorescence in MG63 cells treated with the naked inhibitor, GO-PEI, inhibitor or RNase co-culture (Figure 4).

miR-214 has been reported to promote osteosarcoma cancer cell invasion and proliferation.^{38,39} In this study, we found that GP-inhibitors significantly decreased osteosarcoma cells' invasion ability in transwells and migration ability after wound scratching, resulting in declining tumorigenicity. In accordance with this finding, GP-inhibitors decreased the cancer metastasis-associated genes, twist and N-cadherin, and increased E-cadherin (Figure 5). Furthermore, we investigated whether GP-inhibitors had much stronger inhibition of miR-214 expression than Lipofectamine, suggesting that miR-214 inhibition via GO-PEI effectively degraded or inhibited endogenous miR-214 (Figure 6). Research studies and biomedical database



Figure 8 Antitumor activities of GP-inhibitor in an MG63 xenograft mouse model. (**A**) Representative xenograft tumor tissue images of mice treated with saline, miR-214 inhibitor, GO-PEI or GP-inhibitor complexes (30 μ L). (**B**) Relative changes of tumor volume over time. The values are presented; n=5. (**C**) Relative changes of body weight over time and the values are presented; n=5. (**D**) Representative images of HE staining and immunohistochemical staining of PTEN, caspase 3 and caspase 8 in different treatment groups are shown. Scale bars: 50 μ m. *p < 0.05; **p < 0.01.

analysis demonstrated that PTEN was one of the targets of miR-214 in tumors.⁴⁰⁻⁴² A GP-inhibitor was found to significantly increase PTEN expression, likely because the

miR-214 inhibitor targeted the 3'-UTR of PTEN. Conversely, GP-inhibitors reduced Akt and ERK1/2 expression. Because PTEN is a tumor-suppressor gene that represses the



Figure 9 Organ tissue morphology observation after GP-in the treat of the logical analysis of the liver, spleen, kidney and lung after treatment with saline, miR-214 inhibitor, GO-PEI or GP inhibitor complexes (30 μL) for 20 day. Score bars: 100 μm.

cies,⁴³ PTLA can pathogenesis of various human Laligh negatively regulate the PI32 Akt²⁹ and A K1/2 signaling pathways^{44,45} to specific y inhibit tumorigenesis and contribute to reducing two and N adherin expression.^{46,47} In addition, GP-inhibitors in used approvide cell death and rapy so it by of CDDP (Figures 6 enhanced the emoth arted that miR-214 was an anti-apopand 7). It he been re 2 cells, ¹ however, GP-inhibitors blocked totic factor in cts of oncogenic miR-214 showing cell the downstream apoptosis upregulation. These results indicate that GO-PEI delivered miR-214 inhibitors that successfully degraded endogenous miR-214 and prevented the carcinogenicity of miR-214.

In the tumor xenograft models, we found that GPinhibitor-injected mice showed obvious cancer cell death and significant tumor shrinking. GP-inhibitors not only accumulated in the tumor site but also transported into intratumoral tissues. The nanosize of GO-PEI complexes made them easy to transport in vivo. Because high biocompatibility and stability are required for in vivo application of synthesized nanomaterials, GP-inhibitor complexes with high tumor cell targeting efficacy showed that GO-PEI had conjugation stability with miR-214 inhibitor against blood and body fluid. Here, we showed that miR-214 inhibitors could not only be delivered into osteosarcoma cells by GO-PEI but also specifically target intracellular miR-214, showing an antitumor effect.

Conclusions

High miR-214 expression in cancers has been associated with invasion, metastasis, poor prognosis and weak response to therapy.^{42,48,49} In this work, highly efficient miR-214 inhibitor delivery was shown to inhibit osteosarcoma cell invasion and migration in vitro and suppress tumor progression in vivo. This technology could deliver other therapeutic nucleic acid or nucleic acid inhibitors to treat other tumors or

diseases. Taken together, the GP inhibitor is a potential candidate for antitumor therapy in vivo.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 81703120 and No. 81071751, Beijing, China), Natural Science Foundation of Guangdong Province (2017A030310365, Guangzhou, China), and Guangdong Provincial Medical Research Foundation (A2016360, Guangzhou, China). The authors gratefully acknowledge financial support from China Scholarship Council.

Disclosure

The authors report no conflicts of interest in this work.

References

- Sampson VB, Kamara DF, Kolb EA. Xenograft and genetically engineered mouse model systems of osteosarcoma and Ewing's sarcoma: tumor models for cancer drug discovery. *Expert Opin Drug Discov*. 2013;8(10):1181–1189. doi:10.1517/17460441.2013.817988
- Kushlinskii NE, Fridman MV, Braga EA. Molecular mechanisms and microRNAs in osteosarcoma pathogenesis. *Biochemistry (Mosc)*. 2016;81(4):315–328. doi:10.1134/S0006297916040027
- Ram Kumar RM, Boro A, Fuchs B. Involvement and clinical upper. of microRNA in osteosarcoma. *Int J Mol Sci.* 2016;17(6) 377. doi:10.3390/ijms17060877
- Song YD, Li DD, Guan Y, Wang YL, Zheng J, are a modula s cisplatin sensitivity of osteosarcoma cells three in regula on of an robic glycolysis. *Cell Mol Biol (Noisy-Le-Cond.)*. 2017 (200):75–79. doi:10.14715/cmb/2017.63.9.14
- Wang Z, Cai H, Lin L, Tang M, Confl. Uprega and expression of microRNA-214 is linked to tumor expression and a curse prognosis in pediatric osteosarcoma. *Pedian Blood Cancer*. 2014;61 (2):206–210. doi:10.1002/pc.24763
- Cheng D, Qiu X, Zhuan M, Zhu C, Zou H, Le Z. MicroRNAs with prognostic significant in ostersarcoma: a systemic review and meta-analysis. *On parget.* 21, 7;8(46):81062–81074. doi:10.18632/ oncotarget.19009
- Li Y, Song T, Li Z, et al. Upregulation of miR-214 induced radioresistance of oscillation of workgeting PHLDA2 via PI3K/Akt signalog. Front. Icol. 2019; 298. doi:10.3389/fonc.2019.00298
- Karls TA, Branne TE, Liposome delivery of microRNA-145 to mesench, and stem cells leads to immunological off-target effects mediated by NG-I. *Mol Ther.* 2013;21(6):1169–1181. doi:10.1038/ mt.2013.55
- Wells DJ. Gene merapy progress and prospects: electroporation and other physical methods. *Gene Ther*. 2004;11(18):1363–1369. doi:10. 1038/sj.gt.3302337
- Nayak S, Herzog RW. Progress and prospects: immune responses to viral vectors. *Gene Ther*. 2010;17(3):295–304. doi:10.1038/gt.2009. 148
- Yin L, Song Z, Kim KH, Zheng N, Gabrielson NP, Cheng J. Nonviral gene delivery via membrane-penetrating, mannose-targeting supramolecular self-assembled nanocomplexes. *Adv Mater*. 2013;25 (22):3063–3070. doi:10.1002/adma.201205088
- Feng L, Zhang S, Liu Z. Graphene based gene transfection. Nanoscale. 2011;3(3):1252–1257. doi:10.1039/c0nr00680g

- Liu X, Ma D, Tang H, et al. Polyamidoamine dendrimer and oleic acid-functionalized graphene as biocompatible and efficient gene delivery vectors. ACS Appl Mater Interfaces. 2014;6(11):8173–8183. doi:10.1021/am500812h
- 14. Zhang L, Lu Z, Zhao Q, Huang J, Shen H, Zhang Z. Enhanced chemotherapy efficacy by sequential delivery of siRNA and anticancer drugs using PEI-grafted graphene oxide. *Small.* 2011;7 (4):460–464. doi:10.1002/smll.201001522
- 15. Ma J, Liu R, Wang X, et al. Crucial role of lateral size for graphene oxide in activating macrophages and stimulating pro-inflammatory responses in cells and animals. ACS Nano. 2015;9(10):10498–10515. doi:10.1021/acsnano.5b04751
- 16. Chen GY, Meng CL, Lin KC, et al. Graphene oxide as a chemosensitizer: diverted autophagic flux, enhanced nuclear import, elevated necrosis and improved antitumor effects. *Biomaterials* 10:12–22. doi:10.1016/j. biomaterials.2014.11.034
- Kim H, Namgung R, Singherk, Oh IK, Ku WJ. Graphene oxide-polyethylenimine nanocons byt as a gene convery vector and bioimaging tool. *Bioconic, Chem.* 11;22(12): 58–2567. doi:10. 1021/bc200397j
- Hobel S, Koburger John M, et al. Polyet echimine/small interfering RNA-mediated to eckdor a of vascular endothelial growth factor in vivo exert anti-tume affects symplestically with Bevacizumab. *J Gene Mm* 2010;12(3):2 300 ecdi:10.1002/jgm.1431
- Liu CL, u N. Liu GL, Tiang A. MiR214 promotes osteosarcoma tumor growth all metastasis by decreasing the expression of PTEN. Michael Rep. 2018, 2(4):6261–6266. doi:10.3892/mmr.2015.4197
- Yin K, Liao Q, He Zhong D. Prognostic value of twist and E-cadherin in patients with osteosarcoma. *Med Oncol.* 2012;29 (5):3449–34 doi:10.1007/s12032-012-0317-6
- Hou CH, J. In FL, Hou SM, Liu JF. Cyr61 promotes ithelial esenchymal transition and tumor metastasis of osteosarcoms-cy Raf-1/MEK/ERK/Elk-1/TWIST-1 signaling pathway. *Mol Cancer*. 2014;13:236. doi:10.1186/1476-4598-13-236
- wang X, Sun J, Fu C, Wang D, Bi Z. MicroRNA-214 regulates osteosarcoma survival and growth by directly targeting phosphatase and tensin homolog. *Mol Med Rep.* 2014;10(6):3073–3079. doi:10.3892/mmr.2014.2616
- Chalhoub N, Baker SJ. PTEN and the PI3-kinase pathway in cancer. *Annu Rev Pathol.* 2009;4:127–150. doi:10.1146/annurev.pathol.4.110807. 092311
- Wang X, Jiang X. PTEN: a default gate-keeping tumor suppressor with a versatile tail. *Cell Res.* 2008;18(8):807–816. doi:10.1038/cr.2008.83
- Baker A, Wyatt D, Bocchetta M, et al. Notch-1-PTEN-ERK1/2 signaling axis promotes HER2+ breast cancer cell proliferation and stem cell survival. *Oncogene*. 2018;37(33):4489–4504. doi:10.1038/ s41388-018-0251-y
- 26. Liu JJ, Li Y, Chen WS, et al. Shp2 deletion in hepatocytes suppresses hepatocarcinogenesis driven by oncogenic beta-Catenin, PIK3CA and MET. J Hepatol. 2018;69(1):79–88. doi:10.1016/j.jhep.2018. 02.014
- Chang Z, Xing J, Yu X. Curcumin induces osteosarcoma MG63 cells apoptosis via ROS/Cyto-C/Caspase-3 pathway. *Tumour Biol.* 2014;35 (1):753–758. doi:10.1007/s13277-013-1102-7
- Ding Y, Wang Y, Chen J, et al. p21 overexpression sensitizes osteosarcoma U2OS cells to cisplatin via evoking caspase-3 and Bax/Bcl-2 cascade. *Tumour Biol.* 2014;35(4):3119–3123. doi:10.1007/s13277-013-1404-9
- 29. Yang H, Kong W, He L, et al. MicroRNA expression profiling in human ovarian cancer: miR-214 induces cell survival and cisplatin resistance by targeting PTEN. *Cancer Res.* 2008;68(2):425–433. doi:10.1158/0008-5472.CAN-07-2488
- 30. Zhang L, Gao L, Li Y, et al. Effects of plasmid-based Stat3-specific short hairpin RNA and GRIM-19 on PC-3M tumor cell growth. *Clin Cancer Res.* 2008;14(2):559–568. doi:10.1158/1078-0432.CCR-07-1176

- 31. Tian Y, Guo B, Jia H, et al. Targeted therapy via oral administration of attenuated Salmonella expression plasmid-vectored Stat3-shRNA cures orthotopically transplanted mouse HCC. *Cancer Gene Ther*. 2012;19(6):393–401. doi:10.1038/cgt.2012.12
- Cheever TR, Berkley D, Braun S, et al. Perspectives on best practices for gene therapy programs. *Hum Gene Ther.* 2015;26(3):127–133. doi:10.1089/hum.2014.147
- Yin H, Kanasty RL, Eltoukhy AA, Vegas AJ, Dorkin JR, Anderson DG. Non-viral vectors for gene-based therapy. *Nat Rev Genet.* 2014;15(8):541–555. doi:10.1038/nrg3763
- 34. Sun Q, Wang X, Cui C, Li J, Wang Y. Doxorubicin and anti-VEGF siRNA co-delivery via nano-graphene oxide for enhanced cancer therapy in vitro and in vivo. *Int J Nanomedicine*. 2018;13:3713–3728. doi:10.2147/IJN.S162939
- 35. Yadav N, Kumar N, Prasad P, Shirbhate S, Sehrawat S, Lochab B. Stable dispersions of covalently tethered polymer improved graphene oxide nanoconjugates as an effective vector for siRNA delivery. ACS Appl Mater Interfaces. 2018;10(17):14577–14593. doi:10.1021/acsami.8b0 3477
- 36. Zhu J, Tang A, Law LP, et al. Amphiphilic core-shell nanoparticles with poly(ethylenimine) shells as potential gene delivery carriers. *Bioconjug Chem.* 2005;16(1):139–146. doi:10.1021/bc0498951
- Liang W, Gao B, Fu P, Xu S, Qian Y, Fu Q. The miRNAs in the pathgenesis of osteosarcoma. *Front Biosci.* 2013;18:788–794. doi:10.2741/4142
- 38. Zhang M, Wang D, Zhu T, Yin R. miR-214-5p targets ROCK1 and suppresses proliferation and invasion of human osteosarcoma cells. *Oncol Res.* 2017;25(1):75–81. doi:10.3727/096504016X1471907813 3401
- Xu Z, Wang T. miR-214 promotes the proliferation and invasion of osteosarcoma cells through direct suppression of LZTS1. *Biochem Biophys Res Commun.* 2014;449(2):190–195. doi:10.1016/j.bbrc. 2014.04.140
- Penna E, Orso F, Taverna D. miR-214 as a key hub that controcancer networks: small player, multiple functions. *J Invest Dermatol*. 2015;135(4):960–969. doi:10.1038/jid.2014.479

- 41. Yin Y, Cai X, Chen X, et al. Tumor-secreted miR-214 induces regulatory T cells: a major link between immune evasion and tumor growth. *Cell Res.* 2014;24(10):1164–1180. doi:10.1038/ cr.2014.121
- 42. Yang TS, Yang XH, Wang XD, Wang YL, Zhou B, Song ZS. MiR-214 regulate gastric cancer cell proliferation, migration and invasion by targeting PTEN. *Cancer Cell Int.* 2013;13(1):68. doi:10.1186/ 1475-2867-13-68
- 43. Qi Q, Ling Y, Zhu M, et al. Promoter region methylation and loss of protein expression of PTEN and significance in cervical cancer. *Biomed Rep.* 2014;2(5):653–658. doi:10.3892/br.2014.298
- 44. Jiang X, Li H. Overexpression of LRIG1 regulates PTEN via MAPK/ MEK signaling pathway in esophageal squamous cell carcinoma. *Exp Ther Med.* 2016;12(4):2045–2052. doi:10.3892/etm.2016.3606
- 45. Bai L, Mao R, Wang J, et al. ERK1/2 product proliferation and inhibited apoptosis of human cervical or cer cells a pregulated the expression of c-Fos and c-Jun protein *Med Oncol.* 15;32(3):57. doi:10.1007/s12032-015-0490-5
- Peng B, Zhu H, Leung PC. Chadotroph pleasing formone regulates human trophoblastic cell invasion bia. WIST-induced N-cadherin expression. *Join End Colol Metab*, 2015;100(1):E19–29. doi:10.1210/jc.2014-10.2
- 47. Chetram MA, Hielen CV. Part regulation of ERK1/2 signaling in cancer. J. Republic Signal Transluce Res. 2012;32(4):190–195. doi:10.310200799032012.695798
- 48. Ueda T, Volinia S, Ou pura H, et al. Relation between microRNA expression and prognosis of gastric cancer: a microRNA expression analysis. *Lancet Oncol.* 2010;11 :136–146. doi:0.1016/S1470-2045(09)70343-2
- 49. Lerducci MG, A elli D, Picchio MC, et al. MicroRNA profiling receives that mile 1, miR486 and miR-214 are upregulated and involution of an exervival in Sezary syndrome. *Cell Death Dis.* 2011;2:e151. doi:10.1038/cddis.2011.32

International Journal of Nanomedicine

Dovepress

Publish your work in this journal

The International Journal of Nanomedicine is an international, peerreviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch[®], Current Contents[®]/Clinical Medicine, Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-nanomedicine-journal