## ORIGINAL RESEARCH Long Non-Coding RNA AGAP2-AS1/miR-628-5p/ PTN Axis Modulates Proliferation, Migration, Invasion, and Apoptosis of Glioma Cells

This article was published in the following Dove Press journal: Cancer Management and Research

Yang Yan<sup>1,\*</sup> Yiping Wang<sup>1,\*</sup> Yuxia Liu<sup>2</sup> Tao Chen<sup>3</sup> Yaoli Zhu<sup>4</sup> Huiging Li<sup>5</sup> Fangen Kong <sup>1</sup>Department of Neurosurgery, The Fifth

Affiliated Hospital Sun Yat-Sen University, Zhuhai, Guangdong, People's Republic of China; <sup>2</sup>Department of Gastrointestinal Surgery, The Fifth Affiliated Hospital Sun Yat-Sen University, Zhuhai, Guangdong, People's Republic of China; <sup>3</sup>Department of Spine Surgery, The Fifth Affiliated Hospital Sun Yat-Sen University, Zhuhai, Guangdong, People's Republic of China; <sup>4</sup>Department of Critical Care Medicine, The Fifth Affiliated Hospital Sun Yat-S University, Zhuhai, Guangdong, Peg Republic of China; <sup>5</sup>Department Neurology, The Fifth Affiliated losdital Sun Yat-Sen University, Zh olic of Cha Guangdong, People's Re

\*These authors contribu ally to this work

Correspondence: Fangen Kong Email uyangleyanwlpty@163.com



seen reported h olved in a variety Purpose: Long non-coding RNAs (lncRNAs) have of cancers, including glioma. However, the exact the and inderlying mechanism of lncRNA e not yet een fully elucidated. AGAP2 antisense RNA 1 (AGAP2-AS1) sliom. roR<sup>2</sup> A-628-5p (miR-628-5p) and Methods: The expression levels of P2-AS1, h pleiotrophin (PTN) were measured by quantitive real-time polymerase chain reaction (qRT-PCR). Cell proliferation, apoptosis, migration and invasion were detected by Cell Counting Kit-8 (CCK-8) assay, flow g ometry, transwell as y, respectively. Western blot assay was used to detect the protein vel of PTN. The interaction between miR-628-5p and AGAP2y bioinformates software and confirmed by the dual-luciferase AS1 or PTN was predicted reporter and RNA Immunop. ipitation RIP) assays. Murine xenograft model was established to confirm the b of AGALZ-AS1 in glioma progression in vivo.

S1 ey Results: AGAP2was upregulated in glioma tissues and cells. Knockdown of AGAPAAS1 inh d the proliferation, migration and invasion, but facilitated apoptosis s. Morever, AGAP2-AS1 could directly bind to miR-628-5p and its over-Jma c versed the anti-tumor effect of miR-628-5p restoration on the progression of ression is. In addition, miR-628-5p directly targeted PTN and its inhibition abolished the glio effect of PTN knockdown on the progression of glioma cells. Furthermore, inhibite functioned as a competing endogenous RNA (ceRNA) by sponging miR-AGAP2-A 8-5p to modulate PTN expression. Besides, AGAP2-AS1 depletion reduced tumor growth regulating miR-628-5p and downregulating PTN.

Conclusion: AGAP2-AS1 knockdown suppressed cell proliferation, migration and invasion but promoted cell apoptosis in glioma cells by regulating miR-628-5p/PTN axis, providing novel avenues for treatment of glioma.

Keywords: glioma, AGAP2-AS1, miR-628-5p, PTN, cell progression

#### Introduction

Glioma, a kind of malignant brain tumor, is one of the most deadly cancers in adults.<sup>1</sup> In spite of significant advance has been made in the diagnosis and treatment of glioma, the prognosis is still dismal and the median survival time for high-grade glioma patients is only 10-15 months.<sup>2-4</sup> Hence, elucidating the pathogenic mechanisms at the molecular level and searching effective therapeutic strategies are critical for the treatment of glioma.

Long non-coding RNAs (lncRNAs), more than 200 nucleotides, play essential roles in tumorigenesis in a variety of cancers and without protein-coding ability.<sup>5-7</sup>

CO 0 S C2020 Yan et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

It has been confirmed that the aberrant expression of IncRNAs is tightly associated with diverse biological processes, such as cell differentiation, cell growth, apoptosis, cell cycle, drug resistance, metastasis, and epithelialmesenchymal transition.<sup>8,9</sup> LncRNAs are considered to be critical regulators of tumorigenesis and cancer progression, and more and more lncRNAs have been identified as oncogenes or tumor suppressors.<sup>10</sup> LncRNA AGAP2 antisense RNA 1 (AGAP2-AS1), an antisense lncRNA located at 12q14.1, has been identified as an oncogene in multiple cancers, such as breast cancer,<sup>11</sup> non-small cell lung cancer,<sup>12</sup> and gastric cancer.<sup>13</sup> Additionally, Zheng et al pointed out that AGAP2-AS1 abundance was enhanced in glioma tissues and cells.<sup>14</sup> However, the involvement of AGAP2-AS1 in glioma pathogenesis remains to be explored.

Recent reports have proven that lncRNAs serve as competing endogenous RNA (ceRNA), which interact with microRNAs (miRNAs) and modulate the expression of miRNA target genes.<sup>15</sup> MiR-628-5p is found to be a potential biomarker in several diseases, such as ovarian cancer,<sup>16</sup> prostate cancer,<sup>17</sup> and acute myeloid leukemia.<sup>18</sup> Additionally, miR-628-5p has been reported to be expressed at a low level in the glioma tissues and cell lines.<sup>19</sup> Howev the interaction between AGAP2-AS1 and miR-628-5p has no been reported. Pleiotrophin (PTN), located on chred me 7, is expressed in the brain during embryogenesis 1d com only upregulated in many cancers, include gli Bioinformatics analysis exhibits the ential ding sites between miR-628-5p and AGAP2... or PTN. us, we supposed that AGAP2-AS1 might regulate the progression of glioma through acting a sponge of iR-628-5p to regulate PTN expression in the content study, the abundance of AGAP2-AS1 was menure in gliopra tissues and cells. al functions of GAP2-AS1 in glioma Moreover, the big optosis migration and invasion were further cell growth, investigated. urther plored the ceRNA regulatory network of AO 2-AS1/miR-628-5p/PTN in glioma cells. Collectively, this report showed a new ceRNA regulatory network in glioma.

## Methods

#### Patients' Specimens

Nontumorous brain tissues (NBTs) and glioma tissues were provided by The Fifth Affiliated Hospital Sun Yat-Sen University in this study. NBTs were collected from 14 patients who underwent a partial brain resection because of traumatic brain injury or intracerebral hemorrhage. Glioma clinical tissues were collected from 55 patients with glioma requiring surgical resection. In these tissues, 30 cases were low-grade glioma tissues (LGG; grade II) and 25 cases were high-grade glioma tissues (HGG; grade III and IV). These participants did not receive chemotherapy, radiotherapy or others' therapy before surgery. This study was approved by the Research Ethics Committee of The Fifth Affiliated Hospital Sun Yat-Sen University. All participants signed the informed consent. Fresh samples were promptly frozen in liquid nitrogen and then stored at  $-80^{\circ}$ C before use.

## Cell Culture and Transfection

Two glioma cell lines (2251 and 2N229), were purchased from COBIOER (Nunjne, Coma), and two other glioma cell lines (A172 and SHC-1) were obtained from Cell Bank of the Charge Academ of Sciences (Shanghai, China). Normal hum eastrocytes (NHAs) were brought from conza (Basel, Switerland). These cells were grown in Dulbecco's emodified eagle medium (DMEM; Invegen, Carlstad, CA, USA) containing 10% fetal boving erum (CBS), penicillin (100 U/mL) and streptolingin (100  $\mu$ g/mL) (Gibco, Carlsbad, CA, USA) in a dame ed atmosphere with 5% CO<sub>2</sub> at 37°C.

The small interfering RNA against-AGAP2-AS1 or PTN i-AGAP2-AS1 or si-PTN) and their control (si-control), miR-628-5p mimic or inhibitor (miR-628-5p or miR-628-5p inhibitor) and their control (miR-control or inhibitor-control), and AGAP2-AS1 overexpression vector (AGAP2-AS1) were obtained from GenePharma (Shanghai, China). Lentiviral small hairpin RNA targeting AGAP2-AS1 (sh-AGAP2-AS1) and its control (sh-control) were provided by Ribo Bio Corporation (Guangzhou, China). SHG44 and U251 cells were transfected with oligonucleotides or vectors using the Lipofectamine 3000 reagent (Invitrogen).

## Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

Total RNA from tissue and cell lines was extracted using TRIzol reagent (Invitrogen). The first strand of complementary DNA (cDNA) for AGAP2-AS1 and PTN or miR-628-5p was synthesized with TaqMan Reverse Transcription Kit or TaqMan microRNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA, USA), respectively. Real-time PCR was performed on a 7900HT Fast RT-PCR instrument using SYBR Green PCR Master Mix (Applied Biosystems) and following the primers: AGAP2-AS1 (Forward, 5'-TACCTTGACCTTGCTGCTGCTCC-3'; Reverse, 5'-TGTC CCTTAATGACCCCATCC-3'), miR-628-5p (Forward, 5'-GCTGACATATTTACTAGAG-3'; Reverse, 5'-GAACATG TCTGCGTATCTC-3') PTN (Forward, 5'-AGAAGCAATT TGGCGCGGA-3'; Reverse, 5'-TGCACCACCAACTGCTT AGC-3), GAPDH (Forward, 5'-CGCTCTCTGCTCCTCC TGTTC-3'; Reverse, 5'- ATCCGTTGACTCCGACCT TCAC-3'), U6 (Forward, 5'-CTCGCTTCGGCAGCA CAT'; Reverse, 5'-CGCAAATTCGTGAAGCGTTCCA3'). Analysis of relative AGAP2-AS1, PTN or miR-628-5p expression was evaluated with  $2^{-\Delta\Delta Ct}$  method with normalization to GAPDH or U6, respectively. All experiments are performed three times.

#### **Cell Proliferation Assay**

Cell Counting Kit-8 (CCK-8; Beyotime, Shanghai, China) was used to determine cell proliferation capacity. In brief, SHG44 and U251 cells growing at an exponential rate were seeded in 96-well plates (4000 cells/well). CCK-8 (10  $\mu$ L) solution was added to the culture medium at 0 h, 24 h, 48 h and 72 h after transfection, then incubated for an additional 3 h. After that, the absorbance was evaluated at 450 nm under a microplate reader (Bio-Rad, He CA, USA). All experiments are performed three time

#### Cell Apoptosis Assay

ropidiun Annexin V-fluorescein isothiocyanate (FITC) iodide (PI) apoptosis detection kit couble ning kit (BD Biosciences, Franklin Lakes, N SA) was us yed to detect cell apoptosis. In brief, SHC+ and \$51 cells growing at an exponential rate were secred into 6 cm thes and transfected AP2-AS1, miR-control, miR-628-5p, with si-control, si-A miR-628-5p + AGA 2-AS, si-PTN, or si-PTN + miR-628-5p inhibitor After h sfection or 48 h, SHG44 and s<sup>1</sup> d, and re-suspended in bind-U251cells rere concected, ing bury (500 1) followed by staining with Annexin V-FITC at for 15 min. Finally, cell apoptosis was analyzed by a flor cytometer (BD Biosciences). Non-apoptotic cells were defined as Annexin V-FITC<sup>-</sup>/PI<sup>-</sup>, early apoptotic cells were defined as Annexin V-FITC<sup>+</sup>/PI<sup>-</sup>, late apoptotic/ necrotic cells were defined as Annexin V-FITC<sup>+</sup>/PI<sup>+</sup>, and dead cells were defined as Annexin V-FITC<sup>-</sup>/PI<sup>+</sup>. All experiments are performed three times.

### Transwell Assay

For cell invasion assay, the transfected SHG44 and U251 cells were suspended in serum-free DMEM (100  $\mu L)$  and

placed in the upper chamber of transwell chamber (8  $\mu$ m pore size, Corning Inc., Corning, NY, USA) containing a Matrigel-coated (BD, San Jose, CA, USA), while cell migration assay did not coat with Matrigel. DMEM with 10% FBS (500  $\mu$ L) was added to the lower chamber. Following incubation for 48 h, non-migrated and noninvaded cells were scrubbed by a cotton swab; migrated and invaded cells were fixed with 4% paraformaldehyde and stained with 0.5% crystal violet, and counted by an inverted microscope (Olympus, Tokyo, Japan). All experiments are performed three times.

## Dual-Luciferase Reputter Assa

The wild-type (WT) or outant (AUT) of AGAP2-AS1 or 3'-UTR of PTN containing the public binding sites of miR-628-5p was softhesized and inserted into the pGL3control lucificase reporter vectors (Promega, Madison, WI, USA9 camely whether plasmids (WT-AGAP2-AS1, 3 UTR-WAPTN) or mutant-type plasmids (MUT-AGAP2-AS1, 3'UTA-MUT-PTN). WT or MUT plasmid ansiently co-transfected with miR-628-5p mimic or miRontrol into HG44 and U251 cells for 48 h. Finally, heiferase activity was examined using the Dual-Lucification Assay System (Promega) and normalized to heilla luciferase activity. All experiments are performed three times.

#### RNA Immunoprecipitation (RIP) Assay

EZ-Magna RIP Kit (Millipore) was applied to conduct RIP assay. In brief, SHG44 and U251 cells were lysed by a complete RIP lysis buffer. After that, whole-cell extract (100  $\mu$ L) was incubated by RIP buffer containing magnetic beads coated with human anti-argonaute2 (Ago2) or anti-IgG antibody (as control). Next, samples were digested using the proteinase K. Lastly, immunoprecipitated RNA was isolated, and the expression levels of AGAP2-AS1 and miR-628-5p were examined by qRT-PCR.

#### Western Blot Assay

Total protein from tissues and cells (SHG44 and U251) was extracted using RIPA lysis buffer supplemented with protease (Beyotime) on ice. The protein concentration of the lysates was assessed by a bicinchoninic acid protein assay kit (Tanon, Shanghai, China). Protein from each sample (about 40  $\mu$ g) was subjected to 12% SDS-PAGE (80 V for 30 min and 120 V for 40 min). Subsequently, the gels containing the goal protein were transferred onto a polyvinylidene fluoride (PVDF) membrane (0.2  $\mu$ m, Beyotime). After that, the membranes containing the goal protein were blocked using 5% non-fat milk in Tris-Buffered Saline Tween-20 (TBST), followed by incubation overnight at 4°C with the specific antibodies against PTN (1:2000, ab223675, Abcam, Cambridge, MA, USA) and GAPDH (1:2000, ab37168, Abcam). After washing with TBST, secondary antibody was added to the blots for 2 h. Finally, the blots were detected with enhanced chemiluminescence reagent (Pierce, Rockford, IL, USA). The protein level of PTN was normalized by GAPDH. The intensity of the bands was analyzed by ImageJ software. All experiments are performed three times.

### Tumor Xenograft Model

The animal experiments were performed in accordance with the Guidelines for Care and Use of Laboratory Animals of "National Institutes of Health" and approved by the Animal Experimental Ethics Committee of The Fifth Affiliated Hospital Sun Yat-Sen University. BALB/c nude mice (fiveweek-old, male) were provided by Shanghai Experimental Animal Center (Shanghai, China). The mice were randomly grouped on the basis of body weight on the day of tumor cell inoculation using the stratified randomization method (n=7/ group). U251 cells transfected with sh-NC or sh-AGAP2-AS1 were subcutaneously injected into the dorsal flanks nude mice. We examined the xenograft tumor volume ever week and calculated by the following formula eth × width $^{2}/2$ . After 5 weeks, mice were sacrificed 1 d the t nors were weighed and collected for further analysis

### Statistical Analysis

All experiments were repeated at least 3 topes in this study. Data were expressed as the mean ±standard excition (SD). All statistical analyses were performed using GraphPad version 6.0 software (Graph and software inc., La Jolla, CA,



USA). Student's *t*-test was used to analyze differences between two groups, and the significance of differences among multiple groups was analyzed by one-way analysis of variance (ANOVA) followed by Turkey's post hoc test. P<0.05 was regarded as statistically significant.

## Results

## AGAP2-ASI Expression Was Increased in Glioma Tissues and Cells

To explore the potential role of AGAP2-AS1 in glioma, the qRT-PCR was used to analyze the exp ssion GAP2-AS1 in glioma tissues and cells. The results showed the AGAP2-AS1 expression was increased in LGC sases and GG cases compared with that in NBT Moreover, **N** S constant higher AGAP2-AS1 expression than LGG case. (Figure 1A). er level AGAP2-AS1 was Furthermore, the dive . observed in gline a cell lines. A17, LN229, SHG44 and U251) rather than 1 NHA cells, especially in SHG44 and U251 celle (Figure 1B) Thus, we chose SHG44 and U251 or further investigation. Furthermore, the relationship cells between the relative expression of AGAP2-AS1 and clinicopathengic characeristics of glioma patients was analyzed. As mentary Table 1, the high expression level of shown h P2-AS1 was significantly associated with tumor size and AO grade, but not with age, gender lymphatic metastasis, and differentiation grade. All these results indicated that GAP2-AS1 might play an oncogenic role in glioma.

## Knockdown of AGAP2-AS1 Suppressed Proliferation, Migration and Invasion, but Facilitated Apoptosis in Glioma Cells

To analyze the functional roles of AGAP2-AS1 in glioma, we interfered with endogenous AGAP2-AS1 expression in



Figure I AGAP2-AS1 expression was enhanced in glioma tissues and cells. (A) The level of AGAP2-AS1 was measured in NBTs, LGG and HGG cases by qRT-PCR. (B) The expression of AGAP2-AS1 was determined in glioma cell lines (A172, LN229, SHG44, and U251) and NHA by qRT-PCR. \*P<0.05.

SHG44 and U251 cells via transfection with specific siRNA. The transfection efficiency was evaluated using qRT-PCR. As shown in Figure 2A and B, the expression of AGAP2-AS1 was dramatically decreased in SHG44 and U251 cells transfected with si-AGAP2-AS1 compared with those cells transfected with si-control or NC group (untransfected cells). CCK-8 assay suggested that knockdown of AGAP2-AS1 markedly impaired SHG44 and U251 cell growth (Figure 2C and D). Moreover, the change in the cell apoptosis after silencing AGAP2-AS1 was analyzed using flow cytometry analysis. As presented in Figure 2E and F, silenced AGAP2-AS1 induced the apoptosis of SHG44 and U251 cells. Transwell assay demonstrated that the depletion of AGAP2-AS led to a significant suppression of migration and invasion abilities in SHG44 and U251 cells (Figure 2G and H). All these findings proved that the AGAP2-AS1 was a tumor promoter in glioma progression in vitro.

## AGAP2-AS1 Served as a Molecular Sponge of miR-628-5p in Glioma Cells

It is generally accepted that lncRNAs could function as sponges or decoys of miRNAs to modulate target mRNAs expression. Thus, we intended to explore whether AGAP2-AS1 could modulate gene expression through sponging miRNAs. The starBase v3.0 was employed to search AGAP2-AS1 targets, miR-628-5p was found to potentially bind to AGAP2-AS1 (Figure 3A). Then, a dual-luciferase reporter system was used to determine whether miR-628-5p could directly bind to AGAP2-AS sults showed that overexpression of miR-628-5 remarkably decreased the luciferase activity of WT-AGA, AS1, whe as no obvious change on the luciferar activity of UT-/ AP2-AS1 was found in SHG44 d U251 ells (F e 3B and C). To determine whether ΓA/ -AS1 and miR-628-5p were in induced encine complex (RISC) complex, the same RM



Figure 2 Knockdown of AGAP2-AS1 inhibited glioma cell progression. SHG44 and U251 cells were divided into 3 groups: NC (untransfected cells), si-control and si-AGAP2-AS1. (**A** and **B**) AGAP2-AS1 level was determined by qRT-PCR. (**C** and **D**) Cell proliferation was assessed by CCK-8 assay. (**E** and **F**) Cell apoptosis was analyzed using flow cytometry. The proportion of non-apoptotic cells (left lower quadrant: Annexin V-FITC<sup>-</sup>/PI<sup>-</sup>), early apoptotic cells (right lower quadrant: Annexin V-FITC<sup>+</sup>/PI<sup>-</sup>), late apoptotic/necrotic cells (right upper quadrant: Annexin V-FITC<sup>+</sup>/PI<sup>+</sup>) and dead cells (left upper quadrant: Annexin V-FITC<sup>-</sup>/PI<sup>+</sup>). (**G** and **H**) Transwell was employed to examine cell migration and invasion abilities. \*P<0.05.



Figure 3 AGAP2-AS1 interacted with miR-628-5p. (A) The potential binding sites between neuciferase activity was measured in SHG44 and U251 cells co-transfected with WT- AGAP2-A enrichment of AGAP2-AS1 and miR-628-5p was measured by RIP assay in Subscript U25 transfected with miR-control, miR-628-5p or miR-628-5p + AGAP2-AS1. (F) The number of evaluate cell proliferation. (I) Flow cytometry analysis was used to measure cell to ptosin attert transwell assay. \*P<0.05.

RIP experiments were performed. The rest sho AGAP2-AS1 and miR-628-5p were greatly riched in Ago2-containing beads compared IgG conti group A (Figure 3D and E), suggesting the endo nous interaction between AGAP2-AS1 and AR-628-5p. Be. les, the upregulation of miR-628-5p sulted just significant promotion of miR-628-5p expression, overexpression of AGAP2--hi<sup>1</sup> on ( ure 3F). Moreover, AS1 attenuated expre. miR-62 ently blocked the prolifrestoration or 5p prov tion in but induced apoptosis of eration, mig cells, while these effects were abolished SHG44 and U<sub>2</sub> by upregulating AC P2-AS1 (Figure 3G–K). These results disclosed that AGAP2-AS1 might exert biological function via serving as a sponge of miR-628-5p in glioma cells.

# PTN Was a Direct Target of miR-628-5p in Glioma Cells

To elucidate the underlying mechanism of miR-628-5p exerting its functional effects in glioma, the downstream targets of miR-628-5p were predicted by TargetScan. As

een n. 628-5p and AC P2-ASI were predicted by starBase v3.0. (**B** and **C**) The GAP2-AL or MUT-AL P2-ASI and miR-628-5p or miR-control. (**D** and **E**) The of U251 central action with Ago2 or IgG. (**F**–**K**) SHG44 and U251 cells were n. TopiR-628-5p was detected by qRT-PCR. (**G** and **H**) CCK-8 assay was used to osign attention **K**) The number of migrated and invaded cells was determined by

hown in Figure 4A, putative binding sites between PTN and miR-628-5p, suggesting that PTN might be a target of miR-628-5p. To validate this prediction, we constructed the 3'UTR-WT-PTN or 3'UTR-MUT-PTN luciferase reporter vector and introduced them into SHG44 and U251 cells. Dual-luciferase reporter assay suggested that the luciferase activity was strongly suppressed in SHG44 and U251 cells co-transfected with 3'UTR-WT-PTN and miR-628-5p, while it was not changed in 3'UTR-MUT-PTN group (Figure 4B). In addition, the effect of miR-628-5p on the levels of PTN mRNA and protein in U251 and SHG44 cells was explored. The data showed that the mRNA and protein levels of PTN were decreased after transfection with miR-628-5p mimic (Figure 4C and D), suggesting PTN was negatively regulated by miR-628-5p. Moreover, PTN mRNA and protein levels were notably decreased in SHG44 and U251 cells transfected with si-PTN, while the effect was abated by co-transfection with miR-628-5p inhibitor (Figure 4E and F). Besides, downregulation of miR-628-5p partly reversed anti-proliferation, anti-migration,



Figure 4 MiR-628-5p exerted its anti-tumor role by regulating PTN. (A) The putative binding sites be en miR-6 o and PTN ere predicted by TargetScan. (B) TR-WT-PTN or 3'UTR-MUT-PTN). and Relative luciferase activity was determined in both SHG44 and U251 (cells co-transfected with miR-Sp or miR-con K-628-5p by qRT-PCR and Western blot (C and D) The mRNA and protein expression of PTN were analyzed in SHG44 and U251 cells trap .tea h miR-control analyses, respectively. (E–J) SHG44 and U251 cells were transfected with si-control, si-PTN, si-PTN, miR-628 inhibitor. (**E** and **F**) The mRNA and protein expression of PTN were examined by qRT-PCR and Western blot assays, respectively. (G) Cell proliferation was measured by 8 assay. (H) Cell apoptosis rate was assessed by flow cytometry analysis. (I and J) Transwell assay was used to analyze cell migration and invasi cies. \*P<0.05.

anti-invasion, and pro-apoptosis effects caused by PTN knockdown (Figure 4G–J). Taken together, these date dicated that miR-628-5p exerted its anti-tumor functio via regulating PTN expression in glioma cells.

## PTN Was Regulated by ACAP2-11 hand miR-628-5p

2-AS1 To determine whether fu tioned as a ceRNA of miR-628-5p to regu e PTN expression, SHG44 and U251 celd were transfected with si-control, si-AGAP2-AS1, si GAP2 S1 + inhibitor-control, or si-AGAP2-AS1 + mix 22 p inhibitor. The qRT-PCR and wed at transfection with si-Western bl nys 5 S1 red the NA and protein expression AGAP2 of PTN SF 251 cells, while the decreased expression PTN was increased again by co-transfection with miR-628 inhibitor (Figure 5A–C). Collectively, data revealed that AGAP2-AS1 functioned our as a ceRNA of miR-628-5p to modulate PTN expression.

## Knockdown of AGAP2-AS1 Inhibited Glioma Tumorigenesis in vivo

To confirm the effect of AGAP2-AS1 on glioma in vivo, U251 cells stably transfected with sh-NC or sh-AGAP2-AS1 were subcutaneously injected into the flanks of nude Lice. Knock own of AGAP2-AS1 greatly inhibited tumor volume weight in xenograft model (Figure 6A and B). Idditionally, qRT-PCR analysis presented that knockdown of AGAP2-AS1 led to a significant reduction of AGAP2-AS1 expression, while an increase of miR-628-5p level in excised tumor masses (Figure 6C and D). Furthermore, qRT-PCR and Western blot assays demonstrated that deficiency of AGAP2-AS1 evidently reduced the mRNA and protein expression of PTN (Figure 6E and F). Thus, it was concluded that AGAP2-AS1 inhibition reduced tumor growth by regulating miR-628-5p and PTN in vivo.

## Discussion

Glioma is the most common and lethal brain tumor with an extremely poor prognosis. The aim of this research was to find a novel underlying mechanism in tumorigenesis and progression of glioma. Recently, increasing lncRNAs have been identified to be involved in the occurrence and development of different tumors, including glioma.<sup>22,23</sup> Here, we focused on explaining the biological functions and the potential mechanism of AGAP2-AS1 in the occurrence and progression of glioma.

AGAP2-AS1 has been indicated to be dysregulated in many diseases and participated in multiple cell behaviors, such as proliferation, apoptosis, metastasis, and differentiation.<sup>12,24</sup> Notably, AGAP2-AS1 was confirmed to



Figure 5 PTN expression was regulated by miR-628-5p and AGAP2-AS1 in glioma cells. (A–C) The mRNA and protein expression levels of PTN in SHG44 and U251 transfected with si-control, si-AGAP2-AS1, si-AGAP2-AS1 + inhibitor-control, or si-AGAP2-AS1 + miR-628-5p inhibitor were detected by qRT-PCR and Western blot, respectively. \*P<0.05.





promote cell proliferation in glice a cell via sponging miR-15a/b-5p to regulate HDGF<sup>1/2</sup> in addition, n et al proved that AGAP2-AS1 depletig mited cell growth and metastasis, but accelerated apoptosition gliobestoma (GBM).<sup>25</sup> Moreover, AP2-A level was enhanced Luo et al demonstrated that in GBM and associ d th poor prognosis, and sitive repressed growth and invasion but AGAP2-A depleti in GBM cells through silencing TFPI2.<sup>26</sup> facilitated apo ulatory mechanisms of AGAP2-AS1 in Nevertheless, the glioma are still not well understood. Firstly, we found that AGAP2-AS1 abundance was enhanced in HGG cases compared with NBTs or LGG cases, which was in line with previous studies. Subsequently, we characterized the functional roles of AGAP2-AS1 in glioma by using loss-of-function experiments. The resulted disclosed that AGAP2-AS1 inhibition suppressed the glioma cell growth and metastasis, and contributed to apoptosis. Similarly, in vivo experiments uncovered that AGAP2-AS1 depletion repressed tumor growth. Our findings combined with previous studies demonstrated that AGAP2-AS1 played an oncogenic role in the progression of glioma.

It is widely accepted that lncRNAs commonly modulate genes expression via sharing the same miRNAs with their downstream mRNAs.<sup>27</sup> In this research, starBase 3.0 was employed to search for AGAP2-AS1 targets, and miR-628-5p was found to potentially bind to AGAP2-AS1. Then, dual-luciferase reporter assay suggested that AGAP2-AS1 directly targeted miR-628-5p. MiR-628-5p has been proven to be a tumor suppressor or promoter in different tumors. For example, miR-628-5p has been demonstrated to block the progression of epithelial ovarian cancer via inhibiting FGFR2.<sup>16</sup> Additionally, Wang et al demonstrated that miR-628-5p level was elevated in osteosarcoma and its knockdown blocked the proliferation and metastasis of osteosarcoma cells through targeting IFI44L.<sup>28</sup> Besides, Xie et al found that miR-628-5p repressed glioma cell growth via targeting DDX59.<sup>19</sup> In the current research, we found that restoration of miR-628-5p obviously limited the cell growth, migration and invasion but facilitated apoptosis in glioma cells, while these effects were abolished by upre-gulating AGAP2-AS1. Our findings suggested that miR-628-5p acted as a tumor suppressor in glioma. To clarify how miR-628-5p affected the progression of glioma, potential target was predicted by TargetScan. PTN containing binding sites of miR-628-5p was selected as the target for further analysis.

The emerging evidence has suggested that PTN was abnormally expressed in various cancers.<sup>29,30</sup> Previous studies demonstrated that PTN expression was enhanced in glioma and positively correlated with poor survival, and PTN accelerated migration and proliferation in glioma cells, while its knockdown inhibited tumor growth in vitro and in vivo.<sup>21,31,32</sup> Consistent with these findings, we also proved that downregulation of PTN limited cell proliferation and metastasis but facilitated apoptosis in glioma cells, whereas co-transfection of miR-628-5p abolished these effects. Moreover, the results displayed that PTN expression was positively regulated by AGAP2-AS1 and negatively regulated by miR-628-5p. In addition, deficiency of AGAP2-AS1 resulted in significant promotion of mit 5p expression and reduction of PTN level in tissues om xenograft mice. Therefore, these findings disclosed AGAP2-AS1 functioned as a ceRNA v spon ng mi 628-5p to modulate PTN expression in glioma

#### Conclusion

In conclusion, these findings should that ACAP2-AS1 knockdown blocked the progression of glioma through functioning as ceR1A of piR-628-5p to regulate PTN expression. Hence, the arGAP2-AS1/miR-628-5p/PTN axis might be promising therapidic target for the treatment of groma.

#### Ackno vedgments

The authors puld like to thank the participants in this study.

#### **Author Contributions**

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agreed to be accountable for all aspects of the work.

#### Funding

This project was supported by the Science and Technology Planning Project Foundation of Zhuhai in Guangdong Province (grant no. 20171009E030067); Science and Technology Planning Project Foundation of Zhuhai in Guangdong Province (grant no. 20191210E030073).

#### Disclosure

The authors declare that they have no conflicts of interest in this work.

## References

- Goodenberger ML, Jenkus RB. Geness of a de glioma. *Cancer Genet*. 2012;205(12):1:3-621. doi:10.1015/j.c.eergen.2012.10.009
  Laws ER, Parney & Huang V, et al. Survival following surgery and
- prognostic factors to recently diagnered malignant glioma: data from the stoma outpres project. J Neurosurg. 2003;99 (3):467-1. Noi:10.3171/jn. 001-9.3.0467
- 3. Davis dE. O. blastoma: over dew of disease and treatment. *Clin J Oncol Nurs*. 20 20(5):S2–S8. doi:10.1188/11.CJON.291-297
- Mar JP, Dolece TA, Horbinski C, et al. Epidemiologic and molecular prognostic review of glioblastoma. *Cancer Epidemiol Biomarkers* Prev. 2014;23(10):1985–1996. doi:10.1158/1055-9965. EPI-14-027:
- Li J, Xua, Z, Liu C, Goodenberger ML, Jenkins RB. Long codif RNAs and complex human diseases. *Int J Mol Sci.* 2013;14(9):18790–18808. doi:10.3390/ijms140918790
- Tibb EA, Brown CJ, Lam WL. The functional role of long non-coding RNA in human carcinomas. *Mol Cancer*. 2011;10(1):38. doi:10.1186/1476-4598-10-38
- Zhang X, Sun S, Pu JKS, et al. Long non-coding RNA expression profiles predict clinical phenotypes in glioma. *Neurobiol Dis*. 2012;48(1):1–8. doi:10.1016/j.nbd.2012.06.004
- Wu Q, Xiang S, Ma J, et al. Long non-coding RNA CASC 15 regulates gastric cancer cell proliferation, migration and epithelial mesenchymal transition by targeting CDKN 1A and ZEB 1. *Mol Oncol.* 2018;12(6):799–813. doi:10.1002/1878-0261.12187
- Huang Y, Xiang B, Liu Y, et al. LncRNA CDKN2B-AS1 promotes tumor growth and metastasis of human hepatocellular carcinoma by targeting let-7c-5p/NAP1L1 axis. *Cancer Lett.* 2018;437:56–66. doi:10.1016/j.canlet.2018.08.024
- Tsai M-C, Spitale RC, Chang HY. Long intergenic noncoding RNAs: new links in cancer progression. *Cancer Res.* 2011;71(1):3–7. doi:10.1158/0008-5472.CAN-10-2483
- 11. Dong H, Wang W, Mo S, et al. SP1-induced lncRNA AGAP2-AS1 expression promotes chemoresistance of breast cancer by epigenetic regulation of MyD88. *J Exp Clin Cancer Res.* 2018;37(1):202. doi:10.1186/s13046-018-0875-3
- 12. Li W, Sun M, Zang C, et al. Upregulated long non-coding RNA AGAP2-AS1 represses LATS2 and KLF2 expression through interacting with EZH2 and LSD1 in non-small-cell lung cancer cells. *Cell Death Dis.* 2016;7(5):e2225. doi:10.1038/cddis.2016.126
- 13. Qi F, Liu X, Wu H, et al. Long noncoding AGAP2-AS1 is activated by SP1 and promotes cell proliferation and invasion in gastric cancer. *J Hematol Oncol.* 2017;10(1):48. doi:10.1186/s13045-017-0420-4
- 14. Zheng Y, Lu S, Xu Y, et al. Long non-coding RNA AGAP2-AS1 promotes the proliferation of glioma cells by sponging miR-15a/b-5p to upregulate the expression of HDGF and activating Wnt/β-catenin signaling pathway. *Int J Biol Macromol.* 2019;128:521–530. doi:10.1016/j.ijbiomac.2019.01.121

6067

- Tay Y, Rinn J, Pandolfi PP. The multilayered complexity of ceRNA crosstalk and competition. *Nature*. 2014;505(7483):344–352. doi:10. 1038/nature12986
- Li M, Qian Z, Ma X, et al. MiR-628-5p decreases the tumorigenicity of epithelial ovarian cancer cells by targeting at FGFR2. *Biochem Biophys Res Commun.* 2018;495(2):2085–2091. doi:10.1016/j. bbrc.2017.12.049
- Srivastava A, Goldberger H, Dimtchev A, et al. Circulatory miR-628-5p is downregulated in prostate cancer patients. *Tumor Biol.* 2014;35 (5):4867–4873. doi:10.1007/s13277-014-1638-1
- Favreau AJ, Sathyanarayana P. miR-590-5p, miR-219-5p, miR-15b and miR-628-5p are commonly regulated by IL-3, GM-CSF and G-CSF in acute myeloid leukemia. *Leuk Res.* 2012;36(3):334–341. doi:10.1016/j.leukres.2011.09.027
- Xie P, Wang Y, Liao Y, et al. MicroRNA-628-5p inhibits cell proliferation in glioma by targeting DDX59. *J Cell Biochem*. 2019;120 (10):17293–17302. doi:10.1002/jcb.28991
- Kurtz A, Schulte AM, Wellstein A. Pleiotrophin and midkine in normal development and tumor biology. *Crit Rev Oncog.* 1995;6 (2):151–177.
- 21. Zhang L, Kundu S, Feenstra T, et al. Pleiotrophin promotes vascular abnormalization in gliomas and correlates with poor survival in patients with astrocytomas. *Sci Signal*. 2015;8(406):ra125–ra125. doi:10.1126/scisignal.aaa1690
- 22. Huarte M. The emerging role of lncRNAs in cancer. *Nat Med.* 2015;21(11):1253-1261. doi:10.1038/nm.3981
- 23. Li J, Zhang M, An G, et al. LncRNA TUG1 acts as a tumor suppressor in human glioma by promoting cell apoptosis. *Exp Biol Med.* 2016;241(6):644–649. doi:10.1177/1535370215622708
- 24. Liu Z, Wang Y, Wang L, et al. Long non-coding RNA AGAP2-AS1, functioning as a competitive endogenous RNA, upregulates ANXA11 expression by sponging miR-16-5p and promotes proliferation and metastasis in hepatocellular carcinoma. J Exp Clin Cancer P 2019;38(1):194. doi:10.1186/s13046-019-1188-x

- Tian Y, Zheng Y, Dong X. AGAP2-AS1 serves as an oncogenic lncRNA and prognostic biomarker in glioblastoma multiforme. *J Cell Biochem*. 2019;120(6):9056–9062. doi:10.1002/jcb.28180
- 26. Luo W, Li X, Song Z, et al. Long non-coding RNA AGAP2-AS1 exerts oncogenic properties in glioblastoma by epigenetically silencing TFPI2 through EZH2 and LSD1. *Aging (Albany NY)*. 2019;11 (11):3811–3823. doi:10.18632/aging.102018
- Yoon JH, Abdelmohsen K, Gorospe M. Functional interactions among microRNAs and long noncoding RNAs. *Semin Cell Dev Biol.* 2014;34:9–14. doi:10.1016/j.semcdb.2014.05.015
- Wang JY, Lu SB, Wang JQ. miR-628-5p promotes the growth and migration of osteosarcoma by targeting IFI44L. *Biochem Cell Biol*. 2019. doi:10.1139/bcb-2019-0001
- 29. Feng Z, Gao S, Wu Y, et al. Lung cancer cell migration is regulated via repressing growth factor PTN/RPTP to paling by menin. *Oncogene*. 2010;29(39):5416–5426. doi:10.1038/onr.1010.282
- Riegel AT, Wellstein A. The potent role of the henrin-binding growth factor pleiotrophin in breast can be *Breast Caner Res Treat*. 1994;31(2–3):309–314. doi:10.007/bf000.063
- Powers C, Aigner A, Stoice JE, et al. Pleiothenin chaling through anaplastic lymphoma kip je is rate-builting for guidastoma growth. *J Biol Chem.* 2003;27:10;11:23–14158 doi:10.1074/jbc.M112 354200
- 32. Ulbricht U, Echrich C, Fillbrah, R, et al. RNA interference targeting protein gross, phosphatase acceptor-type protein tyrosine phosphatase β suppreses glioblastoma growth in vitro and in vivo. J Neurophy. 2006;98(5), 197–1506. doi:10.1111/j.1471-4159.2006.

#### **Cancer Management and Research**

#### **Dove**press

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/cancer-management-and-research-journal