

EGCG Upregulates UCP₃ Levels to Protect MIN₆ Pancreatic Islet Cells from Interleukin-1 β -Induced Apoptosis

This article was published in the following Dove Press journal:
Drug Design, Development and Therapy

Xu Jia^{1,*}
Ziren Luo^{2,*}
Ying Gao³
Hua Liu³
Xinghai Liu³
Wenli Mai³ 
Hong Liu³
Qian Zheng³ 

¹Department of Pharmacy, Affiliated Hospital of North Sichuan Medical College, Nanchong 637000, People's Republic of China; ²School of Pharmacy, North Sichuan Medical College, Nanchong 637000, People's Republic of China; ³Department of Physiology, North Sichuan Medical College, Nanchong 637007, People's Republic of China

*These authors contributed equally to this work

Objective: The protective effects of epigallocatechin gallate (EGCG) on interleukin-1 β (IL-1 β)-induced apoptosis were investigated in murine MIN₆ pancreatic β -cells. The role of uncoupling protein-3 (UCP₃) signaling in this process was also explored.

Methods: After treatment with IL-1 β and EGCG, cells were collected and analyzed. Cell viability was measured using the CCK₈ assay and the function of β -cells was evaluated by analyzing insulin secretion. Detection of mitochondrial function in cells was performed by measuring mitochondrial membrane potential, the concentration of ATP and activity of ROS. Apoptosis was analyzed by Hoechst33258 staining and flow cytometry. Expression levels of UCP₃ were interrogated using immunohistochemistry, RT-PCR and Western blotting.

Results: Compared with the control group, IL-1 β treatment (20nM) for 24 h significantly decreased cell viability and insulin secretion, damaged mitochondrial function and increased ROS activity. Results also showed increased apoptosis and a decrease in UCP₃ expression levels ($p < 0.01$). However, treatment with low (1mM) or high (5mM) concentrations of EGCG significantly decreased IL-1 β -induced apoptosis ($p < 0.01$), restored mitochondrial function and subsequently increased UCP₃ levels in IL-1 β -induced β -cells ($p < 0.01$).

Conclusion: These results suggest that EGCG protects against IL-1 β -induced mitochondrial injury and apoptosis in β -cells through the up-regulation of UCP₃.

Keywords: EGCG, pancreatic β -cells, apoptosis, UCP₃

Introduction

Diabetes is a worldwide public health issue. According to the International Diabetes Federation, the number of type 2 diabetes (T2DM) cases in adults ranging from 18 to 99 years of age was 451 million. If this trend continues, this number will rise to 693 million in 2045.¹

Insulin resistance and increased β -cell apoptosis are the main mechanisms of pathogenesis behind T2DM.² High glucose levels, high fat levels and inflammation can result in apoptosis. Studies suggest that reactive oxygen species (ROS) and oxidative stress (OS) are the main culprits of diabetes.³ ROS is mainly produced by the mitochondria, which is generally considered as toxic byproducts of aerobic metabolism and includes superoxide anion radicals (O₂⁻), hydroxyl-free radicals (OH⁻) and hydrogen peroxides (H₂O₂). Apoptosis pathways include the endogenous mitochondrial pathway, endogenous endoplasmic reticulum pathway and exogenous death receptor pathway. In the endogenous mitochondrial pathway, B-cell lymphoma-2 (Bcl-2) family proteins increase mitochondrial permeability by

Correspondence: Qian Zheng
Department of Physiology, North Sichuan Medical College, Nanchong, 637007, People's Republic of China
Email 373969568@qq.com

reducing membrane potential. Reduced membrane potential leads to the destruction of mitochondrial function and structure, an increased production of ROS and reduces antioxidant enzyme activity. Excessive ROS damages the nucleus and mitochondrial DNA. As a consequence, the pro-apoptotic factor cytochrome C (cyt-C) from the mitochondria is released into the cytoplasm and binds with an apoptotic protein activating factor to activate pro-caspase-9 and further activate downstream caspase-3, which mediates apoptosis.^{4,5} Uncoupling proteins (UCPs) are a class of mitochondrial transporters located on the mitochondrial membrane. These transporters include UCP₁, UCP₂, UCP₃, UCP₄, and UCP₅, which can regulate the synthesis of ATP and ROS.⁶ UCPs can affect the survival of β -cells by regulating ATP synthesis, fatty acid release and glucose oxidation, and is closely related to mitochondrial dysfunction mediated by mitochondrial oxidation. Uncoupling protein 3 (UCP₃), a member of the uncoupling protein family, can induce proton leakage across the inner mitochondrial membrane and decreases the electrochemical proton gradient (Δp). In this process, mitochondrial uncoupling generates heat instead of ATP and thereby protects against mitochondrial oxidative damage by reducing the production of ROS.^{7,8} UCP₃ mRNA expression is ubiquitous in vertebrates and can be detected in skeletal muscle, fat, the heart, thymus, spleen and skin cells.⁹ However, in 2008, it was first uncovered that the UCP₃ gene is expressed on the pancreatic islet, which can promote insulin secretion.¹⁰ There is strong evidence that UCP₃ is decreased in the skeletal muscles of T2DM patients.¹¹ Early research identified that fasting blood glucose and insulin levels were reduced in mice overexpressing UCP₃, suggesting that glucose homeostasis may be improved.¹² Diabetes mediates an increase in ROS levels that can lead to impaired endothelial function, decreased smooth muscle in the diabetic corpus cavernosum and increased apoptosis.¹³ Additionally, previous studies have shown that UCP₃ can promote fatty acid oxidation and limit ROS generation.¹⁴ These studies suggested that decreased UCP₃ may be a potential mechanism for T2DM.¹⁵ However, whether UCP₃ affects the onset and development of diabetes is still unknown.

IL-1 β is considered an important endogenous cytokine that can induce β -cell apoptosis and lead to the onset of diabetes by various pathways such as NO,¹⁶ Ca²⁺,¹⁷ mitochondrial membrane potential, ROS¹⁸ and endoplasmic reticulum stress (ERS).¹⁹ The pro-apoptotic effect of IL-1 β was observed in MIN₆ islet cells, which are mouse

tumor cells expressing SV40 large T antigen under the control of an insulin promoter. Therefore, IL-1 β -induced apoptosis has been used to evaluate disease mechanisms and potential treatment strategies for diabetes. EGCG is the most abundant tea-specific catechin and shows evidence in preventing cancer, obesity, diabetes and cardiovascular diseases. More specifically, the treatment of diabetes with EGCG has garnered attention.²⁰ EGCG has an insulin-like effect, where it can regulate blood glucose by improving insulin sensitivity, affecting liver gluconeogenesis and restricting intestinal glucose transport. This mechanism may be attributed to the inhibition of the NF- κ B pathway, limiting the release of pro-inflammatory cytokines and reducing lipid peroxidation in the heart of diabetic animals.¹⁹ In addition, EGCG has strong antioxidant activity, which can prevent or T2DM and complications by scavenging oxygen-free radicals and improving OS. Myung-Kwan Han's study revealed that EGCG can combat IL-1 β -induced RINm5F cell apoptosis. This study also found that this anti-apoptotic mechanism is related to inhibition of NF- κ B and NO.²¹ Whether EGCG protects pancreatic islet cells from interleukin 1 β -induced apoptosis by regulating UCP₃ levels remains to be further studied.

Here, IL-1 β -processed MIN₆ cells were incubated with low and high doses of EGCG to observe the effects of EGCG on β -cell secretion, mitochondrial function and apoptosis. Simultaneously, our study found that UCP₃ was expressed in MIN₆ cells and down-regulated when pro-processed by IL-1 β . Thus, the goal of this work was to reveal whether the protection of EGCG on IL-1 β -induced apoptosis may be related to UCP₃.

Materials and Methods

Cell Culture

MIN₆ cells were purchased from the Baili Company (China) and cultured in High Glucose Dulbecco's minimal essential medium (DMEM) (Hyclone) supplemented with 15% fetal bovine serum (nzk0677, Hyclone), 10000U/mL penicillin (Hyclone) and 1000ug/mL streptomycin (Hyclone) in a 37°C, humidified atmosphere containing 5% CO₂. Culture medium was replaced every 2–3 days and cells were passaged every 5–6 days. Experiments were performed when 28–35th generation cells reached 80% confluency. Briefly, MIN₆ cells (5×10^5 cells/mL) were seeded in 96-well plates overnight and randomly divided into 4 groups (n=6). Then, IL-1 β (20 nM, 792008D, Invitrogen) was used to pre-treat cells for 24h during

experiments except for the blank group. Finally, cells were treated with EGCG (A-0159, YaJi biological) at low (1mM) or high (5mM) concentrations for 24 hours and cultured in a 37°C incubator.

CCK-8 Assay

The viability of MIN₆ cells was evaluated using a Cell Counting Kit-8 (CCK-8) (AR1160-100, Boster). After administration, 96-well cell culture plates were centrifuged (500 r/min, 5min) and the supernatant was removed. Next, cells were resuspended in 200μL serum-free cell culture supplemented with 20ul CCK-8 solution and cultured for 1h at 37°C. A Microplate Reader (Bio-Rad, USA) was used to detect the absorbance of the culture at 450 nm.

Glucose-Stimulated Insulin Secretion (GSIS) Assay

MIN₆ cells were treated with or without high glucose (16.7 mM) culture medium for 2h. Cell culture supernatants were collected to determine the insulin concentration using an ELISA kit (H203, Nanjing Jian Cheng).

Mitochondrial Activity

Mitochondrial membrane potential in MIN₆ cells was measured using the mitochondrial membrane potential assay kit with JC-1 (C2006, Beyotime). JC-1, a fluorochrome, can be accumulated in the mitochondrial matrix to form polymers, which produce a strong red fluorescence (Ex = 585 nm, Em = 590 nm). JC-1 exists in the form of a monomer in the cytoplasm of unhealthy cells and produces green fluorescence (Ex = 514 nm, Em = 529 nm). Quantitative characterization of the mitochondrial membrane potential is $R = \text{red fluorescence} / \text{green fluorescence}$. Briefly, 1×10^4 cells were resuspended in 0.5 mL culture medium, and 0.5 mL of JC-1 staining working solution was added before incubating cells at 37°C for 20 minutes. After centrifugation (4°C, 600r/min, 4min) and discarding the supernatant, JC-1 staining buffer was used to wash the cells two times. Finally, cells were resuspended with buffer and detected by flow cytometry. The ATP assay kit (A095, Nanjing Jian Cheng) and ROS Assay Kit (A087, Nanjing Jian Cheng) were performed according to manufacturer's instructions to further test mitochondrial function.

Apoptosis Assay

To investigate apoptosis of MIN₆ cells, Hoechst 33258 staining (C1017, Beyotime) solution was added to the cells for

a final concentration of 10ug/mL and incubated at 37°C for 15min. Cells were collected and chromatin morphologic changes were observed using fluorescence microscopy (Leica, Germany). Reduced nuclear size, chromatin condensation, intense fluorescence and nuclear fragmentation were indicators of apoptotic cells. Then, we further detected apoptosis using flow cytometry. MIN₆ cells (5×10^5) were plated in a 1.5 mL centrifuge tube and washed twice with cold PBS, suspended in binding buffer and stained by Annexin V-FITC/propidium iodide (PI) (C1056, Beyotime). Cells were incubated for 20 min in the dark (37°C) and the FACSCalibur flow cytometer (Acuuri, USA) was used to observe apoptotic cells within 1 hour. Annexin V-FITC⁺PI⁻: represented early apoptotic cells, Annexin V-FITC⁺PI⁺: represented necrotic cells, Annexin V-FITC⁺PI⁺: represented late apoptotic cells necrotic cells and Annexin V-FITC⁻PI⁻: represented living cells.

Immunohistochemistry

For immunostaining, MIN₆ cells growing on cover glass slides were fixed with 4% paraformaldehyde for 30 min at room temperature, washed 3 times with PBS and then permeabilized for 15 min using 0.5% Triton X-100 in PBS. Cells were washed 3 times with PBS.

Endogenous peroxidase activity was inhibited by immersing sections in methanol with 0.6% H₂O₂ for 30 min at room temperature. After washing with distilled water 3 times and blocking for 30 min at room temperature, diluted primary antibody UCP₃ was added to the cells and incubated at 4°C overnight. After washing twice with PBS, secondary antibody was added and incubated in the darkroom for 30min. At this time, strep avidin-biotin complex (SABC) was added at a proper dilution for 30min (RT) and subsequently washed 4 times with PBS. Colorimetric development of the reaction was performed by incubation in DAB (diaminobenzidine) solution and then washed with water several times. Finally, all specimens were counterstained with hematoxylin solution and mounted with neutral gum. Cells were subjected to microscopy. Brown-yellow staining was considered as positive staining in the cytoplasm as previously described.²² The immunohistochemical score (IHS) was calculated by combining an estimate of the percentage of immunoreactive cells (quantity score=A) with an estimate of the staining intensity (staining intensity score=B). This was scored as follows: For A: no staining was scored as 0, 1~10% of cells stained scored as 1, 11~50% scored as 2, 51~80% scored as 3 and 81~100% scored as 4. B was scored on

a scale of 0 to 3, where 0 (negative), 1 (weak), 2 (moderate) and 3 (strong). $IHS=A \times B$.

Quantitative Real-Time PCR

Total RNA was extracted using the Ultra-pure RNA extraction kit (CW0581, CWbio) according to the manufacturer's instructions. Reverse transcription was performed using the HiFi-MMLV cDNA Kit (CW0744, CWbio) as follows: the mixture was incubated at 37°C for 50 min, then kept at 70°C for 10 minutes and cDNA was stored at -20°C. For qPCR, SYBR green (CW0956, CWbio) was used along with the Real-Time PCR System (ABI7500, USA). Thermal cycles included 10 min at 95°C, 45 cycles of 15 s at 95°C, 60 s at 60°C. Primers included: UCP-3-F: 5'-ACTCCAGCGTCGCC ATCAGG-3'; R: 5'-CTGTAGGCATCCATAGTCCC-3'. GA PDH-F: 5'-TGGAGTCTACTGGCGTCTT-3'; R: 5'-TGTC ATATTCTCGTGGTTCA-3'. Gene expression levels were normalized to Gapdh and relative quantitative calculation analysis was based on the $2^{-\Delta\Delta Ct}$ method.

Western Blotting

To examine the protein expression levels of UCP₃ between different experimental groups, MIN₆ cells were collected after washing twice with PBS and lysed with 200 μ L of lysis buffer containing Phenylmethylsulfonyl fluoride (PMSF) (ST505, Beyotime). Protein concentrations were measured using Coomassie blue staining (P0017B, Beyotime). An equal amount of protein for each sample was denatured at 95°C for 5 minutes in protein-loading buffer, separated by 10% SDS-polyacrylamide gel electrophoresis (PAGE) (80V) and subsequently semi-dry-transferred to polyvinylidene fluoride (PVDF) membranes (Millipore, USA). After PVDF membranes were blocked for 1 hour at room temperature, they were incubated with the following primary antibodies overnight at 4°C: Anti-UCP₃ antibody (ab10985, abcam) (1:300) and mouse mAb-actin (AA128, Beyotime) (1:1000). Finally, membranes were washed with TBST and incubated with an HRP-conjugated goat anti-rabbit (ZB-2301, ZSGB-BIO) or goat anti-mouse (ZB-2305, ZSGB-BIO) secondary antibody (1:3000) for 1 hour at room temperature. Finally, blots were developed with a chemiluminescent HRP substrate (Merk Millipore, USA) and visualized using the Western Lighting-ECL (Fusion solos, China). The absorbance values of target proteins were analyzed using Image J 180.

Statistical Analysis

Data were expressed as mean \pm standard deviation and analyzed using SPSS 15.0 statistical software for all groups

following normal distribution. The *F*-test was used to compare the mean of multiple samples. The Dunnett *t* test and SNK-q test were used for multiple comparisons between multiple sample means. A significant difference was considered when $p < 0.05$.

Results

EGCG Treatment Significantly Improved the Activity and Function of IL-1 β -Stimulated MIN₆ Cells

Consistent with previous reports, we observed that IL-1 β -stimulation reduced cell activity in MIN₆ cells, indicated by significantly decreased OD₄₅₀ values. Co-cultivation with EGCG for another 24 h improved MIN₆ cells activity (Figure 1A). Moreover, IL-1 β pretreatment impaired GSIS (basal and high glucose stimulated) in MIN₆ cells when compared with control group. However, the low and high doses of EGCG treatment significantly increased insulin secretion of IL-1 β -stimulated MIN₆ cells and showed a dose-dependent trend (Figure 1B and C).

EGCG Treatment Recovered Mitochondrial Function and Reduced Oxidative Stress Response of IL-1 β -Stimulated MIN₆ Cells

To further evaluate the effect of EGCG on mitochondrial damage response induced by IL-1 β , we assessed mitochondrial membrane potential and ATP activity. In the IL-1 β -induced MIN₆ cells, a significant decrease in mitochondria membrane potential and ATP activity was observed (Figure 2A–C). Simultaneously, levels of ROS, an oxidative stress response indicator, in IL-1 β -induced MIN₆ cells increased (Figure 2D). EGCG reversed these alterations, dose-dependently increased cell mitochondrial membrane potential, inhibited oxidative stress response and enhanced the activity of ATP (Figure 2D).

EGCG Treatment Decreased Cell Apoptosis in IL-1 β -Stimulated MIN₆ Cells

To confirm the effects of EGCG on IL-1 β -induced apoptosis in MIN₆ cells, Hoechst 33258 staining and flow cytometry were used to examine apoptosis. Microscope analysis revealed condensed and bright apoptotic nuclei were readily observed in cells treated with IL-1 β . However, when treated with EGCG, obvious reduced nuclear condensation in MIN₆ cells was observed (Figure 3), suggesting that




Figure 1 EGCG treatment significantly improved the activity and function of IL-1 β -stimulated MIN₆ cells. MIN₆ cells were stimulated with IL-1 β followed by co-cultivation with low (1mM) or high (5mM) EGCG for 24 h. (A) Cell viability was assessed in each group using the CCK-8 assay. Mitochondrial membrane potential of MIN₆ cells in each group was determined using flow cytometry. Basal insulin (B) and high glucose stimulated insulin (C) levels for group in the cell supernatant were determined using an ELISA kit. Data are representative of 2 independent experiments and presented as mean \pm SD; * p < 0.05, ** p < 0.01. Significant differences were evaluated using one-way ANOVA with a post hoc test (Fisher's least significant difference).

EGCG had a protective effect on IL-1 β -stimulated MIN₆ cells. To quantitatively characterize this observation, we examined apoptotic changes using flow cytometry. Compared with the control, culturing MIN₆ cells in IL-1 β conditions for 24 h markedly increased the fraction of apoptotic cells (Figure 4). As expected, EGCG treatment significantly reduced IL-1 β -induced apoptosis in a dose-dependent manner. These results were consistent with those observed using the Hoechst 33258 assay.

Effect of EGCG on UCP₃ Expression Levels in IL-1 β -Stimulated MIN₆ Cells

UCP₃ is a member of the mitochondrial uncoupling protein family. UCP₃-deficient mice show increased levels of ROS.²³ To assess the potential mediatory role of UCP₃ in IL-1 β -induced apoptosis in MIN₆ cells, UCP₃ protein expression levels were analyzed in cells using immunohistochemistry. UCP₃ protein stained brown and was mainly expressed in the cytoplasm of MIN₆ cells (Figures 5 and 6). EGCG significantly increased UCP₃ protein in MIN₆ cells. Next, both gene and protein expression levels of UCP₃ were detected for each group of MIN₆ cells using RT-PCR and Western blotting, respectively. As expected, when compared to the control group, UCP₃ gene and protein levels were decreased in the IL-1 β -stimulated MIN₆ cells. In

contrast, EGCG treatment significantly increased UCP₃ gene and protein levels (Figure 6). Thus, we speculated that decreased UCP₃ levels under IL-1 β -stimulated conditions could disrupt normal mitochondrial function, thereby resulting in excessive production of ROS and apoptosis, and EGCG reverses this damage through upregulation of UCP₃.

Discussion

Studies have shown that apoptosis is the main method for β -cell death in diabetic patients. IL-1 β can damage β -cell function and mediate apoptosis. Our previous work confirmed that IL-1 β causes β -cell apoptosis in neonatal rats, and this mechanism is related to the NO,²⁴ Ca²⁺ and HO-1/CO pathways.²⁵ A recent study suggested that IL-1 β up-regulates ROS, destroys mitochondrial function and causes apoptosis. John Zeqi Luo et al²⁶ confirmed that mitochondrial membrane potential of INS-1 cells weakens, ATP synthesis decreases and apoptosis increases after IL-1 β treatment. After stimulating with glucose, insulin secretion ability of MIN₆ returned to levels observed in normal islet β -cells. Here, we found that treatment with IL-1 β dramatically increased the apoptosis rate of MIN₆ cells and inhibited cell activity. In addition, insulin secretion function decreased after stimulating with or without glucose. Therefore, we speculated that IL-1 β affected the activity




Figure 2 EGCG treatment recovered mitochondrial function and reduced oxidative stress response in IL-1 β -stimulated MIN₆ cells. MIN₆ cells were stimulated with IL-1 β followed by co-cultivation with low (1mM) or high (5mM) EGCG for 24 h. **(A)** Mitochondrial membrane potential of MIN₆ cells in was determined for each group using flow cytometry. **(B)** The bar chart show a summary of statistical results in **(A)**. The activity of ATP **(C)** and content of ROS **(D)** in MIN₆ cells for each group. Data are representative of 2 independent experiments and presented as mean \pm SD; * p < 0.05, ** p < 0.01. Significant differences were evaluated using one-way ANOVA with a post hoc test (Fisher's least significant difference).




Figure 3 EGCG treatment decreased cell apoptosis in IL-1 β -stimulated MIN₆ cells. MIN₆ cells were stimulated with IL-1 β followed by co-cultivation with low (1mM) or high (5mM) EGCG for 24 h. Morphological analysis of the nuclei were stained using Hoechst 33258. Red arrows indicate apoptotic MIN₆ cells. Representative images for each group (n=6 per group) are shown (magnification \times 400).




Figure 4 EGCG treatment decreased cell apoptosis in IL-1 β -stimulated MIN₆ cells. **(A)** Representative flow cytometry plots show percentages of early apoptotic cells (Annexin V-FITC⁺PI⁻), necrotic cells (Annexin V-FITC⁺PI⁺), late apoptotic cells and necrotic cells (Annexin V-FITC⁺PI⁺) and living cells (Annexin V-FITC⁻PI⁻) for each group of MIN₆ cells receiving specified treatment (n \geq 6 per group). **(B)** The bar charts show a summary of the statistical results in **(A)**. Data are representative of 2 independent experiments and presented as mean \pm SD; *p < 0.05, **p < 0.01. Significant differences were evaluated using one-way ANOVA with a post hoc test (Fisher's least significant difference).




Figure 5 UCP₃ was mainly expressed in the cytoplasm of MIN₆ cells. MIN₆ cells were stimulated with IL-1 β followed by co-cultivation with low (1mM) or high (5mM) EGCG for 24 h. UCP₃ protein levels were examined using immunohistochemistry and were indicated by red arrows. Representative histology images for each group (n=6 per group) are shown (magnification \times 400).

and secretory function of MIN₆ cells and had the ability to induce apoptosis.

Mitochondria are not the only main site of ROS production, but also the first target attacked by ROS. Maladjusted mitochondria causes an accumulation of ROS and reduces antioxidant activity, which further aggravates mitochondrial damage and eventually leads to apoptosis. There is evidence suggesting that UCP₃ inhibits the production of ROS and reduces $\Delta\Psi$, increases proton leaking, affects ATP generation and has an influence on apoptosis.²⁷ Recent studies showed that UCP₃ deficiency impairs myocardial fatty acid oxidation and contractile recovery following ischemia/reperfusion²⁸. In prediabetic subjects and T2DM patients, skeletal muscle UCP₃ content is decreased by almost 50%.²⁹ Our results confirmed that UCP₃ is expressed in MIN₆ cells, however, its expression is significantly reduced after treatment with IL-1 β . Importantly, we observed that ATP synthesis and mitochondrial membrane potential are significantly reduced and increases the synthesis of ROS. Thus, we assumed that IL-1 β -mediated mitochondria disorder is attributed to the down-regulation of UCP₃.

Preventing pancreatic β -cells apoptosis can significantly reduce the incidence of diabetes. Therefore, selecting drugs that inhibit apoptosis is very important. EGCG is a natural catechin distributed in green tea that possesses anti-inflammatory, anticarcinogenic, antioxidative, free-radical scavenging in different cell types and animal models.^{30–32} It has been suggested that the protective effects of green tea may result from antioxidant ingredients, which are molecules containing easily oxidizable phenolic hydroxyl groups. Studies have shown that EGCG protects cell function by reducing oxidative damage and preserving mitochondrial integrity.^{33,34} Animal studies suggested that EGCG can reduce blood glucose in STZ, tetrazolopyrimidine-induced diabetic animals and animal models of spontaneous diabetes.³⁵ Moreover, EGCG can prevent the occurrence of diabetes.³⁶ Liu et al³⁷ emphasized that EGCG has an attenuated effect on myocardial mitochondrial dysfunction in diabetic Goto-Kakizaki rats. Nevertheless, EGCG has distinct effects on different types of cells. For example, H9C2 cells show more stable mitochondrial membrane potential after pre-treatment with EGCG, thereby reducing hypoxia/reoxygenation-induced apoptosis.³⁸ Experiments have confirmed




Figure 6 EGCG significantly increased the gene and protein expression levels of UCP₃ in IL-1 β -stimulated MIN₆ cells. MIN₆ cells were stimulated with IL-1 β followed by co-cultivation with low (1mM) or high (5mM) EGCG for 24 h. **(A)** Summary of the immunohistochemical scores (IHS) for UCP₃ protein levels for each group. **(B)** The gene expression levels of UCP₃ in each group measured by qRT-PCR. **(C)** The protein levels of UCP₃ in each group assessed by Western blotting and normalized to actin. **(D)** Bar chart showing a summary statistical results in **(C)**. Data are representative of 2 independent experiments and presented as mean \pm SD; * p < 0.05, ** p < 0.01, *** p < 0.001. Significant differences were evaluated using one-way ANOVA with a post hoc test (Fisher's least significant difference).

that EGCG protected SH-SY5Y cells against the cytotoxicity induced by lead (Pb) and beta-amyloid peptides (APs) by decreasing OS and preventing apoptosis. This mechanism is related to improving the expression of Bax and BCL₂ and inhibiting annexin V and caspase-3.³⁹ However, Lee et al⁴⁰ found that EGCG promotes 3T3-L1 adipocyte differentiation by increasing ROS production. Our results reflect that EGCG can reduce the apoptosis of MIN₆ cells induced by IL-1 β , enhance cell activity and increase insulin secretion function. These results suggest that EGCG has a protective effect on the function of pancreatic β -cells exposed to IL-1 β . In our study, treatment of EGCG increases UCP₃ and subsequently

increases ATP synthesis and mitochondrial membrane potential. As a result, there are decreased ROS levels observed in IL-1 β -induced MIN₆ cells. These results indicate that EGCG can increase the activity of MIN₆ cells by increasing UCP₃ expression, which stabilizes mitochondrial function by maintaining ATP synthesis and mitochondrial membrane potential.

Conclusion

This study clearly demonstrated that the proinflammatory cytokine IL-1 β significantly decreased the protein and gene expression levels of UCP₃, ATP synthesis and mitochondrial

membrane potential in MIN₆ cells. This work also provided evidence for the first time that EGCG markedly increased protein and gene expression of UCP₃ in IL-1 β -stimulated MIN₆ cells. In conclusion, EGCG has the potential to ameliorate IL-1 β -stimulated cell apoptosis in MIN₆ cells by up-regulating UCP₃ expression and may serve as a therapeutic intervention for metabolic diseases such as diabetes.

Acknowledgments

This work was supported by Key project of Sichuan Province Department of Education, China (17ZA0168, 17ZA0167); Special Project of Strategic Cooperation in Nanchong City, China (18SXH20224)

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The authors do not have potential financial conflicts of interest related to this manuscript. The authors report no conflicts of interest for this work.

References

- Ogurtsova K, Da RFJ, Huang Y, et al. IDF Diabetes Atlas: global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Res Clin Pract.* 2017;128:40–50. doi:10.1016/j.diabres.2017.03.024
- Rhodes CJ. Type 2 diabetes—a matter of beta-cell life and death? *Science.* 2007;307(5708):380–384.
- Giralt M, Villarroja F. Mitochondrial Uncoupling and the Regulation of Glucose Homeostasis. *Curr Diabetes Rev.* 2017;13(4):386–394.
- Chen W, Luo S, Xie P, Hou T, Yu T, Fu X. Overexpressed UCP2 regulates mitochondrial flashes and reverses lipopolysaccharide-induced cardiomyocytes injury. *Am J Transl Res.* 2018;10(5):1347–1356.
- Yang W, Weng J. Early therapy for type 2 diabetes in China. *Lancet Diabetes Endocrinol.* 2014;2(12):992–1002. doi:10.1016/S2213-8587(14)70136-6
- Mailloux RJ, Harper ME. Uncoupling proteins and the control of mitochondrial reactive oxygen species production. *Free Radic Biol Med.* 2011;51(6):1106–1115. doi:10.1016/j.freeradbiomed.2011.06.022
- Cadenas S. Mitochondrial uncoupling, ROS generation and cardioprotection. *Biochim Biophys Acta Bioenerg.* 2018;1859(9):940–950.
- Cadenas S. Mitochondrial uncoupling, ROS generation and cardioprotection. *Biochim Biophys Acta Bioenerg.* 2018;1859(9):940–950. doi:10.1016/j.bbabi.2018.05.019
- Zhao RZ, Jiang S, Zhang L, Yu ZB. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *Int J Mol Med.* 2019;44(1):3–15.
- Li Y, Maedler K, Shu L, Haataja L. UCP-2 and UCP-3 proteins are differentially regulated in pancreatic beta-cells. *PLoS One.* 2008;3(1):e1397. doi:10.1371/journal.pone.0001397
- Schrauwen P, Hesselink MK, Blaak EE, et al. Uncoupling protein 3 content is decreased in skeletal muscle of patients with type 2 diabetes. *Diabetes.* 2001;50(12):2870–2873. doi:10.2337/diabetes.50.12.2870
- Son C, Hosoda K, Ishihara K, et al. Reduction of diet-induced obesity in transgenic mice overexpressing uncoupling protein 3 in skeletal muscle. *Diabetologia.* 2004;47(1):47–54. doi:10.1007/s00125-003-1272-8
- Hirata H, Kawamoto K, Kikuno N, et al. Restoring erectile function by antioxidant therapy in diabetic rats. *J Urol.* 2009;182(5):2518–2525. doi:10.1016/j.juro.2009.07.009
- MacLellan JD, Gerrits MF, Gowing A, Smith PJ, Wheeler MB, Harper ME. Physiological increases in uncoupling protein 3 augment fatty acid oxidation and decrease reactive oxygen species production without uncoupling respiration in muscle cells. *Diabetes.* 2005;54(8):2343–2350. doi:10.2337/diabetes.54.8.2343
- Hesselink MK, Schrauwen P. Towards comprehension of the physiological role of UCP3. *Horm Metab Res.* 2005;37(9):550–554. doi:10.1055/s-2005-870421
- Tannous M, Amin R, Popoff MR, Fiorentini C, Kowluru A. Positive modulation by Ras of interleukin-1 β -mediated nitric oxide generation in insulin-secreting clonal beta (HIT-T15) cells. *Biochem Pharmacol.* 2001;62(11):1459–1468. doi:10.1016/S0006-2952(01)00818-8
- Dula SB, Jecmenica M, Wu R, et al. Evidence that low-grade systemic inflammation can induce islet dysfunction as measured by impaired calcium handling. *Cell Calcium.* 2010;48(2–3):133–142. doi:10.1016/j.ceca.2010.07.007
- Mbaya E, Oules B, Caspersen C, et al. Calcium signalling-dependent mitochondrial dysfunction and bioenergetics regulation in respiratory chain Complex II deficiency. *Cell Death Differ.* 2010;17(12):1855–1866. doi:10.1038/cdd.2010.51
- Verma G, Bhatia H, Datta M. JNK1/2 regulates ER-mitochondrial Ca²⁺ cross-talk during IL-1 β -mediated cell death in RINm5F and human primary β -cells. *Mol Biol Cell.* 2013;24(12):2058–2071. doi:10.1091/mbc.e12-12-0885
- Jang HJ, Ridgeway SD, Kim JA. Effects of the green tea polyphenol epigallocatechin-3-gallate on high-fat diet-induced insulin resistance and endothelial dysfunction. *Am J Physiol Endocrinol Metab.* 2013;305(12):E1444–E1451. doi:10.1152/ajpendo.00434.2013
- Han MK. Epigallocatechin gallate, a constituent of green tea, suppresses cytokine-induced pancreatic beta-cell damage. *Exp Mol Med.* 2003;35(2):136–139. doi:10.1038/emmm.2003.19
- Soslow RA, Dannenberg AJ, Rush D, et al. COX-2 is expressed in human pulmonary, colonic, and mammary tumors. *Cancer Am Cancer Soc.* 2000;89;12:2637–2645.
- Hagen T, Vidal-Puig A. Mitochondrial uncoupling proteins in human physiology and disease. *Minerva Med.* 2015;93(1):41–57.
- Zheng Q, Liu H, Cao DY. Experimental study on the protective effect of Cordyceps mycelium on interleukin 1 β -damaged islet cell. *Chinese J Basic Med Trad Chinese Med.* 2007;014(10):765–767.
- Zheng Q, Liu H, Cao DY. The role of heme oxygenase/carbon monoxide system in anti-IL-1 β -induced islet cell apoptosis treated with fructose 1,6-diphosphate. *Chinese J Applied Physiol.* 2019;4:1254.
- Produit-Zengaffinen N, Davis-Lameloise N, Perreten H, et al. Increasing uncoupling protein-2 in pancreatic beta cells does not alter glucose-induced insulin secretion but decreases production of reactive oxygen species. *Diabetologia.* 2006;50(1):84–93. doi:10.1007/s00125-006-0499-6

27. Brand MD. Uncoupling to survive? The role of mitochondrial inefficiency in ageing. *Exp Gerontol*. 2000;35(6-7):811-820. doi:10.1016/S0531-5565(00)00135-2
28. Edwards KS, Ashraf S, Lomax TM, et al. Uncoupling protein 3 deficiency impairs myocardial fatty acid oxidation and contractile recovery following ischemia/reperfusion. *Basic Res Cardiol*. 2018;113(6):47. doi:10.1007/s00395-018-0707-9
29. Schrauwen P, Mensink M, Schaart G, et al. Reduced skeletal muscle uncoupling protein-3 content in prediabetic subjects and type 2 diabetic patients: restoration by rosiglitazone treatment. *J Clin Endocrinol Metab*. 2006;91(4):1520-1525. doi:10.1210/jc.2005-1572
30. Fang CY, Wu CC, Hsu HY, et al. EGCG inhibits proliferation, invasiveness and tumor growth by up-regulation of adhesion molecules, suppression of gelatinases activity, and induction of apoptosis in nasopharyngeal carcinoma cells. *Int J Mol Sci*. 2015;16(2):2530-2558. doi:10.3390/ijms16022530
31. Yuan JM, Sun C, Butler LM. Tea and cancer prevention: epidemiological studies. *Pharmacol Res*. 2011;64(2):123-135. doi:10.1016/j.phrs.2011.03.002
32. Khan N, Mukhtar H. Multitargeted therapy of cancer by green tea polyphenols. *Cancer Lett*. 2008;269(2):269-280. doi:10.1016/j.canlet.2008.04.014
33. Raza H, John A. In vitro protection of reactive oxygen species-induced degradation of lipids, proteins and 2-deoxyribose by tea catechins. *Food Chem Toxicol*. 2007;45(10):1814-1820. doi:10.1016/j.fct.2007.03.017
34. Meng Q, Velalar CN, Ruan R. Regulating the age-related oxidative damage, mitochondrial integrity, and antioxidative enzyme activity in Fischer 344 rats by supplementation of the antioxidant epigallocatechin-3-gallate. *Rejuvenation Res*. 2008;11(3):649-660. doi:10.1089/rej.2007.0645
35. Ortsater H, Grankvist N, Wolfram S, Kuehn N, Sjöholm A. Diet supplementation with green tea extract epigallocatechin gallate prevents progression to glucose intolerance in db/db mice. *Nutr Metab (Lond)*. 2012;9(1):11. doi:10.1186/1743-7075-9-11
36. Kim SJ, Li M, Jeong CW, et al. Epigallocatechin-3-gallate, a green tea catechin, protects the heart against regional ischemia-reperfusion injuries through activation of RISK survival pathways in rats. *Arch Pharm Res*. 2014;37(8):1079-1085. doi:10.1007/s12272-013-0309-x
37. Liu J, Tang Y, Feng Z, Liu J, Liu J, Long J. (-)-Epigallocatechin-3-gallate attenuated myocardial mitochondrial dysfunction and autophagy in diabetic Goto-Kakizaki rats. *Free Radic Res*. 2014;48(8):898-906. doi:10.3109/10715762.2014.920955
38. Wang W, Huang X, Shen D, Ming Z, Zheng M, Zhang J. Polyphenol epigallocatechin-3-gallate inhibits hypoxia/reoxygenation-induced H9C2 cell apoptosis. *Minerva Med*. 2013;109(2):95-102.
39. Ayyalasomayajula N, Ajumeera R, Chellu CS, Challa S. Mitigative effects of epigallocatechin gallate in terms of diminishing apoptosis and oxidative stress generated by the combination of lead and amyloid peptides in human neuronal cells. *J Biochem Mol Toxicol*. 2014;33(11):e22393.
40. Lee H, Lee YJ, Choi H, Ko EH, Kim JW. Reactive oxygen species facilitate adipocyte differentiation by accelerating mitotic clonal expansion. *J Biol Chem*. 2009;284(16):10601-10609. doi:10.1074/jbc.M808742200

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>