

TRIM66 Overexpression Promotes Glioma Progression and Regulates Glucose Uptake Through cMyc/GLUT3 Signaling

Yuequn Song^{1,2}Lifang Meng³Jian Yu¹Zhi Cao¹Jizhou Sun¹Hongyu Zhao²

¹Department of Neurosurgery, the Fourth Affiliated Hospital of China Medical University, Shenyang, People's Republic of China; ²Department of Neurosurgery, Shengjing Hospital of China Medical University, Shenyang, People's Republic of China; ³Department of Scientific Research, China Medical University, Shenyang, People's Republic of China

Objective: Tripartite motif 66 (TRIM66) is reported to be closely associated with human cancers. However, the roles of TRIM66 in glioma remain unclear. The present study aimed to investigate the clinical significance and biological roles of TRIM66 in human glioma.

Methods: TRIM66 expression in glioma tissues was examined by immunohistochemistry. TRIM66 overexpression and siRNA knockdown were performed in glioblastoma cell lines. CCK8, colony formation assay, transwell assay, Annexin V and JC1 staining, glucose uptake assay, and Western blotting were used to explore the biological roles and potential underlying mechanisms of TRIM66 in glioma progression.

Results: Our results showed that TRIM66 was overexpressed in 52/95 glioma cases. The rates of TRIM66 overexpression in Grade I, Grade II, Grade III, and Grade IV gliomas were 16.6%, 41.3%, 58.6%, and 70.9%, respectively. Oncomine data showed that TRIM66 was upregulated in glioblastoma and oligodendroglioma compared with normal brain tissues. TRIM66 expression was higher in glioblastoma cell lines compared with normal SVG p12 glial cell line. TRIM66 promoted in vitro and in vivo proliferation, invasion, and inhibited temozolomide (TMZ)-induced apoptosis. Notably, TRIM66 increased glucose metabolism by upregulating glucose uptake, glucose consumption, and ATP production. Western blotting showed that TRIM66 positively regulated cMyc and GLUT3. Depletion of cMyc by siRNA abolished the effect of TRIM66 on GLUT3. Chromatin immunoprecipitation (ChIP) assay showed that cMyc could bind to the promoter regions of GLUT3 in glioblastoma cells.

Conclusion: TRIM66 was upregulated in human gliomas, where it promoted cell growth and chemoresistance. Our data also identified novel roles of TRIM66 in glioma progression. TRIM66 upregulates glucose uptake and mitochondrial function through the cMyc/GLUT3 signaling, which makes it a potential therapeutic target.

Keywords: TRIM66, glioma, cMyc, GLUT3

Introduction

Glioma is the most common malignant tumor in the central nervous system. Despite the improvement in earlier diagnosis, surgery, and chemotherapy, the prognosis for this malignancy remains poor.¹ Overexpression of oncogenes plays a vital role in the development and progression of human glioma and identifying molecular targets in tumor progression is an important task.^{2,3}

Tripartite motif (TRIM) superfamily proteins, most of which have E3 ubiquitin ligase activities and characterized by conserved tripartite motif (TRIM), exhibit various cellular functions and biological processes including transcription

Correspondence: Hongyu Zhao
Department of Neurosurgery, Shengjing Hospital of China Medical University, 36 Sanhao St, Heping District, Shenyang, Liaoning, 110004, People's Republic of China
Tel +86 18900912293
Email hongyu_zhao2020@163.com

regulation, intracellular signaling, cell growth, apoptosis, innate immunity, autophagy, developmental processes, and carcinogenesis.^{4–6} Tripartite motif 66 (TRIM66) is a member of the TRIM family which possesses potential functions as a transcriptional repressor.⁷ TRIM66 has emerged as an oncogenic protein that is closely related to multiple malignant tumors. In hepatocellular carcinoma (HCC), overexpression of TRIM66 promoted cell proliferation and invasion.⁸ TRIM66 activated Wnt signaling by increasing glycogen synthase kinase-3 β (GSK-3 β) phosphorylation and β -catenin expression.⁸ TRIM66 deficiency suppressed cell viability, proliferation, migration, and invasion in prostate cancer cells, possibly through regulation of STAT2/IL-2 signaling.⁹ These studies indicated that TRIM66 might function as an oncogenic protein. However, its expression pattern and biological effects in glioma have not been investigated.

Here, we investigated the expression pattern of TRIM66 in glioma tissues. We then explored the biological effects of TRIM66 on proliferation, invasion, and TMZ-induced apoptosis in glioma cells. The possible underlying mechanism was also investigated.

Methods and Materials

Patients and Specimens

This study was approved by the Institutional Review Boards of China Medical University and was conducted following the principles expressed in the Declaration of Helsinki. Informed consent was obtained from patients. The paraffin-embedded breast cancer specimens were obtained from the Pathology archives of the Fourth Affiliated Hospital of China Medical University, which contain specimens no longer required to be maintained.

Immunohistochemical Staining

The glioma tissue samples were fixed with paraformaldehyde and embedded in paraffin. Sections (4 μ m thick) were prepared and the sections were deparaffinized with xylene and blocked for endogenous peroxidase with H₂O₂. The sections were washed with PBS buffer three times and incubated with TRIM66 (1:100, Sigma, USA) primary antibody at 4°C overnight. Subsequently, the sections were treated with Elivision plus Polymer HRP (Mouse/Rabbit) IHC Kit (Maixin Biotech, Fuzhou, China). After incubation, the sections were washed with PBS buffer and stained with DAB plus kit (Maixin Biotech).

Two pathologists (Huihui Xu and Qi Gao) who were blinded to all clinical data independently evaluated all samples and scored for protein immunoreactivity as well as for intensity of positively stained tumor cells. A third pathologist (Yahao Li, Associate professor) reviewed the slide when there was a discrepancy. Staining for each of the proteins was scored using the methods of modified immunoreactive score (IRS).^{10,11} The intensity of TRIM66 staining was scored as 0 negative; 1 weak; 2 moderate/strong. The percentage score was divided as: 0, 0%; 1 1–25%; 2, 26–50%; 3, 51–75% and 4, 76–100%. The two scores were multiplied to get a final staining score from 0–8. Samples with scores <4 were considered as TRIM66 low expression. Samples with scores between 4–8 were considered as TRIM66 high expression. Images were captured using Olympus BX53 microscope and Olympus CellSens software (Tokyo, Japan).

Cell Culture and Transfection

In this study, human glioma cell lines including A172, U87MG, U251, and human normal astrocyte SVG p12 were obtained from American Typical Culture Collection (ATCC, Manassas, VA, USA). A172 and U251 cells were cultured in DMEM medium. U87MG and SVG p12 cells were cultured in EMEM medium. Cells were cultured with mediums supplied with 10% fetal bovine serum (FBS). TRIM66 overexpression plasmid and negative control plasmid were purchased from Origene (Origene, Rockville, USA). Lipofectamine 3000 (Invitrogen, USA) was used for plasmid transfection. TRIM66 knockdown small interfering RNA and the corresponding non-targeting siRNA were obtained from Dharmacon. Dharmafect1 reagent (Horizon, Cambridge, UK) was used for siRNA knockdown.

Western Blot Analysis

Total proteins were harvested and extracted using lysis mixture with phosphatase inhibitor cocktail and protease inhibitor cocktail. Then, the protein samples were quantified and the same amount of proteins were transferred to PVDF membranes (EMD Millipore, MA, USA). The membranes were blocked with BSA at room temperature for 2 hours. Then, the membranes were washed with TTBS buffer and incubated with primary antibodies at 4°C overnight. Primary antibodies used were listed as follows: TRIM66 (1:1000, Sigma, USA), GLUT3 (1:800, Cell signaling technology, USA), cMyc (1:3000, Abcam, UK). Then, membranes were incubated with

HRP-conjugated goat anti-rabbit or goat anti-mouse secondary antibodies for 2 h at 37°C. The immunoreactive bands were detected and reported using SuperSignal West Dura Extended Duration Substrate (Thermo, MA, USA) and Bio-Imaging Systems (DNR, Israel).

Real-Time Fluorescent Quantitative PCR

Total RNA solution of glioma cells was prepared using TRIzol reagent. After quantification, reverse transcription was performed using PrimeScript RT Kit (Takara, Dalian, China). And the reverse transcription reaction was conducted as follows: 37°C for 45 minutes and 85°C for 2 minutes. SYBR select master Mix Kit (Thermo, MA, USA) was used for qPCR, and the reaction was conducted as follows: 96°C for 3 minutes, 40 cycles of 95°C for 20 seconds, and 60°C for 30 seconds. The sequences of primers were listed as follows: TRIM66 for 5'-GAGGAAGCTGCAAAAGAAGGAC-3', rev 5'-TGAGGCGCACA TCTGATACCA-3'; GLUT3 for 5'- CCTTTGGCACTCT CAACCAGC -3', rev 5'- AACCCAGTAGCAGCGGC CAT -3'; GAPDH for 5'-GAAATCCCATCACCATCTTCC AG-3', rev 5'-GAGCCCCAGCCTTCTCCAT-3'. Relative expression of TRIM66 was normalized to GAPDH and calculated according to $2^{-\Delta\Delta ct}$. This experiment was repeated three times.

CCK8 and Colony Formation Assay

The cell viability assay was performed using the Cell Counting Kit-8 solution. Cells were plated in 96-well plates at a concentration of 5000 cells/well. For quantitation of cell proliferation rate, 10 μ L of CCK8 solution was added to each well and incubated for 4 h at 37°C. Optical density was measured at 450nm using a microplate reader.

For colony formation assay, cells were seeded into dishes (about 1000 cells), incubated for 2 weeks, and then washed and stained with Giemsa. Colonies with more than fifty cells were counted using a microscope.

Flow Cytometry

Cells were harvested by trypsinization and centrifuged at 1200 rpm for 10 minutes to pellets. Then, the pellets were washed with PBS buffer (twice, 3 minutes each time). Annexin V/PI kit (BD bioscience, USA) was used for staining. Cells were re-suspended in 500 μ L binding buffer (1 \times). 5 μ L Annexin V-FITC and 5 μ L PI were incubated with the cell suspension in the dark for 15 minutes.

For cell cycle analyses, cells were harvested and washed in ice-cold phosphate-buffered saline (PBS) 3

times. Cells were fixed in 75% ethanol at -20°C overnight. Then, cells were centrifuged at 1200rpm for 10 minutes and the supernatant was discarded. The pellets were washed with ice-cold PBS 3 times (5 minutes each time). Cells were stained with 50 μ g/mL propidium iodide (Sigma-Aldrich, Germany), 10 μ g/mL RNase A and 0.2% Triton X-100 for 30 minutes in the dark. A flow Cytometer was used to analyze the apoptosis rate and cell cycle distribution.

Analysis of Mitochondrial Membrane Potential

Mitochondrial membrane potential was analyzed using JC-1 dye (Cell signaling technology, MA, USA). Cells were collected and then incubated with 1 μ g/mL JC-1 for 30 min at 37°C. The stained cells were centrifuged at 1200 rpm for 10 minutes and washed with PBS 3 times. The cells were re-suspended in 300 μ L PBS buffer. The mitochondrial membrane potential was determined by Flow Cytometer.

Glucose Uptake Assay

Cells were trypsinized from the plate and washed with PBS buffer. Then, cells were centrifuged at 1200 rpm for 5 minutes. The pellets were re-suspended in PBS with 2% FBS (FBS/PBS). The 2-NBDG (Thermo Fisher Scientific, USA) stock solution was added to each cell suspension at a final concentration of 10 μ M and incubated for 30 minutes at 37°C. Cells were collected, washed with ice-cold PBS, and re-suspended in 300 μ L ice-cold FBS/PBS. The fluorescence intensity was measured using Flow Cytometer.

Glucose Consumption and ATP Level Measurement

The supernatants were collected and the glucose consumption was measured using the Glucose Oxidase (GOD) Assay Kit (Applygen Technologies, Beijing, China) according to the manufacturer's instructions. For ATP level measurement, cells were lysed in 200 μ L of ATP Assay Buffer and the lysates were centrifuged at 12,000 rpm for 5 minutes. 100 μ L ATP reaction solution was added into a 96-well plate and 20 μ L samples were added to the ATP reaction solution. A standard curve was performed and the fluorescence was measured using a microplate reader.

Transwell Invasion Assay

Cell invasive capacity was determined by transwell invasion assay using the 24-well transwell chamber (Corning, NY, USA). Matrigel (BD bioscience, CA, USA) was coated in the upper compartment chamber and incubated for four hours. Cells in 100 μ L of serum-free DMEM medium were seeded on the upper chamber, and DMEM medium with 10% FBS was added into the bottom chamber. The cells were incubated for 24 hours; then, the cells on the bottom surface were fixed 4% paraformaldehyde and stained with 0.1% crystal violet for 2 hours.

Chromatin Immunoprecipitation (ChIP)

Cells were cross-linked with formaldehyde and resuspended in lysis buffer. The lysates were sonicated to yield 300–1000 bp DNA fragments. The DNA-protein complex was incubated with cMyc antibody (Abcam, USA) and IgG antibody (Cell signaling technology, USA) overnight at 4°C in the presence of protein A/G beads (Millipore, Hayward, CA, USA). qPCR was performed by using primers for the promoter of the SLC2A3/GLUT3 gene. The primers for ChIP were listed as follows:

SLC2A3/GLUT3 position 1 forward, 5' CACCTCAG CCTCCCAAGTAGC 3'

SLC2A3/GLUT3 position 1 reverse, 5' GCAACACA GTGAGACCCCTT 3'

SLC2A3/GLUT3 position 2 forward, 5' CTCACGC CTGTAATCCCAGCA 3'

SLC2A3/GLUT3 position 2 reverse, 5' GTCTCCA TCTCTGACCTCGT 3'

Nude Mice Xenograft

U251 cell line with stable TRIM66 overexpression was selected using G418 after transfection. BALB/c athymic nude mice were obtained from Slac Laboratory Animals company (Shanghai, China). Xenograft model was established by subcutaneous injection of stable cells (5 million cells/mouse). Tumor size was measured and recorded. Animals were sacrificed and xenograft tumors were removed 25 days after injection.

Statistical Analysis

The data were analyzed using SPSS 22.0 software for Windows (SPSS, Chicago, IL, USA). Data of TRIM66 expression was downloaded from the Oncomine database

and statistically analyzed by Mann–Whitney *U*-test. Overall survival was visualized using Kaplan–Meier curves, and the differences were compared by Log rank test. Comparisons of biological experiments between different groups were performed using the Student's *t*-test. $P < 0.05$ was considered a statistically significant difference.

Results

TRIM66 is Overexpressed in Glioma

We first examined the protein expression of TRIM66 in 95 glioma specimens by immunohistochemistry. Staining intensity and percentage were combined to evaluate TRIM66 expression levels. Normal glial cells showed negative or weak expression (Figure 1A). In contrast, TRIM66 overexpression was observed in 54.7% (52/95) of glioma tissues (Figure 1B–F).

Next, we analyzed the correlation between clinicopathological factors and TRIM66. The results showed that the rate of TRIM66 protein upregulation in Grade I (pilocytic astrocytoma), Grade II (diffuse astrocytoma), Grade III (anaplastic astrocytoma), and Grade IV glioma (glioblastoma multiforme, GBM) were 16.6%, 41.3%, 58.6%, and 70.9%, respectively (Table 1) ($p = 0.0284$). Grade III glioma and glioblastoma multiforme (GBM) showed a significantly higher positive rate compared with Grade I and II gliomas. Since grade I glioma (pilocytic astrocytoma) samples are uncommon compared with higher grade gliomas. The cohort of grade I gliomas is only 6 patients and therefore it might be hard to have a representative TRIM66 expression in this subgroup of patients.

We also analyzed data from the Oncomine database. The Sun Brain Glioblastoma dataset showed TRIM66 mRNA was higher in glioblastoma tissues compared with normal brain tissues (Figure 2A). Sun Brain Oligodendroglioma dataset showed that TRIM66 expression was higher in Oligodendroglioma compared with the normal brain (Figure 2B). The Cancer Genome Atlas (TCGA) glioma cohort showed that glioma patients with high TRIM66 expression have shorter overall survival time compared with those with low TRIM66 expression, although the *p*-value did not reach a statistical significance (Log rank test, $p = 0.056$, Figure 2C). In addition, TCGA dataset analysis showed that TRIM66 DNA copy number was higher in Oligodendroglioma compared with the normal brain (Figure 2D).




Figure 1 Expression of TRIM66 in glioma specimens. (A) Negative TRIM66 staining in normal glial cells. (B) Negative TRIM66 staining in a case of Grade I glioma (pilocytic astrocytoma). (C) Negative TRIM66 staining in a case of Grade II glioma (diffuse astrocytoma). (D) High TRIM66 staining in a case of Grade III glioma (anaplastic astrocytoma). (E) Negative TRIM66 staining in a case of Grade IV glioma (glioblastoma multiforme). (F) High TRIM66 staining in a case of Grade IV glioma (glioblastoma multiforme).

TRIM66 Promotes Proliferation and Invasion

We examined protein and mRNA levels of TRIM66 in GBM cell lines U87MG, U251, A172, and normal glial cell line SVG p12. As shown in [Figure 3A](#), U87 and A172 cell lines showed relatively high endogenous TRIM66 expression compared with SVG p12 cell line. To investigate the biological function of TRIM66, TRIM66 siRNA knockdown and overexpression was performed in A172 and U251 cell line, respectively. The transfection efficiencies were confirmed by Western blotting and RT-qPCR ([Figure 3B](#)).

CCK-8 assays demonstrated that TRIM66 overexpression significantly increased proliferation rate in U251 cell

line while TRIM66 knockdown decreased cellular growth rate in A172 cell line ([Figure 4A](#)). Colony formation assays demonstrated that TRIM66 transfection significantly increased colony numbers in U251 cell line while TRIM66 knockdown decreased colony formation ability in A172 cell line ([Figure 4B](#)). We also evaluated the change in invasiveness after TRIM66 overexpression and knockdown. TRIM66 overexpression increased invasive ability in U251 cells while TRIM66 depletion inhibited invasion in A172 cells ([Figure 4C](#)). We also performed siRNA knockdown in U87MG cell line. Knockdown efficiency was confirmed using Western blot and real-time PCR ([Supplementary Figure 1A](#)). CCK-8 and colony formation

Table I The Relationship Between TRIM66 and Clinical Features of Astrocytoma

Parameters	Number	TRIM66		p value
		Low Expression	High Expression	
Age				
< 60	59	27	32	0.9003
≥ 60	36	16	20	
Gender				
Female	27	12	15	0.9195
Male	68	31	37	
Grading				
Grade I	6	5	1	0.0284
Grade II	29	17	12	
Grade III	29	12	17	
Grade IV	31	9	22	

assays showed that TRIM66 knockdown decreased proliferation rate and colony formation ability ([Supplementary Figure 1B and C](#)). Matrigel invasion assay showed that TRIM66 knockdown inhibited invasion in U87MG cell line ([Supplementary Figure 1D](#)). In addition, To examine the effect of TRIM66 on tumor growth in vivo, we established TRIM66 overexpressing U251 cell line by G418 selection and cells were subcutaneously injected into nude mice. As shown in [Supplementary Figure 2A and B](#), the in vivo growth rate and tumor sizes of TRIM66 overexpressing U251 cells were much larger than control cells, suggesting TRIM66 promoted glioma growth in vivo. We also examined the change in cell cycle transition. As shown in [Figure 5A](#), TRIM66 overexpression significantly increased the S phase percentage and decreased the G1 phase percentage. TRIM66 depletion decreased the S phase percentage and increased the G1 phase percentage. The above results suggested that TRIM66 regulated malignant proliferation and invasion of glioma cells.

TRIM66 Regulates Temozolomide (TMZ)-Induced Apoptosis and Mitochondrial Function

We also explored the role of TRIM66 on chemosensitivity. According to previous studies,^{13–15} temozolomide (100μM) was used to treat glioma cells after TRIM66 plasmid and siRNA transfection. Annexin V/PI staining was used to determine the change of apoptosis. AnnexinV/PI staining showed that TRIM66 depletion

upregulated the levels of TMZ-induced apoptosis after 24h of treatment. TRIM66 overexpression reduced the apoptosis rate, suggesting TRIM66 could decrease chemosensitivity to chemotherapeutic drugs ([Figure 5B](#)). In addition, TRIM66 knockdown also increased apoptosis rate in U87MG cell line ([Supplementary Figure 1E](#)).

Sensitivity to chemotherapy is often regulated by the mitochondrial apoptosis pathway, which could be reflected by mitochondrial membrane potential. Low membrane potential induces green fluorescence after JC-1 staining ([Figure 5C](#)). Our data showed that TRIM66 siRNA increased the percentage of green fluorescence while TRIM66 overexpression decreased green fluorescence, suggesting that TRIM66 plays an important role in maintaining normal mitochondrial membrane potential ([Figure 5C](#)).

TRIM66 Regulates ATP Production and Glucose Uptake

Cancer cells use glucose metabolism to produce ATP, which is essential for the growth and development of chemoresistance. Glucose uptake is an important step in glucose metabolism. Our data showed that TRIM66 overexpression upregulated ATP levels while TRIM66 knockdown decreased ATP levels ([Figure 6A](#)), suggesting TRIM66 is a regulator of energy production in glioma cells. Next, we explored the change of glucose uptake and consumption. Using 2-NBDG uptake assay, we showed that TRIM66 overexpression upregulated glucose uptake while TRIM66 knockdown inhibited the level of glucose uptake ([Figure 6B](#)). As shown in [Figure 6C](#), TRIM66 overexpression increased glucose consumption, while TRIM66 siRNA reduced it.

TRIM66 Regulates cMyc/GLUT3 Signaling

To find out the potential mechanism of TRIM66 activity, we screened several potentially related proteins. Using Western blot, we validated TRIM66 overexpression and downregulation on the immunoblot to confirm TRIM66 level of expression in the cells ([Figure 7A](#)). Our results demonstrated that TRIM66 positively regulated both cMyc and GLUT3 protein expression ([Figure 7A](#)). We also screened the possible change of p-AKT, p-ERK, E-cadherin, ZO-1, beta-catenin, p-c-Met after TRIM66 overexpression and knockdown. However, there was no significant change in these proteins/signaling pathways (data not shown). cMyc has been reported as a transcriptional regulator of glucose metabolism.




Figure 2 TRIM66 in gliomas according to Oncomine and TCGA data. **(A)** Analysis of Sun Brain Glioblastoma data suggested that TRIM66 mRNA was increased in glioblastoma tissues compared with normal brain tissues. **(B)** Analysis of Sun Brain Oligodendroglioma dataset of Oncomine showed that TRIM66 expression was higher in Oligodendroglioma compared with the normal brain. **(C)** Analysis of the TCGA dataset showed that glioma patients with high TRIM66 expression have shorter overall survival time compared with those with low TRIM66 expression, although the p-value did not reach a statistical significance (Log rank test, $p=0.056$). **(D)** TCGA dataset analysis showed that TRIM66 DNA copy number in was higher in Oligodendroglioma compared with the normal brain.




Figure 3 TRIM66 expression in glioma cell lines. (A) Western blot and RT-qPCR results showed that TRIM66 protein was low in normal glial cell line SVG p12. TRIM66 showed high expression in glioblastoma cell lines U87MG and A172. (B) Transfection and siRNA knockdown efficiencies were confirmed by Western blot and RT-qPCR in U251 and A172 cells, respectively. * $p < 0.05$.

To validate the association between cMyc and TRIM66-induced GLUT3 upregulation, cMyc siRNA was used in the U251 cell line, which was co-transfected with the TRIM66 plasmid. As shown in Figure 7B, cMyc knockdown downregulated protein and mRNA levels of GLUT3. cMyc depletion abolished the effect of TRIM66 on GLUT3 upregulation.

We also investigated if cMyc regulated SLC2A3/GLUT3 gene by binding to its promoter in glioma cells. The cMyc matrix and potential binding sites on the

SLC2A3/GLUT3 promoter were predicted by JASPAR (Figure 7C). Binding of cMyc to SLC2A3/GLUT3 promoter was examined using ChIP assays followed by qPCR assay (Figure 7C). These results suggested that TRIM66 transcriptional regulated SLC2A3/GLUT3 through cMyc in glioma cells.

Discussion

Growing evidence showed elevated expression of TRIM66 protein in various cancers. TRIM66 overexpression




Figure 4 TRIM66 regulates proliferation and invasion in glioma cells. **(A)** CCK8 assay results showed TRIM66 overexpression significantly increased proliferation rate in U251 cells while TRIM66 knockdown decreased cellular growth rate in A172 cells. **(B)** Colony formation assay results showed TRIM66 transfection significantly increased colony numbers in U251 cells while TRIM66 knockdown decreased colony numbers in A172 cells. **(C)** TRIM66 overexpression increased the numbers of invading cell number in U251 cells while TRIM66 depletion decreased the numbers of invading number in A172 cells. * $p < 0.05$.




Figure 5 TRIM66 promotes cell cycle, apoptosis and mitochondrial function. **(A)** Flow cytometry showed that TRIM66 overexpression elevated the S phase percentage and decreased the G1 phase percentage in U251 cells. TRIM66 depletion decreased the S phase and increased the G1 phase percentage in A172 cells. **(B)** Annexin V/PI results showed that the TMZ-induced apoptosis rate was decreased in U251 cells transfected with TRIM66. TRIM66 depletion upregulated TMZ-induced apoptosis in A172 cells. **(C)** JC1 staining showed that TRIM66 siRNA increased the percentage of green fluorescence in A172 cells while TRIM66 overexpression decreased green fluorescence in U251 cells. * $p < 0.05$.




Figure 6 TRIM66 regulates ATP production and glucose uptake. **(A)** ATP production assay showed that TRIM66 overexpression upregulated ATP levels while TRIM66 knockdown decreased ATP levels. **(B)** 2-NBDG glucose uptake showed that TRIM66 overexpression increased glucose uptake rate in U251 cells. TRIM66 depletion downregulated glucose uptake rate in A172 cells. **(C)** Glucose consumption assay showed that TRIM66 overexpression increased glucose consumption in U251 cells, while TRIM66 siRNA reduced it in A172 cells. * $p < 0.05$.




Figure 7 TRIM66 regulates GLUT3 through cMyc. **(A)** Western blot results showed that TRIM66 overexpression upregulated TRIM66, GLUT3, and cMyc protein while TRIM66 siRNA knockdown downregulated TRIM66, GLUT3 and cMyc protein. **(B)** cMyc siRNA was transfected with TRIM66 plasmid. cMyc siRNA significantly decreased the expression of cMyc and GLUT3. In cells transfected with cMyc siRNA, the effect of TRIM66 overexpression on GLUT3 protein and mRNA was significantly ameliorated. **(C)** Prediction of cMyc binding sites on SLC2A3/GLUT3 promoter by JASPAR database analysis. ChIP assay showed that cMyc could bind to the SLC2A3/GLUT3 promoter region. * $p < 0.05$.

contributes to osteosarcoma carcinogenesis and indicates poor survival outcomes.¹⁶ TRIM66 expression is upregulated in hepatocellular carcinoma and confers tumorigenicity by regulating GSK3 Wnt signaling.⁸ TRIM66 is overexpressed in NSCLC patients and serves as a prognostic marker.¹⁷ Knockdown of TRIM66 inhibits malignant behavior and epithelial–mesenchymal transition in NSCLC cells.¹⁸ TRIM66 knockdown inhibits tumor growth by regulating JAK2/STAT3 signaling in colorectal cancer cells.¹⁹ However, the expression pattern and potential mechanism of TRIM66 in glioma remain unexplored. In the present study, we demonstrated that TRIM66 was upregulated in 54.7% of glioma and correlated with the higher grade, which was supported by data from the Oncomine database, indicating the potential use of TRIM66 as a clinical indicator of the malignant feature.

TRIM66 has been implicated in a variety of oncogenic processes. CCK-8 and colony formation assays confirmed that ectopically expressed TRIM66 stimulated proliferation, invasion, cell cycle transition. TRIM66 depletion showed the opposite results. The data indicate TRIM66 plays important role in the malignant proliferation of glioma. Furthermore, our data confirmed that TRIM66 overexpression decreased TMZ sensitivity and reduced apoptosis. Mitochondrial function plays an important role in regulating chemo-sensitivity.²⁰ We further showed that TRIM66 was able to maintain mitochondrial membrane potential after TMZ treatment, suggesting TRIM66 could protect mitochondrial function.

Many cancers have long been thought to primarily metabolize glucose for energy production. Both proliferation and chemoresistance in tumor cells rely on energy supply and glucose uptake is the initial step during the process.^{21,22} Our results showed that increased TRIM66 overexpression increased glucose uptake, consumption, and ATP production in glioma cells, indicating that TRIM66 serves as a positive regulator of glucose metabolism. Growing evidence indicates that glucose metabolism plays an important role in the development and growth of glioma.²³ Malignant cells are highly dependent on glycolysis for ATP generation, known as the Warburg effect in which withdrawal of glucose resulted in apoptosis of glioma cells.²⁴ In addition, many report has shown that ATP provided by glycolysis is essential for the development of chemoresistance²⁵ and inhibition of glucose metabolism increased apoptosis in glioma cells.^{24,26} In addition, recent reports showed key regulators of glucose metabolism could control invasion phenotype of glioma

cells.^{27,28} Based on these reports and our results, the positive regulation of TRIM66 on glucose metabolism/ATP production might contribute to its biological roles such as promoting proliferation, invasion and inhibiting apoptosis. Our findings linked the role of TRIM66 on glucose metabolism to its biological functions, which has not been previously reported.

Upregulation of glucose metabolic enzymes plays an important role in elevated glucose consumption and ATP production of cancer cells.^{29,30} After screening several glucose metabolic-related enzymes we found that TRIM66 induced GLUT3 protein, which was an important glucose transporter during the glucose uptake process.³¹ The involvement of GLUT3 in glioma progression has been previously reported.^{32,33} GLUT3 is overexpressed in glioma and correlates with pathological grade.³⁴ GLUT3 upregulation promotes metabolic reprogramming associated with antiangiogenic therapy resistance in glioblastomas.³⁵ A recent study showed that GLUT3 is a druggable target in a subpopulation of glioblastoma.³² Thus, TRIM66 might regulate glucose metabolism through GLUT3 in glioma.

We also found that TRIM66 upregulated cMyc protein expression. cMyc is an important malignant biomarker in glioma and regulates various biological processes such as apoptotic cell death, proliferation, survival, differentiation.^{12,36,37} cMyc is also required for the maintenance of glioma cancer stem cells.³⁸ It has been reported that cMyc could regulate glucose metabolism by activating transcription of related genes.³⁹ Using cMyc siRNA knockdown, we were able to show that cMyc could mediate TRIM66-induced GLUT3 upregulation. This was further supported by ChIP data that cMyc is directly bound to the GLUT3 promoter in glioma cells. Together, these findings showed a strong link among TRIM66, cMyc, and GLUT3 in glioma cells.

In summary, our results reveal that TRIM66 is overexpressed in human glioma. Our data provide evidence that TRIM66 overexpression in glioma plays an important role in proliferation, apoptosis, and glucose uptake, possibly by regulating cMyc/GLUT3 signaling. TRIM66 might function as a novel therapeutic target in gliomas.

Acknowledgments

The present study was funded by Science Research Foundation of Liaoning Provincial Department of Education (FWZR2020015).

Disclosure

The authors declare that they have no competing interests.

References

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin*. 2012;62(1):10–29. doi:10.3322/caac.20138
2. Krupkova O, Loja T, Redova M, et al. Analysis of nuclear nestin localization in cell lines derived from neurogenic tumors. *Tumour Biol*. 2011;32(4):631–639. doi:10.1007/s13277-011-0162-9
3. Zadran S, Amighi A, Otiniano E, Wong K, Zadran H. ENTPD5-mediated modulation of ATP results in altered metabolism and decreased survival in glioblastoma multiforme. *Tumour Biol*. 2012;33(6):2411–2421. doi:10.1007/s13277-012-0505-1
4. Hatakeyama S. TRIM family proteins: roles in autophagy, immunity, and carcinogenesis. *Trends Biochem Sci*. 2017;42(4):297–311. doi:10.1016/j.tibs.2017.01.002
5. Hatakeyama S. TRIM proteins and cancer. *Nat Rev Cancer*. 2011;11(11):792–804. doi:10.1038/nrc3139
6. Jaworska AM, Wlodarczyk NA, Mackiewicz A, Czerwinska P. The role of TRIM family proteins in the regulation of cancer stem cell self-renewal. *Stem Cells*. 2020;38(2):165–173. doi:10.1002/stem.3109
7. McAviera RM, Crawford LJ. TIF1 proteins in genome stability and cancer. *Cancers*. 2020;12(8):2094. doi:10.3390/cancers12082094
8. Fan W, Du F, Liu X. TRIM66 confers tumorigenicity of hepatocellular carcinoma cells by regulating GSK-3 β -dependent Wnt/ β -catenin signaling. *Eur J Pharmacol*. 2019;850:109–117. doi:10.1016/j.ejphar.2019.01.054
9. Cao H, Gao R, Chen L, Feng Y. TRIM66 promotes malignant progression of prostate carcinoma through the JAK/STAT pathway. *FEBS Open Bio*. 2020;10:515–524. doi:10.1002/2211-5463.12798
10. Remmele W, Stegner HE. [Recommendation for uniform definition of an immunoreactive score (IRS) for immunohistochemical estrogen receptor detection (ER-ICA) in breast cancer tissue]. *Der Pathologe*. 1987;8(3):138–140. German.
11. Fedchenko N, Reifensrath J. Different approaches for interpretation and reporting of immunohistochemistry analysis results in the bone tissue - a review. *Diagn Pathol*. 2014;9:221. doi:10.1186/s13000-014-0221-9
12. Annibaldi D, Whitfield JR, Favuzzi E, et al. Myc inhibition is effective against glioma and reveals a role for Myc in proficient mitosis. *Nat Commun*. 2014;5:4632. doi:10.1038/ncomms5632
13. Plummer S, Wallace S, Ball G, et al. A human iPSC-derived 3D platform using primary brain cancer cells to study drug development and personalized medicine. *Sci Rep*. 2019;9(1):1407. doi:10.1038/s41598-018-38130-0
14. Lee SY. Temozolomide resistance in glioblastoma multiforme. *Genes Dis*. 2016;3(3):198–210. doi:10.1016/j.gendis.2016.04.007
15. Agnihotri S, Burrell K, Buczkowicz P, et al. ATM regulates 3-methylpurine-DNA glycosylase and promotes therapeutic resistance to alkylating agents. *Cancer Discov*. 2014;4(10):1198–1213. doi:10.1158/2159-8290.CD-14-0157
16. Chen Y, Guo Y, Yang H, et al. TRIM66 overexpression contributes to osteosarcoma carcinogenesis and indicates poor survival outcome. *Oncotarget*. 2015;6(27):23708–23719. doi:10.18632/oncotarget.4291
17. Ma Y, Dai HY, Zhang F, Zhao D. TRIM66 expression in non-small cell lung cancer: a new predictor of prognosis. *Cancer Biomark*. 2017;20(3):309–315. doi:10.3233/CBM-170207
18. Dai HY, Ma Y, Da Z, Hou XM. Knockdown of TRIM66 inhibits malignant behavior and epithelial-mesenchymal transition in non-small cell lung cancer. *Pathol Res Pract*. 2018;214(8):1130–1135. doi:10.1016/j.prp.2018.06.008
19. He T, Cui J, Wu Y, Sun X, Chen N. Knockdown of TRIM66 inhibits cell proliferation, migration and invasion in colorectal cancer through JAK2/STAT3 pathway. *Life Sci*. 2019;235:116799. doi:10.1016/j.lfs.2019.116799
20. Fu L, Dong Q, He J, et al. SIRT4 inhibits malignancy progression of NSCLCs, through mitochondrial dynamics mediated by the ERK-Drp1 pathway. *Oncogene*. 2017;36(19):2724–2736. doi:10.1038/ncr.2016.425
21. Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*. 2009;324(5930):1029–1033. doi:10.1126/science.1160809
22. Xu M, Chen S, Yang W, et al. FGFR4 links glucose metabolism and chemotherapy resistance in breast cancer. *Cell Physiol Biochem*. 2018;47(1):151–160. doi:10.1159/000489759
23. Bao Z, Chen K, Krepel S, et al. High glucose promotes human glioblastoma cell growth by increasing the expression and function of chemoattractant and growth factor receptors. *Transl Oncol*. 2019;12(9):1155–1163. doi:10.1016/j.tranon.2019.04.016
24. Jelluma N, Yang X, Stokoe D, Evan GI, Dansen TB, Haas-Kogan DA. Glucose withdrawal induces oxidative stress followed by apoptosis in glioblastoma cells but not in normal human astrocytes. *Mol Cancer Res*. 2006;4(5):319–330. doi:10.1158/1541-7786.MCR-05-0061
25. Wang T, Ning K, Sun X, Zhang C, Jin LF, Hua D. Glycolysis is essential for chemoresistance induced by transient receptor potential channel C5 in colorectal cancer. *BMC Cancer*. 2018;18(1):207. doi:10.1186/s12885-018-4123-1
26. Atif F, Yousuf S, Espinosa-Garcia C, Sergeeva E, Stein DG. Progesterone treatment attenuates glycolytic metabolism and induces senescence in glioblastoma. *Sci Rep*. 2019;9(1):988. doi:10.1038/s41598-018-37399-5
27. Miyai M, Kanayama T, Hyodo F, et al. Glucose transporter Glut1 controls diffuse invasion phenotype with perineuronal satellitosis in diffuse glioma microenvironment. *Neuro Oncol Adv*. 2021;3(1):vdaa150. doi:10.1093/oaajnl/vdaa150
28. Abbadi S, Rodarte JJ, Abutaleb A, et al. Glucose-6-phosphatase is a key metabolic regulator of glioblastoma invasion. *Mol Cancer Res*. 2014;12(11):1547–1559. doi:10.1158/1541-7786.MCR-14-0106-T
29. Cho H, Lee YS, Kim J, Chung JY, Kim JH. Overexpression of glucose transporter-1 (GLUT-1) predicts poor prognosis in epithelial ovarian cancer. *Cancer Invest*. 2013;31(9):607–615. doi:10.3109/07357907.2013.849722
30. Brown RS, Wahl RL. Overexpression of Glut-1 glucose transporter in human breast cancer. An immunohistochemical study. *Cancer*. 1993;72(10):2979–2985. doi:10.1002/1097-0142(19931115)72:10<2979::AID-CNCR2820721020>3.0.CO;2-X
31. Barron CC, Bilan PJ, Tsakiridis T, Tsiani E. Facilitative glucose transporters: implications for cancer detection, prognosis and treatment. *Metabolism*. 2016;65(2):124–139. doi:10.1016/j.metabol.2015.10.007
32. Cosset E, Ilmjarv S, Dutoit V, et al. Glut3 addiction is a druggable vulnerability for a molecularly defined subpopulation of glioblastoma. *Cancer Cell*. 2017;32(6):856–868 e855. doi:10.1016/j.ccell.2017.10.016
33. Zheng C, Yang K, Zhang M, et al. Specific protein 1 depletion attenuates glucose uptake and proliferation of human glioma cells by regulating GLUT3 expression. *Oncol Lett*. 2016;12(1):125–131. doi:10.3892/ol.2016.4599
34. Liu Y, Li YM, Tian RF, et al. The expression and significance of HIF-1 α and GLUT-3 in glioma. *Brain Res*. 2009;1304:149–154. doi:10.1016/j.brainres.2009.09.083
35. Kuang R, Jahangiri A, Mascharak S, et al. GLUT3 upregulation promotes metabolic reprogramming associated with antiangiogenic therapy resistance. *JCI Insight*. 2017;2(2):e88815. doi:10.1172/jci.insight.88815
36. Ashrafzadeh M, Zarabi A, Hushmandi K, et al. C-Myc signaling pathway in treatment and prevention of brain tumors. *Curr Cancer Drug Targets*. 2020;20:104–114. doi:10.2174/1568009619666191023100046

37. Shervington A, Cruickshanks N, Wright H, et al. Glioma: what is the role of c-Myc, hsp90 and telomerase? *Mol Cell Biochem.* 2006;283(1–2):1–9. doi:10.1007/s11010-006-2495-z
38. Wang J, Wang H, Li Z, et al. c-Myc is required for maintenance of glioma cancer stem cells. *PLoS One.* 2008;3(11):e3769. doi:10.1371/journal.pone.0003769
39. Osthus RC, Shim H, Kim S, et al. Deregulation of glucose transporter 1 and glycolytic gene expression by c-Myc. *J Biol Chem.* 2000;275(29):21797–21800. doi:10.1074/jbc.C000023200

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.