




Exploring the Prevalence and Patterns of Use of Sleep Aids and Stimulants Among Emergency Physicians and EMS Providers in Saudi Arabia

Osama Y Kentab¹
 Ahmad AAI Ibrahim¹
 Khaled R Soliman¹ 
 Muna Aljahany² 
 Abdulaziz I Alresseeni¹
 Abdulaziz S Algarni¹ 

¹Emergency Department, King Abdullah Bin Abdulaziz University Hospital, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia; ²Clinical Sciences Department, College of Medicine, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia

Purpose: This study explores the prevalence and patterns of the use of pharmacologic sleep aids and stimulants among Saudi Arabia's emergency physicians (EPs) and emergency medical services (EMS) providers.

Patients and Methods: We adopted a descriptive, cross-sectional design. To collect data on the types and frequencies of sleep aids and stimulants used, we distributed a semi-structured, anonymous, web-based questionnaire to registered EPs, paramedics, and emergency medicine technicians (EMTs) in the Saudi Commission for Health Specialties. An internal consistency analysis showed good reliability (Cronbach's alpha=0.667) of the questionnaire. A subscale analysis confirmed the results—alpha values were 0.720 and 0.618 for the use of sleep aids and stimulants, respectively.

Results: Males and females represented 81.8% and 18.2%, respectively, of the valid sample of 669 participants. Respondents aged 25–34, 35–44, and 45–55 years represented 51.9%, 32.7%, and 10.2% of the sample, respectively. Results showed that a majority of the respondents (67.1%) used stimulants. Caffeine was the most common stimulant; caffeine and energy drinks were used by 65.9% and 17.2% of the respondents, respectively. Caffeine, energy drinks, nicotine, and ephedrine were used by 65.9%, 17.2%, 18.5%, and 17.3% of the respondents, respectively. The respondents who used at least one sleeping aid and those using only one and two sleeping aids accounted for 36.6%, 15.6%, and 9.7%, respectively. The most common sleeping aids antihistamines and marijuana were used on most days by 13.4% and 13.3% of the respondents, respectively. The average monthly number of night shifts ($P = 0.025$) significantly influenced sleep aid use. Respondents working in night shifts for 3–5 months or more than 7 days were more likely to use sleeping aids.

Conclusion: Future research should enhance health workers' knowledge of the efficacy and safety of these medications and guide strategies to organize and reduce night shift work.

Keywords: sleep disruption, shift work, circadian rhythm, night shift

Introduction

Working in emergency departments (EDs) represents a unique challenge to physicians, owing to the department's continuous operations. Therefore, it is important to create work schedules that ensure the availability of qualified staff around the clock. This involves deputing staff with an ability to work overnight and in rotating shifts. This nature of job leads to changing sleep patterns and requires the staff to compensate for their sleep debt with daytime sleep. In Saudi Arabia and several other countries, ED physicians work eight hours per shift, and they constantly rotate

Correspondence: Muna Aljahany
 Department of Clinical Sciences, College of Medicine at Princess Nourah Bint Abdulrahman University, P.O. Box 84428, Riyadh, Saudi Arabia
 Tel +966 118233711
 Email msaljahany@pnu.edu.sa

shifts. These clinical schedules cause the staff to experience detrimental effects, including fatigue, decreased alertness, irritability, burnout, decreased memory, and communication problems.^{1,2} In organizations, these effects are reflected through increased rates of absenteeism, accidents, and medical errors.^{3,4}

It has been well-established that the aforementioned negative effects of shift work are attributable to disrupting the circadian rhythms and reducing the daytime recovery sleep of the ED staff.⁵ As per a study in the United States, the night-shift ED residents experienced a higher frequency of difficulties in sleeping, shorter sleep duration, and decreased performance than that of their peers working in other shifts.⁵ Similarly, in Saudi Arabia, the ED physicians who worked in shifts experienced a poorer sleep quality and higher Epworth sleepiness scale scores than that of their non-shift work healthcare professionals.⁶ Other investigations concerning nurses have demonstrated increased risks of reduced periods of sleep, insomnia, and excessive daytime sleepiness and fatigue when doing shift work,⁷ working night shifts,^{8,9} and working a three-shift pattern,¹⁰ respectively.

Given their nature of work, the ED staff should take suitable measures to combat sleep deprivation, circadian rhythm variation, and the possible consequences to their health. Given the lack of notable changes in the medical residents' education system and no reduction in work hours or an abandonment of shift work, there are few solutions to the sleep-related problems. Hence, there is a need to implement strategies facilitating administrative and behavioral changes. In addition to the aforementioned issues, the use of pharmacologic sleep aids has been a cause for significant concern for several years. Although the use of sleep aids is prevalent among the ED staff and may help them counteract the problems associated with their fluctuating and highly demanding schedules, the use of these medications may not be a benign solution. Along with sleep aids, there may be a high likelihood of the on-shift use of stimulants by a considerable proportion of the ED staff.

Studies assessing the use of sleep aids and stimulants among the ED staff are limited by poor response rates^{11,12} or to a single institution's ED.¹³ It must be noted that, to the best of our knowledge, the prevalence of the use of such medication has not yet been investigated in Saudi Arabia. The outcomes of this study will provide significant insights into the current status of sleep patterns and the effect of such medications on the alertness levels. Having

said that, the future research can increase the awareness and knowledge of ED staff on the use of such medications.

We assessed the prevalence and patterns of using sleep aids among ED physicians and EMS providers via a nationwide, web-based survey in the Kingdom of Saudi Arabia. As secondary outcomes, we investigated their use of stimulants and its effect in promoting alertness.

Materials and Methods

Study Design and Setting

Based on the guidelines of the strengthening the reporting of observational studies in epidemiology (STROBE) checklist for cross-sectional studies,¹⁴ we conducted a descriptive, prospective cross-sectional study among all the registered EPs, paramedics, and EMTs in the Saudi Commission for Health Specialties (SCFHS) in Saudi Arabia. Study conducted between February 2020 to March 2021. We invited them to participate in an online, semi-structured questionnaire designed to collect data on their use of sleep aids and stimulants. We excluded the non-ED physicians from the analysis. The study protocol was approved by the Institutional Review Board of the Princess Nourah bint Abdulrahman University.

Study's Tool

The online questionnaire was adapted from a recent survey-based study in Canada.¹³ We obtained an email approval from the authors to use the survey tool. The survey had five main domains comprising 19 items: 1) sociodemographic data (4 items)—age, gender, marital status, and current living situation; 2) practice information (4 items)—work position and experience, work location, and the average number of night shifts in EDs (defined as finishing work after midnight); 3) the perceived effects of working in EDs on sleep (5 items)—the perceived effects of shift work on the ability to sleep, the subjective perceptions of the factors contributing to sleeping difficulties, and the personal perceptions of having insomnia or chronic insomnia; 4) the use of sleep aids (4 items)—the prior and current use of sleep aids, the prescriber of these medications, the reasons attributing to the use of sleep aids; and 5) the use of stimulants (2 items)—the prior and current use of stimulants to maintain wakefulness/alertness during work.

Selection of Participants

The sample comprised all the registered EPs, paramedics, and EMTs in the SCFHS in Saudi Arabia. The sample size

was estimated using the n4Studies application.¹⁵ Using a standard proportion of 0.5, standard error (d) = 0.05, Alpha (α) = 0.05, and $Z(0.975) = 1.959964$, the sample size was estimated to be 341. With a 20% invalid responding rate, the final sample size was determined to be 376.

The questionnaire was uploaded to SurveyMonkey® (www.surveymonkey.com)—a platform that collects participants' responses via customized questionnaires. Participation was entirely voluntary; the participants were informed about the purpose of the study and they provided consent to participate upon the survey's completion, and the study was conducted in accordance with the Declaration of Helsinki. While we did not collect personal information, individual data for each response was coded for subsequent analysis using Microsoft Excel 2016.

Outcome Measures

The primary outcomes represent the proportion of ED staff who had used or were using sleep aids. We also explored the types and frequency of using such medications. The secondary outcomes include the sociodemographic and practice characteristics and the use of stimulants to overcome fatigue during work. We defined insomnia as a condition wherein a person experiences difficulty with the initiation, maintenance, or quality of sleep, despite normal sleep opportunities and times, which eventually leads to daytime functional/cognitive impairment.¹⁶ Chronic insomnia was classified as having insomnia for more than one month.¹⁷ The use of sleep aids/stimulants for more than 6 times per month was defined as frequent use.

Primary Data Analysis

Using a Cronbach's alpha test, we tested the internal consistency of the whole questionnaire and the specific items related to the use of sleep aids and stimulants. The descriptive analysis included frequencies and percentages of all the categorical variables along with their respective 95% confidence intervals (95% CIs). We used the multi-response analysis to analyze items related to the domains of using sleep aids and stimulants (domains 4 and 5). Statistical analysis was performed using the Statistical Package for Social Sciences version 26.0 (SPSS Inc., Chicago, IL, USA).

Internal Consistency of the Questionnaire

An analysis of the internal consistency of all the questionnaire items showed good reliability (Cronbach's

$\alpha=0.667$). A subscale analysis showed similar results, where alpha values were 0.720 and 0.618 for the domains of the use of sleep aids and stimulants, respectively.

Statistical Analysis

Statistical analysis was performed using R 3.6.3. Counts and percentages were used to summarize the distribution of categorical variables, and the mean \pm standard deviation (SD) was used to summarize the distribution of continuous variables. The chi-square test of independence was used to assess the relationship between categorical variables. Hypothesis testing was performed at a 5% level of significance.

Results

Of the 777 respondents, we excluded 30 responses from nurses and administrators. Subsequently, we excluded 78 responses owing to missing data (information limited to the work location). Of the excluded participants, the number of personnel working in and outside the Kingdom of Saudi Arabia accounted for 48 (61.5%) and 30 (38.5%), respectively. Thus, the final sample consisted of 669 respondents working in the Kingdom of Saudi Arabia. The overall response rate was 26%. Males and females represented 81.8% and 18.2% of the study sample, respectively. Respondents aged 25–24 years old represented 51.9% of the study sample. Respondents aged 35–40 years and 45–55 years represented 32.7% and 10.2% of the study sample, respectively. Most of them perceived shift work as negatively affecting their ability to sleep effectively ($n = 560$, 85%).

The percentage of respondents working >7 nights/month, on an average, exceeded more than a quarter of the sample, while 27.7% and 22.6% worked 5–7 nights/month and 5–7 night shifts/month, respectively. We did not observe any statistically significant relationship between the average number of night shifts/month (ordinal) and the number of sleeping aids used ($r = 0.04$, $p = 0.42$) (Table 1).

Results showed that 63.4% ($n = 424$) of the respondents did not use any of the sleeping aids mentioned in the questionnaire. Thus, the number of respondents using at least one, only one, and two sleeping aids accounted for 245 (36.6%), 104 (15.6%), and 65 (9.7%), respectively (Figure 1). Melatonin and antihistamines were used by 16.1% ($n=108$) and 22.4% ($n = 150$) respondents, respectively. Barbiturates ($n = 13$, 1.9%) and marijuana ($n = 17$, 2.5%) were the least used sleeping aids (Figure 2). Based on the available data, the most commonly used sleeping

Table 1 Factors Contributing to Difficulty in Initiating or Maintaining Sleep

	[ALL]	N
	N=659	
Shift work negatively affects the ability to sleep effectively		659
No	99 (15.0%)	
Yes	560 (85.0%)	
Factors contributing to difficulty initiating or maintaining sleep?[†]		
Work hours/demands of work: Yes	299 (44.7%)	669
Work-related emotional stress: Yes	270 (40.4%)	669
Personal or family-related stressors: Yes	174 (26.0%)	669
Family commitments: Yes	147 (22.0%)	669
Circadian misalignment or day/night sleep reversal: Yes	310 (46.3%)	669
Other (please specify):	4 (0.6%)	669
How many night shifts (finishing after midnight) do you work in an average month?		513
>7 nights	167 (32.6%)	
1–3 nights	36 (7.02%)	
3–5 nights	116 (22.6%)	
5–7 nights	142 (27.7%)	
I only work nights	17 (3.31%)	
None	35 (6.82%)	

Note: [†]Percentage for each response was calculated from the total number of respondents.

aids antihistamines and marijuana were used on most days of the week by 13.4% and 13.3% of the respondents, respectively. The least commonly used sleep aid was benzodiazepine, which was used less than once/month, as per 50% of the respondents.

Table 2 showed a statistically significant relationship between gender and the use of sleeping aids ($P = 0.041$). Females were more likely to use sleeping aids (45.1%) than that of males (34.7%). We did not find a statistically significant relationship between age and the use of sleeping aids, but between work area and the use of sleeping aids. The post-hoc comparisons showed that respondents from the Southern province were less likely to use sleeping aids than those from Riyadh ($P = 0.043$). We did not find any statistically significant relationship between the marital status and ED-work duration and the use of sleeping aids, but between the average monthly number of midnight shifts and the use of sleeping aids ($P = 0.025$).

The post-hoc comparisons showed that respondents who worked for 3–5 months or > 7 days in night shifts were more likely to use sleeping aids. There was no statistically significant relationship between work and the use of sleeping aids.

In Table 3, 217 respondents reported the reasons for using pharmacological sleeping aids. The main reason was attributed to the need to reset the circadian rhythms or

natural sleep cycle ($n = 66$, 30.4%). Other reasons included sleeping after ($n = 48$, 22.1%) and before a night shift ($n = 39$, 18%) and regardless of the shifts worked ($n = 47$, 21.7%). The proportion of respondents using OTC medications accounted for less than one-half of the sample ($n = 90$, 42.7%), and the proportion that self-prescribed medications accounted for one-quarter of the sample ($n = 56$, 26.5%). Results showed that only 32.6% of the respondents did not use any stimulants ($n = 218$); the respondents using only one and two stimulants represented 43.8% (less than one-half of the respondents) and 16.9% of the sample, respectively.

Results showed that caffeine and energy drinks were used by 65.9% ($n = 441$) and 17.2% ($n = 115$), respectively, of the respondents. Nicotine and ephedrine were used by 18.5% ($n = 124$) and 17.3% ($n = 116$) of the respondents, respectively. Amphetamines, modafinil, and herbals were the least used stimulants, as per the respondents (Figure 3).

As per Table 4, 82.69% ($n = 86$) of the respondents reported that energy drinks were used once/shift. The proportion of respondents using modafinil once and twice/shift represented 61.9% (more than half of the respondents) and 13.64% of the sample, respectively. Half of the respondents used nicotine at least five times/shift.

Table 5 did not show a statistically significant relationship between the gender and age and the use of stimulants




Figure 1 Sleeping aids used by the respondents.




Figure 2 Frequency of using sleeping aids.

Table 2 Factors Associated with Using at Least One Sleeping Aid

	No N=424	Yes N=245	P-value
Gender			0.041
Female	67 (54.9%)	55 (45.1%)	
Male	357 (65.3%)	190 (34.7%)	
Age (Year)			0.083
less than 24	15 (78.9%)	4 (21.1%)	
25–34	219 (63.1%)	128 (36.9%)	
35–44	128 (58.4%)	91 (41.6%)	
45–54	50 (73.5%)	18 (26.5%)	
55–64	12 (75.0%)	4 (25.0%)	
Work area			0.043
Riyadh	224 (60.4%)	147 (39.6%)	
Eastern province	56 (63.6%)	32 (36.4%)	
Northern province	12 (57.1%)	9 (42.9%)	
Southern province	50 (80.6%)	12 (19.4%)*	
Western province	77 (64.7%)	42 (35.3%)	
Marital status			0.849
Single	141 (65.6%)	74 (34.4%)	
Married	268 (62.2%)	163 (37.8%)	
Divorced	11 (61.1%)	7 (38.9%)	
Widowed	2 (66.7%)	1 (33.3%)	
Job			
Assistant Consultant/Registrar	81 (60.9%)	52 (39.1%)	
Consultant/Associate Consultant	67 (54.0%)	57 (46.0%)	
Emergency Resident in the program/service	65 (63.7%)	37 (36.3%)	
EMS	77 (72.0%)	30 (28.0%)	
Other (please specify)	6 (60.0%)	4 (40.0%)	
Paramedic	126 (66.0%)	65 (34.0%)	
Duration of work in the ER medicine			0.113
<10	277 (61.8%)	171 (38.2%)	
10–20	125 (64.4%)	69 (35.6%)	
>20	22 (81.5%)	5 (18.5%)	
Average number of night shifts (finishing after midnight) per month:			0.025
None	25 (71.4%)	10 (28.6%)	
1–3 nights	11 (52.4%)	10 (47.6%)	
3–5 nights	29 (39.2%)	45 (60.8%)*	
5–7 nights	39 (60.0%)	26 (40.0%)	
>7 nights	86 (51.5%)	81 (48.5%)*	
I only work nights	11 (64.7%)	6 (35.3%)	

Notes: Statistical analysis was performed using the Chi-square test of independence. *Significantly different compared to the first level.

($P > 0.05$ for both comparisons). We found a statistically significant relationship between the work and the use of stimulants ($P < 0.05$).

The post-hoc comparisons showed that respondents from the Western province were more likely to use stimulants than those from Riyadh ($P < 0.05$). We did not find a statistically significant relationship between the marital status and the use of stimulants.

The results showed a statistically significant relationship between the current job and the use of stimulants, with consultants/associate consultants reporting a higher use of stimulants than that of the associated consultants. We found a relationship between the ED-work duration and the stimulant use ($P < 0.05$), with respondents who worked for > 20 years reporting less stimulant use than those who worked for < 10 years.

Discussion

Sleep deprivation resulting from ED shift schedules negatively impacts the clinical and cognitive performance of physicians and non-physicians.^{18,19} Several strategies can be applied to prevent sleep deprivation, such as the appropriate scheduling and sequence of shifts and napping for short periods.²⁰ However, instead of considering these administrative aspects, at an individual level, there has been a prevalence in the use of pharmacologic aids to modulate sleep and wakefulness.

In a recent national wide study, about 15% of the general population of the employee sample reported taking sleep medications in the preceding month, where 10.8% had taken medication only infrequently compared to 36.6% who reported use from our study participants.²¹ In this study, 67.4% had reported use of stimulants at some point of their career. Most of them believed that shift work could have influenced their normal sleep pattern; the most significantly perceived factors of sleep disruption were circadian misalignment, increased work hours, and work-related emotional stress.

The aforementioned results mimic those reported in other studies in the literature. In Canada, the rates of the previous and current use of sleep aids were estimated at 67% and 56%, respectively, among ED physicians.¹³ Similarly, 55.7% of the ED staff reported the use of sleep aids in an early web-based survey in the United States.¹² Concerning their routine use, a study among the staff of a local ED center in the United States revealed that about 38% of the ED residents and nurses have been using pharmacologic sleep aids.²² This reflects the urgent need to reduce the disruptive effects of shift work on the circadian rhythm, which is dependent on the internal clock

Table 3 Reasons for Using Sleep Aids and Supply Sources

	[ALL] N=217	N
Reason for using a pharmacological sleep aid		217
For sleeping, regardless of shifts worked	47 (21.7%)	
For resetting circadian rhythms or the natural sleep cycle	66 (30.4%)	
For sleeping after a night shift	48 (22.1%)	
For sleeping before a night shift	39 (18.0%)	
Other (please specify)	17 (7.83%)	
Supplier for sleeping aid prescriptions		211
Colleague	18 (8.53%)	
Combination: General Practitioner and Colleague	5 (2.37%)	
Combination: General Practitioner and Colleague	7 (3.32%)	
General Practitioner or Family Doctor	26 (12.3%)	
Others (please specify)	9 (4.27%)	
Over-the-counter (OTC) medication	90 (42.7%)	
Self	56 (26.5%)	

controlled by the hypothalamic suprachiasmatic nucleus.²³ In essence, ED shift workers deprived of nighttime sleep cannot get the same amount of rest from the daytime sleep

as that from the nighttime sleep. The repeated or continuous work during night shifts disrupts sleep patterns, which cause a dyssynchronization between chronological


**Figure 3** Stimulants used by the respondents.

Table 4 Frequency of Stimulant Use

	Once/Shift	Twice/Shift	Three Times/Shift	Four Times/Shift	Five Times or More
Caffeine	146 (35.44%)	164 (39.81%)	69 (16.75%)	15 (3.64%)	18 (4.37%)
Energy Drinks (for example: Rockstar and Red Bull)	86 (82.69%)	17 (16.35%)	0 (0.00%)	1 (0.96%)	0 (0.00%)
Amphetamines (for example: Adderall and Dexedrine)	14 (63.64%)	3 (13.64%)	4 (18.18%)	1 (4.55%)	0 (0.00%)
Modafinil (for example: Alertec)	13 (61.90%)	0 (0.00%)	5 (23.81%)	3 (14.29%)	0 (0.00%)
Ephedrine/Ephedra	12 (57.14%)	1 (4.76%)	3 (14.29%)	3 (14.29%)	2 (9.52%)
Nicotine (cigarettes, gum, pipe, and cigars)	17 (14.91%)	15 (13.16%)	18 (15.79%)	11 (9.65%)	53 (46.49%)
Herbals (eg Guarana, Ginseng, and Gotu Kola.)	18 (66.67%)	4 (14.81%)	1 (3.70%)	2 (7.41%)	2 (7.41%)

Notes: Percentage was calculated from the number of respondents who reported the use frequency for each stimulant.

and biological clocks and, in turn, makes it difficult for sleep-deprived individuals to recover effectively from sleep deprivation. Therefore, pharmacological solutions are usually considered by shift workers as an option, including those working in the EDs.

In this analysis, antihistamines, melatonin, marijuana and analgesics were used sleep aids as past or present routine medications. Likewise, Handel et al¹² demonstrated that antihistamines, primarily diphenhydramine, were used frequently by ED residents. Nonetheless, other studies have shown a relative prevalence of other medications. Nonbenzodiazepine hypnotics were the most popular among ED physicians in Canada¹³ and the ED residents of the national American Board of Emergency Medicine.²⁴ Recently, Richards et al²² found marked discrepancies between past and current uses; they showed that the ED staff used diphenhydramine, melatonin, and alcohol in the past, whereas melatonin, diphenhydramine, and doxylamine were routinely used by the participants.

There is a lack of clinical evidence regarding the efficacy and safety of the use of antihistamines among shift work-staff, as indicated by a recent systematic review.²⁵ Antihistamines induce tolerance with repeated use, and they can be associated with adverse events such as dizziness, xerostomia, and post-awakening sedation. However, after night shifts, it has been shown that melatonin improves the length of sleep, but no other sleep quality parameters.²⁶ Other randomized clinical trials have found no significant effects of melatonin on sleep quality, cognitive functions, or night alertness in ED physicians; these studies are also outdated.^{27,28} These inconsistent outcomes indicate the lack of high-quality studies, which necessitates conducting prospective clinical investigations involving different sleep aid medications used by shift work-ED staff.

Concerning stimulants, we found that the sampled participants predominantly used caffeine-based stimulants. This was in agreement with other studies on nightshift workers in healthcare settings.^{11,22,29} This observation was unsurprising since caffeine is the most commonly used psychoactive substance worldwide.³⁰ It increases arousal in a dose-dependent manner, with more significant effects in fatigued than rested individuals.³¹ The non-sleep-deprived and sleep-deprived individuals experience performance benefits, owing to increased alertness, after consuming 30–300 mg and 200–600 mg of caffeine, respectively.^{32,33} Notwithstanding the low-quality evidence, caffeine combined with naps can significantly reduce sleepiness during night shifts, as revealed by a Cochrane systematic review published in 2014.²⁶ Among nightshift workers, the caffeine use is usually associated with improved cognitive performance, mood, and enhanced wakefulness; however, its use may induce tolerance in a significant proportion of individuals.³⁴ We could not assess the effects of caffeine on work performance among ED staff; this could be best assessed by assessing fatigue during shifts.

Limitations

As with any observational study, we could not infer causation owing to the self-reported nature of data. The subjective reporting of data might have also led to underreporting of the prevalence of the use of sleep aids and stimulants. In other words, social desirability bias in the responses of the participants might have resulted in dishonest responses. A recall bias is also evident, where the respondents might have inadvertently under- or over-reported their personal perceptions and uses of medications/active substances. It must be noted that sleep disruption was assessed on the basis of the responses of the participants, instead of the scores of sleep quality and daytime sleepiness. Additionally, work-related

Table 5 Factors Associated with the Use of at Least One Stimulant

	No N=218	Yes N=451	P-value
Gender			0.874
Female	41 (33.6%)	81 (66.4%)	
Male	177 (32.4%)	370 (67.6%)	
Age (Years)			0.161
less than 24	7 (36.8%)	12 (63.2%)	
25–34	109 (31.4%)	238 (68.6%)	
35–44	65 (29.7%)	154 (70.3%)	
45–54	29 (42.6%)	39 (57.4%)	
55–64	8 (50.0%)	8 (50.0%)	
Work area			0.04
Riyadh	130 (35.0%)	241 (65.0%)	
Eastern province	29 (33.0%)	59 (67.0%)	
Northern province	8 (38.1%)	13 (61.9%)	
Southern province	21 (33.9%)	41 (66.1%)	
Western province	27 (22.7%)	92 (77.3%)*	
Current relationship status			0.406
Single	68 (31.6%)	147 (68.4%)	
Married	139 (32.3%)	292 (67.7%)	
Divorced	9 (50.0%)	9 (50.0%)	
Widowed	1 (33.3%)	2 (66.7%)	
Job			< 0.001
Assistant Consultant/Registrar	43 (32.3%)	90 (67.7%)	
Consultant /Associate Consultant	22 (17.7%)	102 (82.3%)*	
Emergency resident in the program/service	40 (39.2%)	62 (60.8%)	
EMS	46 (43.0%)	61 (57.0%)	
Others (please specify)	5 (50.0%)	5 (50.0%)	
Paramedic	62 (32.5%)	129 (67.5%)	
Duration of working in the Emergency Medicine			0.006
<10	147 (32.8%)	301 (67.2%)	
10–20	55 (28.4%)	139 (71.6%)	
>20	16 (59.3%)	11 (40.7%)*	
Average number of night shifts (finishing after midnight) per month:			0.841
None	9 (25.7%)	26 (74.3%)	
1–3 nights	12 (33.3%)	24 (66.7%)	
3–5 nights	30 (25.9%)	86 (74.1%)	
5–7 nights	33 (23.2%)	109 (76.8%)	
>7 nights	41 (24.6%)	126 (75.4%)	
I only work nights	3 (17.6%)	14 (82.4%)	

Notes: Statistical analysis was performed using Chi-square test of independence.

*Significantly different compared to the first level.

stress was not assessed using reliable tools. Finally, although the study's tool was based on another study, and it showed good values of the parameters of the internal consistency, our questionnaire was not validated for use among ED workers, particularly in the national context.

Conclusion

In line with the aforementioned findings and limitations, this study provides recommendations for future investigations. The future research should consider a prospective, randomized design to collect real-time data from cohort groups using different types of sleep aids, in order to assess the efficacy and safety of their use among ED workers, including those working shifts. This will help in identifying robust causal associations of the possible factors that could influence the performance of these medications. The study also recommends formulating administrative strategies that can facilitate a smooth shift transition and reduce the burden of sleep disruption among healthcare professionals. On the basis of clinical, instead of subjective and self-reported, investigations, the future studies should assess the effects of such strategies on the parameters of sleep quality and day-time sleepiness. Finally, in the context of Saudi Arabia, in order to ensure the prudent use of these medications, the future research should promote the awareness and knowledge of ED physicians and nurses regarding the lack of evidence on and the possible side effects of antihistamines. In this context, it must be noted that the ED staff in Saudi Arabia attributed the negative effects of shift work on the sleep quality to circadian misalignment and increased work demands. To offset these effects, 36.6% of the respondents used or have been routinely using sleep aid medications, including antihistamines and melatonin. This use is prevalent, despite the lack of robust clinical evidence on the relationship between the routine use of these medications and their safety and efficacy in promoting sleep. Most of the participants (67.4%) had also used stimulants, mostly caffeine. Future studies should be implemented, at a national level, to assess the clinical significance of sleep aids/stimulants for shift workers. These studies should explore alternative strategies formulated by decision-makers to reduce and organize shift work, mainly the night shifts, and should test the quality of sleep using reliable and effective tools.

Abbreviations

ED, emergency department; EMT, emergency medical technicians; SCFHS, Saudi Commission for Health Specialties.

Funding

This research was funded by the Deanship of Scientific Research at Princess Nourah Bint Abdulrahman

University, Riyadh, Saudi Arabia, through the Fast-track Research Funding Program.

Disclosure

The authors report no conflicts of interest in this work.

References

- Costa C, Mondello S, Micali E, et al. Night shift work in resident physicians: does it affect mood states and cognitive levels? *J Affect Disord*. 2020;272:289–294. doi:10.1016/j.jad.2020.03.139
- Castro EAS, de Almondes KM. Sleep pattern and decision-making in physicians from mobile emergency care service with 12-h work schedules. *Int J Neurosci*. 2018;128:530–539. doi:10.1080/00207454.2017.1400970
- Izadpanah F, Nikfar S, Bakhshi Imcheh F, Amini M, Zargar M. Assessment of frequency and causes of medication errors in pediatrics and emergency wards of teaching hospitals Affiliated to Tehran University of medical sciences (24 hospitals). *J Med Life*. 2018;11:299–305.
- O'Shea J, Vu S, Siegelman J, Heron S, Lall M. "Breaking" the emergency department: does the culture of emergency medicine present a barrier to self-care? *West J Emerg Med*. 2020;21:313–321. doi:10.5811/westjem.2019.10.44584
- Harrison EM, Walbeek TJ, Maggio DG, Herring AA, Gorman MR. Circadian profile of an emergency medicine department: scheduling practices and their effects on sleep and performance. *J Emerg Med*. 2020;58:130–140. doi:10.1016/j.jemermed.2019.10.007
- Alshahrani SM, Baqays AA, Alenazi AA, AlAngari AM, AlHadi AN. Impact of shift work on sleep and daytime performance among health care professionals. *Saudi Med J*. 2017;38:846–851. doi:10.15537/smj.2017.8.19025
- Epstein M, Soderstrom M, Jirwe M, Tucker P, Dahlgren A. Sleep and fatigue in newly graduated nurses—Experiences and strategies for handling shiftwork. *J Clin Nurs*. 2020;29:184–194. doi:10.1111/jocn.15076
- Garde AH, Hansen AM, Hansen J. Sleep length and quality, sleepiness and urinary melatonin among healthy Danish nurses with shift work during work and leisure time. *Int Arch Occup Environ Health*. 2009;82:1219–1228. doi:10.1007/s00420-009-0419-4
- Bazrafshan MR, Moravveji F, Soleymanejad N, et al. Prevalence and effects of sleep disorders among shift-working nurse. *Ann Trop Med Public Health*. 2018;11:13. doi:10.4103/ATMPH.ATMPH_504_17
- Eldevik MF, Flo E, Moen BE, Pallesen S, Bjorvatn B. Insomnia, excessive sleepiness, excessive fatigue, anxiety, depression and shift work disorder in nurses having less than 11 hours in-between shifts. *PLoS One*. 2013;8(8):e70882. doi:10.1371/journal.pone.0070882
- Shy BD, Portelli I, Nelson LS. Emergency medicine residents' use of psychostimulants and sedatives to aid in shift work. *Am J Emerg Med*. 2011;29(9):1034–1036.e1031. doi:10.1016/j.ajem.2010.06.004
- Handel DA, Raja A, Lindsell CJ. The use of sleep aids among emergency medicine residents: a web-based survey. *BMC Health Serv Res*. 2006;6:136. doi:10.1186/1472-6963-6-136
- Francis MN, Wishart IM, Williamson T, Iverach R. Use of pharmacologic sleep aids and stimulants among emergency medicine staff physicians in a Canadian Tertiary care setting: a web-based survey. *Ann Emerg Med*. 2019;73:325–329. doi:10.1016/j.annemergmed.2018.07.006
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Prev Med*. 2007;45(4):247–251. doi:10.1016/j.ypmed.2007.08.012
- Ngamjaras C, Chongsuvivatwong V, McNeil E. n4Studies: sample size calculation for an epidemiological study on a smart device. *Siriraj Med J*. 2016;68:160–170.
- Thomas SJ, Calhoun D. Sleep, insomnia, and hypertension: current findings and future directions. *J Am Soc Hypertens*. 2017;11(2):122–129. doi:10.1016/j.jash.2016.11.008
- Schutte-Rodin S, Broch L, Buysse D, Dorsey C, Sateia M. Clinical guideline for the evaluation and management of chronic insomnia in adults. *J Clin Sleep Med*. 2008;4(05):487–504. doi:10.5664/jcsm.27286
- Persico N, Maltese F, Ferrigno C, et al. Influence of shift duration on cognitive performance of emergency physicians: a Prospective Cross-Sectional Study. *Ann Emerg Med*. 2018;72(2):171–180. doi:10.1016/j.annemergmed.2017.10.005
- Wolf LA, Perhats C, Delao A, Martinovich Z. The effect of reported sleep, perceived fatigue, and sleepiness on cognitive performance in a sample of emergency nurses. *J Nurs Adm*. 2017;47(1):41–49. doi:10.1097/NNA.0000000000000435
- Halm M. Night shift naps improve patient and workforce safety. *Am J Crit Care*. 2018;27:157–160. doi:10.4037/ajcc2018861
- AlShareef SM. Occupational outcomes associated with sleep quality and excessive daytime sleepiness: results from a National Survey. *Nat Sci Sleep*. 2020;12:875–882. doi:10.2147/NSS.S271154
- Richards JR, Stayton TL, Wells JA, Parikh AK, Laurin EG. Night shift preparation, performance, and perception: are there differences between emergency medicine nurses, residents, and faculty? *Clin Exp Emerg Med*. 2018;5:240–248. doi:10.15441/ceem.17.270
- Grimberg A, Kutikov JK. 142 – hypothalamus: neuroendometabolic center. In: Polin RA, Abman SH, Rowitch DH, Benitz WE, Fox WW, editors. *Fetal and Neonatal Physiology (Fifth Edition)*. Elsevier; 2017:1451–1461.e1452.
- McBeth BD, McNamara RM, Ankel FK, et al. Modafinil and zolpidem use by emergency medicine residents. *Acad Emerg Med*. 2009;16(12):1311–1317. doi:10.1111/j.1553-2712.2009.00586.x
- Culpepper L, Wingertzahn MA. Over-the-counter agents for the treatment of occasional disturbed sleep or transient insomnia: a Systematic Review of Efficacy and Safety. *Prim Care Companion CNS Disord*. 2015;17. doi:10.4088/PCC.4015r01798
- Liira J, Verbeek JH, Costa G, et al. Pharmacological interventions for sleepiness and sleep disturbances caused by shift work. *Cochrane Database Syst Rev*. 2014;12(8):Cd009776.
- Jorgensen KM, Witting MD. Does exogenous melatonin improve day sleep or night alertness in emergency physicians working night shifts? *Ann Emerg Med*. 1998;31:699–704. doi:10.1016/S0196-0644(98)70227-6
- Wright SW, Lawrence LM, Wrenn KD, Haynes ML, Welch LW, Schlack HM. Randomized clinical trial of melatonin after night-shift work: efficacy and neuropsychologic effects. *Ann Emerg Med*. 1998;32:334–340. doi:10.1016/S0196-0644(98)70010-1
- Franke AG, Bagusat C, McFarlane C, Tassone-Steiger T, Kneist W, Lieb K. The use of caffeinated substances by surgeons for cognitive enhancement. *Ann Surg*. 2015;261:1091–1095. doi:10.1097/SLA.0000000000000830
- Temple JL, Bernard C, Lipshultz SE, Czachor JD, Westphal JA, Mestre MA. The safety of ingested caffeine: a comprehensive review. *Front Psychiatry*. 2017;8:80. doi:10.3389/fpsy.2017.00080
- Lieberman HR, Agarwal S, Caldwell JA, Fulgoni VL. Demographics, sleep, and daily patterns of caffeine intake of shift workers in a nationally representative sample of the US adult population. *Sleep*. 2020;43(3):zsz240. doi:10.1093/sleep/zsz240
- Carvey CE, Thompson LA, Mahoney CR, Liberman HR. Caffeine: mechanism of action, genetics, and behavioral studies conducted in task simulators and the field. In: Wesensten N, editor. *Sleep Deprivation, Stimulant Medications, and Cognition*. Cambridge, UK: Cambridge University Press; 2012:93–107.

33. Bonnet MH, Arand DL. Studies of fatigue and human performance in the laboratory. In: Sharafkhaneh A, Hirshkowitz M, editors. *Fatigue Management: Principles and Practices for Improving Workplace Safety*. New York: Springer New York; 2018:13–27.
34. Williams GW, Shankar B, Klier EM, et al. Sensorimotor and executive function slowing in anesthesiology residents after overnight shifts. *J Clin Anesth*. 2017;40:110–116. doi:10.1016/j.jclinane.2017.04.002

Open Access Emergency Medicine

Dovepress

Publish your work in this journal

The Open Access Emergency Medicine is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of emergency medicine. The manuscript management system is completely online

and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/open-access-emergency-medicine-journal>