


Clinical and Prognostic Implications of 1p/19q, *IDH*, *BRAF*, *MGMT* Promoter, and *TERT* Promoter Alterations, and Expression of Ki-67 and p53 in Human Gliomas

Zixi Yang¹
Feng Ling¹
Sibei Ruan¹
Jiajia Hu²
Mingxi Tang¹
Xingwang Sun¹
Wenbo Long¹ 

¹Pathology Department of the First Affiliated Hospital, Southwest Medical University, Luzhou, Sichuan, People's Republic of China; ²School of Basic Medicine, Southwest Medical University, Luzhou, Sichuan, People's Republic of China

Background and Objective: Genetic alterations, including *IDH*, *BRAF*, and *TERT* promoter mutations (*IDH*-mu, *BRAF*-mu, *TERT*p-mu, respectively), 1p/19q co-deletion (1p/19q-codel), and *MGMT* promoter methylation (*MGMT*p-M), are correlated with glioma tumor development. Therefore, these genetic alterations could serve as biomarkers for the diagnosis, prognosis, and classification of gliomas, combined with the immunohistochemical markers Ki-67 and p53. However, the correlation between these alterations and the expression of Ki-67 and p53 is poorly understood.

Methods: We analyzed the prevalence and prognosis of these five alterations, as well as Ki-67 and p53 expression, in 103 primary grade II–IV gliomas via fluorescence qPCR, Sanger sequencing, fluorescence in situ hybridization, and immunohistochemistry.

Results: In the 103 cases, *MGMT*p-M was the most common alteration (70.9%), followed by *TERT*p-mu (58.3%), *IDH*-mu (46.6%), 1p/19q-codel (34.0%), and *BRAF*-mu (5.8%). No cases showed quintuple-positive alterations, but 26 cases (25.2%) showed quadruple-positive alterations (*IDH*-mu/*TERT*p-mu/*MGMT*p-M/1p/19q-codel). The percentage of *TERT*p-mu and 1p/19q-codel cases decreased with p53 expression, and the percentage of *IDH*-mu and 1p/19q-codel cases decreased with Ki-67 expression. *IDH*-mu, *MGMT*p-M, and 1p/19q-codel were positive factors for survival rates in glioma patients, while *TERT*p-mu, p53, and Ki-67 positivity were negative factors. Old age, histological grade IV, *IDH*-mu, 1p/19q-codel, Ki-67+, and p53+/Ki-67+ were significantly correlated with overall survival (OS). However, only p53+/Ki-67+ was an independent prognostic factor for OS in the multivariate Cox-model analysis.

Conclusion: *IDH*-mu only and quadruple-positivity were associated with good OS in glioma patients, while *TERT*p-mu only, *TERT*p-mu/*MGMT*p-M and p53+/Ki-67+ were associated with poor prognosis. Combining these genomic alterations and Ki-67/p53 expression should have clinical value in gliomas.

Keywords: human glioma, biomarker, correlation analysis, genetic alteration, survival rate

Introduction

Diffuse gliomas are the most common form of primary malignant brain tumor and affect adults worldwide.¹ Gliomas can be categorized as astrocytomas, oligodendrogliomas, and ependymomas and are graded II–IV according to the World Health Organization (WHO) classification, with the deadliest form being glioblastoma (GBM).² The histopathological classification of diffuse gliomas is influenced by

Correspondence: Wenbo Long
Email wenbolong@swmu.edu.cn

Received: 27 August 2021
Accepted: 8 November 2021
Published: 23 November 2021

Cancer Management and Research 2021:13 8755–8765

8755



© 2021 Yang et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at <https://www.dovepress.com/terms.php> and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (<http://creativecommons.org/licenses/by-nc/3.0/>). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (<https://www.dovepress.com/terms.php>).

interobserver variation arising from morphological classification, which can result in inconsistent histological grading and typing.³ To address this challenge, molecular markers are increasingly being developed and employed to guide the diagnosis and clinical management of gliomas.

Ki-67 and p53 are two well-studied molecules in human cancer. Ki-67 is a marker of cell division and a reliable indicator of tumor cell proliferation that is associated with the histological grade of gliomas. Although there is disagreement regarding the role of Ki-67 as a prognostic marker, many studies have revealed that it predicts poor prognosis in glioma patients.⁴ Although altered expression of the *p53* gene is not necessarily correlated with survival, immunohistochemical expression of p53 is a possible prognostic biomarker in glioma patients.⁵

Besides these traditional immunohistochemical markers, markers based on genomic variation have been developed to facilitate the diagnosis and prognosis of gliomas. Specifically, point mutations in isocitrate dehydrogenase genes (*IDH*-mu), mutations in the promoter region of the telomerase reverse transcriptase gene (*TERT*p-mu), and co-deletion of chromosome arms 1p and 19q (1p/19q-codel) can be used to classify gliomas into five principal groups according to age, survival, and associated genomic alterations.⁶ Although it is unclear how these mutations contribute to tumor development, *IDH*-mu and 1p/19q-codel are used to define oligodendroglioma (O), which is characterized by high interobserver discordance.⁷ Like *IDH*-mu, *TERT*p-mu is frequently observed in gliomas — especially primary GBM tumors — but is less frequent in low grade astrocytoma (A) tumors.⁸ As a result, *TERT*p-mu and *IDH*-mu are used in clinical settings and facilitate the identification of about 80% of GBMs from other tumor types.^{6,8,9} Methylguanine-DNA methyltransferase promoter methylation (*MGMT*p-M) has also been explored to stratify malignant gliomas into subgroups and is used as a prognostic marker for overall survival (OS) and a predictive factor for chemotherapy in GBM patients.¹⁰ The V600 mutation in the v-raf murine sarcoma oncogene homologue B1 (*BRAF*-mu) is not common in adult gliomas but frequently appears in low-grade pediatric gliomas and is highly variable across patient groups.¹¹ Given these confusing features, further research is needed to elucidate the clinical value of the *BRAF* mutation.

To date, not many studies have investigated the relationships among the various glioma markers. Though some

studies have analyzed the correlation among 1p/19q, *IDH*, *TERT*p, *MGMT*p, and *BRAF* alterations in recent years,^{6,8,9,11} there are still some controversial. Moreover, few studies discussed the combination of these alterations with the expression of p53 and Ki-67 in glioma tumors. To address this research gap, we investigated the prevalence of *MGMT*p-M, *BRAF*-mu, *IDH*-mu, *TERT*p-mu, 1p/19q-codel, and immunohistochemical expression of p53 and Ki-67 in 103 patients with grades II–IV primary gliomas. We analyzed the relationships of these molecular markers with clinicopathological characteristics and survival. We also analyzed the clinical significance of combining these markers in predicting the survival of glioma patients. The present study will contribute to an increased understanding of the molecular background of tumor types and support histological classification and prognosis of gliomas based on these markers.

Methods

Patients and Specimens

A total of 103 primary glioma samples were collected from 103 patients diagnosed with primary diffuse gliomas (grade II–IV) at the Affiliated Hospital of Southwest Medical University (Luzhou, China) between June 2017 and June 2020 ([Supplementary Table 1](#)). All patients were chemotherapy-free and targeted drug-naïve at the time of surgery. All patient data in this manuscript were coded and utilized anonymously. This study was conducted in accordance with the Declaration of Helsinki and was approved by the Institutional Research Ethics Committee of the First Affiliated Hospital of Southwest Medical University, China. Written informed consent was obtained from all patients prior to the study.

Glioma samples were formalin-fixed and paraffin-embedded (FFPE), sectioned into 4 µm thick slices, and then permanently fixed on microscope slides. After hematoxylin and eosin (H&E) staining, the slides were reviewed by two pathologists and classified according to the WHO 2016 recommendation.

DNA Extraction

Genomic DNA was extracted from 6 µm thick sections of FFPE tumor tissue using the DNA extraction kit for paraffin embedded tissues (DP331) (Tiangen, Beijing, China) according to the manufacturer's protocol. DNA concentration and quality were measured using a Nano-Drop ND-

1000 spectrophotometer (NanoDrop Technologies, Houston, USA).

IDH1, IDH2, and TERTp Mutations

Mutations in *IDH1* (R132H), *IDH2* (R172K), and *TERTp* (C228T and C250T) were determined by PCR and Sanger sequencing using detection kits (SinoMD Gene Testing Technology Co., Ltd, Beijing, China) with positive and negative controls. Sequences covering mutational hotspots in the *IDH1*, *IDH2*, and *TERT* core promoters were amplified by fluorescence qPCR. Assessments of *IDH1* (R132), *IDH2* (R172), and *TERTp* mutations were conducted using the following primer sets: *IDH1*-F: 5'-CTCCT GATGA GAAGA GGGTT G-3', R: 5'-TGGAA ATTTC TGGGC CATG-3'; *IDH2*-F: 5'-TGGAA CTATC CGGAA CATCC-3', R: 5'-AGTCT GTGGC CTTGT ACTGC; *TERTp*-F: 5'-GGCCG ATTCG ACCTC TCT, R: 5'-AGCAC CTCGC GGTA GGTAG TGG. PCR was performed in a total volume of 25 μ L comprising 2 μ L DNA in solution (10–50 ng/mL), 12.5 μ L *IDH1*, *IDH2*, or *TERTp* PCR reaction mix, 4.5 μ L primers (0.25 nM each), and 6 μ L ddH₂O. Amplification was performed using a LightCycler480 II instrument (Roche) with an initial denaturing step at 95°C for 3 min followed by 40–45 cycles of denaturation at 94°C for 15 s, annealing at 60°C for 45 s, and a final extension at 60°C for 5 min. PCR products were treated with SAP enzyme in a product: enzyme volume ratio of 5:2 at 37°C for 60 min, followed by 80°C for 15 min. After purification, products were subjected to direct sequencing on an ABI 3500 sequencer.

MGMT Promoter Methylation

A total of 1000 ng DNA were subjected to bisulfite treatment using a DNA bisulfite transformation kit (Tiangen, DP215). Methylation of *MGMTp* was determined using a detection kit (SinoMD Gene Testing Technology Co., Ltd, Beijing, China) with positive and negative controls. Primers for PCR amplification of the *MGMT* promoter region were as follows: methylated-F: 5'-GCGTT TCGAC GTTCG TAGGT-3', R: 5'-CACTC TTCCG AAAAC GAAAC G-3'; unmethylated-F: 5'-TGTGT TTTCG ATATG TTGGG ATAGT-3', R: 5'-AACTC CACAC TCTTC CAAAA ACAA-3'. PCR was performed in a total volume of 25 μ L comprising 2 μ L sample or reference DNA in solution, 12.5 μ L *MGMT* PCR reaction mix, 5.5 μ L primers (0.25 nM each), and 5 μ L ddH₂O. The PCR conditions were as follows: a denaturing step at 95°C for 3 min, 45 cycles of denaturation at 94°C for 15 s,

annealing at 60°C for 45 s, and a final extension at 60°C for 5 min. The Ct values of sample DNA amplification (Cts) and reference DNA amplification (Ctr) were recorded. When $\text{Ctr} \leq 35$ and $\text{Cts} \leq 40$, a result with $\text{Cts} - \text{Ctr} \leq 11$ was defined as positive and $\text{Cts} - \text{Ctr} > 11$ as negative; when $\text{Ctr} \leq 35$ but $\text{Cts} > 40$, the result was defined as negative; when $\text{Ctr} > 35$, the result was regarded meaningless.

BRAF V600E Mutation

Mutations in *BRAF* V600E were detected using the *BRAF* V600E fluorescence detection kit (AmoyDx Biotechnology Co., Ltd, 8.0120301X024, Xiamen, China) with positive and negative controls. Samples were amplified in a 50 μ L volume in 96-well plates in duplicate. PCR amplification comprised 45 μ L PCR reaction mix, 0.4 μ L Taq enzymes, and 4.6 μ L DNA (1–10 ng). A LightCycler480 II instrument (Roche) was used to carry out PCR as follows: 95°C for 5 min; 15 cycles of 95°C for 20 s, 64°C for 20 s, and 72°C for 20 s; 30 cycles of 93°C for 25 s, 60°C for 35 s, and 72°C for 20 s; and finally 72°C for 5 min. The Ct values of sample DNA amplification (Cts) and reference DNA amplification (Ctr) were recorded. When Ctr was between 13 and 21, a result with $\text{Cts} < 30$ was defined as positive, while $\text{Cts} \geq 30$ was defined as negative; when $\text{Ctr} > 21$, the result was regarded meaningless or that the sample DNA load should be increased.

Deletion Test of 1p/19q

New 4 μ m sections of the representative tumor areas marked by H&E staining were selected to test the status of 1p/19q. The sections were deparaffinized with xylene, rehydrated with 100%, 90%, and 70% ethanol and water for 5 min each, and then repaired with 1 \times EDTA in a steam cooker. After washing and drying, the sections were subjected to 30 min of pepsin digestion at 37°C. Paired 1p and 19q probes (AnBiPing Pharmaceutical Technology Co., Ltd, F.01081, Guangzhou, China) were added to each slide at a volume of 10 μ L and codenatured with the target DNA at 85°C for 5 min. The slides were incubated overnight at 37°C in a humidified oven and subsequently washed for 10 min with 2 \times SSC. Green and red fluorescent signals were quantified using an Olympus BX60 fluorescence microscope with the appropriate filters. For each slide, a minimum of 100 non-overlapping nuclei was assessed for the numbers of green and red signals.

Immunohistochemistry of Ki-67 and p53

To detect Ki-67 and p53 expression, mouse anti-human monoclonal primary antibodies against Ki-67 and p53 (Maixin Biotechnology Development Co., Ltd, MAB-0672, 1:1000 and MAB-0674, 1:1000, respectively, Fuzhou, China) were used according to the manufacturer's protocol. Each slide was individually reviewed and scored by two independent pathologists. The indexes of Ki-67 and p53 were determined as the percentage of immunostained cells per 200 cells in 5 fields. For analysis, p53 was scored according to Hu et al¹²: negative 0, no positive cells; score 1, weakly positive (<25% positive cells); score 2, moderately positive (25–50% positive cells); and score 3, strongly positive (>50% positive cells). Similarly, we used a 0–3 scoring system for the Ki-67 index: negative 0, <5% positive cells; score 1, weakly positive (5–25% positive cells); score 2, moderately positive (25–50% positive cells); and score 3, strongly positive (>50% positive cells). Controls without the primary antibody and positive control tissues were included in all experiments to ensure quality staining.

Statistical Analysis

The chi-square test or two-tailed Fisher's exact test was used for statistical analyses of categorical variables, such as genetic alterations, immune expression, and clinicopathological features. A *p*-value < 0.05 was considered statistically significant. OS was defined as the first day of surgery until death or the end of follow-up. Survival differences among groups were calculated using the Kaplan–Meier method with the Log rank test. We used the Cox proportional hazards model for univariate and multivariate OS analyses. SPSS software version 19.0 (IBM Armonk, NY, USA) was used for all statistical analyses.

Results

Clinicopathologic Characteristics

A total of 103 patients were included in this study, comprising 45 men (46.7%) and 58 women (56.3%) with a mean age of 48.5 ± 11.3 years (range 17–74 years) and 48.0 ± 14.8 years (range 16–83 years), respectively. There were 84 patients (81.6%) younger than 60 and only 19 patients (18.4%) older than 60. Histological classifications were done according to the 2016 revised criteria of the World Health Organization (WHO) as follows: 17 diffuse A (16.5%) cases, 19 anaplastic astrocytoma (AA) (18.4%) cases, 18 O (17.5%) cases, 12 anaplastic

oligodendroglioma (AO) (11.7%) cases, 1 oligoastrocytoma (OA) (1.0%) cases, and 36 GBM (35.0%) cases. Categorization by WHO histological grade at diagnosis yielded 35 grade II cases (34.0%), 32 grade III cases (31.1%), and 36 grade IV cases (35.0%).

Characteristics of *IDH*, *BRAF*, and *TERT* Mutations, *MGMT* Methylation, and 1p/19q Codeletion in Gliomas

A total of 45 and 3 mutations (Table 1) were observed that produced R132H of the *IDH1* gene and R172K of the *IDH2* gene, respectively, in gliomas. However, no tumors were observed with mutations in both *IDH1* and *IDH2*. The frequencies of *MGMT*-M, *BRAF*-mu, and 1p/19q-codel, but not *TERT*-mu, in *IDH*-mu cases were higher than those in *IDH*-wt cases. Unlike *TERT*-mu, *IDH*-mu, *MGMT*-M, *BRAF*-mu, and 1p/19q-codel mutations were identified with a higher frequency in grade II–III cases and a lower frequency in grade IV cases. Among the 103 cases, 5 (4.9%) showed apparent 1p deletion only, 4 (3.9%) showed apparent 19q deletion only, while 35 (34.0%) showed apparent 1p/19q-codel.

Of the 103 grade II–IV gliomas, there were no cases of quintuple-positive alterations (*IDH*-mu/*BRAF*-mu/*TERT*-mu/*MGMT*-M/1p/19q-codel). Meanwhile, 26 (25.2%), 3 (2.9%), 2 (1.9%), 1 (1.0%), and 12 (11.7%) cases showed *IDH*-mu/*TERT*-mu/*MGMT*-M/1p/19q-codel quadruple-positive alterations, *TERT*-mu/*MGMT*-M/1p/19q-codel triple-positive alterations, *IDH*-mu/*MGMT*-M/1p/19q-codel triple-positive alterations, *IDH*-mu/*TERT*-mu/1p/19q-codel triple-positive alterations, and *IDH*-mu/*MGMT*-M double-positive alterations, respectively.

Characteristics of p53 and Ki-67 Expression in Gliomas

The immunoreactivity of Ki-67, but not p53, significantly increased with increased glioma grade ($P < 0.001$, Table 2). The percentage of *TERT*-mu and 1p/19q-codel cases decreased with p53 immunoreactivity, while the frequencies of *IDH*-mu, *MGMT*-M, and *BRAF*-mu cases were unrelated to p53 immunoreactivity. Comparatively, the percentage of *IDH*-mu and 1p/19q-codel cases decreased with Ki-67 immunoreactivity, whereas the percentages of *MGMT*-M, *TERT*-mu, and *BRAF*-mu cases showed no correlation to Ki-67 immunoreactivity.

Table 1 Correlations Between Clinicopathological Characteristics and Alterations on *IDH1/2*, *MGMTp*, *TERTp*, *BRAF*, and *Ip/19q* in 103 Human Gliomas

	<i>IDH1/2</i> (%)			<i>p</i> value	<i>MGMTp</i> (%)		<i>p</i> value	<i>TERTp</i> (%)		<i>p</i> value	<i>BRAF</i> (%)		<i>p</i> value	<i>Ip/19q</i> (%)				<i>p</i> value
	<i>IDH1-mu</i>	<i>IDH2-mu</i>	<i>Wt</i>		<i>M</i>	<i>UM</i>		<i>Mu</i>	<i>Wt</i>		<i>Mu</i>	<i>Wt</i>		<i>Ip-del</i>	<i>19q-del</i>	<i>Codel</i>	<i>Wt</i>	
Sex																		
Male, n=45	19 (42.2)	1 (2.2)	25 (55.6)	0.887	29 (64.4)	16 (35.6)	0.206	28 (62.2)	17 (37.8)	0.472	2 (4.4)	43 (95.6)	0.598	2 (4.4)	1 (2.2)	14 (31.1)	28 (62.2)	0.770
Female, n=58	26 (44.8)	2 (3.4)	30 (51.7)		44 (75.9)	14 (24.1)		32 (55.2)	26 (44.8)		4 (6.9)	54 (93.1)		3 (5.2)	3 (5.2)	21 (36.2)	31 (53.4)	
Age (years)																		
< 60, n=84	41 (48.8)	3 (3.6)	40 (47.6)	0.044*	57 (67.9)	27 (32.1)	0.157	46 (54.8)	38 (45.2)	0.131	6 (7.1)	78 (92.9)	0.230	5 (5.9)	4 (4.8)	28 (33.3)	47 (56.0)	0.526
≥ 60, n=19	4 (21.1)	0 (0.0)	15 (78.9)		16 (84.2)	3 (15.8)		14 (73.7)	5 (26.3)		0 (0.0)	19 (100)		0 (0.0)	0 (0.0)	7 (36.8)	12 (63.2)	
Histology																		
A, n=17	8 (47.1)	0 (0.0)	9 (52.9)	<0.001***	12 (70.6)	5 (29.4)	0.029*	2 (11.8)	15 (88.2)	<0.001***	3 (17.6)	14 (82.4)	0.304	1 (5.9)	0 (0.0)	2 (11.8)	14 (82.4)	<0.001***
AA, n=19	11 (57.9)	0 (0.0)	8 (42.1)		10 (52.6)	9 (47.4)		6 (31.6)	13 (68.4)		1 (5.3)	18 (94.7)		1 (5.3)	1 (5.3)	3 (15.8)	14 (73.7)	
O, n=18	13 (72.2)	2 (11.1)	3 (16.7)		17 (94.4)	1 (5.6)		17 (94.4)	1 (5.6)		0 (0.0)	17 (94.4)		0 (0.0)	0 (0.0)	16 (88.9)	2 (11.1)	
AO, n=12	8 (66.7)	1 (8.3)	3 (25.0)		11 (91.7)	1 (8.3)		9 (75.0)	3 (25.0)		0 (0.0)	11 (91.7)		0 (0.0)	0 (0.0)	11 (91.7)	1 (8.3)	
OA, n=1	1 (100.0)	0 (0.0)	0 (0.0)		1 (100.0)	0 (0.0)		1 (100.0)	0 (0.0)		0 (0.0)	1 (100.0)		0 (0.0)	0 (0.0)	1 (100.0)	0 (0.0)	
GBM, n=36	4 (11.1)	0 (0.0)	32 (88.9)		22 (61.1)	14 (38.9)		25 (69.4)	11 (30.6)		2 (5.6)	34 (94.4)		3 (8.3)	3 (8.3)	2 (5.6)	28 (77.8)	
Grade																		
Grade II-III, n=67	41 (61.2)	3 (4.5)	23 (34.3)	<0.001***	51 (76.1)	16 (23.9)	0.110	35 (52.2)	32 (47.8)	0.091	4 (6.0)	63 (94.0)	0.932	2 (3.0)	1 (1.5)	33 (49.3)	34 (50.7)	<0.001***
Grade IV, n=36	4 (11.1)	0 (0.0)	15 (78.9)		22 (61.1)	14 (38.9)		25 (69.4)	11 (30.6)		2 (5.6)	34 (94.4)		3 (8.3)	3 (8.3)	2 (5.6)	28 (77.8)	
Tumor location																		
Left, n=49	24 (49.0)	1 (2.0)	24 (49.0)	0.110	39 (79.6)	10 (20.4)	0.064	28 (57.1)	21 (42.9)	0.917	2 (4.1)	47 (95.9)	0.387	3 (6.1)	1 (2.0)	19 (38.8)	26 (53.1)	0.002***
Right, n=43	20 (46.5)	2 (4.7)	21 (48.8)		29 (67.4)	14 (32.6)		26 (60.5)	17 (39.5)		4 (9.3)	39 (90.7)		1 (2.3)	0 (0.0)	15 (34.9)	27 (63.8)	
Others, n=11	1 (9.1)	0 (0.0)	10 (90.9)		5 (45.5)	6 (54.5)		6 (54.5)	5 (45.5)		0 (0.0)	11 (100.0)		1 (9.1)	3 (27.3)	1 (9.1)	6 (54.5)	
<i>IDH1/2</i>																		
<i>Mu</i> , n=48	1	1	1	1	40 (83.3)	8 (16.7)	0.009**	27 (56.3)	21 (43.8)	0.700	2 (4.2)	46 (95.8)	0.502	0 (0.0)	1 (2.1)	29 (60.4)	18 (37.5)	<0.001***
<i>Wt</i> , n=55	1	1	1		33 (60.0)	22 (40.0)		33 (60.0)	22 (40.0)		4 (7.3)	51 (92.7)		5 (9.1)	3 (5.5)	6 (10.9)	41 (74.5)	
<i>MGMTp</i>																		
<i>M</i> , n=73	37 (50.7)	3 (4.1)	33 (45.2)	0.027*	1	1	1	48 (65.8)	25 (34.2)	0.016*	2 (2.7)	71 (97.3)	0.037*	4 (5.5)	2 (2.7)	33 (45.2)	34 (46.6)	0.002***
<i>UM</i> , n=30	8 (26.7)	0 (0.0)	22 (73.3)		1	1		12 (40.0)	18 (60.0)		4 (13.3)	26 (86.7)		1 (3.3)	2 (6.7)	2 (6.7)	25 (83.3)	
<i>TERTp</i>																		
<i>M</i> , n=73	37 (50.7)	3 (4.1)	33 (45.2)	0.865	1	1	0.016*	48 (65.8)	25 (34.2)	1	2 (2.7)	71 (97.3)	0.202	4 (5.5)	2 (2.7)	33 (45.2)	34 (46.6)	<0.001***
<i>UM</i> , n=30	8 (26.7)	0 (0.0)	22 (73.3)		1	1		12 (40.0)	18 (60.0)		4 (13.3)	26 (86.7)		1 (3.3)	2 (6.7)	2 (6.7)	25 (83.3)	
<i>BRAF</i>																		
<i>Mu</i> , n=6	2 (33.3)	0 (0.0)	4 (66.7)	0.759	2 (33.3)	4 (66.7)	0.037*	2 (33.3)	4 (66.7)	0.202	1	1	1	1	0 (0.0)	0 (0.0)	5 (83.3)	0.172

(Continued)

Table 1 (Continued).

	IDH1/2 (%)		p value	MGMTp (%)		p value	TERTp (%)		p value	BRAF (%)		p value	1p/19q (%)				p value
	IDH1-mu	IDH2-mu		M	UM		Mu	Wt		Mu	Wt		1p-del	19q-del	Codel	Wt	
Wt, n=97	43 (44.3)	3 (3.1)	51 (52.6)	71 (73.2)	26 (26.8)		58 (59.8)	39 (40.2)		1	1		4 (4.1)	4 (4.1)	35 (36.1)	54 (55.7)	
1p/19q Codel, n=35	26 (74.3)	3 (8.6)	6 (17.1)	33 (94.3)	2 (5.7)		30 (85.7)	5 (14.3)		0 (0.0)	35 (100.0)	0.070	1	1	1	1	1
Others, n=68	19 (27.9)	0 (0.0)	49 (72.1)	40 (58.8)	28 (41.2)		30 (44.1)	38 (55.9)		6 (8.8)	62 (91.2)		1	1	1	1	1

Notes: Significance analysis was two-tailed, and $p < 0.05$ was considered statistically significant. * at $p < 0.05$ level, ** at $p < 0.01$ level, † no data, for no correlation analysis here.
Abbreviations: IDH1/2, isocitrate dehydrogenase 1 and 2; TERTp, telomerase reverse transcriptase promoter; MGMTp, O6-methylguanine-DNA methyltransferase promoter; 1p/19q, chromosome arms 1p and 19q; BRAF, v-raf murine sarcoma oncogene homologue B1; Mu, mutation; Wt, wild-type; M, methylation; UM, unmethylation; 1p-del, 1p-deletion; 19q-del, 19q-deletion; Codel, codeletion; A, astrocytoma; AA, anaplastic astrocytoma; AO, anaplastic oligodendroglioma; AOA, anaplastic oligoastrocytoma; O, oligodendroglioma; OA, oligoastrocytoma; GM, glioblastoma.

Relationship Between IDH-mu, TERTp-mu, MGMTp-M, 1p/19q-codel, and p53 and Ki-67 Expression with Survival Rate

Kaplan–Meier analysis showed that older patients (>60) and higher histological grade tumors had poorer OS (Figure 1A and B). For the molecular groups, patients with IDH-mu and 1p/19q-codel had significantly increased OS compared with IDH and 1p/19q wild-type patients (Figure 1D and F); OS was comparable between MGMTp-M and MGMTp-UM patients (median 18.0 vs 17.5 months) and between TERTp-mu and TERTp-wt patients (median 17.0 vs 27.0 months) (Figure 1C and E). For p53, OS was lower in p53 positive patients than in negative patients; but for Ki-67, OS was not significantly different between Ki-67 positive patients and negative patients (Figure 1G and H).

Univariate analysis using the Cox regression model showed that old age, high histological grade, IDH-mu, 1p/19q-codel, and positive Ki-67 immunostaining were significantly correlated with OS (Table 3). We combined the two factors of p53 and Ki-67 in the univariate analysis and found that double positivity of p53 and Ki-67 (p53+/Ki-67+) was significantly correlated with OS. However, in the multivariate Cox-model analysis, none of the factors mentioned above except p53+/Ki-67+ was an independent prognostic factor for OS.

Prognostic Impact Analysis of Molecular Subgroups Stratified by MGMT Promoter Methylation, IDH Mutation, TERT Promoter Mutation, 1p/19q Codeletion, Ki-67, and p53 Expression, and Their Combination

In the combined analysis, quadruple positive (MGMTp-M/IDH-mu/TERTp-mu/1p/19q-codel) and Ki-67-/p53+ molecular groups showed the best prognosis for OS in the total 103 glioma cases (Figure 2A–C), while TERTp-mu only and Ki-67+/p53+ groups showed the worst OS. In grade II–III gliomas, Ki-67+/p53+ accounted for 52.2% of total cases. In contrast, in grade IV gliomas, Ki-67+/p53+ accounted for 86.1% of total cases, while the remaining 13.9% of cases were Ki-67-/p53– group. Patients with Ki-67+/p53+ gliomas showed the worst survival rates in both grade II–III and IV gliomas (Figure 2E and H); OS of the TERTp-mu only, MGMTp-M/TERTp-mu and quadruple

Table 2 Ki-67 and p53 Immunoreactivity in 103 Human Gliomas

	p53 Cases, n (%)					Ki-67 Cases, n (%)				
	Score 0	Score 1	Score 2	Score 3	p value	Score 0	Score 1	Score 2	Score 3	p value
Grade II–III, n=67 IV, n=36	20 (29.9) 5 (13.9)	33 (49.3) 16 (44.4)	9 (13.4) 9 (25.0)	5 (7.5) 6 (16.7)	0.101	19 (28.4) 0 (0.0)	46 (68.7) 14 (38.9)	2 (3.0) 20 (55.6)	0 (0.0) 2 (5.6)	<0.001**
IDH Mu, n=48 Wt, n=55	13 (27.1) 12 (21.8)	21 (43.8) 28 (50.9)	7 (14.6) 11 (20.0)	7 (14.6) 4 (7.3)	0.516	13 (27.1) 6 (10.9)	31 (64.6) 29 (42.7)	4 (8.3) 18 (32.7)	0 (0.0) 2 (3.6)	0.003**
MGMTp M, n=73 UM, n=30	19 (26.0) 6 (20.0)	37 (50.7) 12 (40.0)	9 (12.3) 9 (30.0)	8 (11.0) 3 (10.0)	0.200	15 (20.5) 4 (13.3)	42 (57.5) 18 (60.0)	14 (19.2) 8 (26.7)	2 (2.7) 0 (0.0)	0.572
TERTp Mu, n=60 Wt, n=43	16 (26.7) 9 (20.9)	35 (58.3) 14 (32.6)	7 (11.7) 11 (25.6)	2 (3.3) 9 (20.9)	0.003**	9 (15.0) 10 (23.3)	33 (55.0) 27 (62.8)	17 (28.3) 5 (11.6)	1 (1.7) 1 (2.3)	0.211
BRAF Mu, n=6 Wt, n=97	2 (33.3) 23 (23.7)	3 (50.0) 46 (47.4)	0 (0.0) 18 (18.6)	1 (16.7) 10 (10.3)	0.671	3 (50.0) 16 (16.5)	2 (33.3) 58 (59.8)	1 (16.7) 21 (21.6)	0 (0.0) 2 (2.1)	0.230
1p/19q Codel, n=35 Others, n=68	12 (34.3) 13 (19.1)	22 (62.9) 27 (39.7)	0 (0.0) 18 (26.5)	1 (2.9) 10 (14.7)	0.001**	9 (25.7) 10 (14.7)	24 (68.6) 36 (52.9)	1 (2.9) 21 (30.9)	1 (2.9) 1 (1.5)	0.011*

Notes: Significance analysis was two-tailed, and $p < 0.05$ was considered statistically significant. * at $p < 0.05$ level, ** at $p < 0.01$ level.

Abbreviations: IDH, isocitrate dehydrogenase; TERTp, telomerase reverse transcriptase promoter; MGMTp, O6-methylguanine-DNA methyltransferase promoter; 1p/19q, chromosome arms 1p and 19q; BRAF, v-raf murine sarcoma oncogene homologue B1; Mu, mutation; Wt, wild-type; M, methylation; UM, unmethylation; Codel, codeletion.

negative (*MGMTp*-UM/*IDH*-wt/*TERTp*-wt/1p/19q-wt) groups was worse than that of the other groups (Figure 2C, F and I). Quadruple positive status was associated with a good OS no matter the Ki-67 or p53 status in grade II–III gliomas (Figure 2D). However, the quadruple positive group was absent in grade IV gliomas (Figure 2G), while *IDH*-mu was only associated with a good prognosis for OS in both grade II–III and IV gliomas (Figure 2D and G). Out of the 36 grade IV gliomas, only four were *IDH*-mu (11.1%), thus failing to change the poor prognosis of GBMs.

Discussion

The present study investigated whether combined analysis of genetic alterations in *IDH*, *BRAF*, *TERTp*, *MGMTp*, and 1p/19q along with expression changes in p53 and Ki-67 protein is associated with the biological behaviors of gliomas. The results demonstrated significant associations of *IDH*-mu, *BRAF*-mu, *TERTp*-mu, *MGMT*-M, 1p/19q-codel, and p53 and Ki-67 expression with the survival rate of glioma patients.

Previous studies have reported that genetic alterations in *IDH*, *BRAF*, *TERTp*, *MGMTp*, and 1p/19q are associated with histological classification of gliomas.^{6,10,11} *IDH*-mu is present in 50–80% of WHO grade II–III gliomas and secondary GBM but is rare in primary GBM.¹³ Consistent with the reports of these studies, the present study identified 41 *IDH1*-mu and 3 *IDH2*-mu cases in 67 grade II–III gliomas, but only 4 *IDH1*-mu cases in 36 primary GBMs. No cases showed concurrent *IDH1*-mu and *IDH2*-mu, which could be because the two mutations are mutually exclusive.¹⁴ In contrast, we observed a high frequency of *TERTp*-mu and *MGMTp*-M in both grade II–III and IV gliomas, especially oligodendroglial gliomas, in accordance with previous reports. These findings suggest that *TERTp* mutation and *MGMT* methylation are typical molecular characteristics of these tumors.^{15–18} Concurrent loss of 1p/19q and *IDH*-mu has been used to define O tumors and resolve unclear diagnoses due to high inter-observer discordance.^{7,19} The present study demonstrated that 1p/19q-codel is highly concentrated in oligodendrocyte-derived gliomas and strongly associated with *IDH*-mu




Figure 1 Prognostic values of patient age, histological grade, *MGMT* promoter methylation, *IDH* mutation, *TERT* promoter mutation, 1p/19q codeletion, and Ki-67 and p53 expressions by Kaplan-Meier Estimates of Overall Survival in 103 gliomas.

Notes: Older age (>60 years) (A), higher histological grade (B), and overexpression Ki-67 (G) and p53 (H) predicted worse survival rates in glioma patients, while *MGMT* promoter methylation (*MGMTp*-M) (C), *IDH* mutation (*IDH*-mu) (D), *TERT* promoter mutation (*TERTp*-mu) (E), and 1p/19q codeletion (1p/19q-codel) (F) predicted better survival rates. Significance analysis was two-tailed, and $p < 0.05$ was considered statistically significant.

Abbreviations: *IDH*, isocitrate dehydrogenase; *TERTp*, telomerase reverse transcriptase promoter; *MGMTp*, O6-methylguanine-DNA methyltransferase promoter; 1p/19q-codel, chromosome arms 1p and 19q codeletion; *BRAF*, v-raf murine sarcoma oncogene homologue B1; Mu, mutation; Wt, wild-type; M, methylation; UM, unmethylation.

in O tumors. However, 1 AO and 2 O cases were found with intact 1p/19q and *MGMTp*-M, confirming that parts of 1p/19q non-codeleted oligodendroglial gliomas were not astrocytic.²⁰ Moreover, some cases showed conflicting results: two O cases showed *IDH*-wt/1p/19q-wt, but two A, three AA, and two GBM cases showed 1p/19q-codel, suggesting limitations of the existing molecular classification system for tumors with complex karyotypes.²¹

IDH1 mutation is a prognostic indicator of better survival in patients with diffuse gliomas grade II–IV tumors, with the exception of recurrent grade IV tumors.²² Similarly, we found that *IDH*-mu cases showed good prognosis for OS in both grade II–III and IV tumors, which was better than most other molecular subgroups. However, *IDH*-mu frequently appeared along with *TERTp*-mu/*MGMTp*-M/1p/19q-codel (54.2%) and *MGMTp*-M (25.0%) in *IDH* mutated cases. Thus, predicting OS according to *IDH* mutation only without considering other alterations may not be very reliable. Although *TERTp*-mu gliomas were associated with worse survival than *TERTp*-wt gliomas,⁶ patients with *TERTp*-mu/*IDH*-mu gliomas showed better survival than patients with one or none of these two mutations.⁸ In the present study, we did not predict the survival of the patient with *TERTp*-mu/*IDH*-mu glioma, as there was only one such case;

however, we investigated whether *TERTp*-mu is associated with a poor prognosis in gliomas, similar to *MGMTp*-M/*TERTp*-mu. Interestingly, the occurrence of *IDH*-mu was significantly correlated with Ki-67 expression, but not with p53 expression, which is opposite to the finding of *TERTp*-mu, revealing that *IDH* and *TERTp* mutations may influence gliomagenesis through different oncogenic pathways. Ogura et al reported that p53 overexpression is associated with unfavorable outcomes in GBM/AA patients with *IDH1*-wt/*MGMTp*-UM gliomas,²³ suggesting the combined profile of p53 and *IDH* and *MGMTp* status. However, we found that p53 positivity is not significantly correlated with survival in patients with *IDH1*-wt/*MGMTp*-UM gliomas, but that p53+/Ki-67+ predicts unfavorable outcomes in these patients. Thus, the combined assessment of *IDH* and *MGMTp* status with p53 and Ki-67 expression may be more useful for the prognosis of astrocyte-derived gliomas.

p53 expression was previously reported as a potential predictor of OS in glioma patients, but other studies reported contrasting findings.^{23,24} Our study demonstrated that patients with a stronger correlation between positive p53 expression and a worse prognosis than for patients with negative p53 expression. However, p53 expression was not an independent prognostic factor in neither univariate nor multivariate analysis. This finding

Table 3 Clinicopathological and Biological Factors Affecting Overall Survival Rates

	Univariate		p value	Multivariate		p value
	OR	95% CI		OR	95% CI	
Sex (male vs female)	0.640	0.265–1.546	0.321	0.571	0.192–1.698	0.313
Age (≤ 60 y vs > 60 y)	3.300	1.166–9.341	0.025*	3.0182	0.817–11.112	0.098
Histological grade (II–III vs IV)	4.073	1.620–10.240	0.003**	1.034	0.298–3.585	0.958
IDH (wt vs mu)	0.174	0.060–0.509	0.001**	0.394	0.100–1.544	0.181
MGMTp (UM vs M)	0.607	0.239–1.544	0.295	0.834	0.260–2.674	0.760
TERTp (wt vs mu)	1.619	0.646–4.059	0.304	1.431	0.442–4.634	0.550
BRAF (wild-type vs mutated)	0.546	0.061–4.897	0.589	0.727	0.058–9.165	0.805
1p/19q-codel (others vs codel)	0.252	0.079–0.802	0.020*	0.399	0.083–1.905	0.249
p53 (negative vs positive)	3.259	0.890–11.942	0.075	3.074	0.725–13.034	0.128
Ki-67 (negative vs positive)	8.069	1.022–63.698	0.048*	3.400	0.360–32.120	0.286
p53/Ki-67 (others vs both positive)	6.476	1.796–23.353	0.004**	5.173	1.271–21.058	0.022*

Notes: Significance analysis was two-tailed, and $p < 0.05$ was considered statistically significant. * at $p < 0.05$ level, ** at $p < 0.01$ level.

Abbreviations: IDH, isocitrate dehydrogenase; TERTp, telomerase reverse transcriptase promoter; MGMTp, O6-methylguanine-DNA methyltransferase promoter; 1p/19q, chromosome arms 1p and 19q; BRAF, v-raf murine sarcoma oncogene homologue B1; Mu, mutation; Wt, wild-type; M, methylation; UM, unmethylation; Codel, codeletion; OR odds ratio; CI confidence interval.

may be because the cases with strong expression of p53 (p53, 3+) in this study had the highest percentage of IDH-mu (63.6%) but the lowest percentage of TERTp-mu (18.2%) compared to the other groups and thus achieved good OS due to IDH-mu status. Expression of p53 was not an independent prognostic factor for OS, indicating that further research is needed to determine the exact clinical value of p53. Ki-67 is another well-known cancer cell proliferation indicator, and its expression can reflect the proliferation and malignancy of glioma cells.⁴ In the present study, it was observed that most Ki-67+ glioma cases were high-grade and associated with poor OS. However, Ki-67+ was an independent prognostic factor only in the univariate analysis, not in the multivariate analysis. Although Ki-67 may be predictive of poor prognosis, using Ki-67 alone to predict survival in glioma patients may not be reliable.^{4,25} Instead, double positivity of Ki-67 and p53 (Ki-67+/p53+) was investigated as an independent prognostic factor for OS in both the univariate and multivariate analyses. Patients with Ki-67+/p53+ gliomas showed bad prognosis in terms of OS in both grade II–III and IV gliomas, suggesting the need to triage glioma patients based on the molecular status of Ki-67 and p53.

GBM is considered the deadliest primary brain tumor. It is characterized by high frequencies of MGMT and TERT promoter alterations and low frequencies of IDH and 1p/19q mutations in the primary tumor.²⁶ Nevertheless, part of GBMs lack established genetic biomarkers, hence hard to classify into molecular subgroups and prognose OS for

patients. For example, although TERTp and IDH mutations are clinically used to classify most of GBMs, about 20% of GBMs do not fit into this molecularly defined system and are designated IDH-wt/TERTp-wt GBM subgroups.²⁷ In line with previous reports, we found 7 GBMs that were IDH-wt/TERTp-wt, of which 1 case had 1p/19q-codel, 3 cases had triple-negative (IDH-wt/TERTp-wt/1p/19q-wt), and the remaining cases had quadruple-negative (IDH-wt/TERTp-wt/MGMTp-UM/1p/19q-wt). Considering that the majority of GBMs were Ki-67+/p53+ and the rarity of BRAF-mu, Ki-67 and p53 expression and BRAF status is unlikely to aid in distinguishing TERTp/IDH/1p/19q wild-type GBMs from other gliomas. Furthermore, the TERTp/IDH/1p/19q wild-type GBMs cannot be distinguished from other genetic types of GBM by pathological images, highlighting the need to develop new markers to assist in defining subgroup of TERTp/IDH/1p/19q wild-type GBMs.

In conclusion, our findings confirmed existence of complex correlations among TERTp and IDH mutations, MGMTp-M, 1p/19q-codel, and expression of p53 and Ki-67 in human gliomas. BRAF mutation was uncommon in adult astrocytic and oligodendroglial gliomas. IDH-mu only and quadruple-positivity (IDH-mu/TERTp-mu/MGMTp-M/1p/19q-codel) predicted good prognosis for OS in gliomas; TERTp-mu only, TERTp-mu/MGMTp-M and quadruple-negativity predicted poor prognosis, especially in grade II–III gliomas. Ki-67/p53 was an independent prognostic factor for OS, as double positivity predicted poor prognosis in both II–III and IV gliomas. Thus, combining these genome alterations and Ki-67/p53




Figure 2 Kaplan-Meier estimates of overall survival in the glioma molecular groups.

Notes: Survival of the molecular subgroups stratified according to MGMT promoter methylation (MGMTp-M), IDH mutation (IDH-mu), TERT promoter mutation (TERTp-mu), and 1p/19q codeletion (1p/19q-codel) in a total of 103 gliomas (A), comprising 67 grade II–III gliomas (D) and 36 grade IV gliomas (G). 1, quadruple positive (MGMTp-M/IDH-mu/TERTp-mu/1p/19q-codel); 2, IDH-mu only; 3, MGMTp-M/IDH-mu; 4, MGMTp-M only; 5, others; 6, quadruple negative (MGMTp-UM/IDH-wt/TERTp-wt/1p/19q-wt); 7, MGMTp-M/TERTp-mu; 8, TERTp-mu only. Survival of the subgroups stratified according to Ki-67 and p53 overexpression in a total of 103 gliomas (B), comprising 67 grade II–III gliomas (E) and 36 grade IV gliomas (H). Survival of the subgroups stratified according to MGMTp-M, IDH-mu, TERTp-mu, and 1p/19q-codel in a total of 66 Ki-67+/p53+ gliomas (C), comprising 35 Ki-67+/p53+ grade II–III gliomas (F) and 31 Ki-67+/p53+ grade IV gliomas (I). Significance analysis was two-tailed, and $p < 0.05$ was considered statistically significant.

Abbreviations: IDH, isocitrate dehydrogenase; TERTp, telomerase reverse transcriptase promoter; MGMTp, O6-methylguanine-DNA methyltransferase promoter; 1p/19q-codel, chromosome arms 1p and 19q codeletion; BRAF, v-raf murine sarcoma oncogene homologue B1; Mu, mutation; Wt, wild-type; M, methylation; UM, unmethylation.

expression should have clinical value in the diagnosis and prognosis of glioma tumors.

Abbreviations

IDH, isocitrate dehydrogenase; TERTp, telomerase reverse transcriptase promoter; MGMTp, O6-methylguanine-DNA methyltransferase promoter; 1p/19q, chromosome arms 1p and 19q; BRAF, v-raf murine sarcoma oncogene homologue B1; Mu, mutation; Wt, wild-type; M, methylation; UM, unmethylation; 1p-del, 1p-deletion; 19q-del, 19q-

deletion; Codel, codeletion; A, astrocytoma; AA, anaplastic astrocytoma; AO, anaplastic oligodendroglioma; AOA, anaplastic oligoastrocytoma; O, oligodendroglioma; OA, oligoastrocytoma; GM, glioblastoma; OS, overall survival.

Acknowledgments

This study was supported by two joint research projects of Southwest Medical University and Luzhou city (2017LZXNYD-J14 and 2018LZXNYD-ZK22), Natural Science Foundation of Southwest Medical University

(2019ZQN070), as well as the doctoral research initiation fund of the Affiliated Hospital of Southwest Medical University (no. 16238).

Author Contributions

All authors contributed to data analysis, drafting, or revising the article. All authors gave final approval of the manuscript to be published, agreed to the submitted journal, and agreed to be accountable for all aspects of the work.

Disclosure

The authors declare that they have no conflicts of interest in this work.

References

- Labreche K, Kinnersley B, Berzero G, et al. Diffuse gliomas classified by 1p/19q co-deletion, *TERT* promoter and *IDH* mutation status are associated with specific genetic risk loci. *Acta Neuropathol.* 2018;135(5):743–755.
- Pienkowski T, Kowalczyk T, Kretowski A, Ciborowski M. A review of gliomas-related proteins. Characteristics of potential biomarkers. *Am J Cancer Res.* 2021;11(7):3425–3444.
- van den Bent MJ. Interobserver variation of the histopathological diagnosis in clinical trials on glioma: a clinician's perspective. *Acta Neuropathol.* 2010;120(3):297–304.
- Chen WJ, He DS, Tang RX, Ren FH, Chen G. Ki-67 is a valuable prognostic factor in gliomas: evidence from a systematic review and meta-analysis. *Asian Pac J Cancer Prev.* 2015;16(2):411–420.
- Jin Y, Xiao W, Song T, Feng G, Dai Z. Expression and prognostic significance of p53 in glioma patients: a meta-analysis. *Neurochem Res.* 2016;41(7):1723–1731.
- Eckel-Passow JE, Lachance DH, Molinaro AM, et al. Glioma groups based on 1p/19q, *IDH*, and *TERT* promoter mutations in tumors. *N Engl J Med.* 2015;372(26):2499–2508.
- Ammendola S, Caldonazzi N, Simbolo M, et al. H3K27me3 immunostaining is diagnostic and prognostic in diffuse gliomas with oligodendroglial or mixed oligoastrocytic morphology. *Virchows Arch.* 2021. doi:10.1007/s00428-021-03134-1
- Vuong HG, Altibi AMA, Duong UNP, et al. *TERT* promoter mutation and its interaction with *IDH* mutations in glioma: combined *TERT* promoter and *IDH* mutations stratifies lower-grade glioma into distinct survival subgroups-A meta-analysis of aggregate data. *Crit Rev Oncol Hematol.* 2017;120:1–9.
- Killela PJ, Pirozzi CJ, Healy P, et al. Mutations in *IDH1*, *IDH2*, and in the *TERT* promoter define clinically distinct subgroups of adult malignant gliomas. *Oncotarget.* 2014;5(6):1515–1525.
- Schulze Heuling E, Knab F, Radke J, et al. prognostic relevance of tumor purity and interaction with *MGMT* methylation in glioblastoma. *Mol Cancer Res.* 2017;15(5):532–540.
- Galbraith K, Snuderl M. Molecular pathology of gliomas. *Surg Pathol Clin.* 2021;14(3):379–386.
- Hu X, Miao W, Zou Y, Zhang W, Zhang Y, Liu H. Expression of p53, epidermal growth factor receptor, Ki-67 and O6-methylguanine-DNA methyltransferase in human gliomas. *Oncol Lett.* 2013;6(1):130–134.
- Ichimura K, Pearson DM, Kocalkowski S, et al. *IDH1* mutations are present in the majority of common adult gliomas but rare in primary glioblastomas. *Neuro Oncol.* 2009;11(4):341–347.
- Hartmann C, Meyer J, Balss J, et al. Type and frequency of *IDH1* and *IDH2* mutations are related to astrocytic and oligodendroglial differentiation and age: a study of 1010 diffuse gliomas. *Acta Neuropathol.* 2009;118(4):469–474.
- Labussière M, Di Stefano AL, Gleize V, et al. *TERT* promoter mutations in gliomas, genetic associations and clinico-pathological correlations. *Br J Cancer.* 2014;111(10):2024–2032.
- Weller M, Stupp R, Reifenberger G, et al. *MGMT* promoter methylation in malignant gliomas: ready for personalized medicine? *Nat Rev Neurol.* 2010;6(1):39–51.
- Yang P, Cai J, Yan W, et al. Classification based on mutations of *TERT* promoter and *IDH* characterizes subtypes in grade II/III gliomas. *Neuro Oncol.* 2016;18(8):1099–1108.
- Wesseling P, van den Bent M, Perry A. Oligodendroglioma: pathology, molecular mechanisms and markers. *Acta Neuropathol.* 2015;129(6):809–827.
- Yao J, Hagiwara A, Raymond C, et al. Human *IDH* mutant 1p/19q co-deleted gliomas have low tumor acidity as evidenced by molecular MRI and PET: a retrospective study. *Sci Rep.* 2020;10(1):11922.
- Li YX, Shi Z, Aibaidula A, et al. Not all 1p/19q non-codeleted oligodendroglial tumors are astrocytic. *Oncotarget.* 2016;7(40):64615–64630.
- Ballester LY, Huse JT, Tang G, Fuller GN. Molecular classification of adult diffuse gliomas: conflicting *IDH1/IDH2*, *ATRX*, and 1p/19q results. *Hum Pathol.* 2017;69:15–22.
- Tabai Y, Kobayashi K, Saito K, et al. Survival in patients with glioblastoma at a first progression does not correlate with isocitrate dehydrogenase (*IDH1*) gene mutation status. *Jpn J Clin Oncol.* 2021;51(1):45–53.
- Ogura R, Tsukamoto Y, Natsumeda M, et al. Immunohistochemical profiles of *IDH1*, *MGMT* and P53: practical significance for prognosis of patients with diffuse gliomas. *Neuropathology.* 2015;35(4):324–335.
- Levidou G, El-Habr E, Saetta AA, et al. P53 immunoexpression as a prognostic marker for human astrocytomas: a meta-analysis and review of the literature. *J Neuro-Oncol.* 2010;100(3):363–371.
- Alkhaibary A, Alassiri AH, AlSufiani F, Alharbi MA. Ki-67 labeling index in glioblastoma; does it really matter? *Hematol Oncol Stem Cell Ther.* 2019;12(2):82–88.
- Thakkar JP, Dolecek TA, Horbinski C, et al. Epidemiologic and molecular prognostic review of glioblastoma. *Cancer Epidemiol Biomarkers Prev.* 2014;23(10):1985–1996.
- Diplas BH, He X, Brosnan-Cashman JA, et al. The genomic landscape of *TERT* promoter wildtype-*IDH* wildtype glioblastoma. *Nat Commun.* 2018;9(1):2087.

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>