

Effect of interaction of magnetic nanoparticles of Fe_3O_4 and artesunate on apoptosis of K562 cells

Ying Wang¹
Yuxiang Han¹
Yingying Yang¹
Jingci Yang¹
Xiaonan Guo¹
Jingnan Zhang¹
Ling Pan¹
Guohua Xia²
Baohan Chen²

¹Department of Hematology, the 2nd Hospital of Hebei Medical University, People's Republic of China;

²Department of Hematology, the Affiliated Zhongda Hospital, Medical School, Southeast University, Nanjing, People's Republic of China

Abstract: The present study evaluated whether the magnetic nanoparticles of Fe_3O_4 (MNPs- Fe_3O_4) could enhance the activity of artesunate (ART), and to explore its potential mechanisms. Cytotoxicity of the copolymer of ART with MNPs- Fe_3O_4 on K562 cells was detected by MTT assay and the apoptosis rate of K562 cells was measured by flow cytometry. Protein expression levels of bcl-2, bax, bcl-rambo, caspase-3, and survivin in K562 cells were measured by Western blot. After being incubated with the copolymer of ART with MNPs- Fe_3O_4 for 48 hours, the growth inhibition rate of K562 cells was significantly increased compared with that of K562 cells treated with ART alone ($P < 0.05$), and the apoptosis rate of K562 cells was increased significantly compared with that of K562 cells treated with ART alone, suggesting that MNPs- Fe_3O_4 can enhance the activity of ART. Interestingly, the copolymer-induced cell death was attenuated by caspase inhibitor Z-VAD-FMK. Our results also showed that treatment with the copolymer of MNPs- Fe_3O_4 and ART increased the expression of bcl-2, bax, bcl-rambo, and caspase-3 proteins, and decreased the expression of survivin protein in K562 cells compared with ART treatment alone. These results suggest that MNPs- Fe_3O_4 can enhance ART-induced apoptosis, which may be related to the upregulation of bcl-rambo and downregulation of survivin.

Keywords: magnetic nanoparticles, MNPs- Fe_3O_4 , artesunate, K562 cells, apoptosis, bcl-rambo

Introduction

Artesunate (ART), a semi-synthetic derivative of artemisinin extracted from the Chinese herb *Artemisia annua*, is a safe and effective antimalarial drug. Recently, several studies showed that ART could inhibit the growth of various carcinoma cell lines¹⁻³ and suppress tumor cell proliferation by inducing apoptosis,⁴ indicating that ART may be a promising novel candidate for leukemia therapy. However, a variety of lines of evidence indicate that ART inhibited cell proliferation in a dose-dependent manner, and a high concentration of ART induced embryo toxicity,⁵⁻⁷ reversible renal damage toxicity,⁸ and toxicity to bone development,⁹ which limit the usefulness of ART.

To minimize side effects associated with dosage during chemotherapy, a promising approach is to combine a conventional chemotherapy with new strategies to maximize efficacy of tumor treatment through inducing cell apoptosis. Recent studies demonstrate that nanotechnologies and nanoparticles, which have unique peculiarities such as stable biomolecular absorption, high surface energy, and small sizes comparable to those of biomolecules, are becoming a focus for human medical application, including the delivery of antineoplastic agents to cells and tissues.^{10,11} We have recently reported

Correspondence: Ling Pan
Department of Hematology,
The 2nd Hospital of Hebei Medical
University, Hebei Institute of Hematology,
Shijiazhuang 050000,
People's Republic of China
Tel +86 18980601818
Fax +86 02885423921
Email lingpan20002000@yahoo.com.cn

that MNPs-Fe₃O₄, as an anticancer drug deliverer, could enhance the sensitivity of anticancer drugs and reverse the drug resistance of tumor cells.^{12,13}

Apoptosis is the main mechanism of cell death, and is mediated by a cell-intrinsic suicide program, with the relative balance of pro- and anti-apoptotic signaling pathways determining the fate of the cell. There are two main pathways in mammals. One is the death receptor signaling pathway,¹⁴ which is regulated by the IAP family including mainly X-IAP, c-IAP1, c-IAP2, KIAP, and survivin. Another pathway is the mitochondrial signaling pathway, in which bcl-2 plays an important role and bcl-2/bax is the key point of cell apoptosis.¹⁵ Recently, bcl-rambo, a novel member of bcl-2 family, has been shown to induce apoptosis independent of both bcl-2/bax ratio and the death receptor pathway, but this effect can be specifically blocked by the IAP family.¹⁶ The final executor of apoptosis is the caspase family, particularly caspase-3, which is the central proteases for apoptosis.¹⁷

Accordingly, the copolymer of ART with MNPs-Fe₃O₄ may provide a novel clinical candidate for the treatment of leukemia. Here, the sensitivity of K562 cells to this copolymer and its possible underlying mechanisms were investigated.

Materials and methods

Main materials

The following materials were purchased: 3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT) (Amresco, Solon, OH); Z-VAD-FMK (Beyotime Biotechnology); RPMI 1640 medium and fetal calf serum (GIBCO, USA); artesunate (Guilin Pharmaceutical Co., Ltd., China); Annexin V-FITC (Biosea Biotechnology Co., Ltd., Beijing, China); total protein extraction kit P1250 (Applygen Technologies Inc., Beijing, China); BCA protein assay kit (Biosynthesis Biotechnology Co., Ltd., Beijing, China); anti-bcl-2, anti-bax, anti-survivin (Bioworld), anti-bcl-rambo (Biovision), anti-active caspase-3 (Cell Signaling), anti-GAPDH (Santa Cruz, CA), and goat-antirabbit HRP-IgG (Santa Cruz, CA); Western chemiluminescent HRP substrate (Millipore Corporation, Billerica, MA). MNPs-Fe₃O₄ was kindly supplied by the state Key Laboratory of Bioelectronics, Southeast University, Nanjing, China, and was well distributed in 0.01 M phosphate-buffered saline (PBS) by ultrasound to obtain MNPs-Fe₃O₄ colloidal suspension. ART at different concentrations (12.5, 25, 50, 75, and 100 μ M) was conjugated with MNPs-Fe₃O₄, at the molar ratios of 25:1, 50:1, and 100:1 by

mechanical absorption polymerization at 4°C for 12 hours, as previously reported.^{18,19}

Cell line and cell culture

K562 cells, derived from a patient with chronic myeloid leukemia in blast crisis, had been constantly preserved in our laboratory, and were cultured in a flask in RPMI 1640 medium supplemented with 10% fetal calf serum, 100 U/mL penicillin, and 100 μ g/mL streptomycin at 37°C in a humidified 5% CO₂ incubator.

Cytotoxicity assays

K562 cells (2×10^4 cells per well) in logarithmic growth phase were cultured in 96-well flat-bottomed plates in a triplicate pattern and treated with different concentrations of ART, MNPs-Fe₃O₄, or the copolymer of ART with MNPs-Fe₃O₄ in the presence or absence of 40 μ M Z-VAD-FMK for 48 hours. MTT (20 μ L, 5 mg/mL) was added to each well and incubated for 4 hours. Then, 200 μ L of dimethylsulfoxide was added to each well and the plate was vortexed for 10 minutes at 37°C. Finally, the optical density value (A) of each well was measured at a measurement wavelength of 540 nm using a plate reader (Model 550, BIO-RAD, Japan). Cell growth inhibition ratio was calculated as $(1 - A_{540} \text{ of experimental well} / A_{540} \text{ of blank control well}) \times 100\%$. Each assay was repeated at least 3 times.

Apoptosis assay by flow cytometry

Based on the results of cytotoxicity assay, the best combination condition of ART with MNPs-Fe₃O₄ (100:1 M/M) was used to measure the apoptosis rate. Briefly, after being treated with ART, MNPs-Fe₃O₄ or the copolymer of MNPs-Fe₃O₄ and ART for 48 hours, K562 cells were collected, washed twice with ice-cold PBS, and then cells were suspended in 200 μ L of binding buffer and 10 μ L of Annexin V-FITC for 15 minutes in the dark. Thereafter, 300 μ L of binding buffer and 5 μ L of propidium iodide were added to each sample. Finally, the cells were analyzed using BD FACSDiva flow cytometry (BD FACS Canto™ II) with Cell Quest software.

Western blot analysis

After K562 cells were treated as mentioned above, total protein was extracted and quantified according to the manufacturer's protocol. Each equal amount of protein was loaded on sodium dodecyl sulfate polyacrylamide gel, run at 100 volts for 2 hours, and then the protein was transferred to a polyvinylidene fluoride membrane. The membranes were blocked in 5% fat-free milk at room temperature for 2 hours,

and the blots were stained with specific primary antibodies, including anti-bcl-2, anti-bax, anti-survivin, anti-bcl-rambo, anti-active caspase-3, and anti-GAPDH antibodies overnight at 4°C. Then, the membranes were washed and incubated with goat-antirabbit HRP-IgG for 1 hour at room temperature, and imaged with a chemiluminescent substrate. Thereafter, the bound immunoglobulins were removed from the membranes by washing twice with restore western blot stripping buffer, and the signal was detected using the ChemiDoc XR+ (BIO-RAD) with the enhanced chemiluminescence and analyzed by the Image Lab (ECL). GAPDH was used as the internal control.

Statistical analysis

All experiments were repeated 3 times. Data are presented as mean \pm standard deviation and analyzed using the SPSS (Release 13.0; SPSS Inc, Chicago, IL). One-way analysis of variance (ANOVA), 2-sample *t*-test, or rank test was performed to illustrate the significant difference between the treated groups and the control. The difference was considered statistically significant at $P < 0.05$.

Results

Inhibition of cell growth

When K562 cells were treated with the copolymer of ART and MNPs- Fe_3O_4 at the molar ratio of 100:1, the cell viability was decreased significantly compared with that of K562 cells treated with ART alone (0.47 ± 0.11 vs 1.10 ± 0.02) ($P < 0.05$). Although the cell viability of K562 cells was reduced to 0.66 ± 0.03 at a molar ratio of 25:1 and 0.73 ± 0.04 at a molar ratio of 50:1, there were no significant differences between the copolymer and ART alone group ($P > 0.05$), suggesting that the molar ratio of 100:1 was the best condition for copolymerization.

Moreover, when ART at the concentrations of 75, or 100 μM was combined with MNPs- Fe_3O_4 (M/M = 100:1) cell proliferation was remarkably inhibited, especially at the concentration of 100 μM , compared with ART alone ($P < 0.05$) (Figure 1); however there was no significant difference at the concentrations of 12.5, 25, and 50 μM ($P > 0.05$). The cell growth inhibition in the MNPs- Fe_3O_4 group was less than that in the ART group with or without MNPs- Fe_3O_4 . In addition, it was noted that MNPs- Fe_3O_4 alone could slightly inhibit cell proliferation when its concentration was from 0.125 μM to 4 μM , but there was no significant difference ($P > 0.05$).

To further verify that caspase activation is necessary for the copolymer-induced apoptosis, K562 cells were

treated with the copolymer of ART and MNPs- Fe_3O_4 in the presence or absence of general caspase inhibitor Z-VAD-FMK. Cell survival rate was measured to see whether the antiproliferative effect of the copolymer was preventable. As shown in Figure 2, the copolymer-induced cell death was attenuated by Z-VAD-FMK, since cell survival rate in the presence of Z-VAD-FMK was increased more than its absence ($58.73 \pm 0.37\%$ vs $23.67 \pm 1.45\%$) ($P < 0.05$), suggesting that in combination with MNPs- Fe_3O_4 , ART-induced apoptosis proceeded via the caspase-dependent pathway.

Enhancement of apoptosis by the copolymer of ART with MNPs- Fe_3O_4

After being incubated with ART, MNPs- Fe_3O_4 , or the copolymer of ART with MNPs- Fe_3O_4 (M/M = 100:1) for 48 hours, the proportion of apoptotic cells in the copolymer treated group was increased significantly compared with that in 100 μM ART-treated group ($78.02 \pm 10.83\%$ vs $48.90 \pm 14.08\%$) ($P < 0.05$) (Figure 3).

However, there were no significant differences in the apoptosis rate of K562 cells between the copolymer of 12.5 μM ART with 0.125 μM MNPs- Fe_3O_4 -treated group and the 12.5 μM ART-treated group ($28.10 \pm 11.66\%$ vs $17.64 \pm 4.49\%$); between the copolymer of 25 μM ART with 0.25 μM MNPs- Fe_3O_4 -treated group and the 25 μM ART-treated group ($40.54 \pm 13.45\%$ vs $25.76 \pm 8.80\%$); between the copolymer of 50 μM ART with 0.5 μM MNPs- Fe_3O_4 -treated group and the 50 μM ART alone group ($44.95 \pm 13.66\%$ vs $38.98 \pm 14.57\%$); and between the copolymer of 75 μM ART with 0.75 μM MNPs- Fe_3O_4 -treated group and the 75 μM ART alone group ($43.9 \pm 5.53\%$ vs $38.53 \pm 2.10\%$) ($P > 0.05$) (Figures 3 and 4). However, the apoptosis rate in all groups was higher than that in both MNPs- Fe_3O_4 and control groups ($P < 0.05$). These results to some extent indicate that MNPs- Fe_3O_4 may enhance ART-induced cell apoptosis.

Expression of bcl-2, bax, bcl-rambo, caspase-3, and survivin

Western blot assay was performed to trace the changes of apoptosis-related genes on treatment with MNPs- Fe_3O_4 , ART, or the copolymer of ART with MNPs- Fe_3O_4 for 48 hours. It was observed that the expression of survivin was downregulated in the ART group, and this downregulation was even more apparent in the group of copolymer of ART with MNPs- Fe_3O_4 . The levels of bax and bcl-2 were increased in both ART group and the copolymer of ART




Figure 1 The inhibition ratio of K562 treated with ART or the copolymer of ART with MNPs-Fe₃O₄ (100:1) for 48 hours.

Note: * $P < 0.05$, compared with the ART alone.

Abbreviations: MNPs-Fe₃O₄, magnetic nanoparticles of Fe₃O₄; ART, artesunate.




Figure 2 Effect of 1.0 μM MNPs-Fe₃O₄, 100 μM ART, and the copolymer of ART with MNPs-Fe₃O₄ (100:1) on the cell viability in the presence or absence of 40 μM Z-VAD-FMK for 48 hours.

Note: * $P < 0.05$, compared with the absence of Z-VAD-FMK.

Abbreviations: MNPs-Fe₃O₄, magnetic nanoparticles of Fe₃O₄; ART, artesunate.




Figure 3 Apoptosis of K562 cells treated with ART or the copolymer of ART with MNPs- Fe_3O_4 for 48 hours. **A)** Control; **B)** 1 μM MNPs- Fe_3O_4 ; **C)** 100 μM ART; **D)** the copolymer of 100 μM ART with 1 μM MNP- Fe_3O_4 .

Abbreviations: MNPs- Fe_3O_4 , magnetic nanoparticles of Fe_3O_4 ; ART, artesunate.

with MNPs- Fe_3O_4 group compared with control group and MNPs- Fe_3O_4 group (see Figure 5, $P < 0.05$), but there was no obvious difference in the bcl-2/bax ratio. The levels of bcl-rambo and caspase-3 were upregulated when the ART was combined with the MNPs- Fe_3O_4 ($P < 0.05$). These results suggest that the copolymer of ART with MNPs- Fe_3O_4 triggers changes in the expression levels of apoptosis-related genes in K562 cells, among which upregulation of bcl-rambo and downregulation of survivin are the two major alterations.

Discussion

ART could inhibit the proliferation of tumor cells in vitro and in vivo by downregulating the Wnt/ β -catenin pathway.^{4,20} However, toxicity of ART to blood vessels and bone is an obstacle for clinical application of ART, which has limited its widespread application. Nowadays, MNP- Fe_3O_4 as a targeted drug

carrier with target orientation and sustained-release properties has been widely studied in medical research. It has been previously reported that MNP- Fe_3O_4 with anticancer drugs could effectively improve efficiency of chemotherapeutic agents which has a synergistic effect in multidrug resistance.^{12,13}

Our present study demonstrated that the unique MNPs- Fe_3O_4 could increase the biological effects and decrease the side effects of ART. MTT results revealed that after being combined with MNPs- Fe_3O_4 (100:1, M/M), ART showed an enhanced cytotoxic effect on K562 cells; the copolymer of 100 μM ART with 1 μM MNPs- Fe_3O_4 had the most enhanced cytotoxicity among the five groups. Interestingly, the copolymer of ART and MNPs- Fe_3O_4 -induced cell death was attenuated since cell survival rate was increased by Z-VAD-FMK, suggesting that the copolymer of ART and MNPs- Fe_3O_4 -induced apoptosis proceeded via the caspase-dependent pathway.




Figure 4 Apoptosis ratio of K562 cells treated with different concentrations of ART or the copolymer of different concentrations of ART with different concentrations of MNPs-Fe₃O₄.

Notes: * $P < 0.05$, compared with ART alone.

Abbreviations: MNPs-Fe₃O₄, magnetic nanoparticles of Fe₃O₄; ART, artesunate.

Apoptosis, as a process of programmed cell death, is normal for the development and health of multicellular organisms.¹⁴ Apoptotic obstruction is an important event in a tumor genesis, and apoptosis-oriented treatment is common in cancer treatment. Our present studies show the copolymer of 100 μM ART with 1 μM MNPs-Fe₃O₄ could significantly induce K562 cell apoptosis ($P < 0.05$), and the apoptosis induced by ART was dose-dependent, as has been reported in other tumor cells.¹⁻³ These results to some extent indicate that MNPs-Fe₃O₄ can enhance the ART-induced apoptosis.

Apoptosis is the consequence of a series of precisely regulated events that are frequently altered in tumor cells. ART-induced cell apoptosis is apparently involved in the death receptor pathway and the mitochondria.²¹⁻²³ The relative ratios of the various bcl-2 family members such as bcl-2/bax could determine how much cellular stress was needed to induce apoptosis.^{24,25} A recent report showed that bcl-rambo, located in mitochondria, contained conserved bcl-2 homology (BH) motifs 1, 2, 3, 4. But no interaction was found between bcl-rambo with either anti-apoptotic (bcl-2, bcl-XL, bcl-w, A1, MCL-1, E1B-19K, and BHRF1)

or pro-apoptotic (bax, bak, bik, bid, bim, and bad). Its over-expression induces apoptosis by the C-terminal membrane anchor region which was preceded by a unique 250 amino acid insertion containing two tandem repeats. This function was inhibited only by the IAP family.^{15,16} Our results showed that treatment with the copolymer of ART with MNPs-Fe₃O₄ upregulated the expression of bcl-2, bax, bcl-rambo and active caspase-3 protein. It is worth noting that expression of bcl-2 in the copolymer-treated group shows an increasing trend ($P < 0.05$). Although the expression of bax was also increased, the computer-assisted image analysis shows that in combination with MNPs-Fe₃O₄, the ratio of bcl-2/bax in K562 cells was in a slightly ascendant trend compared with ART alone ($P > 0.05$), suggesting that MNPs-Fe₃O₄ enhanced ART-apoptosis through bcl-2/bax-independent mechanism, which differs from the results previously published.^{24,25}

On the other hand, studies showed that Wnt/β-catenin, as an important self-renewal pathway,²⁰ could regulate the transcription of survivin, p53, and c-myc. Overexpression of survivin has been detected in various hematological malignancies, including acute leukemias, myelodysplastic syndrome, chronic myeloid leukemia, and other types of




Figure 5 Expressions of bcl-2, bax, bcl-rambo, survivin, and caspase-3 protein in K562 cells.

Notes: Lane 1) K562 control; Lane 2) 1 μM $\text{MNP}_s\text{-Fe}_3\text{O}_4$; Lane 3) 100 μM ART; Lane 4) the copolymer of 100 μM ART with 1 μM $\text{MNP}_s\text{-Fe}_3\text{O}_4$.

Abbreviations: $\text{MNP}_s\text{-Fe}_3\text{O}_4$, magnetic nanoparticles of Fe_3O_4 ; ART, artesunate.

lymphoid malignancies.^{26–28} Survivin can bind to and potentially inhibit caspases, including caspase-3.²⁹

Caspases are synthesized as proenzymes and can be activated by proteolysis at several sites. The active caspase-3 is a heterotetramer of two large and two small subunits.³⁰ In our present study, the downregulation of survivin expression and a significant increase in the expression of active caspase-3 protein were observed at the same time in the K562 cells treated with the copolymer of $\text{MNP}_s\text{-Fe}_3\text{O}_4$ and ART, suggesting that the potent inhibition of survivin expression could augment the activity of caspase-3, thus substantially enhancing apoptosis of K562 cells induced by the combination of $\text{MNP}_s\text{-Fe}_3\text{O}_4$ with ART.

Conclusion

In summary, these results showed that $\text{MNP}_s\text{-Fe}_3\text{O}_4$ could enhance ART-induced apoptosis in K562 cells, which may be related to the upregulation of bcl-rambo and downregulation

of survivin. It would be worthwhile evaluating the effects of the copolymer on normal cells in vitro for clinical application.

Acknowledgment

This paper was supported by the Open Research Fund of State Key Laboratory of Bioelectronics, Southeast University.

Disclosure

Ying Wang and Yuxiang Han contributed equally to this work. There are no conflicts of interest to report.

References

1. Efferth T, Dunstan H, Sauerbrey A, Miyachi H, Chitambar CR. The anti-malarial artesunate is also active against cancer. *Int J Oncol*. 2001;18: 767–773.
2. Mercer AE, Copple IM, Maggs JL, O'Neill PM, Park BK. The role of heme and the mitochondrion in the chemical and molecular mechanisms of Mammalian cell death induced by the artemisinin antimalarials. *J Biol Chem*. 2011;286:987–996.

3. Du JH, Zhang HD, Ma ZJ, Ji KM. Artesunate induces oncosis-like cell death in vitro and has antitumor activity against pancreatic cancer xenografts in vivo. *Cancer Chemother Pharmacol*. 2010;65:895–902.
4. Li LN, Zhang HD, Yuan SJ, Tian ZY, Wang L, Sun ZX. Artesunate attenuates the growth of human colorectal carcinoma and inhibits hyperactive Wnt/beta-catenin pathway. *Int J Cancer*. 2007;121:1360–1365.
5. Chen HH, Zhou HJ, Wu GD, Lou XE. Inhibitory effects of artesunate on angiogenesis and on expressions of vascular endothelial growth factor and VEGF receptor KDR/flk-1. *Pharmacology*. 2004;71:1–9.
6. Clark RL. Embryotoxicity of the artemisinin antimalarials and potential consequences for use in women in the first trimester. *Reprod Toxicol*. 2009;28:285–296.
7. White TE, Bushdid PB, Ritter S, Laffan SB, Clark RL. Artesunate-induced depletion of embryonic erythroblasts precedes embryo lethality and teratogenicity in vivo. *Birth Defects Res B Dev Reprod Toxicol*. 2006;77:413–429.
8. Li Q, Xie LH, Johnson TO, Si Y, Haeberle AS, Weina PJ. Toxicity evaluation of artesunate and artemisinin in Plasmodium berghei-infected and uninfected rats. *Trans R Soc Trop Med Hyg*. 2007;101:104–112.
9. Clark RL, White TE, A Clode S, Gaunt I, Winstanley P, Ward SA. Developmental toxicity of artesunate and an artesunate combination in the rat and rabbit. *Birth Defects Res B Dev Reprod Toxicol*. 2004; 71:380–394.
10. Fernandez-Pacheco R, Valdivia JG, Ibarra MR. Magnetic nanoparticles for local drug delivery using magnetic implants. *Methods Mol Biol*. 2009;544:559–569.
11. Darton NJ, Hallmark B, Han X, Palit S, Slater NK, Mackley MR. The in-flow capture of superparamagnetic nanoparticles for targeting therapeutics. *Nanomedicine*. 2008;4:19–29.
12. Chen BA, Dai YY, Wang XM, et al. Synergistic effect of the combination of nanoparticulate Fe₃O₄ and Au with daunomycin on K562/A02 cells. *Int J Nanomedicine*. 2008;3:343–350.
13. Chen BA, Lai BB, Cheng J, et al. Daunorubicin-loaded magnetic nanoparticles of Fe₃O₄ overcome multidrug resistance and induce apoptosis of K562-n/VCR cells in vivo. *Int J Nanomedicine*. 2009;4:201–208.
14. Jarpe MB, Widmann C, Knall C, et al. Anti-apoptotic versus pro-apoptotic signal transduction: checkpoints and stop signs along the road to death. *Oncogene*. 1998;17:1475–1482.
15. Oltvai ZN, Millman CL, Korsmeyer SJ. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell*. 1993;74:609–619.
16. Kataoka T, Holler N, Micheau O, et al. Bcl-rambo, a novel Bcl-2 homologue that induces apoptosis via its unique C-terminal extension. *J Biol Chem*. 2001;276:19548–19554.
17. Hengartner MO. The biochemistry of apoptosis. *Nature*. 2000;407: 770–776.
18. Gao H, Wang JY, Shen XZ, Deng YH, Zhang W. Preparation of magnetic polybutylcyanoacrylate nanospheres encapsulated with alicinomy-cin A and its effect on gastric tumor. *World J Gastroenterol*. 2004; 10:2010–2013.
19. Wang X, Zhang R, Wu C, et al. The application of Fe₃O₄ nanoparticles in cancer research: a new strategy to inhibit drug resistance. *J Biomed Mater Res A*. 2007;80:852–860.
20. Hu Y, Chen Y, Douglas L, Li S. Beta-Catenin is essential for survival of leukemic stem cells insensitive to kinase inhibition in mice with BCR-ABL-induced chronic myeloid leukemia. *Leukemia*. 2009;23: 109–116.
21. Li PC, Lam E, Roos WP, Zdzienicka MZ, Kaina B, Efferth T. Artesunate derived from traditional Chinese medicine induces DNA damage and repair. *Cancer Res*. 2008;68:4347–4351.
22. Hamacher-Brady A, Stein HA, Turschner S, et al. Artesunate activates mitochondrial apoptosis in breast cancer cells via iron-catalysed lysosomal reactive oxygen species production. *J Biol Chem*. 2011; 286:6587–6601.
23. Efferth T, Sauerbrey A, Olbrich A, et al. Molecular modes of action of artesunate in tumor cell lines. *Mol Pharmacol*. 2003;64:382–394.
24. Efferth T, Giaisi M, Merling A, Krammer PH, Li-Weber M. Artesunate induces ROS-mediated apoptosis in doxorubicin-resistant T leukemia cells. *PLoS One*. 2007;2:e693.
25. Wu GD, Zhou HJ, Wu XH. Apoptosis of human umbilical vein endothelial cells induced by artesunate. *Vascul Pharmacol*. 2004; 41:205–212.
26. Badran A, Yoshida A, Wano Y, et al. Expression of the anti-apoptotic gene survivin in myelodysplastic syndrome. *Int J Oncol*. 2003; 22:59–64.
27. Tamm I, Kornblau SM, Segall H, et al. Expression and prognostic significance of IAP-family genes in human cancers and myeloid leukemias. *Clin Cancer Res*. 2000;6:1796–1803.
28. Tamm I, Richter S, Oltersdorf D, et al. High expression levels of x-linked inhibitor of apoptosis protein and survivin correlate with poor overall survival in childhood de novo acute myeloid leukemia. *Clin Cancer Res*. 2004;10:3737–3744.
29. Shin S, Sung BJ, Cho YS, et al. An anti-apoptotic protein human survivin is a direct inhibitor of caspase-3 and -7. *Biochemistry*. 2001; 40:1117–1123.
30. Bhardwaj A, Aggarwal BB. Receptor-mediated choreography of life and death. *J Clin Immunol*. 2003;23:317–332.

International Journal of Nanomedicine

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-nanomedicine-journal>

Dovepress

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.