

Lack of association between angiotensin-converting enzyme and dementia of the Alzheimer's type in an elderly Arab population in Wadi Ara, Israel

Abdalla Bowirrat^{1,2,3}

Jing Cui⁴

Kristin Waraska⁵

Robert P Friedland¹⁰

Marlene Oscar-Berman^{2,3,6}

Lindsay A Farrer^{1,6,7,8,9}

Amos Korczyn¹¹

Clinton T Baldwin^{1,5,7}

¹Genetic Program, ²Department of Anatomy and Neurobiology, ³Division of Psychiatry, ⁴Hypertension Section, ⁵Center for Human Genetics, ⁶Department of Neurology, ⁷Department of Genetics and Genomics, Boston University School of Medicine, Boston, MA, USA; ⁸Department of Epidemiology, ⁹Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA; ¹⁰Department of Neurology, Case Western Reserve University, Cleveland, OH, USA; ¹¹Department of Neurology, Tel Aviv University, Tel Aviv, Israel

Abstract: The angiotensin-converting enzyme (ACE), a protease involved in blood pressure regulation, has been implicated as an important candidate gene for Alzheimer's disease (AD). This study investigated whether the ACE gene insertion-deletion (ID) polymorphism is associated with risk of developing dementia of Alzheimer's type (DAT) in an Arab-Israeli community, a unique genetic isolate where there is a high prevalence of DAT. In contrast to several other studies, we found no evidence of an association between this polymorphism and either DAT or age-related cognitive decline (ARCD).

Keywords: angiotensin-converting enzyme, dementia of the Alzheimer's type, age-related cognitive decline, insertion-deletion polymorphism, Arab population

Introduction

Alzheimer's disease (AD) is a catastrophic affliction of the middle and later years of life. Clinically, it is marked by an irreversible decline in mental abilities, including progressive loss of memory, confusion, and dementia, and culminates in childlike helplessness. In the United States, the incidence of AD is exponentially increasing and is projected to affect 8.64 million people by the year 2047 (Grossberg 2003). Median survival times range from 8.3 years for persons diagnosed at age 65, to 3.4 years for persons diagnosed at age 90 (Brookmeyer et al 2002).

Angiotensin-converting enzyme (ACE) catalyzes the conversion of angiotensin I into a physiologically active peptide, angiotensin II. This reaction is an important process in blood pressure regulation. In addition to this well known physiologic role, ACE is able to cleave other proteins, including bradykinin and substance P, and has been shown to inhibit and cleave amyloid β -peptide (A β) aggregation and amyloid plaque formation in vitro (Hu et al 2001). ACE may contribute to the pathogenesis of AD.

To test the hypothesis of association between ACE and DAT, we evaluated an insertion-deletion (ID) polymorphism in the ACE gene as a potential risk factor for dementia of the Alzheimer's type (DAT) in an Arab community in Wadi Ara, Israel. The population comprised 853 persons over the age of 65 (Bowirrat et al 2001; Bowirrat, Friedland, Farrer, et al 2002). Family studies revealed that more than one-third of DAT cases are members of one extended family. We hypothesized that DAT in Wadi Ara may be due to a founder effect enhanced by consanguinity; thus, a specific disease susceptibility allele may be overrepresented in cognitively impaired subjects compared with cognitively healthy residents. Our results suggest that the

Correspondence: Abdalla Bowirrat
Department of Anatomy and
Neurobiology, Laboratory of
Neuropsychology, Boston University
School of Medicine, 715 Albany Street,
L-815 Boston, MA 02118, USA
Tel +1 617 638 4807
Fax +1 617 638 4806
Email bowirrat@bu.edu

ACE-ID polymorphism is not associated with either DAT or age-related cognitive decline (ARCD) in this Arab community.

Methods and results

Subjects were sourced from a prevalence study of dementia conducted in Wadi Ara, a geographically defined area in northern Israel comprising three Arab villages (Bowirrat et al 2001). Among the original list of 853 residents aged 65 years and older (prevalence date, October, 1995), 823 (363 males and 460 females) were available and consented to participate in the study. Only 30 patients (23 females and 7 males) were excluded from the original list of the study (853) because they did not fulfill the diagnostic criteria for DAT and ARCD (they were classified with other medical conditions). Among the excluded females ($n=23$), 19 suffered from vascular dementia (VaD), 1 patient had Parkinson's disease (PD), and 3 had pseudodementia. Among the excluded males ($n=7$), 5 suffered from PD and 2 patients were diagnosed with VaD. The sex-specific prevalence of DAT among males and females was $53/363=14.6\%$ and $115/460=25\%$, respectively; and the sex-specific prevalence of ARCD among males and females was $118/354=33.3\%$ and $236/354=66.6\%$, respectively. These subjects were interviewed and examined by an Arabic-speaking neurologist (AB). Each subject underwent a battery of standard cognitive tests modified to fit the cultural and linguistic characteristics of this community (Bowirrat et al 2001). The diagnosis of DAT and ARCD was determined using DSM-IV criteria (APA 1994). Those with a medical history and laboratory or cognitive test results that suggested the presence of other illnesses, such as vascular dementia, Parkinson's disease, normal-pressure hydrocephalus, or pseudodementia (depression), were excluded. Peripheral blood samples were obtained from the 650 subjects for DNA analysis.

To genotype the ACE-ID polymorphism, 2 polymerase chain reactions (PCRs) were performed. In the first reaction, primers that flank the ID polymorphism were used to amplify DNA from each subject (5'-CTGGAGACC ACTCCCATCCTTCT and 5'-TTGATGAGTTCCAC GTATTTCG). The resulting PCR product was analyzed on a 2% agarose gel for the presence or absence of the 287-bp insertion-deletion polymorphism. Because this reaction frequently fails to detect the insertion variant, a second PCR reaction was performed that specifically amplifies the allele containing the insertion (primers 5'-CTGGGATTA CAGGCGTGATACA and 5'-TTGATGAGTTCCAC

GTATTTCG). The data from both reactions were used to establish the ACE-ID genotype. For statistical analysis, the Hardy-Weinberg equilibrium (HWE) was assessed for each group (DAT, ARCD, normal). Allele and genotype frequencies were compared among groups by a χ^2 test.

The results of the ACE (ID) genotyping (Table 1) showed that the frequency of the I allele was 32% in cognitively normal control, 26% in DAT cases ($p=0.2223$), and 32% in ARCD cases ($p=0.9559$). The genotype frequency distribution in the control groups were also similar to those for DAT cases ($p=0.3731$) and ARCD cases ($p=0.9973$). Therefore, our results suggest that the ID polymorphism within ACE is not associated with DAT or ARCD in this Arab community (Table 1).

Discussion

AD is the most overdiagnosed and misdiagnosed disorder of mental functioning in older adults. Part of the problem, is that many other disorders show symptoms that resemble those of AD. The crucial difference, however, is that many of these disorders – unlike AD – may be stopped, reversed, or cured. Based on these findings, clinical diagnosis of AD has been referred to as “a diagnosis by exclusion”, and one that can only be made in the face of clinical deterioration over time. There is no specific clinical test or finding that is unique to AD. Hence, all disorders that can bring on similar symptoms must be systematically excluded. The “classical” senile plaques and the neurofibrillary tangles seen in an AD

Table 1 Genotypic and allelic frequencies among ARCD, DAT, and cognitively normal control

ACE-ID	Normal	ARCD	DAT
II	8 (7.14)	3 (6.82)	5 (5.88)
DI	56 (50)	22 (50)	35 (41.18)
DD	48 (42.86)	19 (43.18)	45 (52.94)
f(I)	72 (32.14)	28 (31.82)	45 (26.47)

Abbreviations: ACE, angiotensin-converting enzyme; I, insertion allele; D, deletion allele; ARCD, age-related cognitive decline; DAT, dementia of Alzheimer's type; f(I), frequency of the insertion allele.

Table 1a Genotypic p-value

Group	ARCD	Normal
DAT	0.5745	0.3731
ARCD	–	0.9973

Table 1b Allelic p-value

Group	ARCD	Normal
DAT	0.3660	0.2223
ARCD	–	0.9559

brain at autopsy typically are the only definitive diagnosis of the disease.

We did not observe an association between the ACE-ID polymorphism and DAT or ARCD in the elderly Arab population of Wadi Ara. Notably, inherited susceptibility for DAT in Wadi Ara is also not accounted for by the ApoE ϵ 4 allele. Indeed, the frequency of ApoE ϵ 4 among DAT, ARCD, and control was 0.027, 0.033, and 0.029, respectively (Bowirrat et al 2000).

Nonetheless, the relationship between ACE and AD is controversial. Several studies have implicated the ID polymorphism and risk of AD (Narain et al 2000; Cheng et al 2002; Kehoe et al 2003; Elkins et al 2004; Tian et al 2004), while other studies failed to find an association (Myllykangas et al 2000; Taylor et al 2001; Buss et al 2002; Lendon et al 2002; Monastero et al 2002; Harrap et al 2003; Panza et al 2003). Moreover, the pattern of association has been confusing, with some studies showing an association with the I allele (Alvarez et al 1999; Yang et al 2000; Hu et al 2001), while others with the D allele (Amouyel et al 1996; Palumbo et al 1999; Crawford et al 2000; Farrer et al 2000; Richard et al 2001).

The reasons for the disparity in results across studies may be due to differences in ethnic background, different environmental factors, or subject misclassification. To date, few studies have examined the frequency of the ACE genotypes in autopsy-confirmed cases of AD, or investigated the potential effect of these ACE genotypes on the pathology of the disorder. In one autopsy series, the ACE genotype was not associated with the amount of A β ₄₀, A β ₄₂, or pathological tau protein deposited within the frontal cortex, or in the extent of amyloid angiopathy in the brain, whether ApoE ϵ 4 allele was present or not (Lendon et al 2002). The only other autopsy-based study of ACE (based on an elderly Finnish population) also failed to detect an association with ACE (Myllykangas et al 2000). In the Arab community, autopsy is prohibited for religious and traditional reasons. Other limitations of our study are the absence of biomarkers and complete biochemical examinations, and the low number of neuroimaging performed. These helpful clinical techniques are necessary to improve our clinical diagnosis of DAT and are required to differentiate it from similar clinical manifestation that mimic the presentation of dementia.

A possible biological explanation for an association between ACE and AD is still unclear. The insertion allele appears to reduce ACE expression, and DD homozygotes and ID heterozygotes have more circulating ACE than II

homozygotes: 65%, and 31%, respectively (Rieder et al 1999). This increased activity of the D allele may alter A β deposition by a direct proteolytic mechanism (Kehoe et al 2003). Alternatively, ACE may have an indirect role in the progression of AD. The ID polymorphism has also been implicated in various conditions including hypertension (Zaman et al 2002; Hopkins and Hunt 2003), coronary heart disease (Arbustini et al 1995; Kauma et al 1996), cardiomyopathy (Pfohl et al 1998), stroke (de la Torre et al 2002), and juvenile rheumatoid arthritis (Alsaied et al 2003). Other studies don't confirm these associations (see Ueda et al 1995; Hung et al 1999; Harrap et al 2003).

Several studies suggest that there may be a relationship between cardiovascular lesions and the severity and progression of AD (Snowdon et al 1997; Zuliani et al 2001; Bowirrat, Friedland, Korczyn 2002; de la Torre 2002; Korczyn 2002). In summary, in the present study, we found a lack of association between ACE and DAT. Our previous studies have not determined any gene that may explain the high prevalence of DAT in this community, but we have found high prevalence of vascular dementia (VaD) following stroke. Indeed, the prevalence of VaD constitutes 5.9% of the total elderly Arab population in Wadi Ara and about 22% of the total dementia population (Bowirrat, Friedland, Korczyn 2002). This prevalence of VaD (22%) is extremely high compared with VaD patients who accounted for only 6% of all dementia patients identified by Fillit and Hill (2002). Given the role of the ACE gene in vascular disease as well as in AD, the ability to detect association with AD may be complex. On the other hand, and in light of the high frequency of VaD in this population, a new study to evaluate the role of ACE in VaD is essential.

Acknowledgments

This work was supported in part by grants from the National Institute of Health (U01-AG17173), the Institute for the Study of Aging, NIAAA (R37-AA07112), and the US Department of Veterans Affairs.

References

- Alsaied K, Haider MZ, Ayoub EM. 2003. Angiotensin converting enzyme gene insertion-deletion polymorphism is associated with juvenile rheumatoid arthritis. *J Rheumatol*, 30:2705–9.
- Alvarez R, Alvarez V, Lahoz CH, et al. 1999. Angiotensin converting enzyme and endothelial nitric oxide synthase DNA polymorphisms and late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry*, 67:733–6.
- Amouyel P, Richard F, Cotel D, et al. 1996. The deletion allele of the angiotensin I converting enzyme gene as a genetic susceptibility factor for cognitive impairment. *Neurosci Lett*, 217:203–5.

- [APA] American Psychiatric Association. 1994. Diagnostic and statistical manual of mental disorders – Fourth edition (DSM-IV). Washington: APA.
- Arbustini E, Grasso M, Fasani R, et al. 1995. Angiotensin converting enzyme gene deletion allele is independently and strongly associated with coronary atherosclerosis and myocardial infarction. *Br Heart J*, 74:584–91.
- Bowirrat A, Friedland RP, Chapman J, et al. 2000. The very high prevalence of AD in an Arab population is not explained by APOE ϵ 4 allele frequency. *Neurology*, 55:731.
- Bowirrat A, Friedland RP, Farrer L, et al. 2002. Genetic and environmental risk factors for Alzheimer's disease in Israeli Arabs. *J Mol Neurosci*, 19:239–45.
- Bowirrat A, Friedland RP, Korczyn AD. 2002. Vascular dementia among elderly Arabs in Wadi Ara. *J Neurol Sci*, 203–204:73–6.
- Bowirrat A, Treves TA, Friedland RP, et al. 2001. Prevalence of Alzheimer's type dementia in an elderly Arab population. *Eur J Neurol*, 8:119–23.
- Brookmeyer R, Corrada MM, Kawas C. 2002. Survival following a diagnosis of Alzheimer's disease. *Arch Neurol*, 59:1764–7.
- Buss S, Muller-Thomsen T, Hoch C, et al. 2002. No association between DCP1 genotype and late-onset Alzheimer's disease. *Am J Med Gen*, 114:440–5.
- Cheng C-Y, Hong C-J, Lui H-C, et al. 2002. Study of the association between Alzheimer's disease and angiotensin-converting enzyme gene polymorphism using DNA from lymphocytes. *Eur Neurol*, 47:26–9.
- Crawford F, Abdullah L, Schinka J, et al. 2000. Gender-specific association of the angiotensin converting enzyme gene with Alzheimer's disease. *Neurosci Lett*, 280:215–19.
- de la Torre JC. 2002. Alzheimer's disease as a vascular disorder: nosological evidence. *Stroke*, 33:1152–62.
- Elkins JS, Douglas VC, Johnston SC. 2004. Alzheimer's disease risk and genetic variation in ACE. *Neurology*, 62:363–8.
- Farrer LA, Sherbatich T, Keryanov SA, et al. 2000. Association between angiotensin-converting enzyme and Alzheimer's disease. *Arch Neurol*, 57:210–14.
- Fillit H, Hill J. 2002. The costs of vascular dementia: a comparison with Alzheimer's disease. *J Neurol Sci*, 15:203–204:35–9.
- Grossberg GT. 2003. Diagnosis and treatment of Alzheimer's disease. *J Clin Psychiatry*, 64(Suppl 9):3–6.
- Harrap SB, Tzourio C, Cambien F, et al. 2003. The ACE gene I/D polymorphism is not associated with the blood pressure and cardiovascular benefits of ACE inhibition. *Hypertension*, 42:297–310.
- Hopkins PN, Hunt SC. 2003. Genetics of hypertension. *Genet Med*, 5: 413–29.
- Hu J, Igarashi A, Kamata M, et al. 2001. Angiotensin-converting enzyme degrades Alzheimer amyloid β -peptide (A β); retards A β aggregation, deposition, fibril formation; and inhibits cytotoxicity. *J Biol Chem*, 276:47863–8.
- Hung J, McQuillan BM, Nidorf M, et al. 1999. Angiotensin-converting enzyme gene polymorphism and carotid wall thickening in a community population. *Arterioscler Thromb Vasc Biol*, 19:1969–74.
- Kauma H, Paivansalo M, Savolainen MJ, et al. 1996. Association between angiotensin converting enzyme gene polymorphism and carotid atherosclerosis. *J Hypertens*, 14:1183–7.
- Kehoe PG, Katzov H, Feuk L, et al. 2003. Haplotypes extending across ACE are associated with Alzheimer's disease. *Hum Mol Genet*, 12:859–67.
- Korczyn AD. 2002. Mixed dementia – the most common cause of dementia. *Ann N Y Acad Sci*, 977:129–34.
- Lendon CL, Thaker U, Harris JM, et al. 2002. The angiotensin 1-converting enzyme insertion (I)/deletion (D) polymorphism does not influence the extent of amyloid or tau pathology in patients with sporadic Alzheimer's disease. *Neurosci Lett*, 328:314–18.
- Monastero R, Caldarella R, Mannino M, et al. 2002. Lack of association between angiotensin converting enzyme polymorphism and sporadic Alzheimer's disease. *Neurosci Lett*, 335:147–9.
- Myllykangas L, Polvikoski T, Sulkava R, et al. 2000. Cardiovascular risk factors and Alzheimer's disease: a genetic association study in a population aged 85 or over. *Neurosci Lett*, 292:195–8.
- Narain Y, Yip A, Murphy T, et al. 2000. The ACE gene and Alzheimer's disease susceptibility. *J Med Genet*, 37:695–7.
- Palumbo B, Cardini D, Nocentini G, et al. 1999. Angiotensin converting enzyme deletion allele in different kinds of dementia disorders. *Neurosci Lett*, 267:97–100.
- Panza F, Solfrizzi V, D'Introno A, et al. 2003. Shifts in angiotensin I converting enzyme insertion allele frequency across Europe: implications for Alzheimer's disease risk. *J Neurol Neurosurg Psychiatry*, 74:1157–61.
- Pfohl M, Fetter M, Kock M, et al. 1998. Association between angiotensin I-converting enzyme genotypes, extracranial artery stenosis, and stroke. *Atherosclerosis*, 140:161–6.
- Richard F, Fromentin-David I, Ricolfi F, et al. 2001. The angiotensin I converting enzyme gene as a susceptibility factor for dementia. *Neurology*, 56:1593–5.
- Rieder MJ, Taylor SL, Clark AG, et al. 1999. Sequence variation in the human angiotensin converting enzyme. *Nat Genet*, 22:59–62.
- Snowdon DA, Greiner LH, Mortimer JA, et al. 1997. Brain infarction and the clinical expression of Alzheimer's disease. The Nun Study. *JAMA*, 277:813–17.
- Taylor A, Ezquerra M, Bargri G, et al. 2001. Alzheimer's disease not associated with polymorphism in the angiotensinogen and rennin genes. *Am J Med Gen*, 105:761–4.
- Tian J, Shi J, Bailey K, et al. 2004. A polymorphism in the angiotensin 1-converting enzyme gene is associated with damage to cerebral cortical white matter in Alzheimer's disease. *Neurosci Lett*, 354:103–6.
- Ueda S, Weir CJ, Inglis GC, et al. 1995. Lack of association between angiotensin-converting enzyme gene insertion/deletion polymorphism and stroke. *J Hypertens*, 13:1597–601.
- Yang JD, Feng G, Zhang ZX, et al. 2000. Association between angiotensin-converting enzyme gene and late onset Alzheimer's disease in Han Chinese. *Neurosci Lett*, 295:41–4.
- Zaman MA, Oparil S, Calhoun DA, et al. 2002. Drugs targeting the rennin-angiotensin-aldosterone system. *Nat Rev Drug Discov*, 1:621–36.
- Zuliani G, Ble A, Zanca R, et al. 2001. Genetic polymorphisms in older subjects with vascular or Alzheimer's dementia. *Acta Neurol Scand*, 103:304–8.