

A Mini-Review of Diagnostic and Therapeutic Nano-Tools for Pancreatitis

Qixiong Zhang^{1,*}, Shanshan Li^{2,*}, Yang Yu³, Yuxuan Zhu¹, Rongsheng Tong¹

¹Department of Pharmacy, Sichuan Academy of Medical Science & Sichuan Provincial People's Hospital, Personalized Drug Therapy Key Laboratory of Sichuan Province, School of Medicine, University of Electronic Science and Technology of China, Chengdu, 610000, People's Republic of China;

²College of Pharmacy, Southwest Minzu University, Chengdu, 610000, People's Republic of China; ³College of Pharmaceutical Sciences, Southwest University, Chongqing, 400712, People's Republic of China

*These authors contributed equally to this work

Correspondence: Qixiong Zhang, Department of Pharmacy, Sichuan Academy of Medical Science & Sichuan Provincial People's Hospital, Personalized Drug Therapy Key Laboratory of Sichuan Province, School of Medicine, University of Electronic Science and Technology of China, Chengdu, 610000, People's Republic of China, Email qixiongzhong@outlook.com

Abstract: Pancreatitis is an inflammatory reaction of pancreatic tissue digestion, edema, bleeding and even necrosis caused by activation of pancreatin due to various causes. In particular, patients with severe acute pancreatitis (SAP) often suffer from secondary infection, peritonitis and shock, and have a high mortality rate. Chronic pancreatitis (CP) can cause permanent damage to the pancreas. Due to the innate characteristics, structure and location of the pancreas, there is no effective treatment, only relief of symptoms. Especially, AP is an unpredictable and potentially fatal disease, and the timely diagnosis and treatment remains a major challenge. With the rapid development of nanomedicine technology, many potential tools can be used to address this problem. In this review, we have introduced the pathophysiological processes of pancreatitis to understanding its etiology and severity. Most importantly, the current progress in the diagnosis and treatment tools of pancreatitis based on nanomedicine is summarized and prospected.


Keywords: pancreatitis, nanomedicine, diagnosis, therapy

Introduction

Current pathology defines pancreatitis as a disease caused by the digestive action of trypsin, mainly divided into acute pancreatitis (AP) and chronic pancreatitis (CP), which is characterized by varying degrees of edema, bleeding, and acinar and associated pancreatic tissue necrosis.¹⁻⁴ The pathogenesis of AP have been extensively reported but remain unclear.⁵ Gallstones, excessive alcohol intake, drug-induced injury, hypercalcemia, infection, genetics, and autoimmunity can all be listed as risk factors.⁶

AP is characterized by mis-activation and release of trypsinogen that triggers the activation of other digestive enzymes, leading to self-digestion of pancreatic tissue (Figure 1). Mild acute pancreatitis (MAP) when limited inflammation and organ dysfunction are present.⁷ However, when the inflammation expands to the surrounding organs, it evolves into severe acute pancreatitis (SAP), which not only leads to pancreatic necrosis, but also easily leads to multiple organ failure and sepsis. Despite the great advances in intensive care medicine over the past few decades, the mortality rate for patients with SAP is still about 30%.⁸ As the glands are gradually replaced by fibrotic tissue, the glands and ducts atrophy and calcification of proteins in the interstitial tissue and glands and ducts leads to partial or complete obstruction, making pancreatitis prone to progressive progression.⁹ According to statistics, up to 20% of patients with AP will develop to CP.¹⁰ Notably, CP can lead to inflammation, fibrosis, and scarring, which can permanently damage pancreatic tissue and function.⁹ Unfortunately, early diagnosis of CP is difficult due to the lack of typical blood, imaging, or functional biomarkers. Diagnostic tools available include risk factors, clinical history, imaging, and pancreatic function tests. In addition, no specific therapy has been identified to date. The current focus of treatment for CP is to improve quality of life by relieving symptoms. For example, quitting smoking and alcohol or taking non-opioid and

Graphical Abstract



opioid pain relievers for pain. Surgery is performed when there are morphological problems such as catheter abnormalities.

Nanotechnology has been widely used in the field of biomedicine, which has made rapid progress in the prevention, diagnosis and treatment of diseases.¹¹ Nanomedicine focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedicine field.^{12–14} Nanomedicine include nanoparticles, micelles, liposomes and other carriers with drug loading capability, as well as nano-scale drugs composed of some active substances themselves. Nanomedicine have smaller size and larger specific surface area, the ability to load drugs and the unique property that the surface can be modified with other structures.¹⁵ It can be used to improve the effectiveness,




Figure I The molecular and cellular events in acute pancreatitis.

safety, physicochemical properties and pharmacokinetics or pharmacodynamics of drugs. Functionalized nanomedicines have the characteristics of improving the bioavailability of oral drugs in vivo, prolonging the half-life of injectable drugs by reducing immunogenicity, and realizing targeted drug delivery. The use of nanomedicines to treat diseases can reduce the frequency of drug administration, improve the pharmacological effects of drugs and minimize the side effects of drugs, thereby showing better clinical treatment effects and treatment compliance. In addition, nanomedicine can also actively or passively deliver drugs to the target site, which is conducive to the accumulation of drugs in inflammation or tumor microenvironment, so as to play a better therapeutic effect.¹⁶

This review summarizes the application research trends of nanomedicine in pancreatitis from two aspects of treatment and diagnosis, hoping to provide a reference for the research of new nanomedicine for pancreatitis.

Nanomedicine-Based Therapeutic Strategy of Pancreatitis

Nanotechnology has emerged as the most promising approach to pharmaceuticals. The term “nanomedicine” has been defined as regulators take a close look at nano-systems.¹⁷ For example, Doxil (liposomal doxorubicin), Myocet (non-pegylated liposomal doxorubicin), Abraxane (Albumin-based paclitaxel nanoparticles), Genexol-PM (paclitaxel micelles), Marqibo (liposomal vincristine sulfate) are already approved.¹⁸ Currently, commercial formulations for pancreatic disease (pancreatic cancer) are Abraxane and Onivyde (liposomal irinotecan).¹⁸

We know that delivering drugs to the pancreas has some difficulties. Both the pancreas-specific targeting ability and the ability to cross the blood-pancreatic barrier (BPB) need to be improved.¹⁹ In addition, trypsin inhibitors are mostly peptides with short half-life, which affects the therapeutic effect of pancreatitis.²⁰ Finally, the harsh tissue microenvironment of patients also affects drug release and pharmacological activity.²¹

In theory, targeting can be achieved by passive or active targeting approach, which can increase many drugs targeted to the tissue to improve the therapeutic effect. For pancreatitis, passive targeting can be achieved through various abnormal biochemical signals (pH value, reactive oxygen species, abundant digestive enzymes, etc.) and tissue space with enhanced permeability.²² Active targeting can be achieved by specifically targeting overexpressed receptors at the site of inflammation.²³ The following is a review of the development of early diagnosis and targeted therapy for pancreatitis based on nanomedicine from the perspective of different kinds of nanoparticles.

Solid Lipid and Polymer Nanoparticles

Lipid or polymeric nanoparticles are the most widely studied in the field of nanomedicine.²⁴ Not surprisingly, these materials have also been used as nanocarriers in the treatment of pancreatitis. For example, Cervin's group used lipid liquid crystal materials to load somatostatin, a peptide hormone with the potential to treat pancreatitis.²⁵ The nano-somatostatin can increase the effective half-life of this peptide from a few minutes to 1 hour, effectively eliminating its clinical application.

Moreover, polymeric nanoparticles with suitable and tunable particle size range have the effect of extravasation across leaky vasculature and subsequent inflammatory cell-mediated sequestration (ELVIS effect).²⁶ A growing number of studies exploit the ELVIS effect to enable passive targeting of nanoparticles to sites of inflammation and reduced systemic toxicity.^{27–30} In pancreatitis, the nanoparticles with ELVIS effect has shown promise. For example, Yang et al developed a PLGA nanoparticles (CQ/pDNA/PLGA NPs), which co-loaded chloroquine diphosphate (CQ) and pDNA, to achieve targeted delivery to tumors and pancreatitis.³¹ In this study, pDNA compacted by CQ was embedded into PLGA nanoparticles. The CQ/pDNA/PLGA NPs not only improved the transfection efficiency but also enhanced the targeting efficiency of CT26 transplanted tumor. More importantly, CQ/pDNA/PLGA NPs showed excellent targeting ability in *L*-arginine-induced AP model in mice. Impressively, after the first report on PLGA-based AP targeting, the same research group further reported a tamoxifen-loaded PLGA nanoparticles (TAM-NPs) in combination with CQ-loaded liposomes (CQ-LPs) in 2022.³² The follow-up experiment proved that combination therapy achieved relief of AP and even SAP by upregulating IDO signaling pathways in bone marrow derived mesenchymal stem cells. The above two examples illustrated the ELVIS effect of PLGA NPs and their easy uptake by pancreatic macrophages and neutrophils, enabling this kind of delivery system with high therapeutic effect in AP. This property of being taken up by macrophages has also been applied in the precision therapy of pancreatic cancer.³³

Biomacromolecule Nanoparticles

Compared with artificial lipids and synthetic polymers, biological macromolecules (such as proteins, nucleic acids, etc.) have obvious advantages in terms of biocompatibility and safety.³⁴ In addition, some proteins have clear sensitivity to up-regulated amylase, proteases and lipases in pancreatitis lesions, and can be used as responsive drug carriers to achieve effective passive targeting of pancreatitis lesions.³⁵ As shown in Figure 2, Zhao et al developed a primary three-dimensional (3D) structure similar to DNA double helix with bilirubin using silk fibroin as a carrier with the help of multiple intermolecular forces (hydrogen bonds, hydrophobic forces), and collapsed into nanoparticles (BRSNPs).³⁶ In vivo imaging experiments showed that BRSNPs were more efficiently enriched in pancreatic tissue, enabling passive targeting. In acinar cells model and *L*-arginine-induced rat AP model, BRSNPs can release bilirubin under the action of excessive pancreatic enzymes (trypsin) at the AP site. Bilirubin not only directly inhibited cellular mitochondrial ROS generation, but also activated the Nrf2 pathway, increased HO-1, and then inhibited the pro-inflammatory NF- κ B signaling pathway. BRSNPs effectively reduced the levels of various serum biomarkers, including amylase, alanine aminotransferase, aspartate aminotransferase, creatinine, urea nitrogen, etc. It alleviated the oxidative stress state and lipid oxidation of pancreatic tissues, inhibited edema and fibrosis, and finally achieved better AP relief effect than somatostatin or free bilirubin positive treatment group.

In addition, endogenous macromolecules are also ideal carriers. It is known that although carbon monoxide (CO) has anti-inflammatory and antioxidant activities, its potential therapeutic ability for AP is limited in its use in inflammation-related diseases due to disadvantages of toxicity and non-specific distribution.³⁷ However, a blessing in disguise, exploiting the natural binding ability of hemoglobin to CO, Maruyama et al developed a CO-bound hemoglobin vesicles (CO-HbV).³⁸ The authors established a choline-deficient ethionine-supplemented diet-induced SAP mice model. Compared with saline or HbV treatment, CO-HbV obviously attenuated the death of SAP mice. By inhibiting systemic proinflammatory cytokine production, neutrophil infiltration, and oxidative stress status, CO-HbV significantly suppresses AP severity. Notably, the SAP mice also developed secondary multiple organ damage. Interestingly, CO-HbV may be due to inhibition of neutrophil infiltration and oxidative stress damage, as well as moderating multiple organ failure.




Figure 2 (A) Schematic diagram of preparation, targeting and mechanism of bilirubin nanomedicine (BRSNPs). (B) H&E staining sections of pancreatic tissue after BRSNPs treatment. (C) BRSNPs activated Nrf2 signaling pathway, up-regulated SOD-1, and ultimately inhibited the expression level of HO-1 and NF- κ B. Reproduced from Yao Q, Jiang X, Zhai YY, et al. Protective effects and mechanisms of bilirubin nanomedicine against acute pancreatitis. *J Control Release*. 2020;322:312–325. Copyright 2020, with permission from Elsevier.³⁶

In addition to macromolecules such as proteins, nucleic acid is also a qualified carrier construction material.^{39–41} For example, tetrahedral framework nucleic acids, so-called tFNAs, is reported as an emerging class of nanoparticles.⁴² In briefly, tFNAs are self-assembled from four single-stranded DNA (ssDNA) with a three-dimensional cage-like structure.^{43,44} Recent literature indicated that tFNAs can regulate cell behaviors such as cell proliferation, migration, and autophagy.⁴⁵ The tFNAs have also been shown to have excellent anti-inflammatory and anti-apoptotic effects and have been used to treat a variety of diseases: including acute kidney injury,⁴⁶ periodontitis,^{47,48} diabetes,^{49,50} and Sjögren's syndrome.⁵¹ Surprisingly, Yunfeng Lin's team found that tFNAs can inhibit SAP by inhibiting the expression of inflammatory factors in tissues and blood, and at the same time regulating the expression of anti-apoptotic proteins. In addition, tFNA can also inhibit the typical inflammatory manifestations through lymphocyte infiltration, thereby protecting multiple organs such as pancreas, lung, liver and kidney.⁵²

Cell Membrane Camouflaged Nanoparticles

As mentioned above, bio-macromolecular nanoparticles have shown good application effects, which suggested that biomimetic nanoparticles have unique advantages.^{53,54} Cell membrane camouflaged nanoparticles (CMCNPs) are a kind of “Trojan”-like strategy, which is a hot research direction of biomimetic nanomedicine in recent years.⁵⁵ Briefly, the partial or complete membrane of specific cells is extracted and wrapped on the surface of various nanoparticles, so that the nanoparticles inherit some specific functions derived from cells, such as biological activity, homing effect, etc.⁵⁵ Numerous literatures have reported that CMCNPs can be used as excellent therapeutic drug carriers, diagnostic reagent carriers, etc., through the natural homing effect of specific cell membranes on the corresponding lesions, to achieve precise treatment of diseases.^{56–58} We know that pancreatitis is a serious inflammatory disease, and there are a large number of inflammatory cells infiltrating the lesions, such as neutrophils and macrophages.⁵⁹ In view of this, Li Deng's group developed a neutrophil membrane-covered PEG-PLGA nanoparticles (NNPs).⁶⁰ This study found that NNPs can more effectively penetrate the BPB and specifically distribute into pancreatic tissue compared with NPs without cell membrane. Compared with free celastrol (CLT) and NNPs/CLT treatment groups, NNPs/CLT significantly down-regulated serum amylase and myeloperoxidase levels in AP rats. In addition, the use of NNPs also attenuated the systemic toxicity of CLT. Not only that, in vitro imaging experiments also demonstrated that NNPs can be enriched in the lung tissue of AP mice, which is due to a large number of neutrophil infiltration during AP-induced lung injury. Similarly, it has been reported that silk fibroin nanoparticles can precisely deliver the antioxidant ferulic acid to the pancreas after being coated with neutrophil membranes.⁶¹




Figure 3 Macrophage membrane camouflaged PLGA nanoparticles, called as MΦ-NP(L&K), co-loading attractant (Melittin) and inhibitor (MJ-33) of PLA2 to achieve targeted therapy of AP mice. Reproduced from Zhang Q, Zhou J, Zhou J, Fang RH, Gao W, Zhang L. Lure-and-kill macrophage nanoparticles alleviate the severity of experimental acute pancreatitis. *Nat Commun.* 2021;12:4136. Copyright 2021, the Authors. Springer Nature. Creative Commons CC BY.⁶⁵

Meanwhile, macrophage infiltration is also a common and important pathological process of pancreatitis.⁶² Macrophages appear to be a negative factor in the progression of pancreatitis because of excessive recruitment and release of proinflammatory factors.^{63,64} As in Figure 3, Liangfang Zhang's group used autologous macrophage membrane to coating PLGA nanoparticles, supplemented with phospholipase A2 (PLA2) attractant (Melittin) and inhibitor (MJ-33), and prepared a kind of "Lure-and-Kill" macrophage-like nanoparticles (MΦ-NP(L&K)).⁶⁵ The MΦ-NP(L&K) can significantly reduce the expression levels of serum markers (including PLA2, IL-6, TNF- α , IL-1 β) in the MAP mouse model, relieve the edema of pancreatic tissue, and reduce the number of necrotic pancreatic acinar cells. Significantly decreased, reducing the degree of infiltration of CD45-positive lymphocytes in pancreatic tissue. On SAP mice, MΦ-NP (L&K) also achieved excellent therapeutic effect. Moreover, M-NP(L&K) has also been confirmed to have good biocompatibility in vitro and in vivo, which is also the unique advantage of this type of CMCNPs.

Inorganic Nanoparticles

For inflammatory diseases, exogenous materials with antioxidant capacity that mimic endogenous enzymes.⁶⁶ Inorganic nanoparticles are generally smaller in particle size and narrower in size distribution than polymer/lipid nanoparticles, and their surface chemistry is also well suited for ligand conjugation. Therefore, artificial enzymes are the best candidates for treating dysregulated redox homeostasis.^{67–69}

Hu et al reported that a polyvinylpyrrolidone-stabilized MoSe₂ nanoparticle (MoSe₂-PVP NPs) could be easily prepared.⁷⁰ The MoSe₂-PVP NPs effectively simulates various natural enzymes, and eliminates free radicals such as •OH, •O₂[−] and 3-ethylbenzothiazoline-6-sulfonic acid. MoSe₂-PVP NPs significantly increased the survival probability of cells in H₂O₂ and had a significant protective effect on AP animal models.⁷⁰ Another example, Prussian blue, as an ancient dye, is a clinical antidote for radioactive element poisoning such as thallium. Prussian blue nanoparticles (PB




Figure 4 Prussian blue nanoparticles prepared by PVP stabilized Fe[(CN)₆]^{3−}. This artificial nano-enzyme, called the PBzyme, can restore mitochondrial homeostasis by scavenging ROS, reduce pro-inflammatory cytokines IL-6, IL-1 β and TNF- α , and inhibit TLR/NF- κ B signaling pathway, ultimately achieving the remission of acute pancreatitis. Reproduced from Xie X, Zhao J, Gao W, et al. Prussian blue nanozyme-mediated nanoscavenger ameliorates acute pancreatitis via inhibiting TLRs/NF-kappaB signaling pathway. *Theranostics*. 2021;11:3213–3228. This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.⁷⁶

NPs) have stable chemical structure and excellent physical, chemical, optical and magnetic properties. Meanwhile, PB NPs can simulate a variety of antioxidant enzyme activities, which has attracted great attention in the treatment of inflammatory diseases in recent years.^{71–75} As shown in Figure 4, Zheng et al developed a PB NPs (PBzyme) that can act as a nano-enzyme to remove a variety of ROS and pro-inflammatory factors including $\bullet\text{OH}$, $\bullet\text{OOH}$, and H_2O_2 . It reduces necrosis, nucleic acid damage, and peroxidation by downregulating TLR-related NF- κB signaling pathway

Similarly, the rare earth element nanoparticles also have ROS scavenging properties.^{77,78} The nanocerium (NC) developed by Chandraiah Godugu's group can remove excess superoxide anion and hydrogen peroxide in cells by mimicking SOD and catalase activities, thereby alleviating oxidative stress and restoring mitochondrial membrane potential. Furthermore, NC could reduce the expression level of p65-NF- κB and the acetylation of histone H3 in AP mice.⁷⁹ Using the same strategy, in 2019, the same team also reported that yttrium oxide nanoparticles not only inhibited the infiltration of inflammatory cells around damaged pancreatic acinar cells, but also attenuated endoplasmic reticulum stress and the expression of molecular chaperones. Finally, an effective treatment for caerulein-stimulated SAP was achieved.⁸⁰

In addition to the above-mentioned metallic inorganic nanoparticles, non-metallic inorganic nanoparticles represented by selenium can also achieve the purpose of alleviating pancreatitis. Abdel-Hakeem's team restored endocrine and exocrine functions in the pancreas of AP mice using only selenium nanoparticles (10–45 nm) with antioxidant properties.⁸¹

On the other hand, non-metallic nanoparticles represented by porous silica have excellent drug loading capacity. Chitosan oligosaccharides (COS) has been proved to be an antioxidant, but due to its non-specific distribution, it is difficult to exert sufficient effects in vivo.⁸² Zeng et al used porous SiO_2 material to load COS. The complex ($\text{COS}@\text{SiO}_2$) can slowly release COS in pancreatic tissues with acidosis caused by SAP. Accurate release of COS inhibited NF- κB and NLRP3 inflammasome expression by activating Nrf2 pathway, and reduced systemic inflammatory response and oxidative stress indicators in tissues, ultimately alleviating pancreatic and subsequent lung pathological damage in SAP mice.⁸³

But it has also been reported that intravenous injections of commonly used inorganic nanoparticles can speed up the spread of breast cancer cells to other sites, promoting the emergence of new tumors.⁸⁴ This will be a barrier to the clinical application of inorganic nanoparticles.

Liposome, Micelles and Dendrimers

Liposomes, micelles, and dendrimers have also been reported as novel drug carriers for the treatment of pancreatitis. Hsing-wen Sung's research group developed a “transformer”-like nanocarrier system (TLNS). It has been confirmed in vitro that TLNS can undergo structural changes in the intestinal environment and form nanoscale micelles with curcumin (CUR) in the process.⁸⁵ The mechanisms for AP-specific therapy through CUR-loaded micelles is shown (Figure 5). The pancreas of rats treated with TLNS produced about 12 times of CUR signal than that of rats treated with free CUR, possibly improving AP recovery. These results suggest that TLNS can significantly increase drug dissolution in the gut, making oral administration a more effective treatment route for pancreatitis.

Active targeting mediated by specific ligands is more efficient than passive targeting that relies on morphological changes.²³ However, the development of targeted therapies has been hampered by limited treatment options and the lack of molecule-targeted ligands or non-serum-based biomarkers. Kimberly Kelly et al used computation-guided phage display technology to screen five polypeptide ligands selectively for activated pancreatic stellate cells, acinus cells, macrophages, and extracellular matrix in a CP model induced by caerulein.⁸⁶ PEG-DOPC-based liposomes were modified by five ligands to form targeted liposomes, and an optimal target unit (MDLSLKP) was selected by investigating the pharmacokinetic parameters in vivo. The enrichment of optimal liposomes in the inflammatory pancreas increased to 1.3 times compared with the control group. They also observed that apigenin (Api) loaded targeted liposomes (Api-ECM Lip) improved the survival of acinar structures. Api-ECM Lip reduced the expression of collagen and fibronectin in pancreatic tissues of mice with CP by 37% and 33%, respectively. Surprisingly, Api-ECM Lip also alleviated Api-induced hepatic oxidative stress. In another similar study, nano-scale liposomal delivery of caffeic acid




Figure 5 Transformer-like shape-changeable drug-loading system in situ loading curcumin for AP. Oral capsules consist of proton donor DTPA dianhydride, sodium bicarbonate, SDS and curcumin (CUR). When carbon dioxide is released in an acidic environment, the formed bubbles will form CUR-loaded SDS micelles in situ when they cross the water interface composed of SDS and CUR. The micelles can be taken up by M cells in the gut, enter the payer's patch, accumulate in the mesenteric lymph nodes, and then precisely introduced into the pancreatic tissue through the mesenteric-lymphatic system. Reproduced from Chuang EY, Lin KJ, Huang TY, et al. An intestinal "transformers"-like nanocarrier system for enhancing the oral bioavailability of poorly water-soluble drugs. *ACS Nano*. 2018;12:6389–6397. doi:10.1021/acsnano.8b00470.⁸⁵

phenethyl ester, using passive targeting effect, enriched in inflammatory sites, reduced oxidative stress storm by regulating Nrf-2 and NF- κ B signaling pathways, alleviated AP in rats.⁸⁷

In addition, some dendrimers with polyhydroxyl groups have been shown to have antioxidant activity,^{88,89} suggesting their application in the treatment of pancreatitis. As previously reported, two generation 5 (G5) polyamidoamine (PAMAM) dendrimers with hydroxyl or carboxyl, G4.5-COOH and G5-OH were synthesized. In a caerulein-induced AP mouse model, their protective effects were investigated.⁹⁰ G4.5-COOH and G5-OH not only significantly reduced inflammatory storms in the AP mice, but also distinctly inhibited the expression of LPS-induced mouse peritoneal inflammatory macrophages. They also significantly reduced the WBC and monocytes. The in vivo protective effect of G4.5-COOH on AP was better than that of G5-OH. Finally, authors demonstrated that the anti-inflammatory mechanism of G4.5-COOH and G5-OH may be the inhibition of NF- κ B nuclear translocation in macrophages.

To make a long story short, the nanoscale therapeutic strategies for pancreatitis are summarized in Table 1.

Nanomedicine-Based Diagnosis Strategy of Pancreatitis

Current diagnosis of pancreatitis is based on disease symptoms, which is poorly defined and more empirical.² The nanomedicine based on nanotechnology approach provides a new option for the diagnosis of pancreatitis. Given the

Table 1 Reported Nanotechnology-Based Therapeutic Strategy for Experimental Pancreatitis

Category	Purpose	Materials	Drug	Animal Models	Target	Therapeutic Mechanism	Ref.
Solid lipid/polymer nanoparticles	Nanocarrier	Lipid liquid crystal PLGA	Somatostatin Chloroquine diphosphate (CQ) and pDNA	AP <i>L</i> -arginine-induced AP	– Passive targeting (ELVIS effect)	Extending plasma half-lives of drug CQ show synergistic anti-inflammatory effect with the therapeutic gene	[25] [31]
	Nanocarrier	PLGA/liposome	CQ and tamoxifen (TAM)	Caerulein-induced AP and LPS-induce SAP	Passive targeting	CQ/TAM combination and MSCs synergistically up-regulated IDO signaling pathway	[32]
Biomacromolecule nanoparticles	Nanocarrier	Silk fibroin	Bilirubin	<i>L</i> -arginine-induced AP	Passive targeting (enzyme response)	Protect acinar cells by regulating NF- κ B and Nrf2/HO-1 pathways	[36]
	Nanocarrier	Hemoglobin	Carbon monoxide	Choline-deficient ethionine (CDE) diet-induced AP	Bionic targeting	Anti-inflammatory and antioxidation	[38]
	Nanomedicine	Tetrahedral framework nucleic acids	–	2% Sodium taurocholate-induced AP	–	Inhibiting the cytokines in tissues and blood	[52]
Cell membrane camouflaged nanoparticles	Nanocarrier	Neutrophil membrane-coated PEG-PLGA nanoparticles	Celastrol	3% Sodium taurocholate-induce AP	Bionic targeting (Homing)	Down-regulated serum amylase and myeloperoxidase levels	[60]
	Nanocarrier	Neutrophil membrane-coated silk fibroin nanoparticles	Ferulic acid	Not mentioned	Bionic targeting (Homing)	Anti-inflammatory and antioxidation	[61]
	Nanocarrier	Macrophage membrane-coated PLGA nanoparticles	Melittin and MJ-33	Caerulein-induced AP	Bionic targeting (Homing)	While Melittin induces PLA2, MJ-33 inhibits PLA2 activity (PLA2 exacerbates AP)	[65]

(Continued)

Table 1 (Continued).

Category	Purpose	Materials	Drug	Animal Models	Target	Therapeutic Mechanism	Ref.
Inorganic nanoparticles	Nanomedicine	MoSe ₂ -PVP nanoparticles	–	Caerulein-induced AP	–	Effectively mimicked various enzymes and scavenged free radicals	[70]
	Nanomedicine	Prussian blue nanoparticles	–	Caerulein-induced AP	–	Simulate a variety of antioxidant enzyme and inhibit TLRs/NF-κB signaling pathway	[76]
	Nanomedicine	Nanoceria	–	Caerulein-induced AP	–	Reduced NF-κB expression and histone H3 acetylation by mimicking SOD and catalase	[77]
	Nanomedicine	Yttrium oxide nanoparticles	–	Caerulein-induced AP	–	Regulation of the Nrf2/NF-κB pathway to restore mitochondrial and ER homeostasis	[80]
	Nanomedicine	Selenium nanoparticles	–	L-arginine-induced AP	Passive targeting	Anti-inflammatory, antioxidant, and pro-apoptotic actions	[81]
	Nanocarrier	Porous SiO ₂	Chitosan oligosaccharides	Caerulein-induced AP; L-arginine-induced SAP	Passive targeting	Regulating the Nrf2 signaling pathway in AP or SAP mice	[83]
Others	Nanocarrier	Sodium dodecyl sulfate (SDS)-based micelles	Curcumin	L-arginine-induced AP	Passive targeting	Curcumin specifically accumulates AP site through the mesenteric lymphatic system	[85]
	Nanocarrier	Liposome	Apigenin	Caerulein-induced CP	Active targeting (specific peptide as the target unit)	Increased nanoparticle accumulate in inflamed pancreas	[86]
	Nanocarrier	Liposome	Caffeic acid phenethyl ester	L-ornithine	Passive targeting	Modulates Nrf2 and NF-κB Signaling in AP rat	[87]
	Nanomedicine	PAMAM dendrimers with different surface modification groups	–	Caerulein-induced AP	Passive targeting	PAMAM dendrimers reduced the WBC and monocytes, inhibited cytokines and the NF-κB in macrophages	[90]

physiological function of the pancreas, amylase is a major class of serum biomarkers for pancreatitis. A novel synthetic nano-sensor binuclear Pd complex has been reported to be able to sensitively detect α -amylase in serum and urine.⁹¹ The detection limit of this method was (7.4×10^{-10} mol/L) even lower than the α -amylase concentration (3–321 U/L) in different samples from patients with pancreatitis. This method greatly improved the sensitivity (96.88%) and specificity (94.41%) of α -amylase in the early diagnosis of pancreatitis.

Compared with serum biomarkers, in vivo imaging can provide more information for the clinical diagnosis, and magnetic resonance imaging (MRI) is a technique for pancreatitis.⁹² Gadolinium-diethylenetriaminepentaacetic acid (Gd-DTPA) is a common contrast agent for MRI. Although the MRI imaging effect is closely related to the concentration of Gd-DTPA in the lesion, Gd-DTPA lacks the ability to target pancreatic tissue. Some abnormally elevated markers or immune cells in local lesions of pancreatitis provide targets to this problem. As shown in Figure 6, in the case of a large number of macrophages infiltrating in the AP lesion, a mannose-coated nanoliposome containing Gd-DTPA (M-Gd-NL) was previously reported, which was easily phagocytosed by the focal macrophages, thus enhancing the imaging effect of AP. Moreover, since the degree of macrophage infiltration of MAP and SAP is different, the use of M-Gd-NL can further distinguish the severity of AP.⁹³ Similarly, the locally overexpressed lipase in AP is another target. A previous study synthesized Gd-DTPA fatty acid nanoparticles (Gd-DTPA-FA).⁹⁴ Gd-DTPA-FA can be disassembled under the action of excess lipase and release the Gd-DTPA. In the *L*-arginine-induced AP rat model, Gd-DTPA-FA significantly increased T_1 -weighted MRI signal intensity from 1 to 36 hours.

Furthermore, a near-infrared fluorescence and MRI dual-model nanoprobe has been reported recently to realize early visual diagnosis of SAP.⁹⁵ Deng et al prepared a dual-mode nanoprobe (Gd-DTPA-Cy5.5-PsLmAb) by coupling Cy5.5 (a near-infrared fluorescent dye) and P-selectin mAb on DTPA. Take advantage of P-selectin significantly upregulated at the SAP site, this nanoparticles have high spatial resolution and sensitivity in the early stage of SAP animal models. This is important to provide SAP patients with accurate treatment as soon as possible, thereby reducing their mortality.

Factors Affecting the Distribution of Nanomedicines in Pancreatitis Mice

The key to solving the current limitations of the treatment of pancreatitis is the targeting efficiency of nanoparticles, but few literatures have explored the factors affecting the distribution of nanoparticles in animal pancreatitis model.⁹⁶ As we




Figure 6 Mannose-modified gadolinium-containing nanoliposomes enhance local MRI imaging of AP. **(A)** M-Gd-NL is composed of DSPE-PEG-Man, DPPE-DTPA(Gd), and cholesterol stabilizes its structure. **(B)** MRI imaging of M-Gd-NL within 180 minutes of in vivo injection in mice with AP of different severity. **(C)** The comparison of MRI SNR signal and T_1 mapping relaxation time shows that M-Gd-NL can both diagnose MAP and SAP, but the imaging effect of SAP is the best.⁹³ All comparisons were performed between the two groups by one-way analysis of variance with Newman-Keuls posttest. Data are expressed as mean \pm SD (n=5). * $P < 0.05$; ** $P < 0.01$. Reproduced from Tian B, Liu R, Chen S, et al. Mannose-coated gadolinium liposomes for improved magnetic resonance imaging in acute pancreatitis. *Int J Nanomedicine*. 2017;12:1127–1141. Copyright 2017, Dove Medical Press.⁹³

all known, inflammation-related diseases often involve ELVIS effects.⁹⁷ Due to changes in vascular permeability during AP progression, the ELVIS effects may change during progression of AP. Previous studies have explored the effect of nanoparticles size on their behavior in different stages of AP.²¹ After intravenous administration of mesopore silica nanoparticles (MSN) at 60, 150 or 300 nm for 4 h in rats with MAP or SAP. The presence of MSN₁₅₀ in the pancreas was significantly greater than that of MSN₆₀ or MSN₃₀₀ in both MAP and SAP. The content of MSN₁₅₀ in pancreas, intestine and ascites of SAP rats was lower than that of MAP rats, indicating that the targeting performance of MSN in SAP rats was decreased. This may be related to more blood loss and slower blood flow in SAP rats. Such findings can guide us to pay attention to engineering principles when developing therapeutic nanoparticles in pancreatitis.

Conclusion and Prospect

Pancreatitis is an inflammatory disease, especially SAP, which can be life-threatening in severe cases. Without timely diagnosis, CP will seriously affect pancreatic function and induce pancreatic cancer, diabetes and other pancreatic diseases. With the development of pathology, its pathogenesis has been confirmed to be related to premature activation of trypsinogen, calcium overload, pancreatic microcirculation disturbance, NF- κ B pathway activation, leukocyte infiltration, and autophagy damage.

The rapid development of nanomedicine, which focus on targeted delivery and increase local drug accumulation, provides new opportunities for the treatment of pancreatitis. As the most common platforms, lipid and polymer nanoparticles tend to accumulate at sites of inflammation due to their appropriate particle size and ELVIS effect. Camouflaged by inflammatory cell membranes, the nanoparticles can be endowed with the ability to target APs. Therapeutic biomaterials such as antioxidant PAMAM and metallic nanoparticles have also been used to treat acute pancreatitis. We believe that, in accordance with the basic principles of design engineering (such as particle size distribution range, etc.), we should try to integrate the advantages of the above different carriers and propose a nanocarrier with advantages. On the other hand, nanomedicine in the diagnosis of pancreatitis mainly focuses on hypersensitivity detection of serum markers and imaging of pancreatic tissues, while existing studies are limited to amylase and lipase abnormal signals. Therefore, the efforts of researchers related to proteomics, genomics and other emerging disciplines will provide positive help for the development of specific markers for pancreatitis.

Finally, although there are numerous nanomedicine-based strategies for pancreatitis treatment, many experiments have only been carried out in simple animal models, which are insufficient for clinical studies. We believe that the support of pathology and statisticians will also help us to achieve multi-center, large scale clinical trials.

Acknowledgment

This work was supported by grants from the National Natural Science Foundation of China [82204304], the Sichuan Science and Technology Program [2021YJ0128], the Chengdu Science and Technology Program [2022-YF05-01626-SN], the Southwest Minzu University Research Startup Funds [RQD2022033], and the Fundamental Research Funds for the Central Universities [ZYN2022057].

Disclosure

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Mayerle J, Sendler M, Hegyi E, Beyer G, Lerch MM, Sahin-Toth M. Genetics, cell biology, and pathophysiology of pancreatitis. *Gastroenterology*. 2019;156:1951–1968 e1951. doi:10.1053/j.gastro.2018.11.081
2. Tenner S, Baillie J, DeWitt J, Vege SS; American College of Gastroenterology guideline: management of acute pancreatitis. *Am J Gastroenterol*. 2013;108(1400–1415):1416. doi:10.1038/ajg.2013.218
3. Geokas MC, Baltaxe HA, Banks PA, Silva J Jr, Frey CF. Acute pancreatitis. *Ann Intern Med*. 1985;103:86–100. doi:10.7326/0003-4819-103-1-86
4. Whitcomb DC, Frulloni L, Garg P, et al. Chronic pancreatitis: an international draft consensus proposal for a new mechanistic definition. *Pancreatolgy*. 2016;16:218–224. doi:10.1016/j.pan.2016.02.001
5. Sundar V, Senthil Kumar KA, Manickam V, Ramasamy T. Current trends in pharmacological approaches for treatment and management of acute pancreatitis - a review. *J Pharm Pharmacol*. 2020;72:761–775. doi:10.1111/jphp.13229

6. Glasbrenner B, Adler G. Pathophysiology of acute pancreatitis. *Hepatogastroenterology*. 1993;40:517–521.
7. Greenberg JA, Hsu J, Bawazeer M, et al. Clinical practice guideline: management of acute pancreatitis. *Can J Surg*. 2016;59:128–140. doi:10.1503/cjs.015015
8. Portelli M, Jones CD. Severe acute pancreatitis: pathogenesis, diagnosis and surgical management. *Hepatobiliary Pancreat Dis Int*. 2017;16:155–159. doi:10.1016/S1499-3872(16)60163-7
9. Yadav D, Timmons L, Benson JT, Dierkhising RA, Chari ST. Incidence, prevalence, and survival of chronic pancreatitis: a population-based study. *Am J Gastroenterol*. 2011;106:2192–2199. doi:10.1038/ajg.2011.328
10. Singh VK, Yadav D, Garg PK. Diagnosis and management of chronic pancreatitis: a review. *JAMA*. 2019;322:2422–2434. doi:10.1001/jama.2019.19411
11. Bayda S, Adeel M, Tuccinardi T, Cordani M, Rizzolio F. The history of nanoscience and nanotechnology: from chemical-physical applications to nanomedicine. *Molecules*. 2019;25:112. doi:10.3390/molecules25010112
12. Navalakhe RM, Nandedkar TD. Application of nanotechnology in biomedicine. *Indian J Exp Biol*. 2007;45:160–165.
13. Patil-Sen Y. Advances in nano-biomaterials and their applications in biomedicine. *Emerg Top Life Sci*. 2021;5:169–176. doi:10.1042/ETLS20200333
14. Pelaz B, Alexiou C, Alvarez-Puebla RA, et al. Diverse applications of nanomedicine. *ACS Nano*. 2017;11:2313–2381. doi:10.1021/acsnano.6b06040
15. Kim BY, Rutka JT, Chan WC. Nanomedicine. *N Engl J Med*. 2010;363:2434–2443. doi:10.1056/NEJMra0912273
16. Lammers T, Aime S, Hennink WE, Storm G, Kiessling F. Theranostic nanomedicine. *Acc Chem Res*. 2011;44:1029–1038. doi:10.1021/ar200019c
17. Freitas RA Jr. What is nanomedicine? *Nanomedicine*. 2005;1:2–9. doi:10.1016/j.nano.2004.11.003
18. Murday JS, Siegel RW, Stein J, Wright JF. Translational nanomedicine: status assessment and opportunities. *Nanomedicine*. 2009;5:251–273. doi:10.1016/j.nano.2009.06.001
19. Cao X, Hu Y, Luo S, et al. Neutrophil-mimicking therapeutic nanoparticles for targeted chemotherapy of pancreatic carcinoma. *Acta Pharm Sin B*. 2019;9:575–589. doi:10.1016/j.apsb.2018.12.009
20. Brandl T, Simic O, Skaenderup PR, et al. Trypsin inhibitors for the treatment of pancreatitis. *Bioorg Med Chem Lett*. 2016;26:4340–4344. doi:10.1016/j.bmcl.2016.07.029
21. Qiang H, Li J, Wang S, et al. Distribution of systemically administered nanoparticles during acute pancreatitis: effects of particle size and disease severity. *Pharmazie*. 2021;76:180–188. doi:10.1691/ph.2021.0156
22. Chiorean EG, Covelev AL. Pancreatic cancer: optimizing treatment options, new, and emerging targeted therapies. *Drug Des Devel Ther*. 2015;9:3529–3545. doi:10.2147/DDDT.S60328
23. Pearce AK, O'Reilly RK. Insights into active targeting of nanoparticles in drug delivery: advances in clinical studies and design considerations for cancer nanomedicine. *Bioconjug Chem*. 2019;30:2300–2311. doi:10.1021/acs.bioconjchem.9b00456
24. Mehnert W, Mader K. Solid lipid nanoparticles: production, characterization and applications. *Adv Drug Deliv Rev*. 2001;47:165–196. doi:10.1016/S0169-409X(01)00105-3
25. Cervin C, Vandoolaeghe P, Nistor C, Tiberg F, Johnsson M. A combined in vitro and in vivo study on the interactions between somatostatin and lipid-based liquid crystalline drug carriers and bilayers. *Mol Pharm*. 2015;12:2167–2179. doi:10.1021/acs.molpharmaceut.5b00155
26. Gong T, Zhang P, Deng C, Xiao Y, Gong T, Zhang Z. An effective and safe treatment strategy for rheumatoid arthritis based on human serum albumin and Koliphor(R) HS 15. *Nanomedicine*. 2019;14:2169–2187. doi:10.2217/nnm-2019-0110
27. Wang Q, Jiang H, Li Y, et al. Targeting NF- κ B signaling with polymeric hybrid micelles that co-deliver siRNA and dexamethasone for arthritis therapy. *Biomaterials*. 2017;122:10–22. doi:10.1016/j.biomaterials.2017.01.008
28. Xu Y, Mu J, Xu Z, et al. Modular acid-activatable acetone-based ketal-linked nanomedicine by dexamethasone prodrugs for enhanced anti-rheumatoid arthritis with low side effects. *Nano Lett*. 2020;20:2558–2568. doi:10.1021/acs.nanolett.9b05340
29. Pan W, Li Z, Qiu S, et al. Octahedral Pt-MOF with Au deposition for plasmonic effect and Schottky junction enhanced hydrogenothermal therapy of rheumatoid arthritis. *Mater Today Bio*. 2022;13:100214. doi:10.1016/j.mtbio.2022.100214
30. Tan J, Deng Z, Liu G, Hu J, Liu S. Anti-inflammatory polymersomes of redox-responsive polyprodrug amphiphiles with inflammation-triggered indomethacin release characteristics. *Biomaterials*. 2018;178:608–619. doi:10.1016/j.biomaterials.2018.03.035
31. Yang C, Hu T, Cao H, et al. Facile construction of chloroquine containing PLGA-based pDNA delivery system for efficient tumor and pancreatitis targeting in vitro and in vivo. *Mol Pharm*. 2015;12:2167–2179. doi:10.1021/acs.molpharmaceut.5b00155
32. Liu H, Liu S, Song X, et al. Nanoparticle encapsulated CQ/TAM combination harmonizes with MSCs in arresting progression of severity in AP mice through iNOS (IDO) signaling. *Mater Today Bio*. 2022;14:100226. doi:10.1016/j.mtbio.2022.100226
33. Cai H, Wang R, Guo X, et al. Combining gemcitabine-loaded macrophage-like nanoparticles and erlotinib for pancreatic cancer therapy. *Mol Pharm*. 2021;18:2495–2506. doi:10.1021/acs.molpharmaceut.0c01225
34. Zhang Y, Sun T, Jiang C. Biomacromolecules as carriers in drug delivery and tissue engineering. *Acta Pharm Sin B*. 2018;8:34–50. doi:10.1016/j.apsb.2017.11.005
35. Ismail OZ, Bhayana V. Lipase or amylase for the diagnosis of acute pancreatitis? *Clin Biochem*. 2017;50:1275–1280. doi:10.1016/j.clinbiochem.2017.07.003
36. Yao Q, Jiang X, Zhai YY, et al. Protective effects and mechanisms of bilirubin nanomedicine against acute pancreatitis. *J Control Release*. 2020;322:312–325. doi:10.1016/j.jconrel.2020.03.034
37. Stucki D, Stahl W. Carbon monoxide - beyond toxicity? *Toxicol Lett*. 2020;333:251–260. doi:10.1016/j.toxlet.2020.08.010
38. Nagao S, Taguchi K, Sakai H, et al. Carbon monoxide-bound hemoglobin vesicles ameliorate multiorgan injuries induced by severe acute pancreatitis in mice by their anti-inflammatory and antioxidant properties. *Int J Nanomedicine*. 2016;11:5611–5620. doi:10.2147/IJN.S118185
39. Tan X, Jia F, Wang P, Zhang K. Nucleic acid-based drug delivery strategies. *J Control Release*. 2020;323:240–252. doi:10.1016/j.jconrel.2020.03.040
40. Buddolla AL, Kim S. Recent insights into the development of nucleic acid-based nanoparticles for tumor-targeted drug delivery. *Colloids Surf B Biointerfaces*. 2018;172:315–322. doi:10.1016/j.colsurfb.2018.08.057
41. Oliveira C, Ribeiro AJ, Veiga F, Silveira I. Recent advances in nucleic acid-based delivery: from bench to clinical trials in genetic diseases. *J Biomed Nanotechnol*. 2016;12:841–862. doi:10.1166/jbn.2016.2245

42. Zhang T, Tian T, Lin Y. Functionalizing framework nucleic-acid-based nanostructures for biomedical application. *Adv Mater*. 2021;e2107820. doi:10.1002/adma.202107820
43. Zhang T, Cui W, Tian T, Shi S, Lin Y. Progress in biomedical applications of tetrahedral framework nucleic acid-based functional systems. *ACS Appl Mater Interfaces*. 2020;12:47115–47126. doi:10.1021/acsami.0c13806
44. Zhang X, Liu N, Zhou M, Li S, Cai X. The application of tetrahedral framework nucleic acids as a drug carrier in biomedicine fields. *Curr Stem Cell Res Ther*. 2021;16:48–56. doi:10.2174/1574888X15666200422103415
45. Shi S, Li Y, Zhang T, et al. Biological effect of differently sized tetrahedral framework nucleic acids: endocytosis, proliferation, migration, and biodistribution. *ACS Appl Mater Interfaces*. 2021;13:57067–57074. doi:10.1021/acsami.1c20657
46. Zhang Q, Lin S, Wang L, et al. Tetrahedral framework nucleic acids act as antioxidants in acute kidney injury treatment. *Chem Eng J*. 2021;413:127426. doi:10.1016/j.cej.2020.127426
47. Zhou M, Liu NX, Shi SR, et al. Effect of tetrahedral DNA nanostructures on proliferation and osteo/odontogenic differentiation of dental pulp stem cells via activation of the notch signaling pathway. *Nanomedicine*. 2018;14:1227–1236. doi:10.1016/j.nano.2018.02.004
48. Zhou M, Gao S, Zhang X, et al. The protective effect of tetrahedral framework nucleic acids on periodontium under inflammatory conditions. *Bioact Mater*. 2021;6:1676–1688. doi:10.1016/j.bioactmat.2020.11.018
49. Gao S, Zhou M, Li Y, et al. Tetrahedral framework nucleic acids reverse new-onset type 1 diabetes. *ACS Appl Mater Interfaces*. 2021;13:50802–50811. doi:10.1021/acsami.1c16151
50. Li Y, Tang Y, Shi S, et al. Tetrahedral framework nucleic acids ameliorate insulin resistance in type 2 diabetes mellitus via the PI3K/Akt pathway. *ACS Appl Mater Interfaces*. 2021;13:40354–40364. doi:10.1021/acsami.1c11468
51. Gao S, Wang Y, Li Y, et al. Tetrahedral framework nucleic acids reestablish immune tolerance and restore saliva secretion in a sjogren's syndrome mouse model. *ACS Appl Mater Interfaces*. 2021;13:42543–42553. doi:10.1021/acsami.1c14861
52. Wang Y, Li Y, Gao S, Yu X, Chen Y, Lin Y. Tetrahedral framework nucleic acids can alleviate taurocholate-induced severe acute pancreatitis and its subsequent multiorgan injury in mice. *Nano Lett*. 2022;22:1759–1768. doi:10.1021/acs.nanolett.1c05003
53. Oroojalian F, Beygi M, Baradaran B, Mokhtarzadeh A, Shahbazi MA. Immune cell membrane-coated biomimetic nanoparticles for targeted cancer therapy. *Small*. 2021;17:e2006484. doi:10.1002/sml.202006484
54. Beh CY, Prajnamitra RP, Chen LL, Hsieh PC. Advances in biomimetic nanoparticles for targeted cancer therapy and diagnosis. *Molecules*. 2021;26:5052. doi:10.3390/molecules26165052
55. Luk BT, Zhang L. Cell membrane-camouflaged nanoparticles for drug delivery. *J Control Release*. 2015;220:600–607. doi:10.1016/j.jconrel.2015.07.019
56. Xu CH, Ye PJ, Zhou YC, He DX, Wei H, Yu CY. Cell membrane-camouflaged nanoparticles as drug carriers for cancer therapy. *Acta Biomater*. 2020;105:1–14. doi:10.1016/j.actbio.2020.01.036
57. Zhang R, Wu S, Ding Q, et al. Recent advances in cell membrane-camouflaged nanoparticles for inflammation therapy. *Drug Deliv*. 2021;28:1109–1119. doi:10.1080/10717544.2021.1934188
58. Wang H, Liu Y, He R, et al. Cell membrane biomimetic nanoparticles for inflammation and cancer targeting in drug delivery. *Biomater Sci*. 2020;8:552–568. doi:10.1039/C9BM01392J
59. Mayerle J, Dummer A, Sendler M, et al. Differential roles of inflammatory cells in pancreatitis. *J Gastroenterol Hepatol*. 2012;27(Suppl 2):47–51. doi:10.1111/j.1440-1746.2011.07011.x
60. Zhou X, Cao X, Tu H, Zhang ZR, Deng L. Inflammation-targeted delivery of celastrol via neutrophil membrane-coated nanoparticles in the management of acute pancreatitis. *Mol Pharm*. 2019;16:1397–1405. doi:10.1021/acs.molpharmaceut.8b01342
61. Hassanzadeh P, Arbabi E, Rostami F. Coating of ferulic acid-loaded silk fibroin nanoparticles with neutrophil membranes: a promising strategy against the acute pancreatitis. *Life Sci*. 2021;270:119128. doi:10.1016/j.lfs.2021.119128
62. Hu F, Lou N, Jiao J, Guo F, Xiang H, Shang D. Macrophages in pancreatitis: mechanisms and therapeutic potential. *Biomed Pharmacother*. 2020;131:110693. doi:10.1016/j.biopha.2020.110693
63. Allawadhi P, Beyer G, Mahajan UM, Mayerle J. Novel insights into macrophage diversity during the course of pancreatitis. *Gastroenterology*. 2021;161:1802–1805. doi:10.1053/j.gastro.2021.09.049
64. Manohar M, Jones EK, Rubin SJS, et al. Novel circulating and tissue monocytes as well as macrophages in pancreatitis and recovery. *Gastroenterology*. 2021;161:2014–2029 e2014. doi:10.1053/j.gastro.2021.08.033
65. Zhang Q, Zhou J, Zhou J, Fang RH, Gao W, Zhang L. Lure-and-kill macrophage nanoparticles alleviate the severity of experimental acute pancreatitis. *Nat Commun*. 2021;12:4136. doi:10.1038/s41467-021-24447-4
66. Mitchell MJ, Billingsley MM, Haley RM, Wechsler ME, Peppas NA, Langer R. Engineering precision nanoparticles for drug delivery. *Nat Rev Drug Discov*. 2021;20:101–124. doi:10.1038/s41573-020-0090-8
67. Mohammadpour R, Ghandehari H. Mechanisms of immune response to inorganic nanoparticles and their degradation products. *Adv Drug Deliv Rev*. 2022;180:114022. doi:10.1016/j.addr.2021.114022
68. Khan S, Sharifi M, Bloukh SH, Edis Z, Siddique R, Falahati M. In vivo guiding inorganic nanozymes for biosensing and therapeutic potential in cancer, inflammation and microbial infections. *Talanta*. 2021;224:121805. doi:10.1016/j.talanta.2020.121805
69. Roy S, Liu Z, Sun X, et al. Assembly and degradation of inorganic nanoparticles in biological environments. *Bioconjug Chem*. 2019;30:2751–2762. doi:10.1021/acs.bioconjchem.9b00645
70. Xie P, Zhang L, Shen H, et al. Biodegradable MoSe₂-polyvinylpyrrolidone nanoparticles with multi-enzyme activity for ameliorating acute pancreatitis. *J Nanobiotechnology*. 2022;20:113. doi:10.1186/s12951-022-01288-x
71. Gao Y, Yu G, Xing K, et al. Finely tuned Prussian blue-based nanoparticles and their application in disease treatment. *J Mater Chem B*. 2020;8:7121–7134. doi:10.1039/D0TB01248C
72. Zhang W, Hu S, Yin JJ, et al. Prussian blue nanoparticles as multienzyme mimetics and reactive oxygen species scavengers. *J Am Chem Soc*. 2016;138:5860–5865. doi:10.1021/jacs.5b12070
73. Busquets MA, Estelrich J. Prussian blue nanoparticles: synthesis, surface modification, and biomedical applications. *Drug Discov Today*. 2020;25:1431–1443. doi:10.1016/j.drudis.2020.05.014
74. He Q, Yang H, Chen Y, et al. Prussian blue nanoparticles with peroxidase-mimicking properties in a dual immunoassays for glycocholic acid. *J Pharm Biomed Anal*. 2020;187:113317. doi:10.1016/j.jpba.2020.113317

75. Shukla A, Cano-Mejia J, Andricovich J, Burga RA, Sweeney EE, Fernandes R. An engineered Prussian blue nanoparticles-based nanoimmunotherapy elicits robust and persistent immunological memory in a TH-MYCN neuroblastoma model. *Adv Nanobiomed Res.* **2021**;1:2100021. doi:10.1002/anbr.202100021.
76. Xie X, Zhao J, Gao W, et al. Prussian blue nanozyme-mediated nanoscavenger ameliorates acute pancreatitis via inhibiting TLRs/NF-kappaB signaling pathway. *Theranostics.* **2021**;11:3213–3228. doi:10.7150/thno.52010
77. Li H, Wang P, Lin G, Huang J. The role of rare earth elements in biodegradable metals: a review. *Acta Biomater.* **2021**;129:33–42. doi:10.1016/j.actbio.2021.05.014
78. Yang B, Chen Y, Shi J. Reactive oxygen species (ROS)-based nanomedicine. *Chem Rev.* **2019**;119:4881–4985. doi:10.1021/acs.chemrev.8b00626
79. Khurana A, Anchi P, Allawadhi P, et al. Superoxide dismutase mimetic nanoceria restrains cerulein induced acute pancreatitis. *Nanomedicine.* **2019**;14:1805–1825. doi:10.2217/nnm-2018-0318
80. Khurana A, Anchi P, Allawadhi P, et al. Yttrium oxide nanoparticles reduce the severity of acute pancreatitis caused by cerulein hyperstimulation. *Nanomedicine.* **2019**;18:54–65. doi:10.1016/j.nano.2019.02.018
81. Abdel-Hakeem EA, Abdel-Hamid HA, Abdel Hafez SMN. The possible protective effect of Nano-Selenium on the endocrine and exocrine pancreatic functions in a rat model of acute pancreatitis. *J Trace Elem Med Biol.* **2020**;60:126480. doi:10.1016/j.jtemb.2020.126480
82. Yi Z, Luo X, Zhao L. Research advances in chitosan oligosaccharides: from multiple biological activities to clinical applications. *Curr Med Chem.* **2020**;27:5037–5055. doi:10.2174/0929867326666190712180147
83. Mei Q, Deng G, Huang Z, et al. Porous COS@SiO₂ nanocomposites ameliorate severe acute pancreatitis and associated lung injury by regulating the Nrf2 signaling pathway in mice. *Front Chem.* **2020**;8:720. doi:10.3389/fchem.2020.00720
84. Peng F, Setyawati MI, Tee JK, et al. Nanoparticles promote in vivo breast cancer cell intravasation and extravasation by inducing endothelial leakiness. *Nat Nanotechnol.* **2019**;14:279–286. doi:10.1038/s41565-018-0356-z
85. Chuang EY, Lin KJ, Huang TY, et al. An intestinal “transformers”-like nanocarrier system for enhancing the oral bioavailability of poorly water-soluble drugs. *ACS Nano.* **2018**;12:6389–6397. doi:10.1021/acsnano.8b00470
86. Hung J, Awasthi R, Klibanov AL, Kelly KA. Identification of novel ligands for targeted antifibrotic therapy of chronic pancreatitis. *Int J Nanomedicine.* **2021**;16:5495–5512. doi:10.2147/IJN.S318331
87. Shahin NN, Shamma RN, Ahmed IS. A nano-liposomal formulation of caffeic acid phenethyl ester modulates Nrf2 and NF- κ B signaling and alleviates experimentally induced acute pancreatitis in a rat model. *Antioxidants.* **2022**;11:1536. doi:10.3390/antiox11081536
88. Sanz Del Olmo N, Pena Gonzalez CE, Rojas JD, et al. Antioxidant and antibacterial properties of carbosilane dendrimers functionalized with polyphenolic moieties. *Pharmaceutics.* **2020**;12:698. doi:10.3390/pharmaceutics12080698
89. Lee CY, Nanah CN, Held RA, et al. Effect of electron donating groups on polyphenol-based antioxidant dendrimers. *Biochimie.* **2015**;111:125–134. doi:10.1016/j.biochi.2015.02.001
90. Tang Y, Han Y, Liu L, et al. Protective effects and mechanisms of G5 PAMAM dendrimers against acute pancreatitis induced by caerulein in mice. *Biomacromolecules.* **2015**;16:174–182. doi:10.1021/bm501390d
91. Attia MS, Al-Radadi NS. Progress of pancreatitis disease biomarker alpha amylase enzyme by new nano optical sensor. *Biosens Bioelectron.* **2016**;86:413–419. doi:10.1016/j.bios.2016.06.079
92. Hafezi-Nejad N, Singh VK, Fung C, Takahashi N, Zaheer A. MR imaging of autoimmune pancreatitis. *Magn Reson Imaging Clin N Am.* **2018**;26:463–478. doi:10.1016/j.mric.2018.03.008
93. Tian B, Liu R, Chen S, et al. Mannose-coated gadolinium liposomes for improved magnetic resonance imaging in acute pancreatitis. *Int J Nanomedicine.* **2017**;12:1127–1141. doi:10.2147/IJN.S123290
94. Zhang HW, Wang LQ, Xiang QF, et al. Specific lipase-responsive polymer-coated gadolinium nanoparticles for MR imaging of early acute pancreatitis. *Biomaterials.* **2014**;35:356–367. doi:10.1016/j.biomaterials.2013.09.046
95. Long L, Deng L, Wang L, et al. P-selectin-based dual-model nanoprobe used for the specific and rapid visualization of early detection toward severe acute pancreatitis in vivo. *ACS Biomater Sci Eng.* **2020**;6:5857–5865. doi:10.1021/acsbomaterials.0c00596
96. Cai H, Dai X, Guo X, et al. Ataxia telangiectasia mutated inhibitor-loaded copper sulfide nanoparticles for low-temperature photothermal therapy of hepatocellular carcinoma. *Acta Biomater.* **2021**;127:276–286. doi:10.1016/j.actbio.2021.03.051
97. Habtezion A. Inflammation in acute and chronic pancreatitis. *Curr Opin Gastroenterol.* **2015**;31:395–399. doi:10.1097/MOG.0000000000000195

International Journal of Nanomedicine

Dovepress

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine, Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-nanomedicine-journal>