



The Minimal Clinically Important Differences as Well as Statistical Differences of Main Endpoints are Important in Comparing Postoperative Benefits of Different Analgesic Modalities [Letter]

Tian Tian , Xin-Tao Li , Fu-Shan Xue 

Department of Anesthesiology, Beijing Friendship Hospital, Capital Medical University, Beijing, People's Republic of China

Correspondence: Fu-Shan Xue, Department of Anesthesiology, Beijing Friendship Hospital, Capital Medical University, No. 95 Yong-An Road, Xi-Cheng District, Beijing, 100050, People's Republic of China, Tel +86-13911177655, Fax +86-10-63138362, Email xuefushan@aliyun.com; fushanxue@outlook.com

Dear Editor

In a randomized controlled trial including 99 patients who underwent elective laparoscopic bariatric surgery, Sun and colleagues¹ showed that both intravenous infusion of lidocaine (IIL) and ultrasound-guided transverse abdominal plane block (UG-TAPB) provided good postoperative recovery and analgesia, but the IIL resulted in better analgesia at 12 h and 24 h postoperatively compared with UG-TAPB. This study has potential implications, but we would like to remind the readers to pay attention to the clinical significance of their findings.

First, the Quality of Recovery-40 (QoR-40) score at 24 h after surgery was used as the primary endpoint. In the available literature, a 10-point between-group difference in the total QoR-40 scores is generally considered as the minimal clinically important difference.² We noted that the total QoR-40 scores were significantly higher in the patients receiving IIL and UG-TAPB compared with the patients receiving the control intervention, but the net between-group difference in the median of total QoR-40 scores was less than 10. That is, the between-group difference in the quality of postoperative recovery is statistically significant, but its clinical significance is not clearly evident.

Second, this study assessed the visual analog scale (VAS) score of postoperative pain in the resting state, but not the VAS pain score in the active state. In fact, active pain is more severe than resting pain following bariatric surgery, and effective control of active pain after abdominal surgery is very important for the successful use of enhanced recovery after surgery protocols.³ In this study, the VAS resting pain scores at some time-points after surgery were significantly different among the three groups, but the net between-group differences in mean VAS resting pain scores at all time-points postoperatively were less than 1. We would like to remind the readers that the recommended minimal clinically important differences for postoperative pain scores in the available literature are 1.5 in the resting state and 1.8 in the active state when pain is assessed by a 0–10 VAS score.⁴ That is, the improvements in postoperative pain control by IIL and UG-TAPB compared with the control intervention do also not exceed the recommended minimal clinically important differences.

Finally, intravenous dezocine was applied on demand to maintain a VAS score of 4 or less, and dezocine consumption within 24 h postoperatively was significantly decreased in the patients receiving IIL and UG-TAPB compared with the patients receiving the control intervention. However, when between-group differences in postoperative analgesic consumption are compared, it is generally required that the dosage of analgesic used for postoperative pain control should be converted into morphine milligram equivalents in oral or intravenous form.^{3,4} Furthermore, it is commonly recommended that the minimal clinically important difference of morphine milligram equivalents for postoperative pain control is an absolute reduction of 10 mg intravenous morphine in the 24 h.⁴ As the equianalgesic conversion factor of morphine and

dezocine is 1,⁵ the net between-group differences in mean dezocine consumption within 24 h postoperatively in this study are only equivalent to 0.9–5.91 mg intravenous morphine. Accordingly, the real clinical significance of postoperative opioid sparing with IIL or UG-TAPB in this study should be interpreted with caution.

We believe that clarification of the above issues will improve the interpretation of findings in this study.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

All authors have received no financial support and have no potential conflicts of interest in this communication.

References

1. Sun J, Wang S, Wang J, Gao X, Wang G. Effect of intravenous infusion of lidocaine compared with ultrasound-guided transverse abdominal plane block on the quality of postoperative recovery in patients undergoing laparoscopic bariatric surgery. *Drug Des Devel Ther*. 2022;16:739–748. doi:10.2147/DDDT.S356880
2. Kim DH, Oh YJ, Lee JG, Ha D, Chang YJ, Kwak HJ. Efficacy of ultrasound-guided serratus plane block on postoperative quality of recovery and analgesia after video-assisted thoracic surgery: a randomized, triple-blind, placebo-controlled study. *Anesth Analg*. 2018;126(4):1353–1361. doi:10.1213/ANE.0000000000002779
3. Albrecht E, Kirkham KR, Endersby RV, et al. Ultrasound-guided transversus abdominis plane (TAP) block for laparoscopic gastric-bypass surgery: a prospective randomized controlled double-blinded trial. *Obes Surg*. 2013;23(8):1309–1314. doi:10.1007/s11695-013-0958-3
4. Laigaard J, Pedersen C, Rønsbo TN, Mathiesen O, Karlsen APH. Minimal clinically important differences in randomised clinical trials on pain management after total Hip and knee arthroplasty: a systematic review. *Br J Anaesth*. 2021;126(5):1029–1037. doi:10.1016/j.bja.2021.01.021
5. Ye RR, Jiang S, Xu X, Lu Y, Wang YJ, Liu JG. Dezocine as a potent analgesic: overview of its pharmacological characterization. *Acta Pharmacol Sin*. 2021;43(7):1646–1657. doi:10.1038/s41401-021-00790-6

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Drug Design, Development and Therapy 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Drug Design, Development and Therapy editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>

<https://doi.org/10.2147/DDDT.S370327>