LETTER

A Response to Article "Seroprevalence of Hepatitis B Virus, Hepatitis C Virus, Syphilis and Associated Factors Among Female Sex Workers in Gondar Town, Northwest Ethiopia" [Letter]

Rina Isnawati 10*, Widoretno Widoretno 10*, Didik Tulus Subekti 10*

Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency, Cibinong, West Java, Indonesia

*These authors contributed equally to this work

Correspondence: Widoretno Widoretno; Didik Tulus Subekti, Email wido009@brin.go.id; didi028@brin.go.id

Dear editor

We have read and appreciated the article entitled "Seroprevalence of Hepatitis B Virus, Hepatitis C Virus, Syphilis and Associated Factors Among Female Sex Workers in Gondar Town, Northwest Ethiopia".¹ This study draws our attention to providing another insight that we hope will be useful in further studies. The study by Wondmagegn et al is interesting to observe although several other similar studies have been found. The work carried out by the authors has similar ideas to previous publications with the additional observation of serological testing for anti-HCV and Syphilis antibodies.^{2,3} Initially we expected to find a more detailed discussion of the causative mechanism that describes the relationship between risk factors with infection rates and prevalence, unfortunately this is not found in the discussion as in other publications.^{2,3} However, there are some points that can be leveraged to make the study more interesting if some data can be informed or available.

Observation of HBsAg solely is considered inadequate from two points of view. First, HBsAg examination without being followed by other serological tests (HbsAb an HBcAb) could not distinguish whether acute infection, chronic infection, recovery (viral clearance), or co-infection with another HBV serotype.⁴ Clear information in this condition is very useful for elaborating data related to risk factors and infection status of respondents being studied. Second, put alongside the results of the HBsAg test with anti-HCV and syphilis antibody data is not appropriate. Anti-HCV and anti-Syphilis showed that the respondent had been in contact with microorganisms, whether exposure or infection could not be ascertained. Both only provide evidence of developed immunity as a result of past natural infection, post-vaccination or even development of immunity that occurs concurrently with the disease (acute or chronic).⁵ Therefore at least it is highly recommended to do both HBsAg and HBsAb testing simultaneously.

The authors report that HBV, HCV and Syphilis are common occurs via the sexual transmission route in female sex workers (FSWs). Naturally, sexually transmitted infections will increase the risk of multiple infections in respondents. In the publication, authors seems to have been overlooked since multiple infection data such as HBV+HCV+Syphilis, HBV +Syphilis, HBV+HCV and HCV+Syphilis will provide more comprehensive information on actual field situations. This is very useful to provide an overview of the status of respondents in those epidemiological study which is able to capture the actual health status of each FSWs. We therefore suggest that some of the points we have mentioned and criticized above should be considered in future studies.

Disclosure

The authors report no conflicts of interest in this communication.

cc 0 (so 2022 Isnawati et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms by and incorporate the Greative Commons Attribution – Non Commercial (unported, v3.0). License (http://creativecommons.org/licenses/by-nc/3.0/). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

References

- Wondmagegn M, Wondimeneh Y, Getaneh A, Ayalew G. Seroprevalence of hepatitis B virus, hepatitis C virus, syphilis and associated factors among female sex workers in Gondar Town, Northwest Ethiopia. *Infect Drug Resist.* 2022;15:5915–5927. doi:10.2147/IDR.S380952
- 2. Daka D, Hailemeskel G, Fenta DA. Seroprevalence of hepatitis B virus and associated factors among female sex workers using respondent-driven sampling in Hawassa City, Ethiopia. *Infect Drug Resist.* 2021;14:4301–4311. doi:10.2147/IDR.S332333
- Metaferia Y, Ali A, Eshetu S, Gebretsadik D. Seroprevalence and associated factors of human immunodeficiency virus, Treponema pallidum, hepatitis B virus, and hepatitis C virus among female sex workers in Dessie City, Northeast Ethiopia. *Biomed Res Int.* 2021;2021:1–13. doi:10.1155/ 2021/6650333
- 4. Kitt E, Hayes M, Cárdenas AM, Green AM. Interpretation and management of positive anti-hepatitis B core antibody tests in immunocompromised pediatric patients. *Transpl Infect Dis.* 2019;21:e13074. doi:10.1111/tid.13074
- 5. Mast EE, Weinbaum CM, Fiore AE, et al; Advisory Committee on Immunization Practices (ACIP) Centers for Disease Control and Prevention (CDC). A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP) Part 1: immunization of infants, children, and adolescents. *MMWR Recomm Rep.* 2005;54:1–33.

Dove Medical Press encourages responsible, free and frank academic debate. The contentTxt of the Infection and Drug Resistance 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Infection and Drug Resistance editors. While all reasonable steps have been taken to confirm the contentTxt of each letter, Dove Medical Press accepts no liability in respect of the contentTxt of any letter, nor is it responsible for the contentTxt and accuracy of any letter to the editor.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/infection-and-drug-resistance-journal

https://doi.org/10.2147/IDR.S400708

7692 🖪 😏 in 🖸 DovePress