

# Performance of Interferon- $\gamma$ Release Assays in Patients with *Mycobacterium kansasii* Infection

Chun-Hai Gao<sup>1</sup>, Yan-An Zhang<sup>2,3</sup>, Mao-Shui Wang<sup>3-5</sup> 

<sup>1</sup>Department of Clinical Laboratory, Linyi People's Hospital, Linyi, People's Republic of China; <sup>2</sup>Department of Cardiovascular Surgery, Shandong Public Health Clinical Center, Cheeloo College of Medicine, Shandong University, Jinan, People's Republic of China; <sup>3</sup>Shandong Key Laboratory of Infectious Respiratory Disease, Jinan, People's Republic of China; <sup>4</sup>Department of Laboratory Medicine, Shandong Provincial Chest Hospital, Cheeloo College of Medicine, Shandong University, Jinan, People's Republic of China; <sup>5</sup>Department of Laboratory Medicine, Shandong Public Health Clinical Center, Cheeloo College of Medicine, Shandong University, Jinan, People's Republic of China

Correspondence: Yan-An Zhang, Department of Cardiovascular Medicine, Shandong Public Health Clinical Center, 46# Lishan Road, Jinan, 250103, People's Republic of China, Email zhangtbgrou@163.com; Mao-Shui Wang, Department of Laboratory Medicine, Shandong Provincial Chest Hospital, 46# Lishan Road, Jinan, 250013, People's Republic of China, Email wangmaoshui@gmail.com

**Objective:** To evaluate the performance of interferon- $\gamma$  release assays (IGRAs) in patients with *Mycobacterium kansasii* infection.

**Methods:** Consecutive patients between May 2012 and June 2021 who had positive for mycobacterial cultures and who underwent IGRAs (T-SPOT.TB or QuantiFERON-TB Gold [QFT-G]) were included in the analysis. The IGRA positivity rates among patients with *M. kansasii* isolates were then calculated. If *M. kansasii* was identified in at least two sputum samples or in sterile samples, *M. kansasii* disease was then diagnosed. Otherwise, colonisation was considered.

**Results:** During the study period, 54 patients with *M. kansasii* infection underwent T-SPOT.TB (n=48) or QFT-G (n=6) assays. The mean age was 44.1 $\pm$ 13.4 years, 85.2% (46/54) were male. Eight patients were diagnosed with *M. kansasii* disease and another 46 patients were considered to have colonisation. Twenty-four patients (T-SPOT.TB, n=23; QFT-G, n=1) were positive for IGRAs, for an overall rate of 44.4% (24/54; T-SPOT.TB, 47.9% [23/48]; QFT-G, 16.7% [1/6]) for IGRAs, 25.0% (2/8) for *M. kansasii* disease, and 47.8% (22/46) for colonisation.

**Conclusion:** Positive IGRA rates were relatively low in patients with *M. kansasii* infection. More efforts are required to improve the performance of IGRAs in diagnosing *M. kansasii* infection.

**Keywords:** interferon-gamma release assay, tuberculosis, *M. kansasii*, diagnosis, infection

## Introduction

Interferon- $\gamma$  (IFN- $\gamma$ ) release assays (IGRAs) are designed to identify tuberculosis (TB) infections. Currently, two commercial assays are available; each uses different platforms to measure the IFN- $\gamma$  production stimulated by early secretory antigenic target (ESAT)-6 and culture filtrate protein (CFP)-10.<sup>1</sup> The QuantiFERON-TB assay (QFT; Qiagen, Hilden, Germany) is an overnight whole-blood enzyme-linked immunosorbent assay (ELISA) for quantitation of IFN- $\gamma$  in the supernatant, whereas the T-SPOT.TB (Oxford Immunotec, Oxford, UK) measures the number of mononuclear cells secreting IFN- $\gamma$  via an overnight enzyme-linked immunospot assay. In general, an IGRA is sensitive for detecting TB infection, with a sensitivity of 0.94 for active TB and 0.91 for latent TB infection.<sup>2</sup> However, IGRAs have shown limitations in borderline intervals, as well as in consistency. For example, caution should be taken when interpreting QFT results close to the assay cut-off, where assay conversion/reversion is likely to be observed in serial testing.<sup>3</sup> For Chinese village doctors, the concordance between the T-SPOT and QFT-Plus assays was 88.93% (Kappa coefficient: 0.73), and this remains unsatisfactory.<sup>4</sup>

For many decades, the tuberculin skin test (TST) was the only test available for detecting TB infection. However, the TST has several well-known disadvantages, including cross-reactivity with *Bacillus Calmette-Guérin* (BCG) and non-tuberculous mycobacteria (NTM), poor adherence due to return for results and wide variations observed in reading the results. IGRAs have recently been recommended in clinical practice and have several advantages for the diagnosis of TB

infection<sup>1,5–7</sup>. Compared to the TST, IGRAs have several advantages, such as being unaffected by BCG vaccination, IGRAs may produce fewer false positive reactions due to exposure to environmental mycobacteria. Compared with the TST, IGRAs are also less time consuming, because only a single patient visit is required and the results can be available within 24 hours.

NTMs are opportunistic pathogens and are widely distributed in the environment. Current, evidence suggests that the isolation rate of NTMs has been increasing in recent years.<sup>8</sup> In addition, human immunodeficiency virus (HIV) infection is a known risk factor for mycobacterial disease (both TB and NTM infections),<sup>9</sup> and its epidemic increases the complexity of the situation regarding NTM diseases. *Mycobacterium kansasii* is one of the most common NTM species, accounting for about 10% of NTM species isolated from clinical samples.<sup>10–12</sup> Until now, the diagnosis of NTM disease has mostly relied on microbiological evidence showing positive at least two separate sputum samples.<sup>13</sup> This diagnostic criterion is very specific and may contribute to a significantly missed diagnosis of *M. kansasii* infection. Therefore, more efforts aimed at improving the clinical and radiologic items of the guidelines would improve the effective diagnosis of NTM disease.<sup>13</sup> NTMs like *M. kansasii* that shares the RD1 region and show expression of ESAT-6 and CFP-10, can give rise to positive IGRA results.<sup>14</sup> Although the IGRA results for *M. kansasii* infection were reported in our previous study, the sample size (n=4) was too small to provide full details of the IGRA performance.<sup>15</sup> Therefore, the present study adopted a larger sample size to evaluate the performance of IGRAs in patients with *M. kansasii* infection and to assess the actual role of IGRAs in detecting *M. kansasii* infections.

## Methods

This study was conducted at Shandong Provincial Chest Hospital and conformed to the tenets of the Declaration of Helsinki. The study protocol was approved by the Ethics Committee of Shandong Provincial Chest Hospital. Due to the retrospective nature of this investigation and the anonymous nature of the data collection, written informed consent was waived by that Ethics Committee. Between May 2012 and June 2021, consecutive patients who were positive for mycobacterial culture and underwent IGRAs (T-SPOT.TB or QuantiFERON-TB Gold [QFT-G]) were included in the analysis. The patient data were retrieved from the laboratory information system (Ruimei, Shanghai, China). Other clinical characteristics, such as age, sex and HIV status, were also collected.

Sputum samples were cultured using Löwenstein-Jensen medium method and corresponding quality control was conducted as previously reported.<sup>10</sup> *M. kansasii* was identified using a Mycobacteria Identification Array Kit (CapitalBio, Beijing, China) or by sequencing 16S rDNA.<sup>10</sup> If *M. kansasii* was identified in at least two sputum samples or in sterile samples, *M. kansasii* disease was then diagnosed.<sup>16,17</sup> Otherwise, colonisation was considered.

The T-SPOT.TB and QFT-G IGRAs were performed according to the manufacturer's instructions. A positive T-SPOT.TB test result was defined using a cut-off value of at least 6 spots. For QFT-G results, the manufacturer's indicated cut-off of more than 0.35 IU/mL was considered a positive test result.

Continuous variables were presented as mean  $\pm$  standard deviation (SD) and categorical variables were described as count and percentages.

## Results

During the study period, 25,617 mycobacterial strains were found in the clinical samples. Of these, 1107 (4.3%) strains were identified as NTM species, including 110 strains isolated from outpatients and 997 strains from 828 inpatients. The 997 strains were identified as *M. intracellulare* (n=662, 66.4%), *M. kansasii* (n=111, 11.1%), *M. chelonae* (n=103, 10.3%), *M. fortuitum* (n=39, 3.9%), *M. gordonae* (n=20, 2.0%), *M. avium* (n=20, 2.0%), and others (n=42, 4.2%). A total of 111 *M. kansasii* isolates were collected from 97 inpatients.

Of the 97 patients, 54 underwent T-SPOT.TB (n=48) or QFT-G (n=6). The mean age was 44.1 $\pm$ 13.4 years, 85.2% (46/54) of them were male, and 97.6% (40/41) were HIV-negative. Of the 54, 8 were diagnosed with *M. kansasii* disease and the remaining 46 patients were considered to have colonisation. In total, 24 patients (T-SPOT.TB, n=23; QFT-G, n=1) were positive for IGRAs and the positive IGRA rate in patients with *M. kansasii* was calculated as 44.4% (24/54; T-SPOT.TB, 47.9% [23/48]; QFT-G, 16.7% [1/6]). The positivity rates were further calculated as 25.0% (2/8) for *M. kansasii* disease and 47.8% (22/46) for colonisation. Table 1 shows the characteristics of the included patients.

**Table I** The IGRA Results and Clinical Characteristics of Patients with *M. kansasii* Infection

	IGRA (+)	IGRA (-)	Total
N	24 (44.4%)	30 (55.6%)	54
Age (years)	42.9±10.4	45.1±15.5	44.1±13.4
Male, %	87.5% (21/24)	83.3% (25/30)	85.2% (46/54)
HIV (-)	100% (16/16)	96.0% (24/25)	97.6% (40/41)
Infection			
<i>M. kansasii</i> diseases	2 (8.3%)	6 (20.0%)	8 (14.8%)
Colonization	22 (91.7%)	24 (80.0%)	46 (85.2%)
IGRA results	T-SPOT.TB (n=23), QFT (n=1)	T-SPOT.TB (n=25), QFT (n=5)	T-SPOT.TB (n=48), QFT (n=6)

**Abbreviations:** IGRA, interferon-gamma release assay; HIV, human immunodeficiency virus; QFT, QuantiFERON-TB Gold.

## Discussion

A significant IFN- $\gamma$  production in response to a high conservation antigen of *M. tuberculosis* was previously observed in individuals with NTM infection.<sup>18</sup> Hence, positive IGRA results are expected among patients with *M. kansasii* infection, as both the ESAT-6 and CFP-10 antigens are found in *M. kansasii*.<sup>19,20</sup> In this study, assessment of the IGRA performance in patients with *M. kansasii* infection yielded an overall positivity rate of 44.4%. Our data suggested that the T cell response to antigens of *M. kansasii* is relatively weak and that the diagnostic value of IGRAs may be limited for *M. kansasii* infection.

However, in China, the use of IGRAs has several advantages in the diagnosis of *M. kansasii* infection. First, if anti-TB therapy fails, a positive IGRA result would increase the suspicion of *M. kansasii* infection and provide a rapid diagnosis.<sup>14</sup> Second, *M. kansasii* is usually considered pathogenic, as it readily causes disease.<sup>21</sup> A previous study demonstrated that a single positive culture for *M. kansasii* in the proper context could provide sufficient evidence for treatment initiation.<sup>22</sup> Hence, single isolation and a positive IGRA may be appropriate criteria for the diagnosis of *M. kansasii* disease. Third, *M. intracellulare*, *M. kansasii* and *M. fortuitum* are the three most common NTM species in China.<sup>8,10</sup> These epidemiological characteristics which were similar to our reports, raise several clinical implications for IGRA identification of *M. kansasii* infection. For example, IGRAs have the ability to discriminate *M. kansasii* infections from infections by other NTMs (such as *M. intracellulare* and *M. fortuitum*, which lack ESAT-6 and CFP-10).<sup>14</sup> A positive IGRA could increase the certainty of a diagnosis of *M. kansasii* infection.

At present, several studies with small sample sizes have been performed to evaluate of IGRA performance in *M. kansasii* infection. Kobashi Y et al evaluated the clinical usefulness of the QFT-2G test in patients with NTM disease and found a positive response rate of 52% for QFT-2G test in 33 patients with *M. kansasii* disease.<sup>23</sup> In a preliminary study conducted in Poland, positive IGRA results were found in 22% (2/9) of patients with *M. kansasii* infection.<sup>24</sup> In a Danish study, only two patients with *M. kansasii* infection were tested using QFT-G, but both were positive for QFT-G.<sup>25</sup> In Japan, a retrospective study showed that patients with *M. kansasii* infection had positive IGRA rates ranging between 25.9% and 33.3%.<sup>26</sup> By contrast, due to the lack of ESAT-6 and CFP-10, the IGRA positivity rate was significantly lower for infection with *M. avium* complex (1–2%),<sup>23,27</sup> which is a major source of NTM infection.<sup>23,28</sup>

The response to ESAT-6 and CFP-10 antigens is weaker in patients with *M. kansasii* infection than in those with TB infection. One reason may be that differences in ESAT-6 and CFP-10 between *M. kansasii* and *M. tuberculosis* have been reported for nucleotide sequences (10–12%) and amino acid sequences (<5%); therefore, these differences may attenuate the response. Indirect evidence has indicated attenuation of the T cell response in TB patients due to the differences in TB-ESAT-6 and *M. kansasii*-ESAT-6.<sup>29</sup> Another reason may be that the production capacity of IFN- $\gamma$  is lower in NTM patients than in TB patients.<sup>30</sup>

Impairment of the IFN- $\gamma$  responses dependent on stimulation has been observed in patients with NTM infection,<sup>31</sup> suggesting an immunologic imbalance in NTM infection. Future studies should re-evaluate the optimal cut-off values of IGRAs for the diagnosis of *M. kansasii* infection with respect to CFP-10 and ESAT-6.<sup>32,33</sup> Recently, a highly specific protein that can stimulate specific IFN- $\gamma$  secretion has been designed for the diagnosis of *M. abscessus* infection. This novel assay has a moderate AUC of 0.773 for diagnosis,<sup>34</sup> suggesting antigens originating from *M. kansasii* could also be used to improve the IGRA results for *M. kansasii* infection.

Our study had several limitations. First, it was conducted in a high-TB burden country, so latent TB infection may affect our results. Second, the distribution of NTM species a geographical variation; therefore, the epidemiological characteristics could have a significant effect on IGRA usefulness. Third, although the study had a small sample size, it remains one of the largest studies compared with others. Fourth, the IGRAs are designed for the evaluation of TB infection; therefore, further studies are required to evaluate the IFN- $\gamma$  response to antigens from NTM for the diagnosis of NTM diseases.<sup>34</sup> Fifth, two kinds of commercial IGRAs were used in this study, and this could have introduced an additional bias due to the inherent differences between them, such as the TB antigens (T-SPOT.TB assays, RD-1 antigens; QFT assays, TB 7.7 and RD-1 antigens),<sup>35</sup> technical platforms (T-SPOT.TB uses effector T cells; QFT determines IFN- $\gamma$  levels), specimens used (QFT uses whole blood cells; T-SPOT.TB uses mononuclear cells), indeterminate results (these are more common in QFT),<sup>36</sup> and diagnostic performance.<sup>37,38</sup> Therefore, interpreting the results requires caution.

In conclusion, our data demonstrated that a significant proportion of *M. kansasii* infections generate positive results for IGRAs. Infection by *M. kansasii* could therefore induce IFN- $\gamma$  production with the TB antigens. However, further studies are required to address this more completely, and to evaluate the IFN- $\gamma$  responses to *M. kansasii* antigens for the diagnosis of *M. kansasii* infections.

## Data Sharing Statement

The raw data supporting the conclusions of this article will be made available by the authors without undue reservation.

## Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

The study was supported by the Young Taishan Scholars Program of Shandong Province (2021).

## Disclosure

The authors have no competing interests in this work.

## References

1. Mazurek GH, Jereb J, Vernon A., et al. Updated guidelines for using Interferon Gamma Release Assays to detect Mycobacterium tuberculosis infection - United States, 2010. *MMWR Recomm Rep*. 2010;59:1–25.
2. Sotgiu G, Saderi L, Petruccioli E, et al. QuantiFERON TB Gold Plus for the diagnosis of tuberculosis: a systematic review and meta-analysis. *J Infect*. 2019;79:444–453. doi:10.1016/j.jinf.2019.08.018
3. Nemes E, Rozot V, Geldenhuys H, et al. Optimization and Interpretation of Serial QuantiFERON Testing to Measure Acquisition of Mycobacterium tuberculosis Infection. *Am J Respir Crit Care Med*. 2017;196:638–648. doi:10.1164/rccm.201704-0817OC
4. Zhang H, Xin H, Wang D, et al. Serial testing of Mycobacterium tuberculosis infection in Chinese village doctors by QuantiFERON-TB Gold Plus, QuantiFERON-TB Gold in-Tube and T-SPOT.TB. *J Infect*. 2019;78:305–310. doi:10.1016/j.jinf.2019.01.008
5. Sosa LE, Njie GJ, Lobato MN, et al. Tuberculosis Screening, Testing, and Treatment of U.S. Health Care Personnel: recommendations from the National Tuberculosis Controllers Association and CDC, 2019. *MMWR Morb Mortal Wkly Rep*. 2019;68:439–443. doi:10.15585/mmwr.mm6819a3
6. Santin M, Garcia-Garcia JM, Rigau D, et al. Executive Summary of the Guidelines for the Use of interferon-gamma Release Assays in the Diagnosis of Tuberculosis Infection. *Arch Bronconeumol*. 2016;52:477–481. doi:10.1016/j.arbres.2016.02.020

7. Getahun H, Matteelli A, Abubakar I, et al. Management of latent *Mycobacterium tuberculosis* infection: WHO guidelines for low tuberculosis burden countries. *Eur Respir J*. 2015;46:1563–1576. doi:10.1183/13993003.01245-2015
8. Xu D, Han C, Wang MS, Wang JL. Increasing prevalence of non-tuberculous mycobacterial infection from 2004–2009 to 2012–2017: a laboratory-based surveillance in China. *J Infect*. 2018;76:422–424. doi:10.1016/j.jinf.2017.12.007
9. Corbett EL, Churchyard GJ, Clayton T, et al. Risk factors for pulmonary mycobacterial disease in South African gold miners. A case-control study. *Am J Respir Crit Care Med*. 1999;159:94–99. doi:10.1164/ajrcm.159.1.9803048
10. Jing H, Wang H, Wang Y, et al. Prevalence of nontuberculous mycobacteria infection, China, 2004–2009. *Emerg Infect Dis*. 2012;18:527–528. doi:10.3201/eid1803.110175
11. Yu X, Liu P, Liu G, et al. The prevalence of non-tuberculous mycobacterial infections in mainland China: systematic review and meta-analysis. *J Infect*. 2016;73:558–567. doi:10.1016/j.jinf.2016.08.020
12. Tan Y, Deng Y, Yan X, et al. Nontuberculous mycobacterial pulmonary disease and associated risk factors in China: a prospective surveillance study. *J Infect*. 2021;83:46–53. doi:10.1016/j.jinf.2021.05.019
13. Daley CL, Iaccarino JM, Lange C, et al. Treatment of nontuberculous Mycobacterial Pulmonary Disease: an Official ATS/ERS/ESCMID/IDSA Clinical Practice Guideline. *Clin Infect Dis*. 2020;71:905–913. doi:10.1093/cid/ciaa1125
14. Kuznetcova TI, Sauty A, Herbot CP. Uveitis with occult choroiditis due to *Mycobacterium kansasii*: limitations of interferon-gamma release assay (IGRA) tests (case report and mini-review on ocular non-tuberculous mycobacteria and IGRA cross-reactivity). *Int Ophthalmol*. 2012;32:499–506. doi:10.1007/s10792-012-9588-3
15. Wang MS, Wang JL, Wang XF. The performance of interferon-gamma release assay in nontuberculous mycobacterial diseases: a retrospective study in China. *BMC Pulm Med*. 2016;16:163. doi:10.1186/s12890-016-0320-3
16. Daley CL, Iaccarino JM, Lange C, et al. Treatment of nontuberculous mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Eur Respir J*. 2020;56:905–913. doi:10.1183/13993003.00535-2020
17. He Y, Zhang YA, Wang MS. Stool culture for diagnosis of nontuberculous mycobacteria pulmonary disease: an indirect evidence. *J Infect*. 2021;83(5):607–635. doi:10.1016/j.jinf.2021.08.002
18. Lin MY, Reddy TB, Arend SM, et al. Cross-reactive immunity to *Mycobacterium tuberculosis* DosR regulon-encoded antigens in individuals infected with environmental, nontuberculous mycobacteria. *Infect Immun*. 2009;77:5071–5079. doi:10.1128/IAI.00457-09
19. Arend SM, Andersen P, van Meijgaarden KE, et al. Detection of active tuberculosis infection by T cell responses to early-secreted antigenic target 6-kDa protein and culture filtrate protein 10. *J Infect Dis*. 2000;181:1850–1854. doi:10.1086/315448
20. Scherrer S, Landolt P, Friedel U, Stephan R. Distribution and expression of *esat-6* and *cfp-10* in non-tuberculous mycobacteria isolated from lymph nodes of slaughtered cattle in Switzerland. *J Vet Diagn Invest*. 2019;31:217–221. doi:10.1177/1040638718824074
21. van Ingen J, Bendien SA, de Lange WC, et al. Clinical relevance of non-tuberculous mycobacteria isolated in the Nijmegen-Arnhem region, The Netherlands. *Thorax*. 2009;64:502–506. doi:10.1136/thx.2008.110957
22. Jankovic M, Sabol I, Zmak L, et al. Microbiological criteria in non-tuberculous mycobacteria pulmonary disease: a tool for diagnosis and epidemiology. *Int J Tuberc Lung Dis*. 2016;20:934–940. doi:10.5588/ijtld.15.0633
23. Kobashi Y, Mouri K, Yagi S, et al. Clinical evaluation of the QuantiFERON-TB Gold test in patients with non-tuberculous mycobacterial disease. *Int J Tuberc Lung Dis*. 2009;13:1422–1426.
24. Augustynowicz-Kopeck E, Siemion-Szczepniak I, Zabost A, et al. Interferon Gamma Release Assays in Patients with Respiratory Isolates of Non-Tuberculous Mycobacteria - a Preliminary Study. *Pol J Microbiol*. 2019;68:15–19. doi:10.21307/pjm-2019-002
25. Hermansen TS, Thomsen VO, Lillebaek T, Ravn P. Non-tuberculous mycobacteria and the performance of interferon gamma release assays in Denmark. *PLoS One*. 2014;9:e93986. doi:10.1371/journal.pone.0093986
26. Sato R, Nagai H, Matsui H, et al. Interferon-gamma release assays in patients with *Mycobacterium kansasii* pulmonary infection: a retrospective survey. *J Infect*. 2016;72:706–712. doi:10.1016/j.jinf.2016.03.011
27. Siegel SAR, Cavanaugh M, Ku JH, Kawamura LM, Winthrop KL. Specificity of QuantiFERON-TB Plus, a New-Generation Interferon Gamma Release Assay. *J Clin Microbiol*. 2018;1:56. doi:10.1128/JCM.00629-18
28. Kobashi Y, Obase Y, Fukuda M, Yoshida K, Miyashita N, Oka M. Clinical reevaluation of the QuantiFERON TB-2G test as a diagnostic method for differentiating active tuberculosis from nontuberculous mycobacteriosis. *Clin Infect Dis*. 2006;43:1540–1546. doi:10.1086/509327
29. Hur YG, Crampin AC, Chisambo C, et al. Identification of immunological biomarkers which may differentiate latent tuberculosis from exposure to environmental nontuberculous mycobacteria in children. *Clin Vaccine Immunol*. 2014;21:133–142. doi:10.1128/CCI.00620-13
30. Greinert U, Schlaak M, Rusch-Gerdes S, Flad HD, Ernst M. Low in vitro production of interferon-gamma and tumor necrosis factor-alpha in HIV-seronegative patients with pulmonary disease caused by nontuberculous mycobacteria. *J Clin Immunol*. 2000;20:445–452. doi:10.1023/a:101038700000000000
31. Rae W, Gao Y, Eren E, Doffinger R, Marshall B, Williams AP. Differential Impairment of Interferon-gamma Responses in Two Cases of Pulmonary Nontuberculous Mycobacterial Disease. *Case Reports Immunol*. 2016;2016:9165641. doi:10.1155/2016/9165641
32. Elnaggar MM, El-Naggar MM, Abdellazeq GS, et al. Development of an improved ESAT-6 and CFP-10 peptide-based cytokine flow cytometric assay for bovine tuberculosis. *Comp Immunol Microbiol Infect Dis*. 2015;42:1–7. doi:10.1016/j.cimid.2015.07.005
33. Yang C, Luo X, Fan L, Sha W, Xiao H, Cui H. Performance of Interferon-Gamma Release Assays in the Diagnosis of Nontuberculous Mycobacterial Diseases-A Retrospective Survey From 2011 to 2019. *Front Cell Infect Microbiol*. 2020;10:571230. doi:10.3389/fcimb.2020.571230
34. Steindor M, Stehling F, Olivier M, et al. Species-Specific Interferon-Gamma Release Assay for the Diagnosis of *Mycobacterium abscessus* Complex Infection. *Front Microbiol*. 2021;12:692395. doi:10.3389/fmicb.2021.692395
35. van Zyl-Smit RN, Lehloenyia RJ, Meldau R, Dheda K. Impact of correcting the lymphocyte count to improve the sensitivity of TB antigen-specific peripheral blood-based quantitative T cell assays (T-SPOT.®) TB and QFT-GIT. *J Thorac Dis*. 2016;8:482–489. doi:10.21037/jtd.2016.02.65
36. Du F, Xie L, Zhang Y, et al. Prospective Comparison of QFT-GIT and T-SPOT.TB Assays for Diagnosis of Active Tuberculosis. *Sci Rep*. 2018;8:5882. doi:10.1038/s41598-018-24285-3
37. Chee CB, Gan SH, Khinmar KW, et al. Comparison of sensitivities of two commercial gamma interferon release assays for pulmonary tuberculosis. *J Clin Microbiol*. 2008;46:1935–1940. doi:10.1128/JCM.02403-07
38. Chee CB, KhinMar KW, Gan SH, et al. Tuberculosis treatment effect on T-cell interferon-gamma responses to *Mycobacterium tuberculosis*-specific antigens. *Eur Respir J*. 2010;36:355–361. doi:10.1183/09031936.00151309



**Infection and Drug Resistance****Dovepress****Publish your work in this journal**

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>