

# Efficacy and Safety of Secukinumab in Elderly Patients with Moderate to Severe Plaque-Type Psoriasis: Post-Hoc Analysis of the SUPREME Study

Marina Talamonti<sup>1</sup>, Filomena Russo<sup>2</sup>, Giovanna Malara<sup>3,4</sup>, Katharina Hansel<sup>5</sup>, Manuela Papini<sup>6</sup>, Angelo Cattaneo<sup>7</sup>, Aurora Parodi<sup>8</sup>, Andrea Chiricozzi<sup>9,10</sup>, Piergiorgio Malagoli<sup>11</sup>, Federico Bardazzi<sup>12</sup>, Valeria Brazzelli<sup>13</sup>, Paolo Dapavo<sup>14</sup>, Paolo Gisondi<sup>15</sup>, Cristina Zane<sup>16</sup>, Concetta Potenza<sup>17</sup>, Franca Cantoresi<sup>18</sup>, Maria Concetta Fagnoli<sup>19</sup>, Sara Trevisini<sup>20</sup>, Pina Brianti<sup>21</sup>, Leonardo Pescitelli<sup>22</sup>, Giovanni Gigante<sup>23</sup>, Marta Bartezaghi<sup>23</sup>, Luisa Caputo<sup>23</sup>, Elisabetta Aloisi<sup>23</sup>, Antonio Costanzo<sup>24,25</sup>

On behalf of the SUPREME Study Group

<sup>1</sup>Dermatology, University of Rome Tor Vergata, Rome, Italy; <sup>2</sup>Department of Medical, Surgical and Neurological Science, Dermatology Section, University of Siena, S. Maria alle Scotte Hospital, Siena, Italy; <sup>3</sup>Dermatology Unit, Hospital "Bianchi Melacrino Morelli", Reggio, Calabria, Italy; <sup>4</sup>Department of Dermatology, Papardo Hospital, Messina, Italy; <sup>5</sup>Section of Dermatology, Department of Medicine and Surgery, University of Perugia, Perugia, Italy; <sup>6</sup>Dermatologic Clinic of Terni, Department of Medicine and Surgery, University of Perugia, Perugia, Italy; <sup>7</sup>U.O. Dermatologia, Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Milano, Italy; <sup>8</sup>Di.Sal. Section of Dermatology, Ospedale Policlinico San Martino, University of Genova, Genova GE, 16132, Italy; <sup>9</sup>UOC di Dermatologia, Dipartimento di Scienze Mediche e Chirurgiche, Fondazione Policlinico Universitario A. Gemelli - IRCCS, Rome, Italy; <sup>10</sup>Dermatologia, Dipartimento di Medicina e Chirurgia Traslazionale, Università Cattolica del Sacro Cuore, Rome, Italy; <sup>11</sup>Psocare Unit, I.R.C.C.S. Policlinico San Donato, Milano, Italy; <sup>12</sup>Dermatology Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy; <sup>13</sup>Dermatology, Fondazione IRCCS Policlinico San Matteo and University of Pavia, Pavia, Italy; <sup>14</sup>Department of Biomedical Science and Human Oncology, Second Dermatologic Clinic, University of Torino, Torino, Italy; <sup>15</sup>Section of Dermatology and Venereology, Department of Medicine, University of Verona, Verona, Italy; <sup>16</sup>Dermatology Department, ASST Spedali Civili di Brescia, University of Brescia, Brescia, Italy; <sup>17</sup>Dermatology Unit "Daniele Innocenzi", Department of Medico-Surgical Sciences and Biotechnologies, Faculty of Pharmacy and Medicine, Sapienza University of Rome – Polo Pontino, Latina, Italy; <sup>18</sup>Dermatology Unit, Department of Medicine, University of Roma, Roma, Italy; <sup>19</sup>Dermatology, Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, L'Aquila, Italy; <sup>20</sup>Department of Dermatology, University of Trieste, Trieste, Italy; <sup>21</sup>Dermatology and Cosmetology Unit - San Raffaele Hospital, Milan, Italy; <sup>22</sup>Department of Health Sciences, Dermatology Clinic, University of Firenze, Firenze, Italy; <sup>23</sup>Novartis Farma SpA, Origgio, Italy; <sup>24</sup>Unit of Dermatology, IRCCS Humanitas Research Hospital, Milano, Italy; <sup>25</sup>Department of Biomedical Sciences, Humanitas University, Milano, Italy

Correspondence: Marina Talamonti, Dermatology-Department of Systems Medicine, University of Rome Tor Vergata, PTV – Policlinico Tor Vergata, V. le Oxford 81, Rome, 00133, Italy, Tel +39 0620902743, Fax +39 0620902742, Email [talamonti.marina@gmail.com](mailto:talamonti.marina@gmail.com)

**Purpose:** Secukinumab is a fully human monoclonal antibody that inhibits interleukin (IL)-17A approved for the treatment of moderate to severe plaque psoriasis in adults and children. We compared the efficacy and safety of secukinumab in patients aged < 65 years (adult patients) versus patients aged ≥ 65 years (elderly patients) in a post-hoc analysis of the SUPREME study.

**Patients and Methods:** Patients with moderate to severe plaque psoriasis received subcutaneous secukinumab 300 mg per week for the first 5 weeks, then 300 mg per month. We compared the following outcomes in patients aged ≥ 65 years vs < 65 years: baseline characteristics; PASI50/75/90/100 response rates (improvements ≥ 50%/75%/90%/100% in Psoriasis Area and Severity Index (PASI) from baseline); changes in Dermatology Life Quality Index (DLQI); Hospital Anxiety and Depression Scale (HAD-A, HAD-D) score changes; treatment-emergent adverse events (TEAEs).

**Results:** Secukinumab was slightly less effective in elderly patients than in adult patients (response rates at week 16: PASI90, 69.4% vs 80.9%,  $p = 0.4528$ ; PASI100, 44.4% vs 56.7%,  $p = 0.8973$ ). Elderly and adult patients showed a similar time course of changes in absolute PASI scores. Patients aged ≥ 65 years had a statistically significantly lower improvement in quality of life (mean DLQI reduction) than patients aged < 65 years at week 16 [ $-5.4 (\pm 4.3)$  vs  $-8.8 (\pm 6.9)$ ,  $p = 0.0065$ ] and at week 24 [ $-5.3 (\pm 4.4)$  vs  $-9.2 (\pm 7.1)$ ,  $p = 0.0038$ ]. Secukinumab treatment resulted in comparable mean reductions in anxiety and depression scores in both cohorts at 24 weeks [HAD-A,  $-1.3 (\pm 3.3)$  vs  $-2.1 (\pm 3.8)$ ,  $p = 0.9004$ ; HAD-D,  $-1.0 (\pm 3.3)$  vs  $-1.5 (\pm 3.1)$ ,  $p = 0.4598$ ]. The frequency of TEAEs in the two cohorts was similar (16.7% vs 14.6%,  $p = 0.7391$ ).

**Conclusion:** Secukinumab is a valid option for the management of moderate to severe psoriasis in elderly patients.

**Keywords:** psoriasis, biologics, interleukin-17A, secukinumab, elderly

## Introduction

Psoriasis is a common inflammatory, immune-mediated skin disease; its prevalence tends to increase with age.<sup>1–3</sup> Due to the steadily increasing number of people aged  $\geq 65$  years worldwide, the population of geriatric patients with psoriasis is expanding.<sup>4–6</sup> The management of psoriasis in the elderly can be difficult as the presence of comorbidities and the resulting use of concomitant medications, organ dysfunction, impaired immune system, and altered pharmacokinetics complicate the balancing of benefits and adverse effects of anti-psoriasis treatment.<sup>5,7,8</sup> The use of topical medications is complex in this age group and the compliance to topical therapies is generally low. Also, patients aged  $\geq 65$  years are usually excluded from clinical studies and, therefore, evidence-based guidance for dermatologists is lacking.

The introduction of targeted systemic therapies with biologics has considerably improved the management of moderate to severe psoriasis.<sup>9</sup> Secukinumab, a fully human monoclonal antibody that inhibits interleukin (IL)-17A is approved by the European Medicines Agency (EMA) for the treatment of moderate to severe plaque psoriasis in adults and children, psoriatic arthritis, axial spondyloarthritis (ankylosing spondylitis and non-radiographic axial spondyloarthritis), and juvenile idiopathic arthritis (enthesitis-related arthritis and juvenile psoriatic arthritis). Evidence of its effectiveness in elderly patients with psoriasis is limited.<sup>5</sup>

The recent SUPREME study (NCT02394561) was performed to determine whether the presence of the allele HLA-Cw6 within the chromosomal psoriasis susceptibility gene 1 (PSORS1) influences the response to treatment with secukinumab.<sup>10,11</sup> The study showed that secukinumab ensures high rates of skin clearance regardless of HLA-Cw6 status.<sup>10,11</sup> Here we present the results of a post-hoc analysis of the SUPREME study comparing the efficacy and safety of secukinumab in patients aged  $< 65$  years (adult patients) versus patients aged  $\geq 65$  years (elderly patients).

## Materials and Methods


The design of the SUPREME study has been described in detail in a previous report.<sup>10</sup> Briefly, SUPREME was a 24-week, phase IIb, multicentre, prospective study conducted across 50 centers in Italy, with an extension period of up to 72 weeks, and involving patients aged  $\geq 18$  years with moderate to severe plaque psoriasis. Eligible patients were treated with subcutaneous secukinumab 300 mg per week for the first 5 weeks, followed by a maintenance period of 300 mg per month.

The post-hoc analysis of SUPREME presented here compared the following outcomes in patients aged  $\geq 65$  years vs patients aged  $< 65$  years: baseline characteristics; PASI50/75/90/100 response rates (improvements  $\geq 50\%/75\%/90\%/100\%$  in Psoriasis Area and Severity Index (PASI) from baseline); changes in Dermatology Life Quality Index (DLQI); Hospital Anxiety and Depression Scale (HAD-A, HAD-D) score changes; treatment-emergent adverse events (TEAEs). Results are reported as absolute numbers and percentages or mean  $\pm$  standard deviation (SD) as appropriate. Significance tests for differences between patient groups included *t*-test, Wilcoxon test, chi-square test, or Fisher exact test as appropriate. A nominal *p*-value of  $< 0.05$  was considered statistically significant. A logistic regression model was used to evaluate difference in PASI response (taking into account the following covariates: height, weight, waist, age at psoriasis diagnosis and presence of metabolic syndrome), while Kaplan-Meier survival curve and Log rank test was used for time to reach PASI responses. All statistical analyses were performed using SAS software 9.2 (SAS Institute, Cary, NC, US).


## Results

Of the 433 patients treated in the SUPREME study, 36 (8.3%) were aged  $\geq 65$  years. Mean age was 69.2 ( $\pm 4.2$ ) years and 43.1 ( $\pm 11.5$ ) years in the elderly and adult cohorts, respectively. In both cohorts, males were predominant [26/36 (72.2%) and 284/397 (71.5%), respectively]. Elderly patients had a higher mean waist circumference [101.9 ( $\pm 14.1$ ) vs 96.1 ( $\pm 15.1$ ) cm; *p* = 0.0336], were older at diagnosis [45.5 ( $\pm 14.6$ ) vs 25.8 ( $\pm 12.8$ ) years old; *p* < 0.0001] and suffered more frequently of comorbidities (83.3% vs 60.2%; *p* = 0.0061), in particular metabolic syndrome (41.7% vs 14.4%; *p* < 0.0001).

(A)



(B)



**Figure 1** Efficacy of treatment with secukinumab in patients with moderate to severe psoriasis stratified by age < 65 years and ≥ 65 years. **(A)** PASI90 response rates during treatment. **(B)** Absolute PASI scores during treatment.

**Table 1** Summary of TEAEs

Overall Analysis	Age < 65 Years		Age ≥ 65 Years		Total	
	(N=397)		(N=36)		(N=433)	
	n	%	n	%	n	%
Number of TEAEs	579	–	83	–	662	–
Number of patients with TEAEs	215	54.6	17	47.22	232	53.58
Number of patients with serious TEAEs	28	7.05	3	8.33	31	7.16
Number of patients with TEAEs related to study drug	69	17.38	7	19.44	76	17.55
Number of patients with TEAEs leading to discontinuation of study drug	27	6.80	3	8.33	30	6.93
Deaths	1	0.25	1	2.78	2	0.46

Abbreviation: TEAEs, treatment-emergent adverse events.

Significantly fewer older patients were current smokers (27.8% vs 48.6%,  $p = 0.0012$ ). Mean PASI score at baseline was numerically lower for elderly patients [ $19.5 (\pm 7.7)$  vs  $21.3 (\pm 9.7)$ ;  $p = 0.2587$ ] who showed, however, a significantly greater rate of prior therapies with anti-TNF $\alpha$  biologics (44.4% vs 28.2%;  $p = 0.0410$ ). In both cohorts, biologics were discontinued mostly due to lack or loss of efficacy. Of note, most elderly patients (75%) interrupted biologic therapy for loss of efficacy.

Especially in terms of PASI90 and PASI100 responses, secukinumab was slightly less efficacious in elderly patients than in adult patients (response rates at week 16: PASI90, 69.4% vs 80.9%,  $p = 0.4528$  (Figure 1A); PASI100, 44.4% vs 56.7%,  $p = 0.8973$ ). The time to first PASI90 response was similarly short in both cohorts (median time to PASI90 response, 61 vs 58 days, respectively; log-rank, 0.1761). Elderly and adult patients showed a similar time course of changes in absolute PASI scores (Figure 1B).

Elderly patients had a statistically significantly lower improvement in quality of life (mean DLQI reduction) than adult patients at week 16 [ $-5.4 (\pm 4.3)$  vs  $-8.8 (\pm 6.9)$ ,  $p = 0.0065$ ] and at week 24 [ $-5.3 (\pm 4.4)$  vs  $-9.2 (\pm 7.1)$ ,  $p = 0.0038$ ]. At week 16, significantly fewer older patients achieved DLQI 0/1 than adult patients (51.5% vs 71.7%,  $p = 0.0157$ ). Secukinumab treatment resulted in comparable mean reductions in anxiety and depression scores in both cohorts at 24 weeks [HAD-A,  $-1.3 (\pm 3.3)$  vs  $-2.1 (\pm 3.8)$ ,  $p = 0.9004$ ; HAD-D,  $-1.0 (\pm 3.3)$  vs  $-1.5 (\pm 3.1)$ ,  $p = 0.4598$ ].

The frequency of TEAEs in the two cohorts was similar (respectively, 16.7% vs 14.6%,  $p = 0.7391$ ) (Table 1).

## Discussion

This post-hoc analysis of the SUPREME study provides a glimpse of the characteristics of elderly patients with moderate to severe plaque psoriasis and the outcomes of systemic treatment with secukinumab. Elderly patients accounted for less than 10% of the SUPREME population confirming the limited presence of patients aged  $\geq 65$  years in clinical studies. The baseline characteristics of this subgroup were in line with those reported in other studies evaluating older patients, including the higher frequency of comorbidities, lower frequency of current smoking, and later disease onset.<sup>4,8,12,13</sup> Secukinumab was efficacious and well tolerated through week 24 regardless of age. PASI90 and PASI100 response rates tended to be lower in patients aged  $\geq 65$  years than in younger patients, but the differences between the two cohorts were not statistically significant. Improvements in quality of life during treatment were also less pronounced in elderly patients.

Few studies have addressed the efficacy and safety of secukinumab in psoriasis patients aged  $\geq 65$  years. A pooled analysis of three Phase III trials evaluated secukinumab after patient stratification by age and found equivalent efficacy of this anti-IL17 in patients aged  $\geq 65$  years and in younger patients.<sup>12</sup> The rapidity and duration of response were also similar between older and younger patients, in agreement with our observations.<sup>12</sup> A retrospective analysis investigating

the effectiveness of anti-IL17 therapies for psoriasis in elderly patients confirmed the efficacy results reported in clinical trials.<sup>13</sup> Notably, elderly and younger patients had a similar profile of adverse events, despite the higher frequency of comorbidities of the elderly subgroup.<sup>13</sup> The safety findings of our analysis confirm this observation.

The fact that the improvement of the quality of life of the elderly patients treated with secukinumab in the SUPREME study was significantly inferior to that experienced by patients aged < 65 years may reflect the greater disease and treatment burden on elderly patients. At the same time, it highlights the need for tools able to describe more adequately outcomes relevant to the older population. As pointed out by a recent survey among psoriasis patients from different age groups, greater efforts should be devoted to the individualized management of elderly with psoriasis, because their needs differ substantially from those of younger patients with psoriasis.<sup>4,14</sup> A personalized approach to elderly patients with psoriasis may reduce the disease burden and improve treatment outcomes.

Our analysis is limited by its post-hoc nature and the small number of elderly patients in the SUPREME study.

## Conclusion

We believe that this analysis provides useful information on the use of secukinumab in patients aged  $\geq 65$  years and confirms that this anti-IL17 agent is a valid option for the management of moderate to severe psoriasis in elderly patients.

## Data Sharing Statement

The statistically analyzed data is available in the text, and the detailed data will be available on request from the corresponding author.

## Ethics Statement

All patients provided written informed consent before enrolment into the study. The study was approved by all competent Ethics Committees and regulatory authorities ([Supplementary Table 1](#)). The trial was conducted in accordance with the Declaration of Helsinki and Good Clinical Practice, and in compliance with all federal, local and regional requirements.

## Acknowledgments

Study group: Paolo Amerio; Anna Balato; Luca Bianchi; Claudio Bonifati; Cinzia Buligan; Martina Burlando; Giacomo Caldarola; Piergiacomo Calzavara Pinton; Anna Campanati; Alessandra Capo; Maurizio Congedo; Andrea Conti; Francesco Cusano; Stefano Dastoli; Clara De Simone; Micol Del Giglio; Luisa Di Costanzo; Vito Di Lernia; Sergio Di Nuzzo; Daniele Dusi; Gabriella Fabbrocini; Rosaria Fidanza; Chiara Franchi; Francesca Gaiani; Marco Galluzzo; Alberto Ghilardi; Giampiero Girolomoni; Federica Giuliani; Dario Graceffa; Francesco Loconsole; Valentina Mastrandrea; Matteo Megna; Santo Raffaele Mercuri; Giuseppe Micali; Maria Letizia Musumeci; Luigi Naldi; Alessandra Narcisi; Ylenia Natalini; Annamaria Offidani; Diego Orsini; Gianluca Pagnanelli; Annalisa Patrizi; Monica Pau; Giovanni Pellacani; Ketty Peris; Severino Persechino; Stefano Piaserico; Lucia Pietroleonardo; Francesca Prignano; Marco Romanelli; Franco Rongioletti; Roberta Scuderi; Riccardo Sirna; Nevena Skroza; Giuseppe Stinco; Luca Stingeni; Marina Venturini; Leonardo Zichichi; Antonio Zini.

## Funding

The SUPREME study was supported by Novartis Pharma AG.

## Disclosure

M.T. has nothing to disclose; F.R. acted as speaker and consultant for Sanofi, AbbVie and Novartis, outside the submitted work; G.M. has nothing to disclose; K.H. reports personal fees from AbbVie, Almirall, Ganassini, LEO Pharma, Novartis, and Sanofi, outside the submitted work; M.P. has nothing to disclose; A.Ca. has nothing to disclose; A. P. has nothing to disclose; A.Ch. served as advisory board member and consultant and has received fees and speaker's honoraria or has participated in clinical trials for AbbVie, Almirall, Incyte, LEO Pharma, Lilly, Janssen, Novartis, Pfizer, and Sanofi Genzyme; P.M. has nothing to disclose; F.B. has nothing to disclose; V.B. has nothing to disclose. P.D. has nothing to disclose; P.G. reports payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or

educational events for AbbVie, Ammirall, Amgen, Janssen, LEO Pharma, Eli Lilly, Novartis, Pierre Fabre, Pfizer, Sandoz, Sanofi, UCB. C.Z. has nothing to disclose; C.P. has nothing to disclose; F.C. has nothing to disclose; MC.F. has served on advisory boards, received honoraria for lectures and research grants from Amgen, Ammirall, AbbVie, BMS, Galderma, Kyowa Kirin, Leo Pharma, Pierre Fabre, UCB, Lilly, Pfizer, Janssen, MSD, Novartis, Sanofi-Regeneron, Sun Pharma.; S. T. acted as a speaker for AbbVie, Novartis and Janssen; P.B. has nothing to disclose; L.P. has nothing to disclose; G.G., M.B., L.C., and E.A. are employees of Novartis. AC received speaker's/consultancy honoraria or grants for research from AbbVie, Ammirall, Amgen, LEO Pharma, Novartis, Janssen, UCB, and Lilly. The authors report no other conflicts of interest in this work.

## References

1. Griffiths CEM, Armstrong AW, Gudjonsson JE, et al. Psoriasis. *Lancet*. 2021;397(10281):1301–1315. doi:10.1016/S0140-6736(20)32549-6
2. Michalek IM, Loring B, John SM. A systematic review of worldwide epidemiology of psoriasis. *J Eur Acad Dermatol Venereol*. 2017;31:205–212. doi:10.1111/jdv.13854
3. Iskandar IYK, Parisi R, Griffiths CEM, et al. Systematic review examining changes over time and variation in the incidence and prevalence of psoriasis by age and gender. *Br J Dermatol*. 2021;184:243–258. doi:10.1111/bjd.19169
4. van Winden MEC, Haar ELM, Groenewoud HMM, et al. Disease and treatment characteristics in geriatric psoriasis: a patient survey comparing age groups. *Acta Derm Venereol*. 2020;100(14):adv00215. doi:10.2340/00015555-3569
5. Megna M, Camela E, Cinelli E, Fabbrocini G. Real-life efficacy and safety of secukinumab in elderly patients with psoriasis over a 2-year period. *Clin Exp Dermatol*. 2020;45:848–852. doi:10.1111/ced.14258
6. Bakirtzi K, Sotiriou E, Papadimitriou I, et al. Elderly patients with psoriasis: long-term efficacy and safety of modern treatments. *J Dermatol Treat*. 2022;33:1339–1342. doi:10.1080/09546634.2020.1809623
7. Balato N, Patruno C, Napolitano M, et al. Managing moderate-to-severe psoriasis in the elderly. *Drugs Aging*. 2014;31(4):233–238. doi:10.1007/s40266-014-0156-6
8. Phan C, Sigal M-L, Estève E, et al. Psoriasis in the elderly: epidemiological and clinical aspects, and evaluation of patients with very late onset psoriasis. *J Eur Acad Dermatol Venereol*. 2016;30:78–82. doi:10.1111/jdv.12850
9. Sbidian E, Chaimani A, Afach S, et al. Systematic pharmacological treatments for chronic plaque psoriasis: a network meta-analysis. *Cochrane Database Syst Rev*. 2020. doi:10.1002/14651858.CD011535.pub3
10. Costanzo A, Bianchi L, Flori ML, et al. Secukinumab shows high efficacy irrespective of HLA-Cw6 status in patients with moderate-to-severe plaque-type psoriasis: SUPREME study. *Br J Dermatol*. 2018;179:1072–1080. doi:10.1111/bjd.16705
11. Costanzo A, Russo F, Galluzzo M, et al. Secukinumab exhibits sustained and stable response in patients with moderate-to-severe psoriasis: results from the SUPREME study. *Acta Derm Venereol*. 2021;101:adv00576. doi:10.2340/00015555-3816
12. Körber A, Papavassilis C, Bhosekar V, Reinhardt M. Efficacy and safety of secukinumab in elderly subjects with moderate to severe plaque psoriasis: a pooled analysis of phase III studies. *Drugs Aging*. 2018;35(2):135–144. doi:10.1007/s40266-018-0520-z
13. Phan C, Beneton N, Dalaunay J, et al. Effectiveness and safety of anti-interleukin-17 therapies in elderly patients with psoriasis. *Acta Derm Venereol*. 2020;100:adv00316. doi:10.2340/00015555-3678
14. van Winden MEC, Haar ELM, Groenewoud JMM, Kerkhof PCM, Jong EMGJ, Lubeek SFK. Quality of life, treatment goals, preferences and satisfaction in older patients with psoriasis: a patient survey comparing age groups. *Br J Dermatol*. 2021;184(4):759–762. doi:10.1111/bjd.19665

Clinical, Cosmetic and Investigational Dermatology

Dovepress

**Publish your work in this journal**

Clinical, Cosmetic and Investigational Dermatology is an international, peer-reviewed, open access, online journal that focuses on the latest clinical and experimental research in all aspects of skin disease and cosmetic interventions. This journal is indexed on CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dermatology-journal>