

Response to Article “Antipyretic Potential of 80% Methanol Extract and Solvent Fractions of *Bersama Abyssinica* Fresen. (Melianthaceae) Leaves Against Yeast-Induced Pyrexia in Mice” [Letter]

Putri Reno Intan^{1,*}, Ariyani Noviantari^{1,*}, Sukmayati Alegantina^{2,*}

¹Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency (BRIN), Cibinong Science Center, Cibinong - Bogor, West Java, Indonesia; ²Research Center for Pharmaceutical Ingredients and Traditional Medicine, Research Organization for Health, National Research and Innovation Agency (BRIN), Cibinong Science Center, Cibinong - Bogor, West Java, Indonesia

*These authors contributed equally to this work

Correspondence: Putri Reno Intan, Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency (BRIN), Genomic Building, Cibinong Science Center, Jalan Raya Bogor Km. 46, Cibinong – Bogor, West Java, 16911, Indonesia, Email putri.reno.intan@brin.go.id

Dear editor

We are grateful to the authors for publishing their research “Antipyretic Potential of 80% Methanol Extract and Solvent Fractions of *Bersama abyssinica* Fresen. (Melianthaceae) Leaves Against Yeast-Induced Pyrexia in Mice” in the Journal of Experimental Pharmacology 2023:15 81–91.¹ Understanding the efficacy of *B. abyssinica*'s leaf extract and solvent fractions in vivo requires an understanding of this information. We would to communicate our thoughts on the test of preparation and extraction of plant material method used against yeast-induced pyrexia in mice in this study.

In this study, pyrexia or fever was induced by injecting a suspension of 30% w/v yeast extract powder, at a dose of 3 g/kg or 10 mL/kg, in 0.9% normal saline beneath the nape of the neck. After 18 hours, the rectal temperature was recorded. The temperature was recorded at 0.5, 1, 1.5, 2, 2.5, and 3 hours after dosing with crude extract and solvent fractions. The previous experiment had similar methods.² In another experiment, the temperature was recorded at 30 minutes, 1 hour, 2 hours, 3 hours, and 24 hours after treatment.³

Preparing medicinal plants for experimentation is the first and most important stage in producing high-quality research results. Before beginning the process, it involves extracting and determining the quantity and quality of bioactive components.⁴ The next thing that needs to be done according to WHO standards for herbal medicines is potentially hazardous contaminants and residues. There are potential sources for pollutants and residues, as well as the stages of production where they might be visible. Some of them are regarded as inevitable contaminants or herbal medicinal residues. Physical, chemical, and biological pollutants are the two categories of contaminants found in herbal medicines. There could be significant residues of several agrochemicals and organic solvents in herbal medications.⁵

Because of this, the test is potentially hazardous contaminants and residues on herbal medicine may also be evaluated to be used in the near future to complete methods in preparation of medicinal plants. In general, this study still bears faithfully to the initial principle of “Extraction of Plant Material” that will be used as an antipyretic in mice in vivo and may be further explored and evaluated to address the suggestions stated above.

Acknowledgments

It is appropriate to express appreciation to Bantayehu Addis Tegegne and Agumas Alemu Alehegn for their good teamwork in this study. All authors would also like to thank Dr. Sunarno and all researchers in the Center for Biomedical Research, National Research and Innovation Agency (BRIN) for their ongoing assistance.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Tegegne BA, Alehegn AA. Antipyretic potential of 80% methanol extract and solvent fractions of *Bersama abyssinica* Fresen. (Melianthaceae) leaves against yeast-induced pyrexia in mice. *J Exp Pharmacol*. 2023;15:81–91. doi:10.2147/JEP.S390825
2. Yimer T, Emiru YK, Kifle ZD, Ewunetei A, Adugna M, Birru EM. Pharmacological evaluation of antipyretic and antioxidant activities of 80% methanol root extract and derived solvent fraction of *Echinops kebericho* M. (Asteraceae) in mice model. *Biomed Res Int*. 2021;2021:1–8. doi:10.1155/2021/6670984
3. Reza MS, Jashimuddin M, Ahmed J, et al. Pharmacological investigation of analgesic and antipyretic activities of methanol extract of the whole part of *Aeginetia indica*. *J Ethnopharmacol*. 2021;271:113915. doi:10.1016/j.jep.2021.113915
4. Abubakar A, Haque M. Preparation of medicinal plants: basic extraction and fractionation procedures for experimental purposes. *J Pharm Bioallied Sci*. 2020;12:1–10. doi:10.4103/jpbs.JPBS_175_19
5. World Health Organization. *Guidelines for Assessing Quality of Herbal Medicines with Reference to Contaminants and Residues*. World Health Organization; 2007.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Experimental Pharmacology 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Experimental Pharmacology editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Experimental Pharmacology

Dovepress

Publish your work in this journal

The Journal of Experimental Pharmacology is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of laboratory and experimental pharmacology. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-experimental-pharmacology-journal>

<https://doi.org/10.2147/JEP.S413591>