Open Access Full Text Article

Pomegranate Juice Diminishes the Mitochondria-Dependent Cell Death and NF-kB Signaling Pathway Induced by Copper Oxide Nanoparticles on Liver and Kidneys of Rats [Erratum]

Hassanen EI, Tohamy AF, Issa MY, Ibrahim MA, Farroh KY, Hassan AM. Int J Nanomedicine. 2019;14:8905-8922.

An error during the preparation of Figures 6 and 7 led to the inadvertent creation of duplicate regions within multiple images from these figures on pages 8916 and 8917, respectively. The journal wishes to apologise for this error. The correct versions of Figures 6 and 7 are as follows.

© 2023 Hassanen et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms. work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/twrms.php).



Figure 6 Immunohistochemical expression of caspase-3 and NF-κB protein in the liver sections in different groups showing (**A** and **B**) mild to negative Caspase-3 and NF-κB protein expression in control negative group. (**C** and **D**) Strong positive caspase-3 and extensive nuclear translocation of NF-κB protein within hepatocytes in the group intoxicated with CuO-NPs. (**E** and **F**) Moderate positive caspase-3 and NF-κB protein expression in the group pretreated with 1 mL/kg bwt PJ. (**G** and **H**) Mild to negative caspase-3 and NF-κB protein expression in the group pretreated with 1 mL/kg bwt PJ. (**G** and **H**) Mild to negative caspase-3 and NF-κB protein expression in the group pretreated with 1 mL/kg bwt PJ.



Figure 7 Immunohistochemical expression of caspase-3 and NF-κB protein in the kidney tissue sections in different groups showing (**A** and **B**) mild to negative caspase-3 and NF-κB protein expression in control negative group. (**C** and **D**) Moderate positive caspase-3 and extensive nuclear translocation of NF-κB protein within the renal tubular epithelial cells in the group intoxicated with CuO-NPs. (**E** and **F**) Mild to moderate positive caspase-3 and NF-κB protein expression in group pretreated with 1 mL/ kg bwt PJ. (**G** and **H**) Mild to negative caspase-3 and NF-κB protein expression in group pretreated with 1 mL/ kg bwt PJ. (**G** and **H**) Mild to negative caspase-3 and NF-κB protein expression in group pretreated with 1 mL/ kg bwt PJ.

International Journal of Nanomedicine

Dovepress

1851

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch[®], Current Contents[®]/Clinical Medicine, Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http:// www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-nanomedicine-journal

f 🄰 in 🕨 DovePress