

# The PCQ-Infertility Revised: A New Digital Instrument to Measure Treatment Satisfaction of Fertility Patients

Lotte van der Kolk<sup>1</sup>, Ellen Smit<sup>2</sup>, José Bloemer<sup>1</sup>, Lise Marit van Wijk<sup>2</sup>

<sup>1</sup>Institute for Management Research, Radboud University, Nijmegen, the Netherlands; <sup>2</sup>Ferring B.V., Hoofddorp, the Netherlands

Correspondence: Ellen Smit, Polarisavenue 130, Hoofddorp, 2132 JX, the Netherlands, Tel +31 6 208 72 493, Email ellen.smit@ferring.com

**Background:** One of the key dimensions of healthcare quality is patient-centeredness, which represents how well healthcare is geared towards patients' needs and wishes. Many questionnaires that measure the patient-centeredness are long and complicated, eliciting non-response or careless responding. Moreover, responses to some commonly used questionnaires are difficult to interpret. The Patient-Centeredness Questionnaire-Infertility (PCQ-Infertility) is used to measure the patient's experience of fertility healthcare quality. The aim of this study was to improve the PCQ-Infertility to allow large-scale clinical implementation.

**Methods:** The study was performed in three parts. First, shortcomings of the original PCQ-Infertility were identified by evaluation of expert opinions. Second, the number of items were reduced, and items were rescaled and rephrased. Third, 844 patients filled in the original PCQ-Infertility and 260 patients filled in the revised PCQ-Infertility and reliability analyses were performed. In addition, a confirmatory factor analysis was performed on the revised PCQ-Infertility.

**Results:** The number of items in the revised questionnaire was reduced by 24% (from 51 to 39 items), which increased the internal consistency and reliability. The reliability analyses and confirmatory factor analysis indicated high consistency and convergent validity in all seven dimensions (accessibility, information, communication, patient involvement, respect for patient's values, continuity and transition, and competence) of the revised PCQ-Infertility.

**Conclusion:** The revised PCQ-Infertility is a more valid and reliable instrument than the original PCQ-Infertility, easier to interpret and shorter. Therefore, large-scale clinical implementation and data analysis are now possible, giving the opportunity for fertility care professionals to evaluate and improve their healthcare.

**Keywords:** PCQ-Infertility, PREM, fertility, questionnaire, patient-centeredness, patient satisfaction

## Plain Language Summary

It is of great importance to keep improving the quality of healthcare. One important way to do this, is by understanding the needs and wishes of patients. For patients undergoing fertility treatment, a questionnaire was developed that helps clinics to understand which aspects of the fertility care should be improved. This questionnaire was, however, rather long and the questions were difficult to understand. In this study, we show how the questionnaire was improved by removing 24% of questions and rephrasing difficult to understand questions in a way they could be interpreted more easily. We also show that the new questionnaire can capture the patients' needs and wishes in fertility care more accurately than the original questionnaire. Therefore, this new questionnaire can be used in many fertility clinics and will help improve the quality of fertility care.

## Background

Patient-centeredness, which represents how well healthcare is geared towards patients' needs and wishes, is acknowledged as one of the six key aspects of healthcare.<sup>1,2</sup> Within the fertility care, most healthcare professionals are aware of the meaning and importance of patient-centeredness, but they do not, however, always consistently undertake the necessary steps to improve their patient-centered care.<sup>3,4</sup> Important obstacles are the lack of detailed, concrete feedback from patients to understand their specific needs and the healthcare providers lack of motivation to change practice.<sup>4</sup> As

a result, healthcare professionals may not be aware of which limits their ability to improve the care they provide. From a patient's perspective, not only medical skills are considered important, but also respect, coordination, accessibility, information, comfort, support, partner involvement, having a lead physician, and a good attitude of and a good relationship with fertility clinic staff.<sup>5-7</sup> Several of these aspects can and should be improved.<sup>8</sup> Personalized treatment benefits the patients and reduces the substantial physical and emotional burden of fertility treatment.<sup>9</sup> This may reduce the high dropout rate in fertility care.<sup>8,10</sup>

Monitoring patient-centeredness by measuring patients' experiences via surveys, is a valuable tool to improve fertility care.<sup>5,11</sup> Questionnaires that measure the outcome of the given care (eg the amount of physical pain and quality of life patients experience) are Patient Reported Outcome Measures (PROMs). Patient Reported Experience Measures (PREMs) measure patient satisfaction with a health service and capture overall patient experience including the process of care (eg communication and information provision).<sup>12</sup> In 2010, a PREM was developed based on focus group interviews and a literature review in the Netherlands to measure patient-centeredness within the infertility field, the so called Patient-Centeredness Questionnaire-Infertility (PCQ-Infertility).<sup>13</sup> This original PCQ-Infertility (PCQ-original, consisting of 51 items) measures patient-centered care by assessing seven dimensions: accessibility, information, communication, patient involvement, respect for patient's values, continuity and transition, and competence.<sup>13</sup> The PCQ-original showed a high internal consistency and good construct validity.<sup>13,14</sup> In addition, it has been successfully validated in six countries (the Netherlands, Slovakia, Slovenia, Portugal, Iran, and New-Zealand)<sup>13-19</sup> and it is currently used in clinical practice in countries such as Germany and the Netherlands. However, important limitations of the PCQ-original lies in its format. As a paper-based questionnaire it is less likely to be filled in and data processing is labor-intensive. Furthermore, it's lengthy and probably due to the diversity in answering categories and the presence of dichotomous items there is a relatively low reliability of dimensions such as "continuity and transition".<sup>13,19</sup> The response options in the PCQ-original are diverse and include open-ended, multiple-choice, dichotomous (yes/no), rating scales (0-10), and 3-, 4-, and 5-point-Likert response scales. Data analysis is therefore unnecessary complicated and could be improved by consistently applying the same response scales to acquire more reliable and easier to interpret results.

Despite its limitations, the PCQ-original discriminates well on the quality of patient-centered healthcare between fertility clinics and hospitals, making it suitable for benchmarking.<sup>13</sup> It contains many attributes, such as the "Hospitals Experience Questionnaire (American Hospital CAHPS) and the Hospital Comparison System questionnaires (developed by the Consumers' Association and NIVEL measuring hospital service from a patient's perspective) that are required in Consumer Quality Indexes (CQI), a type of patient satisfaction instrument widely used in the Netherlands.<sup>20,21</sup> However, despite their popularity, CQIs have been criticized for having too many items, being costly to administer, and for the results being difficult to interpret.<sup>22</sup> These problems also apply to the PCQ-original. The complicated and long questionnaires could demotivate the patients resulting in nonresponse or careless responding.<sup>23-26</sup>

We sought to overcome the shortcomings of the PCQ-original regarding length, interpretability and reliability, and describe the development and validation based upon the essential standards for patient-outcome measures of a revised version of the PCQ-Infertility (PCQ-revised),<sup>27</sup> which is now available for large-scale clinical implementation.

## Materials and Methods

### Data Collection

#### Sample Population

Women (aged 18-45, single, or in a same sex or heterosexual partnerships) receiving fertility treatment in the Netherlands were invited to fill in the PCQ-original or the PCQ-revised (together with a partner if applicable) by either invitation cards available at the reception or via digital invitation (eg e-mail) which is sent out by the clinic. It was estimated that the response rate was ~50%, but no exact numbers could be collected due to the nature of the invitation procedure. Patients who completed the PCQ-Infertility (either original or revised) consent that data is anonymously collected and used for scientific research. This study was conducted according to the guidelines laid down in the Declaration of Helsinki.

## Collection of Feedback on the Original PCQ-Infertility by Fertility Experts and Patients

A medical advisor from Ferring B.V. consulted three fertility healthcare experts from Dutch clinics (one fertility doctor and two nurses specialized in fertility), and the Dutch patient association for people with fertility problems, Freya, to get insights into commonly experienced issues with the use of the PCQ-original by performing semi-structured interviews. Conversations were held 1:1 on location, and a second Ferring B.V. representative was involved to note the mentioned issues and shortcomings of the PCQ-original, and the possible solutions on how to improve the PCQ-original. Questions such as “Which feedback do patients give after filling in the PCQ-Infertility?”, “Which problems do you encounter with the PCQ-Infertility?”, and “How do you think the PCQ-Infertility could be improved?” were asked. Based on these conversations, commonly mentioned issues were noted and concrete shortcomings and solutions to improve the PCQ-original were formulated. In addition, all three fertility healthcare experts and a representative from Freya were asked to critically evaluate the PCQ-original. They were specifically asked to pinpoint essential items, and to give suggestions for improvement regarding the phrasing and wording of items and answers. The acquired insights and the written feedback served as a base to reduce and rephrase items in the PCQ-revised. The development and validation of the PCQ-revised was based upon previously published guidelines for patient-outcome measures.<sup>27</sup>

## Collection of Original PCQ-Infertility Output

LiveResearch, a Dutch research institute that specializes in measuring Customer (Patient) experience, and fertility departments of four fertility care clinics in the Netherlands supplied us with an anonymized dataset of questionnaires completed voluntarily by 844 patients receiving fertility treatment (Materials and methods, [Sample Population](#)). Approval by an institutional ethics committee to use this dataset was not required according to the Dutch law, since no patient-sensitive data was collected, data was untraceable, and patients participated voluntarily. Data were collected between 2017 and 2019. The dataset consisted of responses to the questions in the PCQ-original, collected in an excel database that was cleaned from duplicates.<sup>13</sup>

## Collection of Revised PCQ-Infertility Output

The same four hospitals that provided the anonymized dataset of the PCQ-original were contacted to participate in this study, in which the PCQ-revised was distributed among new patients. Two hospitals, Nij Geertgen and Hospital Gelderse Vallei, agreed upon participation. The PCQ-revised was completed by 260 new patients who had not previously completed the PCQ-original, of whom 230 came from Nij Geertgen and 30 came from Hospital Gelderse Vallei. Data collection took place between August 2019 and October 2019 in collaboration with LiveResearch.

## Data Analyses

Data was exported to SPSS for statistical analysis (Statistical Package for Social Sciences: version 27 for Mac, SPSS Inc., Chicago, IL, USA). For all analyses, normality of data and the proper assumptions such as frequency, kurtosis, skewness, and missing data were checked and statistical significance was set at a  $p$ -value of  $< 0.05$ . SPSS Amos was used to perform confirmatory analyses.

## Psychometric Analyses

First, reliability of the PCQ-original was tested using Cronbach's alpha for each dimension. Values  $>0.70$  were considered acceptable. Based on the SPSS output of the analysis, an item was flagged if the Cronbach's alpha increased after deletion of the item. Next, the flagged items for deletion were revised with the requirements of the Patient-Centeredness Quality Index (PCQ index), an index widely used in the Netherlands for measurement of patient satisfaction, and items which were considered essential were not deleted.<sup>20</sup> Flagged items for deletion were also cross-referred with the requirements from Freya and deletion only took place if the question was not considered essential. Freya requested items Q4, Q24 and Q26 to remain in the questionnaire, although deletion would have improved the Cronbach's alpha of the corresponding dimensions. In addition, face validity was applied by fertility healthcare experts to determine which items could be deleted and to decide which items should be reformulated. Reliability analysis was then performed on the PCQ-revised and values  $>0.70$  were considered acceptable.

We performed a confirmatory factor analysis to determine convergent and discriminant validity of the PCQ-revised.<sup>13</sup> We assessed the model fit by the Comparative Fit Index (CFI) and the Root Mean Square Error of Approximation (RMSEA). The CFI needs to be  $> 0.9$  to indicate a good model fit and the RMSEA needs to be  $< 0.08$  for a good model fit. We calculated standardized factor loadings for each item in relation to the corresponding dimension and considered loadings of  $> 0.40$  to be sufficient.<sup>28</sup> Convergent validity was determined by calculation of Average Variance Extracted (AVE) scores, with a score  $> 0.5$  indicating the presence of convergent validity. Discriminant validity was tested with the Fornell and Larcker Criterion, with a criterion of  $> 0.5$ .<sup>29</sup>

## Results

We divided our study in three parts. First, we interviewed fertility health care experts and the Dutch patient association Freya to identify shortcomings of the PCQ-original. Second, based on the received feedback, and reliability and validity analyses of the PCQ-original, the scale was revised to make the PCQ-revised parsimonious. Third, we validated the PCQ-revised with the PCQ-original by performing reliability and confirmatory factor analyses.

### Identifying Shortcomings of the Original PCQ-Infertility by Expert Opinion

The most commonly mentioned shortcomings in the PCQ-original as indicated by the consulted fertility health care experts and the Dutch patient association Freya, are the high number of items, the difficulty in interpreting the patient's responses due to differences in answering scale, and inconsistent wording in the items (Table 1). A summary of the shortcomings mentioned for the PCQ-original and their possible solutions are listed in Table 1.

### Revision of the Original PCQ-Infertility

To improve the response rate, interpretability and the length of the PCQ-original, first the number of items were revised based upon reliability analysis, face validity, and predefined criteria (Materials and Methods, Data Analyses).

The PCQ-original comprises of 51 items in total: four background items (Q1-Q4) in the section "Background Information", 46 patient's experience items (Q5-Q50) divided into seven dimensions and one item in the section "To conclude" (Q51, Table 2).<sup>13</sup> In total, 13 items were removed, as can be seen in Table 2. Most of the items were removed from the section patient's experience (12/46; 26%).

In the last section of the PCQ-original "To conclude", the item "What is the probability of recommending this hospital / clinic?" was added (Table 2, "To conclude"). The final PCQ-revised comprises of 39 items: three items in the "Background information" section, 34 items in the main section, and two items "To conclude". The PCQ-revised is 24% shorter than the PCQ-original (from 51 to 39 items).

After reducing the number of items all items were rephrased. Negatively phrased items were rephrased into positively phrased items, as positively phrased items were expected to be less confusing and more consistent, which would increase the reliability of the results.<sup>30</sup> Items that could be considered as vague or ambiguous were also rephrased. The rephrased items are shown in Table 2.

The biggest change that was implemented in the PCQ-revised was the conversion of the dichotomous yes/no into a 5-point Likert scale. For Q35 the dichotomous (yes/no) response scales remained in place and the response scales for items with multiple choice or interval (0–10) scales (Q2, Q3, Q16, Q49, Q50, and Q51) were also not changed.

**Table 1** Summary of Shortcomings of the PCQ-Original and Proposed Solutions as Indicated by Fertility Health Care Experts and the Dutch Patient Association Freya

Shortcomings	Solution
(1) Low response rate	Reduction of number of items
(2) Interpretability	Reduction of number of items Proper choice of wording Consistency in nature of the item Application of metric response scales
(3) Size	Reduction of number of items

**Table 2** PCQ-Original and PCQ-Revised Items and Answer Types

Original Item	Answer Type (n)	Deleted	Revised Item	Answer Type (n)
<b>Background Information</b>				
1. For which hospital are you filling in this questionnaire?	Open	Yes	–	–
2. What is your highest successfully completed education?	Multiple choice (9)	No	1. What is your highest completed level of education?	Multiple choice (8)
3. Which treatment are you undergoing or did you finish?	Multiple choice (6)	No	2. What treatment are you currently undergoing or have you just completed?	Multiple choice (5)
4. Are you currently pregnant?	Dichotomous (Yes/No)	No	3. Are you currently pregnant?	Dichotomous (Yes/No)
<b>Dimension “Accessibility”</b>				
5. How often did you get someone to speak to you right away when calling the fertility department?	Likert-scale (4)	No	4. When I called the clinic, I was able to speak to someone without waiting.	Likert-scale (5) + “N/a”
6. When having items, was it a problem to reach the treatment team (by phone or mail)?	Likert-scale (4)	No	5. If I had any questions, I could easily reach the Reproductive Medicine Department (by email or phone).	Likert-scale (5) + “N/a”
<b>Dimension “Information”</b>				
7. Have you received information on how to reach the treatment team outside of office hours (when having urgent items or issues)?	Dichotomous (Yes/No)	No	6. I received clear information on how to reach the Reproductive Medicine Department outside of office hours.	Likert-scale (5)
8. In addition to oral information, have you also received written information?	Likert-scale (3)	No	7. I received written and/or digital information in addition to oral information.	Likert-scale (5)
9. Was the information about your planned examinations clear?	Likert-scale (4)	No	8. The explanations about examinations that were to be carried out on me were clear and understandable.	Likert-scale (5)
10. Have you been informed about the different treatment options?	Likert-scale (3)	Yes	-	-
11. Was the information about your planned treatment clear?	Likert-scale (4)	No	9. The explanations about the treatment I was to receive were clear and understandable.	Likert-scale (5)
12. Have you received an overview with the schedule of your course of treatment?	Dichotomous (Yes/No)	No	10. I received a clear and understandable treatment overview with time planning.	Likert-scale (5)
13. Have the potential side effects of prescribed medicine been explained?	Likert-scale (4)	Yes	-	-
14. Were the instructions clear on how to inject the medication?	Likert-scale (5)	No	11. I received clear and understandable instructions on how to use the fertility drugs (instructions on how to take or inject them).	Likert-scale (5) + “N/a”
15. Did the treatment team inform you about the possibility to receive support from a social worker or psychologist?	Likert-scale (3)	No	26. I received sufficient information from the treatment team about the possibilities of receiving psychological help from a psychologist or social worker.	Likert-scale (5)
16. Have you missed receiving information from a nurse? If so, at which moment(s)?	Multiple choice (7) Multiple answers are allowed	No	12. I lacked explanation from members of the treatment team at the following times (multiple answers possible)	Multiple choice (7) Multiple answers allowed
17. Were there evaluation talks to look back on your course of treatment?	Likert-scale (4)	Yes	-	-
<b>Dimension “Communication”</b>				
18. Were your practitioners honest and clear about what you could expect from the healthcare?	Likert-scale (4)	Yes	-	-
19. Have the results of your research been discussed with you?	Likert-scale (3)	Yes	-	-

(Continued)

**Table 2** (Continued).

Original Item	Answer Type (n)	Deleted	Revised Item	Answer Type (n)
<b>Dimension “Communication”</b>				
20. How often did the doctor listen to you attentively?	Likert-scale (4)	No	13. The doctor listened to me attentively.	Likert-scale (5)
21. How often did the doctor take you seriously?	Likert-scale (4)	No	14. The doctor took me seriously.	Likert-scale (5)
22. How often did the doctor take time for you?	Likert-scale (4)	No	15. The doctor took time for me.	Likert-scale (5)
23. How often did you have the idea that people were talking about you instead of with you?	Likert-scale (4)	Yes	-	-
24. Was the treatment team prepared to talk to you about mistakes or things that did not go well?	Dichotomous (Yes/No)	No	16. The treatment team was willing to talk to me about mistakes and things that did not go well.	Likert-scale (5) + “N/a”
<b>Dimension “Patient involvement”</b>				
25. How often stood the doctor open for your opinion and ideas?	Likert-scale (4)	No	17. The doctor was open to my opinion and ideas.	Likert-scale (5)
26. How often were you granted the opportunity to ask the doctor items?	Likert-scale (4)	No	18. I was given the opportunity to ask the doctor questions.	Likert-scale (5)
27. Was there - if you had the wish to - The opportunity to co-decide on your treatment?	Likert-scale (4)	No	19. I was given the opportunity to have a say in my treatment.	Likert-scale (5)
<b>Dimension “Respect for patient”'s values’</b>				
28. Did you have access to your own medical file?	Likert-scale (4)	No	20. I had access to my patient file during my course of treatment.	Dichotomous (Yes/No)
29. How often did the doctor show interest in your personal situation?	Likert-scale (3)	No	21. The doctor was interested in my personal situation.	Likert-scale (5)
30. How often did the doctor understand your emotions and current situation?	Likert-scale (4)	No	22. The doctor understood my emotions.	Likert-scale (5)
31. Did the nurses show empathy for your situation?	Likert-scale (4)	No	23. The nurses were understanding of my personal situation.	Likert-scale (5)
32. Did the treatment team also pay attention to your partner?	Likert-scale (4)	No	24. The treatment team also responded to my partner.	Likert-scale (5) + “N/a”
33. How often during the treatment(s) was there personal supervision by the nurses?	Likert-scale (4)	Yes	-	-
34. Did the treatment team pay attention to the potential emotional consequences of fertility issues?	Likert-scale (5)	No	25. The treatment team addressed the possible emotional consequences of fertility problems.	Likert-scale (5)
<b>Dimension “Continuity and transition”</b>				
35. Have you been appointed a single person of contact, who you – in case you have items or issues – can always get in contact with?	Dichotomous (Yes/No)	No	27. A contact person in the team was designated whom I could contact at any time (eg a doctor or nurse(s)).	Dichotomous (Yes/No)
36. How many different doctors are or have been involved in your treatment(s)?	Likert-scale (3)	Yes	-	-
37. Did you have a single doctor (head practitioner) with whom you had the most important conversations and with whom you made decisions?	Likert-scale (3)	No	28. I had my own doctor (primary caregiver) with whom I had the most important conversations and made decisions.	Dichotomous (Yes/No)
38. How often did you have your appointments with the same doctor?	Likert-scale (4)	Yes	-	-

(Continued)



Table 2 (Continued).

Original Item	Answer Type (n)	Deleted	Revised Item	Answer Type (n)
<b>Dimension “Continuity and transition”</b>				
39. How often did you have to repeat your story for different doctor?	Likert-scale (4)	No	29. I only had to tell my story once within the treatment team.	Likert-scale (5)
40. How often did you receive opposing information or advice?	Likert-scale (4)	No	30. I received clear and understandable information or recommendations from those treating me.	Likert-scale (5)
41. Did practitioners contradict each other when talking about policy?	Likert-scale (4)	Yes	-	-
<b>Dimension “Competence”</b>				
42. How often did the treatment team use specific terminology without explaining the meaning?	Likert-scale (4)	Yes	-	-
43. How often was your doctor well-prepared if you came to see him / her for an appointment?	Likert-scale (3)	No	31. The doctor was well prepared when I went to see him/her for an appointment.	Likert-scale (5)
44. Did the doctor(s) make a professional impression on you?	Likert-scale (4)	No	32. The doctor(s) made a professional impression on me.	Likert-scale (5)
45. How often did the treatment team work carelessly?	Likert-scale (4)	No	33. The treatment team worked carefully.	Likert-scale (5)
46. How often did the organization go smoothly (well organized) at the polyclinic?	Likert-scale (4)	No	34. The organisation at the fertility centre/ clinic went smoothly (well organised).	Likert-scale (5)
47. How long did you usually have to wait in the waiting room?	Likert-scale (4)	Yes	-	-
<b>Remaining Items</b>				
48. How often did you have to deal with a waiting time longer than 3 weeks when making an appointment with the doctor?	Likert-scale (4)	No	35. I never had to wait longer than 3 weeks when I wanted to make an appointment with the doctor.	Likert-scale (5)
49. How much time was there in between your 1st visit to the hospital and the moment you received your treatment plan?	Multiple choice (4)	No	36. Between my first visit to the fertility centre/clinic and receiving a treatment plan, there were...	Multiple choice (5)
50. On average, how long did you have to “unnecessarily” wait before you were able to start with the next treatment? (eg due to summer break)	Multiple choice (5)	No	37. How long did you have to wait on average before you could start another treatment due to organisational reasons (for example, due to a waiting list or summer break)?	Multiple choice (5) + “N/a”
<b>To Conclude</b>				
51. What is your rating for the overall fertility healthcare in your hospital?	Interval scale 0–10 (11)	No	38. How likely are you to recommend fertility treatment at your fertility centre or clinic to others?	Interval scale (10)
-	-	-	39. What grade would you give the overall fertility treatment at your fertility centre or clinic?	Interval scale (10)

**Notes:** Van Empel, Inge W.H.; Aarts, Johanna W.M., Measuring patient-centredness, the neglected outcome infertility care: a random multicentre validation study, Human Reproduction, 2010, 25, 10, 2516-2526, by permission of Oxford University Press.<sup>13</sup>

## Testing and Validation of the Revised PCQ-Infertility

### Reliability of the Original and the Revised Version of the PCQ-Infertility

Reliability analysis was performed on the PCQ-original, which was completed by 844 patients, and on the PCQ-revised, which was completed by 260 new patients who did not previously complete the PCQ-original (Materials and Methods, [Psychometric Analyses](#)). All items from the “Background Information” section were excluded from the reliability analysis,

since they either have multiple-choice or dichotomous (yes/no) response scales, which are not eligible for calculation of reliability statistics. The same accounts for Q7, Q12, Q16, Q24, and Q35 of the main section of the PCQ-original. No reliability analysis was performed on the section “Remaining Items”, as these items do not belong to a dimension.

In contrast to the PCQ-original, no Cronbach alpha value was  $< 0.5$  in the PCQ-revised, which indicates that all dimensions are at least reliable to an acceptable standard (Table 3). Importantly, the reduction in number of items did not harm the internal consistency and reliability. For the dimension “Respect for patient’s values”, a higher Cronbach’s alpha was reached when Q24 was deleted (data not shown). However, Freya indicated that this item is essential and should therefore not be deleted from the questionnaire.

### Confirmatory Factor Analysis

As the reliability was high in the PCQ-revised, a confirmatory factor analysis was performed to confirm content and discriminant validity of the dimensions, and if the items fit in the previously defined dimensions of the PCQ-original<sup>13</sup> (Table 4). The confirmatory factor analysis showed good model fit according to the root mean square error of approximation (RMSEA) of 0.072, and the comparative fit index (CFI) was 0.881, which was only slightly below 0.90.<sup>31,32</sup> In five out of seven dimensions, convergent validity was present as shown with an average variance extracted (AVE) scores  $> 0.5$  Table 4. Only the dimensions “Accessibility” (AVE = 0.437) and “Information” (AVE = 0.434) had an AVE  $< 0.5$ . Lower factor loadings of Q6 (0.305) and Q7 (0.413) could explain the slight lack of convergent validity in the dimension “Information”. This suggest that these items relate better to a separate latent construct. The dimension “Accessibility” might benefit from

**Table 3** Reliability Analyses of the PCQ-Original and the PCQ-Revised

Dimension	Cronbach's $\alpha$ PCQ-Original n = 844	Cronbach's $\alpha$ PCQ-Revised n = 260
Accessibility	0.481	0.582
Information	0.701	0.759
Communication	0.893	0.863
Patient involvement	0.748	0.777
Respect for patient's values	0.848	0.860
Continuity and transition	0.718	0.697
Competence	0.717	0.811

**Table 4** Confirmatory Factor Analysis of the PCQ-Revised

Dimension	Item	Loading	Cronbach's $\alpha$	AVE
Accessibility	4. When I called the clinic, I was able to speak to someone without waiting.	0.594	0.582	0.437
	5. If I had any questions, I could easily reach the Reproductive Medicine Department (by email or phone).	0.722		
Information	6. I received clear information on how to reach the Reproductive Medicine Department outside of office hours.	0.305	0.759	0.434
	7. I received written and/or digital information in addition to oral information.	0.413		
	8. The explanations about examinations that were to be carried out on me were clear and understandable.	0.881		
	9. The explanations about the treatment I was to receive were clear and understandable.	0.898		
	10. I received a clear and understandable treatment overview with time planning.	0.528		
	11. I received clear and understandable instructions on how to use the fertility drugs (instructions on how to take or inject them).	0.515		

(Continued)



**Table 4** (Continued).

Dimension	Item	Loading	Cronbach's $\alpha$	AVE
Communication	13. The doctor listened to me attentively.	0.930	0.863	0.667
	14. The doctor took me seriously.	0.926		
	15. The doctor took time for me.	0.832		
	16. The treatment team was willing to talk to me about mistakes and things that did not go well.	0.502		
Patient involvement	17. The doctor was open to my opinion and ideas.	0.667	0.777	0.510
	18. I was given the opportunity to ask the doctor questions.	0.810		
	19. I was given the opportunity to have a say in my treatment.	0.655		
Respect for patient's values	21. The doctor was interested in my personal situation.	0.775	0.860	0.511
	22. The doctor understood my emotions.	0.780		
	23. The nurses were understanding of my personal situation.	0.711		
	24. The treatment team also responded to my partner.	0.718		
	25. The treatment team addressed the possible emotional consequences of fertility problems.	0.752		
	26. I received sufficient information from the treatment team about the possibilities of receiving psychological help from a psychologist or social worker.	0.519		
Continuity and transition	29. I only had to tell my story once within the treatment team.	0.671	0.697	0.512
	30. I received clear and understandable information or recommendations from those treating me.	0.737		
Competence	31. The doctor was well prepared when I went to see him/her for an appointment.	0.644	0.811	0.517
	32. The doctor(s) made a professional impression on me.	0.789		
	33. The treatment team worked carefully.	0.793		
	34. The organisation at the fertility centre/clinic went smoothly (well organised).	0.636		

**Abbreviations:** Loading, factor loading; AVE, average variance extracted.

the inclusion of extra items for more variance explained, as factor loadings showed that the included items do load well onto the dimension. Discriminant validity for all dimensions was confirmed by the Fornell and Larcker criterion.<sup>29</sup>

## Discussion

Our study showed that the PCQ-original was improved by thorough evaluation of the number of items, the phrasing and framing of both items and answers, and by implementation of consistent 5-point-Likert answering scales, while keeping the original seven dimensions. We provide a fully revised PCQ-Infertility that is shorter, more comprehensible, and more reliable than the original PCQ-Infertility ([Supplementary Data](#) for the final PCQ-revised). This should improve response rates and offer easier data analysis and data interpretation for healthcare professionals.

The reduction in number of items and the changes we made in wording increased internal consistency on five dimensions while keeping it stable on two dimensions. Two dimensions had a Cronbach's alpha lower than 0.7, indicating that for these two dimensions, it is also important to evaluate the individual item scores. Importantly, the overall reliability of the revised questionnaire was improved. We expect that the response rate of the PCQ-revised will be higher than the PCQ-original because it will take much less time to complete. In addition, the rephrased items and answers are easier to comprehend, which helps patients to communicate their experiences better. It is likely that this will lead to a reduction in careless responses and improve the measurement of patients' experience of fertility health-care quality.

Another important improvement we made to the PCQ-original was the implementation of 5-point Likert scales for all items that originally had 3-, 4-point Likert scales or dichotomous (yes/no) answering scales. This increases ease of use of the scores and reduces data analysis time. Raw scores of the PCQ-revised can be used to calculate Quality

Improvement (QI) scores to measure patient-centeredness and using 5-point Likert scales increases reliability of such scores.<sup>13</sup>

Our study is not without limitations. This study included only patients from Dutch fertility clinics and the PCQ-revised was validated in a smaller patient population than the PCQ-original. Patients who filled in the PCQ-revised came from two fertility clinics, while the PCQ-original was completed by patients coming from four fertility clinics, causing a possible bias. It is therefore important to validate the PCQ-revised in a larger, more diverse patient population coming from different fertility clinics. It is also important to note that we decided to keep items in the questionnaire upon request of Freya, although deletion of these items would have increased reliability of the dimensions. The reliability of the dimension “Continuity and transition” was previously reported to be less reliable in the PCQ-original,<sup>13,19</sup> and deleting items in this dimension did not improve the reliability in the PCQ-revised. However, the Cronbach’s  $\alpha$  was just below 0.7, indicating that minor adaptations could improve the reliability. The reliability of the dimension “Accessibility” was significantly improved in the PCQ-revised compared to the PCQ-original, but it was still below 0.7. Future research should focus on improving the reliability of these two dimensions.

The PCQ-revised is digitalized and can be completed within ten minutes, which fits better in the current day-to-day practice than the paper version of the PCQ-original. This is an important step towards the implementation of a standardized evaluation system that can help fertility health care experts improving their health care. The PCQ-revised is now implemented in more than thirty Dutch fertility clinics. They can access the PCQ-revised outcome data from their clinic via a dashboard, perform data extraction and evaluate the results directly. The dashboard shows the quality improvement (QI) scores of each domain, which gives them the opportunity to immediately act upon the findings, ultimately leading to an improvement of the quality of fertility care. The next steps are to validate the PCQ-revised in other countries and to evaluate patient satisfaction in fertility care with this new digital measurement tool on an even large scale.

## Conclusion

The PCQ-revised, measuring patient satisfaction in fertility care, is more valid, reliable, and shorter than the PCQ-original. The questions are easier to comprehend, leading to a higher quality of answers. In addition, improved answering scales can make data analysis easier and faster. The PCQ-revised is digitalized and currently implemented in more than thirty Dutch clinics, indicating that it fits the current day-to-day practice and that it helps fertility health care experts to keep improving the quality of their patient-centered care.

## Abbreviations

PCQ, Patient-Centeredness Questionnaire; PROM, Patient Reported Outcome Measure; PREM, Patient Reported Experience Measure; CQI, Consumer Quality Index; CFI, Comparative Fit Index; RMSEA, Root Mean Square Error of Approximation; AVE, Average Variance Extracted; N/A, Not Applicable.

## Data Sharing Statement

All data generated or analysed during this study are included in this published article and its [Supplementary Information](#).

## Ethics Approval and Consent to Participate

All participants were informed about the study and voluntarily participated. According to Dutch law, approval by an institutional ethics committee was not required for this study as data were anonymously collected and not traceable.

## Acknowledgments

We would like to thank all patients who voluntarily completed the PCQ-Infertility, the healthcare professionals who are actively involved in the invitation of patients, C. Ijlstra from LiveResearch for her assistance in data extraction, and J.M. L. Knijnenburg from Freya, drs. C. Leendertz from Hospital De Gelderse Vallei, the Netherlands, and J. Germs and K. Feicke from the University Medical Center Groningen (UMCG), the Netherlands for their expert insights.

## Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

The writing and publication of this work was supported by Ferring B.V., the Netherlands.

## Disclosure

J. Bloemer received a financial compensation from Ferring BV for reviewing this article. The authors report no other conflicts of interest in this work.

## References

1. Institute of Medicine (US) Committee on Quality of Health Care in America. Crossing the quality chasm: a new health system for the 21st century. *BMJ*. 2001;2001:323.
2. Keirns CC, Goold SD. Patient-centered care and preference-sensitive decision making. *JAMA*. 2009;302:1805–1806. doi:10.1001/jama.2009.1550
3. Aarts JWM, Faber MJ, van Empel IWH, et al. Professionals' perceptions of their patients' experiences with fertility care. *Human Reprod*. 2011;26:1119–1127. doi:10.1093/humrep/der054
4. Huppelschoten AG, Aarts JWM, van Empel IWH, et al. Feedback to professionals on patient-centered fertility care is insufficient for improvement: a mixed-method study. *Fertil Steril*. 2013;99:1419–1427. doi:10.1016/j.fertnstert.2012.12.024
5. Dancet EAF, Nelen WL, Sermeus W, et al. The patients' perspective on fertility care: a systematic review. *Hum Reprod Update*. 2010;16:467–487. doi:10.1093/humupd/dmq004
6. Blyth E. Fertility patients' experiences of cross-border reproductive care. *Fertil Steril*. 2010;94:e11–e15. doi:10.1016/j.fertnstert.2010.01.046
7. Van Empel IWH, Hermens RP, Akkermans RP, et al. Organizational determinants of patient-centered fertility care: a multilevel analysis. *Fertil Steril*. 2011;95:513–519. doi:10.1016/j.fertnstert.2010.08.021
8. Van Empel IWH, Nelen WL, Tepe ET, et al. Weaknesses, strengths and needs in fertility care according to patients. *Human Reprod*. 2010;25:142–149. doi:10.1093/humrep/dep362
9. Miles A, Mezzich J. The care of the patient and the soul of the clinic: person-centered medicine as an emergent model of modern clinical practice. *Int J Person Center Med*. 2011;1. doi:10.5750/ijpcm.v1i2.61
10. Verberg MFG, Eijkemans MJC, Heijnen EM, et al. Why do couples drop-out from IVF treatment? A prospective cohort study. *Human Reprod*. 2008;23:2050–2055. doi:10.1093/humrep/den219
11. Gerteis M, Edgman-Levitan S, Daley J, Delbanco TL. *Through the Patient's Eyes*. Jossey-Bass; 1993.
12. Weldring T, Smith SMS. Article commentary: Patient-Reported Outcomes (PROs) and Patient-Reported Outcome Measures (PROMs). *Health Services Insights*. 2013;6:61–68. doi:10.4137/HSI.S11093
13. Van Empel IWH, Aarts JWM, Cohlen BJ, et al. Measuring patient-centredness, the neglected outcome in fertility care: a random multicentre validation study. *Human Reprod*. 2010;25:2516–2526. doi:10.1093/humrep/deq219
14. Pedro J, Canavarro MC, Boivin J, Gameiro S. Positive experiences of patient-centred care are associated with intentions to comply with fertility treatment: findings from the validation of the Portuguese version of the PCQ-infertility tool. *Human Reprod*. 2013;28:2462–2472. doi:10.1093/humrep/det259
15. Aarts JWM, Huppelschoten AG, van Empel IWH, et al. How patient-centred care relates to patients quality of life and distress: a study in 427 women experiencing infertility. *Human Reprod*. 2012;27:488–495. doi:10.1093/humrep/der386
16. Gameiro S, Canavarro MC, Boivin J. Patient centred care in infertility health care: direct and indirect associations with wellbeing during treatment. *Patient Educ Couns*. 2013;93:646–654. doi:10.1016/j.pec.2013.08.015
17. Karajičić S. *Towards Patient-Centered Infertility Health Care: Case Study Slovak Republic*. Health Policy Institute; 2014.
18. Dizaj AVT, Ezabadi Z, Takbiri A, Baniyadi A. Infertility care in Royan Institute: a patient centeredness analysis. *Payesh*. 2016;15:325–332.
19. Mourad SM, Curtis C, Gudex G, et al. Measuring patient-centredness in publicly funded fertility care: a New Zealand validation and international comparison of the Patient-Centred Questionnaire-Infertility. *Aust N Z J Obstet Gynaecol*. 2019;59:265–271. doi:10.1111/ajo.12869
20. Delnoij D, Hendriks M. De CQ-index: het meten van klantervaringen in de zorg. *TSG*. 2008;86:440–446. doi:10.1007/BF03082141
21. Sixma H, Spreeuwenberg P, Zuidgeest M, Rademakers J. CQ-index Ziekenhuisopname: meetinstrumentontwikkeling. *Nivel*. 2009;2009:254.
22. Bast B. CQ-index is duur en ineffectief om cliënttevredenheid te verbeteren. *Zorgvisie*. 2011;2011:256.
23. Credé M. Random responding as a threat to the validity of effect size estimates in correlational research. *Educ Psychol Meas*. 2010;70:596–612. doi:10.1177/0013164410366686
24. Meade AW, Craig SB. Identifying careless responses in survey data. *Psychol Methods*. 2012;17:437–455. doi:10.1037/a0028085
25. van Sonderen E, Sanderman R, Coyne JC. Ineffectiveness of reverse wording of questionnaire items: let's learn from cows in the rain. *PLoS One*. 2013;8:1–7.
26. Weijters B, Baumgartner H. Misresponse to reversed and negated items in surveys: a review. *J Market Res*. 2012;49:737–747. doi:10.1509/jmr.11.0368
27. Reeve BB, Wyrwich KW, Wu AW, et al. ISOQOL recommends minimum standards for patient-reported outcome measures used in patient-centered outcomes and comparative effectiveness research. *Qual Life Res*. 2013;22:1889–1905. doi:10.1007/s11136-012-0344-y

28. Bollen KA. *Structural Equations with Latent Variables*. John Wiley & Sons, Inc.; 1989; doi:10.1002/9781118619179
29. Fornell C, Larcker DF. Evaluating structural equation models with unobservable variables and measurement error. *J Market Res*. 1981;18:39–50. doi:10.1177/002224378101800104
30. Suárez-Alvarez J, Pedrosa I, Lozano LM, et al. Using reversed items in Likert scales: a questionable practice. *Psicothema*. 2018;30:149–158. doi:10.7334/psicothema2018.33
31. Browne MW, Cudeck R. Alternative ways of assessing model fit. *Sociol Methods Res*. 1993;21:230–258. doi:10.1177/0049124192021002005
32. Jöreskog KG, Sörbom D. *LISREL 8: Structural Equation Modeling with the SIMPLIS Command Language*. Ill: Scientific Software International; 1998.

#### Patient Related Outcome Measures

Dovepress

### Publish your work in this journal

Patient Related Outcome Measures is an international, peer-reviewed, open access journal focusing on treatment outcomes specifically relevant to patients. All aspects of patient care are addressed within the journal and practitioners from all disciplines are invited to submit their work as well as healthcare researchers and patient support groups. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/patient-related-outcome-measures-journal>