

Viewpoint on “The Frail Scale – A Risk Stratification in Older Patients with Acute Coronary Syndrome” [Response to Letter]

Hung Manh Pham^{1,2}, Anh Phuong Nguyen^{1,2}, Huong Thi Thu Nguyen^{3,4}, Tam Ngoc Nguyen^{3,4}, Thanh Xuan Nguyen^{3,4}, Thu Thi Hoai Nguyen^{3,4}, Huong Thi Thanh Nguyen^{5,6}, Anh Trung Nguyen^{3,4}, Quang Ngoc Nguyen^{1,2}, Giang Song Tran¹, Huyen Thi Thanh Vu^{3,4}

¹Vietnam National Heart Institute, Bach Mai Hospital, Hanoi, 100000, Vietnam; ²Department of Cardiology, Hanoi Medical University, Hanoi, 100000, Vietnam; ³Department of Geriatrics, Hanoi Medical University, Hanoi, 100000, Vietnam; ⁴Scientific Research Department, National Geriatric Hospital, Hanoi, 100000, Vietnam; ⁵Physiology Department, Hanoi Medical University, Hanoi, 100000, Vietnam; ⁶Dinh Tien Hoang Institute of Medicine, Hanoi, 100000, Vietnam

Correspondence: Anh Phuong Nguyen, Vietnam National Heart Institute, Bach Mai Hospital, Hanoi, 100000, Vietnam, Tel +84 989314464, Email npa6@bachmai.edu.vn

Dear editor

We would like to thank doctor Xinyue Li for comments on our paper and the Editor for the opportunity to respond.^{1,2} Here is point-by-point response to the author's comments and concerns.

Firstly, we discuss issues related to renal disease and frailty in elderly patients with acute coronary syndromes. As mentioned by the author, there is an association between patients with kidney disease and frailty in the elderly, not in the group of patients with acute coronary syndrome. Based on studies in the literature, we also find that chronic kidney disease (CKD) and acute kidney injury (AKI) in elderly patients with acute coronary syndromes are truly interesting.^{3,4} However, this issue is not within the scope of our research, especially during the COVID pandemic, when testing conditions are limited. Therefore, we save this issue for the following studies. Secondly, the category of BMI in Table 1, the proportion of obese patients is too low 1/116 (0.86%), so we do not analyze it separately, but group it together with the overweight group. Thirdly, in-hospital adverse outcome is composite event, so it includes major bleeding and stroke events in the NACE.

About some statistical analysis, firstly, in Table 3, we agree that presenting the reference set as 1 as suggested would be clearer and more accessible. Secondly, the regression analysis presented in Table 4, as the number of variables included increased, the OR values increased. In statistics, when increasing the number of variables in a multivariable logistic regression analysis, the OR can increase or decrease, depending on the relationship and impact between the variables. So, our results are really appropriate because these results further show the truly relationship between frailty and NACE after adjusting for the impact of related factors.

In conclusion, we greatly appreciate the author's comments and suggestions. We fully agree with author about the potential role of the simple FRAIL score in risk stratification for older patients with ACS in the coronary care setting and that it is important to consider the need for evidence and extensive experience among different populations because the incidence of these different patients.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Xinyue L. Viewpoint on “The frail scale – a risk stratification in older patients with acute coronary syndrome”. *J Multidiscip Healthc*. 2023;16:1937–1938. doi:10.2147/JMDH.S429655

2. Pham HM, Nguyen AP, Nguyen HTT, et al. The frail scale - a risk stratification in older patients with acute coronary syndrome. *J Multidiscip Health c.* **2023**;16:1521–1529. doi:10.2147/JMDH.S409535
3. Hashimoto Y, Ozaki Y, Kan S, et al.; J-MINUET Investigators. Impact of chronic kidney disease on in-hospital and 3-year clinical outcomes in patients with acute myocardial infarction treated by contemporary percutaneous coronary intervention and optimal medical therapy- insights from the J-MINUET Study. *Circ J.* **2021**;85(10):1710–1718. doi:10.1253/circj.CJ-20-1115
4. Chalikias G, Serif L, Kikas P, et al. Long-term impact of acute kidney injury on prognosis in patients with acute myocardial infarction. *Int J Cardiol.* **2019**;283:48–54. doi:10.1016/j.ijcard.2019.01.070

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Multidisciplinary Healthcare 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Multidisciplinary Healthcare editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>

<https://doi.org/10.2147/JMDH.S432083>