

LETTER

Suicidality and Its Association with Stigma in Clinically Stable Patients with Schizophrenia in Rural China [Letter]

Riya Gosrani^{1,2}

UCL Medical School, University College London, London, UK; ²School of Anthropology and Museum Ethnography, University of Oxford, Oxford, UK

Correspondence: Riya Gosrani, UCL Medical School, 74 Huntley St, London, WCIE 6DE, UK, Email riya.gosrani.17@ucl.ac.uk

Dear editor

I recently read with keen interest the article by Wang et al that investigates the relationship between suicidality and stigma among clinically stable patients living with schizophrenia in China's rural Anhui Province. 1 Given that suicide is a leading cause of death among individuals with schizophrenia, elucidating associated risk factors represents a crucial area for research.² Whilst I commend the authors for their contributions on this important subject, I believe there are some aspects of the study that warrant further reflection. As a final year medical student with a background in medical anthropology, I aim to provide some pertinent insights and recommendations in this letter.

My initial concern revolves around the consistent use of the term "commit suicide" throughout the paper. Considering the focus of the study, the role of language in either perpetuating or alleviating stigma cannot be overstated. Given the historical and present-day criminalisation of suicide in different parts of the world, the word "commit" inadvertently insinuates undertones of misconduct.³ I believe that it is vital for medical professionals to be acutely aware of their language choices. Opting for terms that are less stigmatising holds the potential to transform perceptions, making a world of difference.³ I would thus propose replacing "committed suicide" with a more neutral phrase such as "died by suicide".

Secondly, the authors' decision to use the Social Impact Scale (SIS) to assess stigma among Chinese patients diagnosed with schizophrenia merits discussion. Psychometric tools such as the SIS, originally developed in English, may not resonate universally across varied cultural landscapes. Whilst high Cronbach's alpha values suggest internal consistency, this does not necessarily ensure the suitability of the SIS in diverse settings. Creating tools that are culturally appropriate and tailored to address specific manifestations of stigma are crucial for designing effective interventions. Accordingly, I advocate for a robust validation of the SIS within this specific demographic to further substantiate the study's findings.

Furthermore, the standards employed in the study to determine clinically stable patients are debatable, describing this as less than a 50% change in psychiatric medication over the past three months. Considering the authors' own observation that individuals in rural areas face limited access to medical resources, I urge a re-evaluation of these criteria. This is because a lack of regular medical check-ups can lead to misleading indications regarding a patient's clinical stability.

Lastly, the authors highlighted that patients displaying negative schizophrenia symptoms sometimes offer selfassessments that do not mirror the actual situation. I would argue that this potential discrepancy is not limited to patients with negative symptoms alone. Given the neurocognitive deficits often associated with positive symptoms of schizophrenia,² these individuals can struggle to accurately gauge their cognitive and functional capacities.⁵ To mitigate inconsistencies in self-assessments across a patient cohort exhibiting both positive and negative symptoms, incorporating feedback from collateral informants could enhance confidence in the results.

Gosrani **Dove**press

In conclusion, I sincerely appreciate the authors' contributions to this journal and trust that my comments will be taken in the constructive spirit in which they are offered.

Disclosure

The author reports no conflicts of interest in this communication.

References

- 1. Wang R, Zheng S, Ouyang X, et al. Suicidality and its association with stigma in clinically stable patients with schizophrenia in rural China. Psychol Res Behav Manag. 2023;16:1947-1956. doi:10.2147/prbm.s413070
- 2. Grover LE, Jones R, Bass NJ, McQuillin A. The differential associations of positive and negative symptoms with suicidality. Schizophr Res. 2022;248:42-49. doi:10.1016/j.schres.2022.07.016
- 3. Sommer-Rotenberg D. Suicide and language. Can Med Assoc J. 1998;159(3):239-240.
- 4. Weiss MG, Jadhav S, Raguram R, Vounatsou P, Littlewood R. Psychiatric stigma across cultures: local validation in Bangalore and London. Anthropol Med. 2001;8(1):71-88. doi:10.1080/13648470120063906
- 5. Morgan O, Strassnig MT, Moore RC, et al. Accuracy of immediate self-assessment of neurocognitive test performance: associations with psychiatric diagnosis and longitudinal psychotic symptoms. J Psychiatr Res. 2022;156:594-601. doi:10.1016/j.jpsychires.2022.10.069

Dove Medical Press encourages responsible, free and frank academic debate. The contentTxt of the Psychology Research and Behavior Management 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Psychology Research and Behavior Management editors. While all reasonable steps have been taken to confirm the contentTxt of each letter, Dove Medical Press accepts no liability in respect of the contentTxt of any letter, nor is it responsible for the contentTxt and accuracy of any letter to the editor.

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/psychology-research-and-behavior-management-journal





