

Quantifying the Adverse Effects of Long COVID on Individuals' Health After Infection: A Propensity Score Matching Design Study

Yudong Miao¹, Ruizhe Ren¹, Zhanlei Shen¹, Yi Li¹, Wanliang Zhang¹, Junwen Bai¹, Dongfang Zhu¹, Jingbao Zhang¹, Dan Guo^{1,2}, Clifford Silver Tarimo³, Rongmei Liu⁴, Qiuping Zhao⁴, Jianping Hu⁵, Xue Zhou⁶, Wenying Dong⁷

¹College of Public Health, Zhengzhou University, Zhengzhou, People's Republic of China; ²Department of Neurology, Henan Provincial People's Hospital, Zhengzhou, People's Republic of China; ³Department of Science and Laboratory Technology, Dar es Salaam Institute of Technology, Dar es Salaam, Tanzania; ⁴Henan Key Laboratory for Health Management of Chronic Diseases, Central China Fuwai Hospital, Zhengzhou, People's Republic of China; ⁵Henan Medical Communication and Project Forward Center, Zhengzhou, People's Republic of China; ⁶College of Health Management, Mudanjiang Medical University, Mudanjiang, People's Republic of China; ⁷Department of Hypertension, Henan Provincial People's Hospital, Zhengzhou, People's Republic of China

Correspondence: Rongmei Liu, Assistant research fellow, Fuwai Central China Cardiovascular Hospital, Zhengzhou University Affiliated Central China Fuwai Hospital, #1, Fuwai Avenue, Zhengdong New District, Zhengzhou, 450014, People's Republic of China, Tel +8615038204525, Email liuyongmei0203@163.com

Objective: To evaluate the prevalence and influencing factors of long COVID, and measure the difference in health status between long COVID and non-long COVID cases.

Methods: A cross-sectional survey was conducted from February 1 to 8, 2023, using a stratified random sampling method in four regions (eastern [Changzhou], central [Zhengzhou], western [Xining] and northeastern [Mudanjiang]) of China. The survey collected COVID-19 patients' socio-demographic characteristics and lifestyles information. The scores of lifestyles and health status range from 5 to 21 and 0 to 100 points, respectively. The criteria of "persistent health problems after 4 weeks of COVID-19 infection" issued by the US Centers for Disease Control and Prevention was used to assess long COVID. Multiple linear regression was used to analyze the influencing factors of the health. The bootstrap method was used to analyze the lifestyles' mediating effect. Propensity score matching (PSM) was used to evaluate the net difference in health scores between long COVID and non-long COVID cases.

Results: The study included 3165 COVID-19 patients, with 308 (9.73%) long COVID cases. The health score of the long COVID cases (74.79) was lower than that of the non-long COVID cases (81.06). After adjusting for potential confounding variables, we found that never focused on mental decompression was a common risk factor for the health of both groups. Lifestyles was the mediating factor on individuals' health. After PSM, the non-long COVID cases' health scores remained higher than that of long COVID cases.

Conclusion: The proportion of long COVID cases was low, but they were worse off in health. Given the positive moderating effect of healthy lifestyles on improving the health of long COVID cases, healthy lifestyles including mental decompression should be considered as the core strategy of primary prevention when the epidemic of COVID-19 is still at a low level.

Keywords: long COVID, lifestyles, self-reported health status, propensity score matching, effect of mediation

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), a novel virus, has caused a worldwide pandemic of coronavirus disease 2019 (COVID-19). Although the World Health Organization has declared that "COVID-19 no longer constitutes a Public health Emergency of International concern",¹ but this does not mean that COVID-19 infection is over. COVID-19 is still circulating at a low level, and data suggest that by April 2023, approximately 764 million people worldwide had contracted the COVID-19 virus.² A large proportion of infected people can return to their pre-infection life after recovery.³ However, one study has shown that COVID-19 infection can cause sequelae in the immune system, blood system, lung system, cardiovascular system, gastrointestinal system, musculoskeletal system, nervous system, and

other aspects.⁴ These sequelae may persist after infection and may recur after cure.⁵ This disease, which manifests as persistent symptoms that affect some people after infection, is known as long COVID and affects people's health.⁶

Currently, there is no universally known definition of long COVID. In this study, the definition of long COVID as defined by the Centers for Disease Control and Prevention of the United States of America was adopted: the condition in which an individual has persistent symptoms or health problems for at least 4 weeks (28 days) after COVID-19 infection.⁷ Persistent symptoms include a range of physical and mental (psychological) symptoms,⁸ including fatigue,⁹ headache, joint pain,¹⁰ respiratory symptoms,¹¹ cognitive impairment and impaired mental health.¹² The persistent effects of these symptoms can lead to poorer health and reduced quality of life.¹³ One study shows that at least 65 million people are infected with long COVID-19 worldwide,¹⁴ but due to lack of surveillance and reporting, the actual number of infected people may be much higher than reported. Previous studies have revealed risk factors for long COVID-19, such as being female,^{15,16} the number of symptoms during the infection,¹⁷ education level, smoking,^{18,19} sleep,^{20,21} and psychological factors.²²

In February 2023, China announced the implementation of “Class B epidemic and B management” control measures for COVID-19 to reduce the negative impact of large-scale lockdown.²³ This policy indicates that the Chinese government will no longer adopt strict quarantine measures for COVID-19 infected people and close contacts, no longer divide high-risk areas and low-risk areas, and the population in each area can move freely. As a result of this policy, the risk of COVID-19 infection has increased nationwide, and acute cases have been treated in a timely and effective manner. However, the health status of long COVID cases also requires urgent attention. The purpose of this study is to explore the health status and influencing factors of individuals with and without long COVID after the social distancing strategy has ended in China, and to explore the effects of long COVID infection on the health of individuals and the regulatory effect of lifestyle to develop targeted healthy lifestyle guidance and policy recommendations to improve the health status of long COVID cases.

Methods

Procedures and Participants

From February 1 to 8, 2023, a cross-sectional study was conducted in the eastern (Changzhou), central (Zhengzhou), western (Xining) and northeastern (Mudanjiang) areas of China mainland using a stratified random sampling method (see [Figure S1](#)). Within each city, more than two rural areas and two urban areas were randomly selected. A cross-sectional online survey was conducted among adults aged 18 years and above in each household to obtain a sample of 5780 eligible participants.

In this study, a total of 3165 patients infected with COVID-19 were enrolled. The sample exclusion criteria were as follows: minors aged less than 18 years, participants with incomplete information, were not infected with COVID-19, and individuals who were not residents of China mainland.

We performed 1:1 nearest neighbor matching between long COVID cases and non-long COVID cases, and a total of 281 pairs of samples were matched. The participant flow chart is shown in [Figure 1](#).

Ethics Approval

Participants were informed of the benefits and risks of participating in this study, and they provided informed consent. All data were used merely for research purposes. This study was approved by the Life Science Ethics Review Committee of Zhengzhou University (2021–01-12-05).

Assessments

In the questionnaire, participants were asked about the post-COVID-19 symptoms they experienced after testing negative for COVID-19. Additionally, they were asked to indicate the duration of these symptoms in terms of the number of days. Participants who self-reported symptoms lasting for more than 4 weeks (28 days) were categorized as long COVID cases based on the measurement criteria used in the study. The health status of participants was assessed using a questionnaire that asked them to rate their current health status on a scale from 0 to 100. Higher scores on the scale indicated a better health status.

We used 13 covariates (sociodemographic factors and health factors), including gender (male, female), age (18–29, 30–39, 40–49, 50–59, ≥60), marital status (married, other), education level (high school or less, bachelor's degree or more), religious affiliation (atheist, others), ethnic (Han, Minority), annual household income (CNY) (<20,000, 20,000–49,999,




Figure 1 The flowchart of participants selection of this study.

50,000–99,999, 100,000–199,999, $\geq 200,000$), BMI (<18.50 , 18.50 – 24.99 , 25.00 – 29.99 , ≥ 30.00), smoking (smoking, quit, never), drinking (drinking, quit, never), wearing masks (always, often, sometimes, seldom, never), sufficient sleeping (always, often, sometimes, seldom, never), and mental decompression (always, often, sometimes, seldom, never).

To verify the mediating effect of smoking, drinking, wearing masks, sufficient sleeping, and mental decompression on the health status of individuals affected by long COVID, we assigned scores to each aspect. Smoking status was scored as 1 point for current smokers, 2 points for those who had quit smoking, and 3 points for individuals who had never smoked. Similarly, drinking status was scored as 1 point for current drinkers, 2 points for ex-drinkers, and 3 points for those who had never consumed alcohol. The frequency of wearing masks, sufficient sleeping, and mental decompression were evaluated and assigned scores based on the following scale: always = 5 points, often = 4 points, sometimes = 3 points, seldom = 2 points, and never = 1 point. The overall scores for these factors ranged from 5 to 21, reflecting the cumulative assessment of participants' behaviors in these areas.

Statistical Analysis

The rank-sum test of two independent samples and the Kruskal–Wallis test were used to analyze the differences in health status scores between the long COVID cases and non-long COVID cases. The frequencies and proportions of categorical variables were calculated, along with their corresponding health status scores while continuous variables were categorized into quartiles, and the frequencies and proportions within each quartile were reported along with the respective health status scores. The factors influencing the health status score were analyzed by using linear regression models. A bias-corrected non-parametric percentile bootstrap method was used to verify the mediating effect of lifestyle on the

impact of long COVID on the health status of patients. To minimize the potential influence of confounding factors, propensity score matching was employed. Covariate selection was carried out using probit regression, and the propensity score was calculated through probit regression as well. Subsequently, 1:1 nearest neighbor matching was conducted between the long COVID and non-long COVID populations among the 3,165 individuals, ensuring a balanced comparison between the two groups while addressing potential biases. All analyses were performed by using STATA17.0 and SPSS27.0 statistical software, and $P < 0.05$ was considered statistically significant.

Results

Characteristics, Lifestyles and Health Status Scores of Long COVID and Non-Long COVID Cases

The basic characteristics, lifestyles, and health status scores of the participants are presented in Table 1. A total of 3165 COVID-19 patients were included in this study, of which 308 (9.73%, 95% CI: 8.70~10.76) were long COVID cases and 2857 (90.27%, 95% CI: 89.24~91.30) were non-long COVID cases. The overall health status score of COVID-19 patients was 80.45 (95% CI: 79.91~80.99). The health status score of long COVID cases was 74.79 (95% CI: 72.75~76.82), which was lower than that of non-long COVID cases 81.06 (95% CI: 80.50~81.61).

The findings show that among long-COVID cases, lower education level (68.33, 95% CI: 62.87~73.8), higher annual income level of 50,000~99,999 yuan (68.46, 95% CI: 63.73~73.19), never felt sufficient sleep (60.50, 95% CI: 20.16~100.84) and those who never paid attention to mental decompression (61.05, 95% CI: 51.66~70.44) had the lowest health status score. Among non-long COVID cases, the elderly (76.73, 95% CI: 75.45~78.01), lower education level (79.69, 95% CI: 78.78~80.6), non-married (80.71, 95% CI: 80.12~81.29), those who with religious belief (80.56, 95% CI: 76.10~85.02) had the lowest health status score. Those who never drank alcohol (81.52, 95% CI: 80.86~82.19) and those who always wore masks (82.97, 95% CI: 82.03~83.91) had the highest health status scores. In addition, people who always slept enough (81.00, 95% CI: 76.55~85.45; 85.01, 95% CI: 83.85~86.16) and those who always had mental decompression (92.60, 95% CI: 86.99~98.21; 86.95, 95% CI: 85.23~88.68) had higher health status scores.

Influencing Factors of Health Status of Long COVID and Non-Long COVID Cases

After accounting for confounding factors, the results of the linear regression model revealed that religious belief, annual household income, education level, wearing masks, smoking, drinking, and mental decompression (all $P < 0.05$) were significantly associated with the health status of long COVID cases, and the influencing factors of the health status score of non-long COVID cases were age, religious belief, BMI, annual household income, sufficient sleep and mental decompression (all $P < 0.05$), as shown in Figure 2 (see Table S1).

Atheism was associated with a higher health status score among long COVID cases ($\beta = 11.69$, 95% CI: 0.25~23.13), while those with a bachelor's degree or higher had a better health status score ($\beta = 7.44$, 95% CI: 1.14~13.74). People who seldom used masks had lower health scores ($\beta = 7.33$, 95% CI: -14.29~-0.37). Among non-long COVID cases, the health status score of the elderly (≥ 60 years old) was lower ($\beta = -10.30$, 95% CI: -12.91~-7.70), and the health status score of atheism was higher ($\beta = 4.12$, 95% CI: 1.09~7.14). People with normal BMI (18.5~25.0) had higher health status score ($\beta = 4.03$, 95% CI: 0.9~7.15), and those who never felt sufficient sleep had lower health status score ($\beta = -15.8$, 95% CI: -26.72~-4.88).

Moreover, among long COVID cases, those who had quit smoking ($\beta = 10.92$, 95% CI: 4.23~17.62) and those who had never smoked ($\beta = 12.83$, 95% CI: 3.83~21.83) had higher health status scores than smokers. Ex-drinkers ($\beta = -14.28$, 95% CI: -22.68~-5.88) and never-drinkers ($\beta = -5.56$, 95% CI: -11.01~-0.10) had lower health status scores than drinkers, however, there was no statistically significant difference in non-long COVID cases (all $P > 0.05$). In both groups, those who never paid attention to mental decompression had the lowest health status score ($\beta = -25.40$, 95% CI: -38.82~-11.98; $\beta = -4.86$, 95% CI: -7.51~-2.21), which represented the worst health status.

Mediating Effect Analysis of Lifestyle Scores

As shown in Figure 3, long COVID significantly positively predicted health status scores ($c = 6.2734$, $P < 0.001$) and negatively predicted lifestyle scores ($a = -0.3542$, $P < 0.05$). When both long COVID and lifestyle were considered in the

Table 1 The Characteristics, Lifestyles and Health Status Scores of the Study Participants

Variables	Total %	Health Status Scores (95% CI)	P value ^a	Long COVID			Non-long COVID		
				(N=308) %	Health Status Scores (95% CI)	P value ^b	(N=2857) %	Health Status Scores (95% CI)	P value ^c
Total	3165 (100)	80.45 (79.91~80.99)		308 (9.73)	74.79 (72.75~76.82)		2857 (90.27)	81.06 (80.50~81.61)	
Sex			0.413			0.697			0.300
Male	1484 (46.89)	80.30 (79.51~81.08)		123 (39.94)	74.19 (70.94~77.43)		1361 (47.64)	80.85 (80.05~81.65)	
Female	1681 (53.11)	80.58 (79.83~81.33)		185 (60.06)	75.18 (72.57~77.8)		1496 (52.36)	81.25 (80.47~82.02)	
Age			0.0001			0.253			0.0001
18~29	357 (11.28)	85.29 (83.67~86.92)		19 (6.17)	77.73 (67.70~87.77)		338 (11.83)	85.71 (84.10~87.33)	
30~39	890 (28.12)	81.13 (80.06~82.19)		94 (30.52)	73.37 (69.56~77.18)		796 (27.86)	82.05 (80.97~83.12)	
40~49	627 (19.81)	81.18 (79.95~82.42)		78 (25.32)	77.79 (73.97~81.60)		549 (19.22)	81.66 (80.33~82.96)	
50~59	671 (21.20)	80.03 (79.02~81.05)		65 (21.10)	74.08 (70.07~78.09)		606 (21.21)	80.67 (79.65~81.70)	
≥60	620 (19.59)	76.39 (75.14~77.64)		52 (16.88)	72.63 (67.45~77.81)		568 (19.88)	76.73 (75.45~78.01)	
Education level			0.0001			0.017			<0.001
High school or below	1182 (37.35)	79.06 (78.13~79.98)		66 (21.43)	68.33 (62.87~73.8)		1116 (39.06)	79.69 (78.78~80.6)	
Bachelor or above	1983 (62.65)	81.27 (80.61~81.95)		242 (78.57)	76.55 (74.48~78.62)		1741 (60.94)	81.94 (81.24~82.63)	
Marital status			<0.001			0.380			<0.001
Married	283 (89.42)	80.15 (79.59~80.72)		28 (91.23)	71.37 (62.52~80.23)		2549 (89.22)	83.96 (82.21~85.70)	
Other	335 (10.58)	82.94 (81.16~84.73)		27 (8.77)	75.11 (73.05~77.18)		308 (10.78)	80.71 (80.12~81.29)	
Ethnic			0.732			0.730			0.842
Han	3094(97.76)	80.30 (76.01~84.58)		303 (98.38)	74.75 (72.70~76.81)		2791 (97.69)	81.07 (80.51~81.63)	
Minority	71(2.24)	80.45 (79.91~81.00)		5 (1.62)	76.8 (60.02~93.58)		66 (2.31)	80.56 (76.10~85.02)	
Religion			0.008			0.354			0.011
Atheist	303 (95.96)	75.74 (72.30~79.18)		29 (96.75)	75.07 (73.05~77.09)		2739 (95.87)	81.25 (80.69~81.81)	
Others	128 (4.04)	80.65 (80.10~81.19)		10(3.24)	66.4 (48.82~83.98)		118 (4.13)	76.53 (73.12~79.95)	
Annual household income (CNY)			0.185			0.034			0.404
<20,000	246 (7.78)	77.94 (75.63~80.25)		18 (5.84)	76.72 (70.17~83.28)		228 (7.98)	78.04 (75.60~80.48)	
20,000~49,999	510 (16.11)	80.13 (78.69~81.57)		35 (11.36)	74.6 (68.77~80.43)		475 (16.63)	80.54 (79.06~82.02)	
50,000~99,999	725 (22.91)	79.29 (78.05~80.53)		78 (25.32)	68.46 (63.73~73.19)		647 (22.65)	80.60 (79.36~81.83)	
100,000~199,999	847 (26.76)	81.31 (80.32~82.29)		85 (27.60)	77.01 (72.93~81.09)		762 (26.67)	81.78 (80.80~82.77)	
≥200,000	837 (26.45)	81.51 (80.61~82.42)		92 (29.87)	77.78 (74.94~80.62)		745 (26.08)	81.97 (81.08~82.92)	
BMI			0.098			0.206			0.281
<18.50	105 (3.32)	77.90 (74.12~81.69)		15 (4.87)	74.6 (62.67~86.53)		90 (3.15)	78.46 (74.49~82.42)	
18.50~24.99	199 (63.13)	81.01 (80.35~81.66)		19 (63.96)	75.87 (73.47~78.27)		1801 (63.04)	81.57 (80.90~82.24)	
25.00~29.99	897 (28.34)	80.02 (78.98~81.06)		73 (23.70)	74.95 (71.04~78.85)		824 (28.84)	80.46 (79.39~81.54)	
≥30	165 (0.05)	77.67 (74.92~80.43)		23 (0.07)	65.13 (55.19~75.07)		142 (4.97)	79.70 (77.07~82.34)	
Wearing masks			0.0001			0.118			0.0001
Always	120 (38.17)	82.16 (81.25~83.06)		15 (49.35)	76.52 (73.67~79.37)		1056 (36.96)	82.97 (82.03~83.91)	
Often	151 (47.93)	79.87 (79.11~80.63)		12 (40.91)	74.33 (71.39~77.26)		1391 (48.69)	80.38 (79.60~81.16)	
Seldom	440 (13.90)	77.74 (76.30~79.17)		30 (9.74)	67.93 (59.46~76.40)		410 (14.35)	78.46 (77.07~79.84)	

(Continued)

Table 1 (Continued).

Variables	Total %	Health Status Scores (95% CI)	P value ^a	Long COVID			Non-long COVID		
				(N=308) %	Health Status Scores (95% CI)	P value ^b	(N=2857) %	Health Status Scores (95% CI)	P value ^c
Smoking			0.069			0.300			0.116
Smoke	623 (19.68)	80.21 (78.97~81.46)		43 (13.96)	68.40 (61.16~75.63)		580 (20.30)	81.09 (79.89~82.28)	
Quit	199 (6.29)	78.15 (75.89~80.41)		28 (9.09)	74.10 (67.69~80.52)		171 (5.99)	78.81 (76.40~81.21)	
Never	134 (42.43)	80.71 (80.08~81.33)		23 (7.95)	76.03 (73.88~78.17)		2106 (73.71)	81.23 (80.58~81.88)	
Drinking			0.008			0.091			0.016
Drink	857 (27.08)	80.26 (79.22~81.30)		76 (24.68)	76.82 (72.26~81.37)		781 (27.34)	80.60 (79.54~81.65)	
Quit	173 (5.47)	76.14 (73.35~78.92)		23 (7.47)	67.26 (59.21~75.31)		150 (5.25)	77.50 (74.58~80.42)	
Never	213 (67.46)	80.87 (80.23~81.52)		20 (67.86)	74.88 (72.55~77.20)		1926 (67.41)	81.52 (80.86~82.19)	
Sufficient sleeping			0.0001			0.0016			0.0001
Always	681 (21.52)	84.74 (83.62~85.86)		46 (14.94)	81.00 (76.55~85.45)		635 (22.23)	85.01 (83.85~86.16)	
Often	155 (49.04)	81.24 (80.52~81.96)		11 (36.04)	76.93 (73.64~80.21)		1441 (50.44)	81.57 (80.84~82.30)	
Sometimes	727 (22.97)	77.43 (76.28~78.57)		10 (34.09)	73.89 (70.83~76.94)		622 (21.77)	78.02 (69.96~75.49)	
Seldom	196 (6.2)	71.14 (68.48~73.80)		44 (14.29)	65.68 (58.83~72.53)		152 (5.32)	72.72 (69.96~75.49)	
Never	9 (0.28)	66.33 (52.08~80.59)		2 (0.65)	60.5 (20.16~100.84)		7 (0.25)	68.00 (51.77~84.23)	
Mental decompression			0.0001			0.0001			0.0001
Always	315 (0.10)	87.13 (85.45~88.81)		10 (3.25)	92.6 (86.99~98.21)		305 (10.68)	86.95 (85.23~88.68)	
Often	850 (0.27)	82.40 (81.47~83.34)		51 (16.56)	80.80 (77.02~84.58)		799 (27.97)	82.51 (81.55~83.47)	
Sometimes	1014 (32.04)	79.39 (78.46~80.31)		13 (42.21)	75.48 (72.71~78.24)		884 (30.94)	79.96 (78.99~80.94)	
Seldom	728 (23.00)	78.02 (76.84~79.19)		97 (31.49)	71.69 (67.64~75.74)		631 (22.09)	78.99 (77.80~80.18)	
Never	258 (8.15)	76.86 (74.62~79.11)		20 (6.49)	61.05 (51.66~70.44)		238 (8.33)	78.19 (75.96~80.42)	

Notes: 95% CI, confidence interval. CNY, China Yuan, The exchange rate, 1 dollar =7.005 CNY. ^aDifferences between categories within each variable in total samples. ^bDifferences between categories within each variable in long COVID samples. ^cDifferences between categories within each variable in non-long COVID samples. Bolded values indicate that in this context, P value < 0.05 and the results are meaningful.




Figure 2 Factors influencing the health status of long COVID and non-long COVID cases.

Notes: 95% CI, confidence interval. CNY, China Yuan, The exchange rate, 1 dollar=7.055 CNY. β , Adjusted for sex, age, ethnic, marriage, religion, BMI, Annual household income, education level, wearing masks, smoking, drinking, sufficient sleeping, mental decompression.

model, the 95% CI did not include 0, suggesting that the mediating effect of lifestyle was confirmed. Furthermore, the results indicated that there was a significant correlation between long COVID and health status scores ($c'=6.0486$, $P<0.001$), as well as between lifestyle and health status scores ($b=-0.6349$, $P<0.001$). This suggests that lifestyle partially mediated the relationship between long COVID and health status scores, with a mediating effect estimated to be 0.2249 (95% CI: 0.077~0.410) (see [Table S2](#)).

Results of Propensity Score Matching

A total of 281 pairs of 562 individuals were matched from the population sample of 3165. The covariates included education level, wearing masks, and sufficient sleeping (see [Table S3](#)). Based on the matched data, the differences in the health status scores between long and non-long COVID cases are shown in [Figure 4](#). After PSM matching, the health




Figure 3 The mediating effect of lifestyle scores.

Notes: a long COVID predicts lifestyle scores. b lifestyle scores predict health status scores. c long COVID predicts health status scores. c' long COVID and lifestyle scores co-predict health status scores.

status score of long COVID cases 73.43 (95% CI: 72.29~76.57) was still lower than that of non-long COVID cases 79.03 (95% CI: 77.22~80.83). Before and after matching, the difference in the health status score between the long COVID cases and non-long COVID cases was 5.59, as shown in Figure 4. (Common support domain of PSM for long COVID and non-long COVID cases is shown in Figure S2).

The above results can be used to form a possible mechanistic pathway for all the findings of this study, which can be seen in detail in Figure 5.




Figure 4 The health status scores between long COVID and non-long COVID cases pre- and post-PSM.

Notes: 95% CI, confidence interval. PSM, Propensity Score Matching.




Figure 5 The possible mechanistic pathway for the findings of this study. 95% CI, confidence interval.

Discussion

Primary Findings

Our results showed that the incidence of what we termed long COVID was at a low level. The overall health status score of the patients was 80.45, but the health status scores of long COVID cases was lower than that of non-long COVID cases. Even after propensity score matching, long COVID cases still had lower health status scores than non-long COVID cases. We found that the lifestyles of COVID-19 patients, including wearing masks, smoking, drinking, sufficient sleeping and mental decompression, was associated with their health status. Healthy lifestyles play a moderating role in improving the health status of long COVID cases.

Comparison with Previous Studies

One study showed that the prevalence of long COVID was between 10% and 15%,²⁴ and a previous study from China showed that 8.89% of infected individuals self-reported long COVID symptoms,²⁵ which is basically consistent with the results of our study (9.73%). However, other studies have reported different prevalence rates. For example, a study from Italy showed that 87.4% of people who recovered from COVID-19 reported at least one persistent symptom (such as asthma and fatigue),⁹ a study from the United States showed that 14.7% of patients had varying degrees of prolonged COVID-19 symptoms,²⁶ and a study from the Faroe Islands showed that 39% of patients developed long COVID symptoms.²⁷ This difference in research results is caused by a variety of factors, mainly because the definition of long COVID has not been formally determined, and the definitions and standards adopted by different studies are inconsistent. In addition, the different treatment statuses received by the hospitalized and non-hospitalized populations,²⁸ the difference in vaccination,²⁹ and the different immune response of the people³⁰ also lead to the inconsistent results. Moreover, our questionnaires were self-reported by infected persons, who tended to report the symptoms that boded them most, and were not professionally validated by health care providers. The patients may also attribute symptoms to their illness rather than their infection, and recall bias can occur when reporting the number of days. Therefore, it can be speculated that our survey results are conservative and may be at the lower limit of prevalence. However, we do not think this discrepancy seriously undermines the value of our findings, because most persistent symptoms are subjective, such as fatigue, and are difficult to verify objectively, even when assessed by a health care professional.

This study suggests that the effects of long COVID on the health status of infected individuals are common and persistent, which is consistent with the findings of previous studies.^{31,32} The health status of COVID-19 patients differs significantly by having long COVID, with long COVID cases having worse health status, which is similar to the findings of previous studies.^{18,33,34} The main reason is that the long-term persistent symptom cluster caused by long COVID includes respiratory sequelae, mental health disorders, cardiovascular diseases,³⁵ gastrointestinal diseases, fatigue, hypertension,³⁶ musculoskeletal pain and anemia,³⁷ which disrupt almost all aspects of daily life of infected people. In the absence of effective treatment, the persistence of sequelae reduces the quality of life of COVID-19 patients and

negatively impacts their health status.³⁸ Another reason for the worse health status of long COVID cases may be that compared with acutely infected patients, long COVID cases are not prioritized as a group that does not need to be hospitalized for dynamic monitoring, resulting in their lack of close monitoring or guidance during the recovery process.³⁹ Moreover, given the huge scale of the COVID-19 epidemic, even if the proportion of long COVID cases is very low, it will cause a significant burden of long-term illness among COVID-19 patients.⁴⁰

In this study, we found that the health status of long COVID cases was associated with smoking, drinking, and mental decompression. Compared with smokers, ex-smokers and never-smokers had better health status, which is consistent with a previous finding.³⁹ This may be because smoking is a potential risk factor for long COVID,⁴¹ which can worsen the health status of the infected person. Since SARS-CoV-2 is mainly transmitted from person to person through respiratory droplets,⁴² smoking can increase the burden on the respiratory system and lungs and the symptoms of virus infection, weaken the immune and cardiovascular systems of infected people, increase their susceptibility to various health complications,¹⁸ and have a negative impact on the health status of infected individuals.

Among long COVID cases, ex-drinkers and those who never drank did not show better health than drinkers. We consider that the most important reason for this result may be due to the infection itself because such an association was not found in non-long COVID cases. Moreover, alcohol has both advantages and disadvantages for human health. Although studies have shown that heavy drinking or frequent alcohol abuse is positively correlated with many diseases including cardiovascular and cerebrovascular diseases, which is not conducive to personal health,^{43,44} there is also evidence that compared with moderate drinkers, non-drinkers have worse health conditions.⁴⁵

The health status of non-long COVID cases is related to sufficient sleeping, and less adequate sleep was associated with poorer health status, consistent with a previous study.⁴⁶ This may be due to physical discomfort caused by symptoms during the acute infection phase⁴⁷ and the negative effects of restrictive measures including “social distancing strategy” to contain the spread of infection.

We did not find gender differences in the health status of the COVID-19 patients, but we found that people who never paid attention to mental decompression had the worst health status, regardless of whether they were long COVID cases or not. This is consistent with the results of a previous study.²² This may be due to the disruptive social changes caused during the COVID-19 epidemic, which exacerbated the mental stress of individuals, and the increase in public restrictions during the epidemic, the pressure of hospital medical care, the fear of infecting others, and the stigma increased the psychological stress of COVID-19 patients, which had a considerable negative impact on their health status.^{48,49}

Although any one of these factors can reduce health status individually, the combination of all these factors is most likely to have an additive effect,⁵⁰ suggesting a link between lifestyle and the health status of patients. Despite the current help of vaccines and pharmaceutical interventions, the actual route of infection in humans cannot be fully simulated,⁵¹ so adherence to a healthy lifestyle is crucial. The results of this study could show that a healthy lifestyle has a positive effect on the health of patients, especially mental health support for individuals who recover from COVID-19, can help regulate the harm of long COVID to the health of individuals.⁵²

Strengths and Limitations

This study has several strengths. First, this is the first national survey after the “Class B epidemic and B management” policy was adjusted in mainland China. The timeliness of the findings helps to deepen the understanding of the long-term harm of COVID-19 infection. Given the significant mediating effect, healthy lifestyles should be considered a cost-effective approach to cope with the continued low level of the COVID-19 epidemic. Second, the PSM method was used to control for confounding factors and ensure the reliability and robustness of the absolute difference in health status between long COVID and non-long COVID cases in this study, which was better than previous similar studies.

This study has several limitations. First, COVID-19 infection and health status scores were based on self-reports, which caused recollection and subjectivity bias. Second, the factors affecting the health status of COVID-19 patients are complex, which means that in addition to the positive mediating effect of lifestyle identified in this study, there are other unknown covariates to be uncovered. Finally, although PSM was used in this research to avoid the influence of some confounding variables, the confounding effect of unknown covariates was difficult to eliminate.

Conclusion

The proportion of long COVID cases was relatively low in all infected individuals. Education level, wearing masks, smoking, drinking and mental decompression were influencing factors. The health status of long COVID cases is significantly worse, and it is necessary to strengthen the follow-up of this population. Given the positive moderating role of healthy lifestyles in improving the health status of long COVID cases, healthy lifestyles including mental decompression should still be considered the most active prevention strategy at the current low epidemic level of COVID-19.

Data Sharing Statement

The data underlying this article will be shared on reasonable request to the corresponding author.

Ethics Statement

The study complied with the Declaration of Helsinki. Participants were informed of the benefits and risks of participating in this study, and they provided informed consent. All data were used merely for research purposes. This study was approved by the Life Science Ethics Review Committee of Zhengzhou University (2021-01-12-05).

Acknowledgments

All authors thank all participants involved in the study.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the Major Program of National Fund of Philosophy and Social Science of China [Grant No.21BGL222]; the Health Medical Collaborative Innovation Program of Zhengzhou [Grant No.20XTZX05015]; National Natural Science Foundation of China [Grant No.82270463].

Disclosure

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Burki T. WHO ends the COVID-19 public health emergency. *Lancet Respir Med*. 2023;(23):S2213–2600. doi:10.1016/S2213-2600(23)00217-5
2. Weekly epidemiological update on COVID-19 - 27 April 2023. Available from: <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19—27-april-2023>. Accessed March 14, 2024.
3. Zhang L, Lei J, Zhang J, et al. Undiagnosed Long COVID-19 in China among non-vaccinated individuals: identifying persistent symptoms and impacts on patients' health-related quality of life. *J Epidemiol Glob Health*. 2022;12:560–571. doi:10.1007/s44197-022-00079-9
4. Sukocheva OA, Maksoud R, Beeraka NM, et al. Analysis of post COVID-19 condition and its overlap with myalgic encephalomyelitis/chronic fatigue syndrome. *J Adv Res*. 2022;40:179–196. doi:10.1016/j.jare.2021.11.013
5. Soriano JB, Murthy S, Marshall JC, et al. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect Dis*. 2022;22:e102–e107. doi:10.1016/S1473-3099(21)00703-9
6. Alwan NA. Surveillance is underestimating the burden of the COVID-19 pandemic. *Lancet Lond Engl*. 2020;396:e24. doi:10.1016/S0140-6736(20)31823-7
7. DeVries A, Shambhu S, Sloop S, Overhage JM. One-year adverse outcomes among us adults with post-COVID-19 condition vs those without COVID-19 in a large commercial insurance database. *JAMA Health Forum*. 2023;4:e230010. doi:10.1001/jamahealthforum.2023.0010
8. Aiyegbusi OL, Hughes SE, Turner G, et al. Symptoms, complications and management of long COVID: a review. *J R Soc Med*. 2021;114:428–442. doi:10.1177/01410768211032850
9. Carfi A, Bernabei R, Landi F, & Gemelli Against COVID-19 Post-Acute Care Study Group. Persistent Symptoms in Patients After Acute COVID-19. *JAMA*. 2020;324:603–605. doi:10.1001/jama.2020.12603

10. Carvalho-Schneider C, Laurent E, Lemaigen A, et al. Follow-up of adults with noncritical COVID-19 two months after symptom onset. *Clin Microbiol Infect off Publ Eur Soc Clin Microbiol Infect Dis.* **2021**;27:258–263.
11. Tenforde MW. Symptom duration and risk factors for delayed return to usual health among outpatients with COVID-19 in a multistate health care systems network - United States, march-June 2020. *MMWR Morb Mortal Wkly Rep.* **2020**;69:993–998.
12. Halpin SJ, McIvor C, Whyatt G, et al. Postdischarge symptoms and rehabilitation needs in survivors of COVID-19 infection: a cross-sectional evaluation. *J Med Virol.* **2021**;93:1013–1022. doi:10.1002/jmv.26368
13. B L, Il K. Health-related quality of life mildly affected following COVID-19: a retrospective pre-post cohort study with a propensity score-matched control group. *J Gen Intern Med.* **2022**;37:1.
14. Av B, Skr van Z, Tc OH, Jgm R. Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study. *Lancet Lond Engl.* **2022**;400:1.
15. Hastie CE, Lowe DJ, McAuley A, et al. Outcomes among confirmed cases and a matched comparison group in the Long-COVID in Scotland study. *Nat Commun.* **2022**;13:5663. doi:10.1038/s41467-022-33415-5
16. Sneller MC, Liang CJ, Marques AR, et al. A longitudinal study of COVID-19 sequelae and immunity: baseline findings. *Ann Intern Med.* **2022**;175:969–979. doi:10.7326/M21-4905
17. Xie Y, Bowe B, Al-Aly Z. Burdens of post-acute sequelae of COVID-19 by severity of acute infection, demographics and health status. *Nat Commun.* **2021**;12:6571. doi:10.1038/s41467-021-26513-3
18. Wong MC-S, Huang J, Wong -Y-Y, et al. Epidemiology, symptomatology, and risk factors for long COVID symptoms: population-based, multicenter study. *JMIR Public Health Surveill.* **2023**;9:e42315. doi:10.2196/42315
19. M W, Elliott J, Chadeau-Hyam M, et al. Persistent COVID-19 symptoms in a community study of 606,434 people in England. *Nat Commun.* **2022**;13.
20. Ahmed GK, Khedr EM, Hamad DA, et al. Long term impact of Covid-19 infection on sleep and mental health: a cross-sectional study. *Psychiatry Res.* **2021**;305:114243. doi:10.1016/j.psychres.2021.114243
21. Salfi F, Amicucci G, Corigliano D, et al. Poor sleep quality, insomnia, and short sleep duration before infection predict long-term symptoms after COVID-19. *Brain Behav Immun.* **2023**;112:140–151. doi:10.1016/j.bbi.2023.06.010
22. Penninx BWJH, Benros ME, Klein RS, Vinkers CH. How COVID-19 shaped mental health: from infection to pandemic effects. *Nat Med.* **2022**;28:2027–2037. doi:10.1038/s41591-022-02028-2
23. Xiao Y, Chen -T-T, Liu L, Zong L. China ends its zero-COVID-19 policy: new challenges facing mental health services. *Asian J Psychiatr.* **2023**;82:103485. doi:10.1016/j.ajp.2023.103485
24. Nalbandian A, Desai AD, Wan EY. Post-COVID-19 Condition. *Annu Rev Med.* **2023**;74:55–64. doi:10.1146/annurev-med-043021-030635
25. Cai J, Lin K, Zhang H, et al. A one-year follow-up study of systematic impact of long COVID symptoms among patients post SARS-CoV-2 omicron variants infection in Shanghai, China. *Emerg Microbes Infect.* **2023**;12:2220578. doi:10.1080/22221751.2023.2220578
26. Perlis RH, Santillana M, Ognyanova K, et al. Prevalence and correlates of long COVID symptoms among US adults. *JAMA Network Open.* **2022**;5:e2238804. doi:10.1001/jamanetworkopen.2022.38804
27. Petersen MS, Kristiansen MF, Hanusson KD, et al. Prevalence of long COVID in a national cohort: longitudinal measures from disease onset until 8 months' follow-up. *Int J Infect Dis IJID off Publ Int Soc Infect Dis.* **2022**;122:437–441.
28. Kayaaslan B, Eser F, Kalem AK, et al. Post-COVID syndrome: a single-center questionnaire study on 1007 participants recovered from COVID-19. *J Med Virol.* **2021**;93:6566–6574. doi:10.1002/jmv.27198
29. Antonelli M, Penfold RS, Merino J, et al. Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study. *Lancet Infect Dis.* **2022**;22:43–55. doi:10.1016/S1473-3099(21)00460-6
30. Abdalla M, El-Arabey AA, Jiang X. What are the challenges faced by COVID-19 vaccines? *Expert Rev Vacc.* **2022**;21:5–7. doi:10.1080/14760584.2022.2008245
31. Al-Aly Z, Agarwal A, Alwan N, Luyckx VA. Long COVID: long-term health outcomes and implications for policy and research. *Nat Rev Nephrol.* **2023**;19:1–2. doi:10.1038/s41581-022-00652-2
32. El-Arabey AA, Abdalla M. Cardiovascular complications in the post-acute COVID-19 syndrome: a novel perspective down the road. *Travel Med Infect Dis.* **2022**;49:102421. doi:10.1016/j.tmaid.2022.102421
33. Valent A, Dudoignon E, Ressaire Q, Dépret F, Plaud B. Three-month quality of life in survivors of ARDS due to COVID-19: a preliminary report from a French academic centre. *Anaesth Crit Care Pain Med.* **2020**;39:740–741. doi:10.1016/j.acepm.2020.10.001
34. Moreno-Pérez O, Merino E, Leon-Ramirez J-M, et al. Post-acute COVID-19 syndrome. Incidence and risk factors: a Mediterranean cohort study. *J Infect.* **2021**;82:378–383. doi:10.1016/j.jinf.2021.01.004
35. Abdalla M, El-Arabey AA, Gai Z. Interplay between LPL and GPIHBP1 in COVID-19 patients: a possible mechanism for post-recovery cardiomyopathy. *Hum Cell.* **2023**;36:2270–2272. doi:10.1007/s13577-023-00953-z
36. M A, Aa E-A, Z G. Hypertension is still a moving target in the context of COVID-19 and post-acute COVID-19 syndrome. *J Med Virol.* **2023**;95:1.
37. Zarei M, Bose D, Nouri-Vaskeh M, et al. Long-term side effects and lingering symptoms post COVID-19 recovery. *Rev Med Virol.* **2022**;32:e2289. doi:10.1002/rmv.2289
38. Zeng N, Zhao Y-M, Yan W, et al. A systematic review and meta-analysis of long term physical and mental sequelae of COVID-19 pandemic: call for research priority and action. *Mol Psychiatry.* **2023**;28:423–433. doi:10.1038/s41380-022-01614-7
39. Subramanian A, Nirantharakumar K, Hughes S, et al. Symptoms and risk factors for long COVID in non-hospitalized adults. *Nat Med.* **2022**;28:1706–1714. doi:10.1038/s41591-022-01909-w
40. Thompson EJ, Williams DM, Walker AJ, et al. Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records. *Nat Commun.* **2022**;13:3528. doi:10.1038/s41467-022-30836-0
41. Reddy RK, Charles WN, Sklavounos A, et al. The effect of smoking on COVID-19 severity: a systematic review and meta-analysis. *J Med Virol.* **2021**;93:1045–1056. doi:10.1002/jmv.26389
42. Meyerowitz EA, Richterman A, Gandhi RT, Sax PE. Transmission of SARS-CoV-2: a review of viral, host, and environmental factors. *Ann Intern Med.* **2021**;174:69–79. doi:10.7326/M20-5008

43. Grønbaek M. The positive and negative health effects of alcohol- and the public health implications. *J interna med.* **2009**;265:407–420. doi:10.1111/j.1365-2796.2009.02082.x
44. Hendriks HFJ. Alcohol and human health: what is the evidence? *Annu Rev Food Sci Technol.* **2020**;11:1–21. doi:10.1146/annurev-food-032519-051827
45. Baumeister SE, Schumann A, Nakazono TT, et al. Alcohol consumption and out-patient services utilization by abstainers and drinkers. *Addict Abingdon Engl.* **2006**;101:1285–1291. doi:10.1111/j.1360-0443.2006.01538.x
46. Huang C, Huang L, Wang Y, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet Lond Engl.* **2023**;401:e21–e33. doi:10.1016/S0140-6736(23)00810-3
47. Zhang J, Xu D, Xie B, et al. Poor-sleep is associated with slow recovery from lymphopenia and an increased need for ICU care in hospitalized patients with COVID-19: a retrospective cohort study. *Brain Behav Immun.* **2020**;88:50–58. doi:10.1016/j.bbi.2020.05.075
48. Shi L, Lu Z-A, Que J-Y, et al. Prevalence of and risk factors associated with mental health symptoms among the general population in China during the coronavirus disease 2019 pandemic. *JAMA Network Open.* **2020**;3:e2014053. doi:10.1001/jamanetworkopen.2020.14053
49. Yuan K, Gong Y-M, Liu L, et al. Prevalence of xlnl1 stress disorder after infectious disease pandemics in the twenty-first century, including COVID-19: a meta-analysis and systematic review. *Mol Psychiatry.* **2021**;26:4982–4998. doi:10.1038/s41380-021-01036-x
50. Malik P, Patel K, Pinto C, et al. Post-acute COVID-19 syndrome (PCS) and health-related quality of life (HRQoL)-A systematic review and meta-analysis. *J Med Virol.* **2022**;94:253–262. doi:10.1002/jmv.27309
51. El-Arabey AA, Abdalla M. In the face of the future, what do we learn from COVID-19? *Hum Vaccines Immunother.* **2021**;17:4119–4120. doi:10.1080/21645515.2021.1963174
52. Wang S, Li Y, Yue Y, et al. Adherence to healthy lifestyle prior to infection and risk of post-COVID-19 condition. *JAMA Intern Med.* **2023**;183:232–241. doi:10.1001/jamainternmed.2022.6555

Risk Management and Healthcare Policy

Dovepress

Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/risk-management-and-healthcare-policy-journal>