

# Association Between TCBI (Triglycerides, Total Cholesterol, and Body Weight Index) and Stroke-Associated Pneumonia in Acute Ischemic Stroke Patients

Yufeng Liu<sup>1,\*</sup>, Yan Chen<sup>2,\*</sup>, Zhongwen Zhi<sup>1,\*</sup>, Ping Wang<sup>1</sup>, Mengchao Wang<sup>1</sup>, Qian Li<sup>1</sup>, Yuqian Wang<sup>1</sup>, Liandong Zhao<sup>1</sup>, Chun Chen<sup>1</sup>

<sup>1</sup>Department of Neurology, The Affiliated Huai'an Hospital of Xuzhou Medical University and The Second People's Hospital of Huai'an, Huai'an, Jiangsu, 223002, People's Republic of China; <sup>2</sup>Department of Neurological Medicine, Siyang Hospital of Traditional Chinese Medicine, Siyang, Jiangsu, 223700, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Chun Chen; Liandong Zhao, Department of Neurology, The Affiliated Huai'an Hospital of Xuzhou Medical University and The Second People's Hospital of Huai'an, No. 62, South Huaihai Road, Huai'an, Jiangsu, 223002, People's Republic of China, Email hongzecc@163.com; zld202109@163.com

**Purpose:** Stroke-associated pneumonia (SAP) usually complicates stroke and is linked to adverse prognoses. Triglycerides, total cholesterol, and body weight index (TCBI) is a new and simple calculated nutrition index. This study seeks to investigate the association between TCBI and SAP incidence, along with its predictive value.

**Patients and Methods:** Nine hundred and sixty-two patients with acute ischemic stroke were divided into SAP group and Non-SAP group. The TCBI was divided into three layers: T1, TCBI < 948.33; T2, TCBI 948.33–1647.15; T3, TCBI > 1647.15. Binary Logistic regression analysis was used to determine the relationship between TCBI levels and the incidence of SAP. Furthermore, restricted cubic splines (RCS) analysis was utilized to evaluate the influence of TCBI on the risk of SAP.

**Results:** TCBI in the SAP group was markedly lower compared to that in the Non-SAP group ( $P < 0.001$ ). The Logistic regression model revealed that, using T3 layer as the reference, T1 layer had the highest risk for SAP prevalence (OR = 2.962, 95% CI: 1.600–5.485,  $P = 0.001$ ), with confounding factors being controlled. The RCS model found that TCBI had a linear relationship with SAP ( $P$  for nonlinear = 0.490,  $P$  for overall = 0.004). Moreover, incorporating TCBI into the A<sup>2</sup>DS<sup>2</sup> (Age, atrial fibrillation, dysphagia, sex, and severity) model substantially enhanced the initial model's predictive accuracy.

**Conclusion:** Low TCBI was associated with a higher risk of SAP. In clinical practice, TCBI has shown predictive value for SAP, contributing to early intervention and treatment of SAP.

**Keywords:** acute ischemic stroke, stroke-associated pneumonia, nutritional status, TCBI

## Introduction

Stroke stands as a primary global cause of mortality and chronic disability, with clinical research indicating that the majority of stroke-related fatalities are not directly due to the stroke itself, but rather its complications. Stroke-associated pneumonia (SAP) constitutes a frequent complication among acute ischemic stroke (AIS) patients, with an incidence rate ranging from 7% to 38%.<sup>1,2</sup> SAP detrimentally impacts patient neurological recovery, significantly alters prognosis, and is a major mortality factor in stroke cases.<sup>3,4</sup> Research has associated factors such as age, dysphagia, severity of stroke, inflammatory response, and stroke-induced immunosuppressive syndrome with an increased risk of SAP.<sup>5,6</sup> The early detection of potential patients holds clinical significance for implementing surveillance and prevention measures. However, relying on a single biomarker presents limited predictive capability regarding disease outcome. Several risk

stratification scoring systems, including the Friedland pneumonia prediction score, A<sup>2</sup>DS<sup>2</sup> (Age, atrial fibrillation, dysphagia, sex, and severity) score, Kwon pneumonia score, and Preventive Antibiotics in Stroke Study pneumonia Rule, etc.,<sup>7–9</sup> have been developed to forecast the incidence of SAP events. The A<sup>2</sup>DS<sup>2</sup> score, determined by factors including age, dysphagia, gender, atrial fibrillation, and stroke severity, is the prevalent risk assessment tool in clinical settings, corroborated across various external cohorts.<sup>10–12</sup>

Studies have shown a correlation between nutritional status and stroke-related adverse outcomes and complications.<sup>13,14</sup> Patients exhibiting malnutrition at baseline exhibited a higher propensity for developing SAP.<sup>15</sup> Therefore, insights into malnutrition assessment could facilitate the early detection of AIS patients with a high risk of SAP. TCBI (triglycerides, total cholesterol, and body weight index) is a new simple calculated nutritional index,<sup>16</sup> encompassing serum triglycerides (TG), total cholesterol (TC), and body weight (BW). Previous research has demonstrated that TCBI serves as both a nutritional and prognostic marker in patients suffering from coronary artery disease, acute decompensated heart failure, and hemodynamic instability within intensive care settings.<sup>16–18</sup> In the Chinese population with H-type hypertension, there exists a negative correlation between TCBI and stroke incidence.<sup>19</sup> In patients with AIS, TCBI has been found to be associated with adverse outcomes.<sup>20</sup>

Currently, research on the connection between TCBI and SAP is still lacking. Therefore, this study hypothesizes that a lower TCBI is associated with an increased incidence of SAP events in patients with AIS. If so, we will evaluate whether TCBI is valuable in predicting SAP incidence and improving the predictive effectiveness of traditional A<sup>2</sup>DS<sup>2</sup> model.

## Materials and Methods

### Study Design and Population

Patients with AIS who were admitted to the Department of Neurology at the Affiliated Huai'an Hospital of Xuzhou Medical University between March 2021 and May 2022 were prospectively enrolled in the study. This study was conducted in strict accordance with the ethical guidelines of the Declaration of Helsinki and was approved by the Ethics Committee of Affiliated Huai'an Hospital of Xuzhou Medical University (Approval number: HEYLL202137). Written informed consent was obtained from each participant.

The inclusion criteria were as follows: (1) Confirmed by computed tomography or magnetic resonance imaging to meet the diagnostic criteria for AIS; (2) The time from onset to admission is less than 7 days; (3) Age  $\geq 18$  years old.

The exclusion criteria were as follows: (1) Transient ischemic attack; (2) Active infection or prophylactic use of antibiotics within 2 weeks prior to admission; (3) Combined with a history of central nervous system diseases, such as brain trauma, cerebral hemorrhage or hydrocephalus; (4) The presence of swallowing dysfunction before stroke; (5) Patients with severe liver and kidney dysfunction (Child-Pugh C or serum creatinine  $>177$   $\mu\text{mol/L}$ ); (6) Patients with diseases of the blood system, cancer or receiving immunosuppressive therapy; (7) Clinical data are incomplete.

### Data Collection

General data of stroke patients were collected, including demographic characteristics: gender, age, height and weight. Previous medical histories include: hypertension, diabetes, coronary heart disease, atrial fibrillation, smoking, drinking and a history of stroke. Stroke-related data: National Institutes of Health Stroke Scale (NIHSS) at admission and stroke subtypes (based on TOAST classification). Treatment received during hospitalization: endovascular treatment, intravenous thrombolysis therapy and drug use. Laboratory data include: hemoglobin, albumin, total cholesterol (TC), triglycerides (TG), low-density lipoprotein cholesterol (LDL-C) and high-density lipoprotein cholesterol (HDL-C). Swallowing function data collection: The swallowing function of patients was evaluated by modified water swallowing test within 24 hours after admission. A<sup>2</sup>DS<sup>2</sup> scoring model score collection: This is a model for predicting clinical risk of SAP based on factors such as age, dysphagia, male, atrial fibrillation, and stroke severity (NIHSS) on a scale of 0 to 10. The A<sup>2</sup>DS<sup>2</sup> score has been validated in multiple external cohorts.<sup>10–12</sup>

## Diagnosis of SAP

According to the revised standards of the Centers for Disease Control and Prevention, patients diagnosed with SAP are those who experience lower respiratory tract infections within the first seven days following an acute stroke.<sup>21</sup>

## Assessment of TCBI

Fasting blood samples were taken within 24 hours after admission for TG, TC and other indicators. The formula includes TG, TC and BW,  $TCBI = TG \text{ (mg/dL)} \times TC \text{ (mg/dL)} \times BW \text{ (kg)} / 1000$ .<sup>16</sup>

## Statistical Analysis

Data conforming to normal distribution were expressed as mean  $\pm$  standard deviation (SD), two independent sample *t*-test was used for inter-group comparison, median and quartile [M (P25-P75)] description was used for non-normal distribution data, and non-parametric rank sum test (Mann–Whitney *U*-test) was used for inter-group comparison. Count data were represented as frequency and percentage (n,%), with the Pearson Chi-square test applied for group comparisons. To assess differences among multiple groups, we conducted either a one-way ANOVA or the Kruskal–Wallis test for parametric and non-parametric data, respectively. Pairwise comparisons were performed on variables that differed between groups, with *P*-values adjusted using the Bonferroni correction method. Due to the severely skewed distribution of TCBI, Log transformation was used in part of the statistical analysis. According to the sample size of this study and TCBI, the patients were divided into three groups on average (Tertile 1 < 948.33, Tertile 2 948.33–1647.15, Tertile 3 > 1647.15). A Binary Logistic regression model was used to examine the relationship between TCBI and SAP. Variables with *P* less than 0.05 in univariate logistic regression analysis were included in multivariate regression analysis. Adjusted variables included age, coronary heart disease, atrial fibrillation, TOAST classification, NIHSS at admission, dysphagia, endovascular treatment, anticoagulant drug use, hemoglobin and albumin. Restricted cubic spline models were used to provide and explore the shape of the association between TCBI and SAP, fitting a restricted cubic spline function with 4 knots (at the 5th, 35th, 65th, and 95th percentiles). The Receiver operating characteristic (ROC) curve was used to estimate the optimal threshold for biomarkers to predict the occurrence of SAP in AIS patients. Area Under the Receiver Operating Characteristic Curve (AUROC), sensitivity, specificity, positive prediction rate and negative prediction rate were calculated. The optimal threshold was identified using the Youden index. The AUROC efficacy of TCBI was compared with other biomarker values by Z-test.


SPSS 25.0 (IBM Corporation, IL, USA), MedCalc 20.0 (MedCalc Software LTD., Ostend, Belgium), R Programming Language 4.3.2 (Vienna, Austria), and GraphPad Prism 9 (GraphPad Software, USA) were used for all study data analysis and image processing. All *P* values were two-tailed, and statistical significance was defined by the *P* value < 0.05.

## Results

### Baseline Characteristics

This study screened 1075 AIS patients hospitalized within seven days of symptom onset for analysis. Exclusions were 21 patients with transient ischemic attacks, 42 with fever or active infections pre-admission, 25 with severe hepatic and renal insufficiency, 9 with hematological diseases and tumors, and 16 with incomplete data. Ultimately, 962 patients were enrolled, with 128 developing SAP, corresponding to an incidence rate of 13.3% (Figure 1). Of the total cohort, 369 (38.4%) were females. The median age was 68 (interquartile range, 59–77). The median TCBI was 1259.03 (interquartile range, 812.18 to 1942.30).

As shown in Table 1, patients in the SAP group were older, had a lighter body weight, had higher NIHSS at admission, had higher rates of coronary heart disease, atrial fibrillation, cardioembolism and dysphagia, and were more likely to receive endovascular treatment and anticoagulant drug use. Compared with Non-SAP group, SAP group patients had lower levels of hemoglobin, albumin, TC and TG, and higher A<sup>2</sup>DS<sup>2</sup> score. The TCBI in SAP group was significantly lower than that in Non-SAP group (All *P* < 0.05). In this study, there were no significant differences between the two groups in gender, height, hypertension, diabetes, smoking, drinking, history of stroke, intravenous thrombolysis, antiplatelet drug use, lipid-lowering drug use, LDL-C and HDL-C (All *P* > 0.05) (Table 1).



**Figure 1** Patient flowchart.

**Abbreviations:** SAP, Stroke-associated pneumonia.

## Association between TCBI and stroke-Associated Pneumonia

According to the quantile of TCBI, all patients were divided into three groups: Tertile 1 < 948.33, Tertile 2 948.33–1647.15, and Tertile 3 > 1647.15. In the different TCBI level groups, there was a significant difference in the proportion of SAP ( $P < 0.001$ ) (Table 2). In addition, Table 2 shows significant differences in age, height, body weight, hypertension, coronary heart disease, atrial fibrillation, TOAST classification, NIHSS at admission, endovascular treatment, intravenous thrombolysis, antiplatelet drug use, anticoagulant drug use, hemoglobin, albumin, TC, TG, LDL-C, HDL-C, and A<sup>2</sup>DS<sup>2</sup> score among the three groups (All  $P < 0.05$ ) (Table 2).

**Table 1** Comparison of Baseline Data Between SAP Group and Non-SAP Group

Variables	All Patients n = 962	Non-SAP n = 834	SAP n = 128	P value
<b>Demographic information</b>				
Female, n (%)	369 (38.4)	316 (37.9)	53 (41.4)	0.446
Age (years), median (IQR)	68 (59–77)	67 (58–76)	73 (64–81)	<0.001
Height (m), mean ± SD	1.65±0.08	1.65±0.08	1.65±0.07	0.488
Body weight (kg), mean ± SD	68.08±11.22	68.73±11.18	63.88±10.61	<0.001
Hypertension, n (%)	795 (82.6)	691 (82.9)	104 (81.3)	0.656
Diabetes, n (%)	331 (34.4)	285 (34.2)	46 (35.9)	0.696
Coronary heart disease, n (%)	144 (15.0)	117 (14.0)	27 (21.1)	0.037
Atrial fibrillation, n (%)	99 (10.3)	70 (8.4)	29 (22.7)	<0.001
Smoking, n (%)	233 (24.2)	199 (23.9)	34 (26.6)	0.507
Drinking, n (%)	177 (18.4)	157 (18.8)	20 (15.6)	0.384
History of stroke, n (%)	178 (18.5)	148 (17.7)	30 (23.4)	0.123

(Continued)

Table 1 (Continued).

Variables	All Patients n = 962	Non-SAP n = 834	SAP n = 128	P value
<b>Clinical features</b>				
TOAST classification, n (%)				<0.001
LAA	636 (66.1)	545 (65.3)	91 (71.1)	
CE	100 (10.4)	74 (8.9)	26 (20.3)	
SAO	226 (23.5)	215 (25.8)	11 (8.6)	
NIHSS at admission (score), median (IQR)	2 (1–5)	2 (1–4)	9 (3–17)	<0.001
Dysphagia, n (%)	135 (14.0)	87 (10.4)	48 (37.5)	<0.001
<b>Treatment, n (%)</b>				
Endovascular treatment	55 (5.7)	24 (2.9)	31 (24.2)	<0.001
Intravenous thrombolysis	127 (13.2)	105 (12.6)	22 (17.2)	0.153
Antiplatelet drug use	884 (91.9)	771 (92.4)	113 (88.3)	0.108
Anticoagulant drug use	116 (12.1)	91 (10.9)	25 (19.5)	0.005
Lipid-lowering drug use	946 (98.3)	821 (98.4)	125 (97.7)	0.518
<b>Laboratory data</b>				
Hemoglobin (g/L), median (IQR)	141 (129–153)	142 (130–153)	137 (122–148)	0.002
Albumin (g/L), mean $\pm$ SD	39.55 $\pm$ 4.28	39.78 $\pm$ 4.19	38.07 $\pm$ 4.60	<0.001
TC (mmol/L), median (IQR)	4.36 (3.61–5.02)	4.40 (3.67–5.06)	4.10 (3.31–4.69)	<0.001
TG (mmol/L), median (IQR)	1.25 (0.94–1.78)	1.31 (0.95–1.84)	1.08 (0.82–1.37)	<0.001
LDL-C (mmol/L), mean $\pm$ SD	2.66 $\pm$ 0.89	2.68 $\pm$ 0.87	2.53 $\pm$ 1.04	0.076
HDL-C (mmol/L), median (IQR)	1.12 (1.02–1.32)	1.12 (1.02–1.32)	1.13 (1.02–1.36)	0.761
TCBI, median (IQR)	1259.03 (812.18–1942.30)	1322.76 (859.93–2046.42)	898.54 (606.86–1396.16)	<0.001
A <sup>2</sup> DS <sup>2</sup> (score), median (IQR)	1 (1–3)	1 (1–2)	5 (3–7)	<0.001

**Abbreviations:** SD, standard deviation; IQR, interquartile range; SAP, Stroke-associated pneumonia; LAA, Large artery atherosclerosis; CE, Cardioembolism; SAO, Small-artery occlusion; NIHSS, National Institutes of Health Stroke Scale; TC, Total cholesterol; TG, Triglycerides; LDL-C, Low-density lipoprotein cholesterol; HDL-C, High-density lipoprotein cholesterol; TCBI, Triglycerides, total cholesterol, and body weight index.

Table 2 Baseline Characteristics of Patients According to TCBI Tertiles

Variables	TCBI Tertiles			P value
	T1 (<948.33) n=320	T2 (948.33–1647.15) n=321	T3 (>1647.15) n=321	
<b>Demographic information</b>				
Female, n (%)	134 (41.9)	124 (38.6)	111 (34.6)	0.163
Age (years), median (IQR)	73 (64–81)	66 (59–76)	64 (55–72)	<0.001
Height (m), mean $\pm$ SD	1.64 $\pm$ 0.08	1.65 $\pm$ 0.08	1.66 $\pm$ 0.07	<0.001
Body weight (kg), mean $\pm$ SD	63.50 $\pm$ 9.72	68.73 $\pm$ 11.02	72.01 $\pm$ 11.21	<0.001
Hypertension, n (%)	251 (78.4)	267 (83.2)	277 (86.3)	0.030
Diabetes, n (%)	99 (30.9)	108 (33.6)	124 (38.6)	0.115
Coronary heart disease, n (%)	68 (21.3)	51 (15.9)	25 (7.8)	<0.001
Atrial fibrillation, n (%)	59 (18.4)	25 (7.8)	15 (4.7)	<0.001
Smoking, n (%)	69 (21.6)	84 (26.2)	80 (24.9)	0.371
Drinking, n (%)	49 (15.3)	65 (20.2)	63 (19.6)	0.214
History of stroke, n (%)	67 (20.9)	54 (16.8)	57 (17.8)	0.372
<b>Clinical features</b>				
TOAST classification, n (%)				<0.001
LAA	201 (62.8)	206 (64.2)	229 (71.3)	

(Continued)

**Table 2** (Continued).

Variables	TCBI Tertiles			P value
	T1 (<948.33) n=320	T2 (948.33–1647.15) n=321	T3 (>1647.15) n=321	
CE	56 (17.5)	24 (7.5)	20 (6.2)	0.001
SAO	63 (19.7)	91 (28.3)	72 (22.4)	
NIHSS at admission (score), median (IQR)	3 (1–7)	2 (1–4)	2 (1–4)	
Dysphagia, n (%)	49 (15.3)	41 (12.8)	45 (14.0)	0.651
<b>Treatment, n (%)</b>				
Endovascular treatment	24 (7.5)	23 (7.2)	8 (2.5)	0.009
Intravenous thrombolysis	49 (15.3)	28 (8.7)	50 (15.6)	0.015
Antiplatelet drug use	275 (85.9)	302 (94.1)	307 (95.6)	<0.001
Anticoagulant drug use	61 (19.1)	32 (10.0)	23 (7.2)	<0.001
Lipid-lowering drug use	316 (98.8)	314 (97.8)	316 (98.4)	0.644
<b>Laboratory data</b>				
Hemoglobin (g/L), median (IQR)	135 (121–147)	142 (129.5–155)	145 (135–156)	<0.001
Albumin (g/L), mean $\pm$ SD	38.70 $\pm$ 4.46	39.81 $\pm$ 4.16	40.14 $\pm$ 4.09	<0.001
TC (mmol/L), median (IQR)	3.60 (3.04–4.20)	4.36 (3.73–4.97)	5.00 (4.49–5.67)	<0.001
TG (mmol/L), median (IQR)	0.84 (0.71–1.02)	1.24 (1.10–1.50)	2.04 (1.68–2.86)	<0.001
LDL-C (mmol/L), mean $\pm$ SD	2.20 (1.62–2.64)	2.70 (2.20–3.19)	3.02 (2.63–3.64)	<0.001
HDL-C (mmol/L), median (IQR)	1.16 (1.05–1.38)	1.14 (1.02–1.35)	1.07 (0.99–1.22)	<0.001
A <sup>2</sup> DS <sup>2</sup> (score), median (IQR)	2 (1–4)	1 (1–3)	1 (1–3)	<0.001
SAP, n (%)	69 (21.6)	39 (12.1)	20 (6.2)	<0.001


**Abbreviations:** SD, standard deviation; IQR, interquartile range; TCBI, Triglycerides, total cholesterol, and body weight index; LAA, Large artery atherosclerosis; CE, Cardioembolism; SAO, Small-artery occlusion; NIHSS, National Institutes of Health Stroke Scale; TC, Total cholesterol; TG, Triglycerides; LDL-C, Low-density lipoprotein cholesterol; HDL-C, High-density lipoprotein cholesterol; SAP, Stroke-associated pneumonia.

As indicated in the forest plot in [Figure 2](#), with the highest tertile serving as the reference after adjusting for the confounders such as age, coronary heart disease, atrial fibrillation, TOAST classification, NIHSS at admission, dysphagia, endovascular treatment, anticoagulant drug use, hemoglobin and albumin, the lowest tertile of TCBI (<948.33) was substantially related to a greater risk of SAP (odds ratio [OR] = 2.962, 95% confidence interval [CI]: 1.600–5.485,  $P$  = 0.001), while the middle tertile of TCBI was not significantly associated with the prevalence of SAP (OR = 1.858, 95% CI = 0.970–3.559,  $P$  = 0.062). When taken as a continuous variable, TCBI was still related to SAP in the logistic regression analysis, with the confounders being controlled (unadjusted: OR = 0.127, 95% CI = 0.062–0.260,  $P$  < 0.001; adjusted: OR = 0.223, 95% CI = 0.094–0.528,  $P$  = 0.001). [Figure 2](#) also showed that age, NIHSS at admission, dysphagia, endovascular treatment, and albumin were also independent predictors of SAP (All  $P$  < 0.05) ([Figure 2](#)). In addition, the RCS curve revealed a linear relationship between TCBI and SAP, after adjusting for confounding variables ( $P$  for nonlinear = 0.490,  $P$  for overall = 0.004) ([Figure 3](#)).

## Predictive Value of TCBI for Stroke-Associated Pneumonia

Results from the ROC analysis indicated that the optimal cutoff value of TCBI for predicting SAP was 1112.37, with an AUROC of 0.664 (95% CI = 0.633–0.694). The sensitivity and specificity were 65.62% and 60.91%, respectively. The positive predictive rate and the negative predictive rate were 20.49% and 92.03%, respectively. Compared to other biomarkers, TCBI demonstrated superior predictive capabilities for SAP over TC (AUROC = 0.599, 95% CI = 0.568–0.630; TCBI vs. TC,  $P$  = 0.002) and TG (AUROC = 0.628, 95% CI = 0.597–0.659; TCBI vs. TG,  $P$  = 0.003). However, there was no significant difference when compared with BW (AUROC = 0.623, 95% CI = 0.591–0.654; TCBI vs. BW,  $P$  = 0.206) ([Table 3](#) and [Figure 4](#)).





**Figure 2** Forest plot of odds ratios for SAP.


**Abbreviations:** LAA, Large artery atherosclerosis; CE, Cardioembolism; SAO, Small-artery occlusion; NIHSS, National Institutes of Health Stroke Scale; TCBI, Triglycerides, total cholesterol, and body weight index.

The AUROC of A<sup>2</sup>DS<sup>2</sup> model was 0.819 (95% CI = 0.794–0.843). The modified A<sup>2</sup>DS<sup>2</sup> model combining A<sup>2</sup>DS<sup>2</sup> model with TCBI has an AUROC of 0.838 (95% CI = 0.813–0.860). The difference in prediction efficiency between the modified A<sup>2</sup>DS<sup>2</sup> model and the original A<sup>2</sup>DS<sup>2</sup> model was statistically significant ( $P = 0.004$ ) (Table 3 and Figure 4).

## Discussion

To our knowledge, this study is the first to investigate the relationship between TCBI and SAP. Following adjustment for potential confounders, results indicated that lower baseline TCBI levels were associated with an increased risk of SAP occurrence. There was a linear dose–response relationship between TCBI and SAP. Additionally, our results again confirmed the predictive efficacy of the A<sup>2</sup>DS<sup>2</sup> model for SAP. Moreover, incorporating TCBI with the conventional A<sup>2</sup>DS<sup>2</sup> model significantly enhanced its accuracy, suggesting TCBI's potential as a valuable biomarker for SAP in clinical settings.

Prior research indicates a high prevalence of malnutrition among stroke patients. Individuals with AIS presenting normal nutritional levels can still exhibit malnutrition due to factors such as advanced age, dysphagia, immunosuppression, cognitive impairments, and restricted mobility. Additionally, acute-phase malnutrition is believed to correlate with a diminished functional prognosis and an elevated mortality rate from all causes.<sup>22</sup> New evidence shows that initiating enteral nutrition early and integrating probiotic therapy enhances nutritional status, diminishes systemic inflammation, bolsters immune function, and decreases the incidence of infectious complications among stroke patients.<sup>23</sup> Hence, assessing nutritional status suitably is crucial for effective risk stratification and treatment planning. Several nutritional assessment instruments, such as the Nutritional Risk Screening 2002 and the Malnutrition Universal Screening Tool, have been employed to identify malnutrition in AIS patients.<sup>24</sup> These indicators consist of subjective parameters, for



**Figure 3** Association of TCBI with risk of stroke-associated pneumonia. Odds ratios and 95% confidence intervals derived from restricted cubic spline regression, with knots placed at the 5th, 35th, 65th, and 95th percentiles of the distribution of Log TCBI. The reference point is the median of Log TCBI. Odds ratios were adjusted for the same variables as model in [Figure 2](#).

instance, recent weight changes, which are arbitrary. In addition, certain AIS patients manifesting symptoms like decreased consciousness and aphasia are unable to participate in scale assessments, rendering these instruments unsuitable for universal application in the AIS population. Consequently, there is a necessity for more objective and broadly applicable methodologies to evaluate patients' nutritional status.


The TCBI is a novel and simple nutritional tool that reflects the intrinsic metabolism of fat, cholesterol, carbohydrates, and sugars, primarily by simple multiplication of body weight and variables that reflect lipid metabolism.<sup>17</sup> The TCBI was initially developed to evaluate the nutritional status of individuals afflicted with atherosclerotic cardiovascular disease. It has been observed that populations suffering from coronary artery disease with lower TCBI scores experience increased rates of all-cause mortality, cardiovascular mortality, and cancer mortality.<sup>16</sup> Recent studies indicate that TCBI could serve as a significant prognostic nutrition-related marker in critically ill cardiac patients needing mechanical circulatory support, experiencing acute decompensated heart failure, suffering from acute myocardial infarction, or undergoing transcatheter aortic valve replacement.<sup>17,18,25,26</sup>

**Table 3** Comparison of Predictive Power of TCBI Vs Other Indicators in the Prediction of SAP

Variables	AUROC (95% CI)	Cutoff Value	Youden Index	Sensibility (%)	Specificity (%)	Positive Prediction Rate (%)	Negative Prediction Rate (%)	P value
TCBI	0.664 (0.633–0.694)	1112.37	0.2654	65.62	60.91	20.49	92.03	Reference
BW	0.623 (0.591–0.654)	65.30	0.1964	64.84	54.80	18.04	91.04	0.206
TC	0.599 (0.568–0.630)	4.42	0.1767	68.75	48.92	17.05	89.85	0.002
TG	0.628 (0.597–0.659)	1.37	0.2321	76.56	46.64	18.05	92.84	0.003
A <sup>2</sup> DS <sup>2</sup>	0.819 (0.794–0.843)	2	0.5108	75.78	75.30	32.01	95.30	<0.001
A <sup>2</sup> DS <sup>2</sup> +TCBI	0.838 (0.813–0.860)	0.1418	0.5485	73.44	81.41	37.75	95.23	<0.001

**Abbreviations:** SAP, stroke-associated pneumonia; AUROC, Area Under the Receiver Operating Characteristic Curve; TCBI, Triglycerides, total cholesterol, and body weight index; BW, Body weight; TC, Total cholesterol; TG, Triglyceride.





**Figure 4** Comparison of AUROC of TCBI with other biomarkers for predicting SAP.

**Abbreviations:** BW, Body weight; TC, Total cholesterol; TG, Triglycerides; TCBI, Triglycerides, cholesterol, body weight index; SAP, Stroke-associated pneumonia.

In a Chinese cohort with H-type hypertension, a higher stroke incidence was observed among individuals with low TCBI levels. Multivariable-adjusted analyses indicated a negative association between TCBI and stroke prevalence.<sup>19</sup> This may be attributed to the fact that malnutrition can induce inflammation and oxidative stress.<sup>27,28</sup> An investigation involving 9708 participants indicates that TCBI levels correlate with both short-term and long-term functional disability, as well as all-cause mortality, following a stroke.<sup>20</sup> Despite the growing interest, research on the association between TCBI and stroke remains sparse. Consequently, we initiated a prospective study to assess the relationship between TCBI and SAP. The findings supported our preliminary hypothesis: patients exhibiting lower TCBI levels exhibited a higher propensity for developing SAP, a relationship that persisted as significant even after adjusting for potential confounders. Numerous investigations have demonstrated a correlation between early malnutrition and an elevated risk of SAP. For the prompt identification of patients at high risk, objective nutritional assessment tools, including the Geriatric Nutrition Risk Index (GNRI), the Controlling Nutritional Status (CONUT), and the Prognostic Nutritional Index (PNI), are recommended.<sup>15,29</sup> However, these three scoring indicators are not widely used in clinical practice due to the complexity of calculation methods and parameters. The advantages of TCBI are easy to obtain, simple to calculate, objective and unrestricted in AIS population. Furthermore, TCBI showed significant predictive value for SAP. Our findings support the use of TCBI as a simple nutritional marker for the surveillance of SAP occurrences.

The underlying mechanism between lower TCBI and increased SAP risk is not fully understood. Weight loss is regarded as a risk factor for SAP.<sup>6</sup> Body weight and body mass index (BMI) commonly quantify muscle and fat mass, and high BMI (overweight or obese) has been well-acknowledged as a risk factor for AIS.<sup>30</sup> Nevertheless, research has demonstrated a linear correlation between elevated BMI and improved survival rates in AIS patients,<sup>31,32</sup> a phenomenon termed the “obesity paradox”. Therefore, relying solely on body weight or BMI to forecast adverse stroke outcomes is contentious. Circulating lipid levels are indicators of nutritional status and inflammation, TG and TC can reflect the nutritional status of patients to a certain extent.<sup>33,34</sup> Prior research indicates an association between dual X-ray absorptiometry (the gold standard for assessing nutritional status) and GNRI.<sup>35,36</sup> Furthermore, multiple evidence suggests a moderately positive correlation between TCBI and GNRI,<sup>16–18</sup> indicating that TCBI may serve as an indirect marker of nutritional status. Hence, interpreting the relationship between TCBI and SAP incidence through the lens of malnutrition is justifiable. The association between malnutrition and immune function is well-documented. Malnutrition may lead to reduced lymphocyte production, thereby impairing immune function,<sup>37,38</sup> and immune dysfunction significantly contributes to the pathology of SAP events. Malnutrition may also contribute to both the development and worsening of SAP by inducing inflammation and oxidative

stress.<sup>27,28</sup> Additionally, based on the correlation between inflammatory indicators and nutritional scores, malnutrition is thought to reflect a high inflammatory and metabolic state of the body, which also contributes to the development of SAP.<sup>15</sup>

However, our study has several limitations. Firstly, considering the results from a single-center cohort, a larger sample size from prospective registries across multiple regions and countries is necessary to generalize the findings. Secondly, TCBI was only measured at admission, preventing the assessment of its dynamic changes and their correlation with SAP throughout the hospitalization period. Thirdly, the study did not include an evaluation of other nutritional indices like CONUT, GNRI, or PNI, nor did it explore the comparative relevance and applicability of TCBI against these tools. Lastly, the absence of data on functional outcomes and mortality precluded analysis of the relationship between TCBI, SAP, and stroke outcomes, which needs to be clarified in further studies.

## Conclusion

In Conclusion, our study demonstrates that low TCBI was associated with a higher risk of SAP. T1 layer (TCBI < 948.33) had the highest risk for SAP prevalence. Moreover, integrating TCBI into traditional A<sup>2</sup>DS<sup>2</sup> model significantly improves its predictive performance. TCBI measured for the first time after admission can be used for early identification of patients at high risk of SAP. Timely attention and effective intervention may improve the prognosis of patients.

## Acknowledgments

The authors thank the study participants and their relatives as well as the clinical staff for their support and contributions to this study.

## Funding

Huai'an municipal Science and Technology Program Project (HAB202322).

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Kishore AK, Vail A, Chamorro A, et al. How is pneumonia diagnosed in clinical stroke research? A systematic review and meta-analysis. *Stroke*. 2015;46(5):1202–1209. doi:10.1161/STROKEAHA.114.007843
2. Badve MS, Zhou Z, van de Beek D, et al. Frequency of post-stroke pneumonia: systematic review and meta-analysis of observational studies. *Int J Stroke*. 2019;14(2):125–136. doi:10.1177/1747493018806196
3. de Montmollin E, Ruckly S, Schwebel C, et al. Pneumonia in acute ischemic stroke patients requiring invasive ventilation: impact on short and long-term outcomes. *J Infect*. 2019;79(3):220–227. doi:10.1016/j.jinf.2019.06.012
4. Li J, Zhang P, Wu S, et al. Stroke-related complications in large hemisphere infarction: incidence and influence on unfavorable outcome. *Ther Adv Neurol Disord*. 2019;12:1756286419873264. doi:10.1177/1756286419873264
5. Eltringham SA, Kilner K, Gee M, et al. Impact of dysphagia assessment and management on risk of stroke-associated pneumonia: a systematic review. *Cerebrovasc Dis*. 2018;46(3–4):99–107. doi:10.1159/000492730
6. Patel UK, Kodumuri N, Dave M, et al. Stroke-associated pneumonia: a retrospective study of risk factors and outcomes. *Neurologist*. 2020;25(3):39–48. doi:10.1097/NRL.0000000000000269
7. Gong S, Zhou Z, Zhou M, et al. Validation of risk scoring models for predicting stroke-associated pneumonia in patients with ischaemic stroke. *Stroke Vasc Neurol*. 2016;1(3):122–126. doi:10.1136/svn-2016-000025
8. Kishore AK, Vail A, Bray BD, et al. Clinical risk scores for predicting stroke-associated pneumonia: a systematic review. *Eur Stroke J*. 2016;1(2):76–84. doi:10.1177/2396987316651759
9. Hotter B, Hoffmann S, Ulm L, et al. External validation of five scores to predict stroke-associated pneumonia and the role of selected blood biomarkers. *Stroke*. 2021;52(1):325–330. doi:10.1161/STROKEAHA.120.031884
10. Hoffmann S, Malzahn U, Harms H, et al. Development of a clinical score (A2DS2) to predict pneumonia in acute ischemic stroke. *Stroke*. 2012;43(10):2617–2623. doi:10.1161/STROKEAHA.112.653055
11. Smith CJ, Bray BD, Hoffman A, et al. Can a novel clinical risk score improve pneumonia prediction in acute stroke care? A UK multicenter cohort study. *J Am Heart Assoc*. 2015;4(1):e001307. doi:10.1161/JAHA.114.001307
12. Song X, He Y, Bai J, et al. A nomogram based on nutritional status and A(2)DS(2) score for predicting stroke-associated pneumonia in acute ischemic stroke patients with type 2 diabetes mellitus: a retrospective study. *Front Nutr*. 2022;9:1009041. doi:10.3389/fnut.2022.1009041
13. Hao R, Qi X, Xia X, et al. Malnutrition on admission increases the in-hospital mortality and length of stay in elder adults with acute ischemic stroke. *J Clin Lab Anal*. 2022;36(1):e24132. doi:10.1002/jcla.24132
14. Mehta A, De Paola L, Pana TA, et al. The relationship between nutritional status at the time of stroke on adverse outcomes: a systematic review and meta-analysis of prospective cohort studies. *Nutr Rev*. 2022;80(12):2275–2287. doi:10.1093/nutrit/nuac034

15. Li D, Liu Y, Jia Y, et al. Association between malnutrition and stroke-associated pneumonia in patients with ischemic stroke. *BMC Neurol.* **2023**;23(1):290. doi:10.1186/s12883-023-03340-1
16. Doi S, Iwata H, Wada H, et al. A novel and simply calculated nutritional index serves as a useful prognostic indicator in patients with coronary artery disease. *Int J Cardiol.* **2018**;262:92–98. doi:10.1016/j.ijcard.2018.02.039
17. Minami-Takano A, Iwata H, Miyosawa K, et al. A novel nutritional index serves as a useful prognostic indicator in cardiac critical patients requiring mechanical circulatory support. *Nutrients.* **2019**;11(6):1420. doi:10.3390/nu11061420
18. Ishiwata S, Yatsu S, Kasai T, et al. Prognostic effect of a novel simply calculated nutritional index in acute decompensated heart failure. *Nutrients.* **2020**;12(11):3311. doi:10.3390/nu12113311
19. Shi Y, Wang X, Yu C, et al. Association of a novel nutritional index with stroke in Chinese population with hypertension: insight from the China H-type hypertension registry study. *Front Nutr.* **2023**;10:997180. doi:10.3389/fnut.2023.997180
20. Zhang G, Pan Y, Zhang R, et al. A novel nutritional index and adverse outcomes in ischemic stroke: results from the third China National Stroke Registry. *Nutr, Metab Cardiovasc Dis.* **2022**;32(6):1477–1484. doi:10.1016/j.numecd.2022.02.015
21. Smith CJ, Kishore AK, Vail A, et al. Diagnosis of stroke-associated pneumonia: recommendations from the pneumonia in stroke consensus group. *Stroke.* **2015**;46(8):2335–2340. doi:10.1161/STROKEAHA.115.009617
22. Sabbouh T, Torbey MT. Malnutrition in stroke patients: risk factors, assessment, and management. *Neurocrit Care.* **2018**;29(3):374–384. doi:10.1007/s12028-017-0436-1
23. Chen X, Hu Y, Yuan X, et al. Effect of early enteral nutrition combined with probiotics in patients with stroke: a meta-analysis of randomized controlled trials. *Eur J Clin Nutr.* **2022**;76(4):592–603. doi:10.1038/s41430-021-00986-3
24. Zhang M, Ye S, Huang X, et al. Comparing the prognostic significance of nutritional screening tools and ESPEN-DCM on 3-month and 12-month outcomes in stroke patients. *Clin Nutr.* **2021**;40(5):3346–3353. doi:10.1016/j.clnu.2020.11.001
25. Kim HR, Kang MG, Kim K, et al. Comparative analysis of three nutrition scores in predicting mortality after acute myocardial infarction. *Nutrition.* **2021**;90:111243. doi:10.1016/j.nut.2021.111243
26. Sudo M, Shamekhi J, Aksoy A, et al. A simply calculated nutritional index provides clinical implications in patients undergoing transcatheter aortic valve replacement. *Clin Res Cardiol.* **2023**. doi:10.1007/s00392-023-02220-5
27. Ikizler TA, Cano NJ, Franch H, et al. Prevention and treatment of protein energy wasting in chronic kidney disease patients: a consensus statement by the international society of renal nutrition and metabolism. *Kidney Int.* **2013**;84(6):1096–1107. doi:10.1038/ki.2013.147
28. Ciancarelli I, Morone G, Iosa M, et al. Influence of oxidative stress and inflammation on nutritional status and neural plasticity: new perspectives on post-stroke neurorehabilitative outcome. *Nutrients.* **2022**;15(1):108. doi:10.3390/nu15010108
29. Dai C, Yan D, Xu M, et al. Geriatric Nutritional Risk Index is related to the risk of stroke-associated pneumonia. *Brain Behav.* **2022**;12(8):e2718. doi:10.1002/brb3.2718
30. Mitchell AB, Cole JW, McArdle PF, et al. Obesity increases risk of ischemic stroke in young adults. *Stroke.* **2015**;46(6):1690–1692. doi:10.1161/STROKEAHA.115.008940
31. Kim BJ, Lee SH, Jung KH, et al. Dynamics of obesity paradox after stroke, related to time from onset, age, and causes of death. *Neurology.* **2012**;79(9):856–863. doi:10.1212/WNL.0b013e318266fad1
32. Liu Z, Sanossian N, Starkman S, et al. Adiposity and outcome after ischemic stroke: obesity paradox for mortality and obesity parabola for favorable functional outcomes. *Stroke.* **2021**;52(1):144–151. doi:10.1161/STROKEAHA.119.027900
33. Reuben DB, Ix JH, Greendale GA, et al. The predictive value of combined hypoalbuminemia and hypocholesterolemia in high functioning community-dwelling older persons: macArthur studies of successful aging. *J Am Geriatr Soc.* **1999**;47(4):402–406. doi:10.1111/j.1532-5415.1999.tb07230.x
34. Zhang Z, Pereira SL, Luo M, et al. Evaluation of blood biomarkers associated with risk of malnutrition in older adults: a systematic review and meta-analysis. *Nutrients.* **2017**;9(8):829. doi:10.3390/nu9080829
35. Chen SC, Chung WS, Wu PY, et al. Associations among geriatric nutrition risk index, bone mineral density, body composition and handgrip strength in patients receiving hemodialysis. *Nutrition.* **2019**;65:6–12. doi:10.1016/j.nut.2019.02.013
36. Mizuiri S, Nishizawa Y, Doi T, et al. Association and predictive value of geriatric nutritional risk index, body composition, or bone mineral density in haemodialysis patients. *Nephrology (Carlton).* **2021**;26(4):341–349. doi:10.1111/nep.13826
37. Ignacio de Ulibarri J, González-Madroño A, de Villar NG, et al. CONUT: a tool for controlling nutritional status. First validation in a hospital population. *Nutr Hosp.* **2005**;20(1):38–45.
38. Gu A, Malahias MA, Strigelli V, et al. Preoperative malnutrition negatively correlates with postoperative wound complications and infection after total joint arthroplasty: a systematic reviewing and meta-analysis. *J Arthroplasty.* **2019**;34(5):1013–1024. doi:10.1016/j.arth.2019.01.005

## Clinical Interventions in Aging

Dovepress

## Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>