

Prevalence and Molecular Characteristics of Enterococci Isolated from Clinical Bovine Mastitis Cases in Ningxia [Letter]

Fitrine Ekawasti¹, Putri Reno Intan², Riza Zainuddin Ahmad¹

¹Research Center for Veterinary Science, Research Organization for Health, National Research and Innovation Agency (BRIN), Bogor, West Java, Indonesia; ²Research Center for Biomedical, Research Organization for Health, National Research and Innovation Agency (BRIN), Bogor, West Java, Indonesia

Correspondence: Riza Zainuddin Ahmad, Research Center for Veterinary Science, Research Organization for Health, National Research and Innovation Agency (BRIN), Genomic Building, Cibinong Science Center, Jalan Raya Bogor Km. 46, Cibinong, Bogor, West Java, 16911, Indonesia, Email rizamiko@yahoo.co.id

Dear editor

With grateful curiosity, we have intently studied an original article titled “Prevalence and Molecular Characteristics of Enterococci Isolated from Clinical Bovine Mastitis Cases in Ningxia” by Liu et al.¹ This is a very valuable study because it focuses on the investigation of the prevalence, antimicrobial resistance, and virulence characteristics of Enterococci isolated from clinical bovine mastitis in Ningxia Hui Autonomous Region, China, is important as it provides valuable insight into the incidence rate of Enterococci in clinical bovine mastitis cases which is an opportunistic mastitis pathogen present in the environment. *Enterococcus* spp. may work as a reservoir of antimicrobial-resistance genes, allowing them to be transmitted to humans through the food chain.²

The researcher in this paper used the culturing milk samples on sheep's blood agar for bacteria identification. Enterococcus bacteria had been found to grow, which were identified based on morphological features, then confirmed by PCR. In fact, several studies have shown that *Enterococcus* spp., *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus uberis*, *Escherichia coli*, and *Klebsiella pneumoniae* are the pathogen types that are most frequently found in cases of mastitis.³ In this study, the detection and sequencing of Enterococcus DNA are very important to confirming bacterial infections that cause mastitis, but we suggest applying the Real-Time Polymerase Chain Reaction (RT-PCR) technique, which has expanded into multiplex real-time PCR (rPCR) to diagnose specific mastitis infections. Bacterial culture and isolate identification are considered the gold standard in mastitis diagnosis but are time-consuming and produce many culture negative samples. Multiplex rPCR has proven to be extremely reliable in the diagnosis of mastitis since it can detect a huge number of bacteria in a relatively short time when compared to traditional methods, and mixed-infection mastitis pathogens are also more easily identified.^{4,5} Identification is also possible for clinical pathogen samples that fail to grow in culture. The low concentration of bacteria in milk has been suggested as one of the reasons for the absence of growth in bacterial culture.⁴ Even though there may have been other bacteria in the milk samples in this study that caused an increase in mastitis. Rapid and accurate identification of mastitis pathogens is important for early disease control.

In addition, antibiotic treatment has been utilized to manage mastitis. However, greater antimicrobial exposure in the environment may result in a rise in the number of multidrug resistant organisms. Antimicrobial resistance (AMR) is a topic that represents the One Health approach because it affects humans, the environment, and animals. Therefore, in the future, it is critical to investigate the AMR problem in dairy products regarding the incidence of mastitis through a One Health approach. Given the perspectives we presented, there is potential to improve the findings of this study for future research.

Acknowledgments

Acknowledgements are extended to Dr. Harimurti Nuradji at the Research Center for Veterinary Science, Research Organization for Health, National Research and Innovation Agency (BRIN) for their continuous support.

Disclosure

The authors report no conflicting interests in this communication.

References

1. Liu J, Liang Z, Zhongla M, et al. Prevalence and molecular characteristics of enterococci isolated from clinical bovine mastitis cases in Ningxia. *Infect Drug Resist.* 2024;17:2121–2129. doi:10.2147/IDR.S461587
2. Kim HJ, Youn HY, Kang HJ, et al. Prevalence and virulence characteristics of enterococcus faecalis and enterococcus faecium in bovine mastitis milk compared to bovine normal raw milk in South Korea. *Animals.* 2022;12(11):1407. doi:10.3390/ani12111407
3. Kabelitz T, Aubry E, van Vorst K, Amon T, Fulde M. The role of streptococcus spp. in bovine mastitis. *Microorganisms.* 2021;9(7):1497. doi:10.3390/microorganisms9071497
4. Keane OM, Budd KE, Flynn J, McCoy F. Increased detection of mastitis pathogens by real-time PCR compared to bacterial culture. *Vet Rec.* 2013;173(11):268. PMID: 23976784. doi:10.1136/vr.101598
5. Kahya Demirebilek S, Yıldız M, Akkoç A, Mutlu AM, Ardıçlı Ö, Aner H. Comparison of bacteriological culture method and multiplex real-time PCR for detection of mastitis. *Res Vet Sci.* 2024;172:105237. doi:10.1016/j.rvsc.2024.105237

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Infection and Drug Resistance 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Infection and Drug Resistance editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

<https://doi.org/10.2147/IDR.S482116>