

# Altered Heart Rate Variability During Mobile Game Playing and Watching Self-Mobile Gaming in Individuals with Problematic Mobile Game Use: Implications for Cardiac Health

Shih-Ching Chin<sup>1,\*</sup>, Yun-Hsuan Chang<sup>2-6,\*</sup>, Chih-Chun Huang<sup>6,7</sup>, Ting-Hsi Chou<sup>1</sup>, Chieh-Liang Huang<sup>8</sup>, Hsiu-Man Lin<sup>9</sup>, Marc N Potenza<sup>10</sup>

<sup>1</sup>Department of Psychology, Asia University, Taichung, Taiwan; <sup>2</sup>Institute of Gerontology, College of Medicine, National Cheng Kung University, Tainan, Taiwan; <sup>3</sup>Institute of Behavioral Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan; <sup>4</sup>Department of Psychology, National Cheng Kung University, Tainan, Taiwan; <sup>5</sup>Institute of Genomics and Bioinformatics, College of Life Sciences, National Chung Hsing University, Taichung, Taiwan; <sup>6</sup>Department of Psychiatry, National Cheng Kung University Hospital, Douliou Branch, Yunlin, Taiwan; <sup>7</sup>Department of Psychiatry, College of Medicine, National Cheng Kung University, Tainan, Taiwan; <sup>8</sup>Department of Psychiatry, Tsaotun Psychiatric Center, Ministry of Health and Welfare, Nantou, Taiwan; <sup>9</sup>Department of Child and Adolescent Development and Mental Health, China Medical University Children's Hospital, Taichung, Taiwan; <sup>10</sup>Psychiatry, Child Study and Neuroscience, Center of Excellence in Gambling Research, Yale School of Medicine, New Haven, CT, USA

\*These authors contributed equally to this work

Correspondence: Yun-Hsuan Chang, Licensed Clinical Psychologist, Institute of Gerontology, College of Medicine; Department of Psychology, National Cheng Kung University, No. 1., University Road, Tainan, Taiwan, Email z11108051@ncku.edu.tw; yh.chang.snoopy@gmail.com

**Introduction:** The surge in mobile gaming, fueled by smartphone and internet accessibility, lacks a comprehensive understanding of physiological changes during gameplay.

**Methods:** This study, involving 93 participants (average age 21.75 years), categorized them into Problematic Mobile Gaming (PMG) and non-problematic Mobile Gaming (nPMG) groups based on Problematic Mobile Gaming Questionnaire (PMGQ) scores. The PMGQ is a 12-item scale developed in Taiwan to assess symptoms of problematic mobile gaming. The research delved into heart rate variability (HRV) alterations during real-time mobile gaming and self-gaming video viewing.

**Results:** Results showed that the PMG group significantly presents a lower root mean square of successive differences (RMSSD), and High Frequency (lnHF) than does the nPMG group ( $F=4.73$ ,  $p=0.03$ ;  $F=10.65$ ,  $p=0.002$ , respectively) at the baseline. In addition, the PMG group significantly displayed elevated HF and low-frequency to high-frequency (LF/HF) in the mobile-gaming ( $F=7.59$ ,  $p=0.007$ ;  $F=9.31$ ,  $p=0.003$ ) condition as well as in the watching self-gaming videos ( $F=9.75$ ,  $p=0.002$ ;  $F=9.02$ ,  $p=0.003$ ) than did the nPMG.

**Conclusion:** The study suggests targeted interventions to mitigate autonomic arousal, offering a potential avenue to address adverse effects associated with problematic mobile gaming behavior. The PMG group displayed increased craving scores after real-time mobile gaming and watching self-gaming video excerpts, unlike the nPMG group. Elevated LF/HF ratios in frequent gaming cases heightened autonomic arousal, presenting challenges in relaxation after mobile gaming. These findings contribute to a nuanced understanding of the complex interplay between mobile gaming activities, physiological responses, and potential intervention strategies.

**Keywords:** addictive behaviors, video games, internet addiction, autonomic nervous system, craving, heart rate variability, self-regulation

## Introduction

With the widespread adoption of smartphones, gaming has shifted to mobile devices. People use smartphones for gaming to relax and cope with negative emotions, potentially leading to problematic usage.<sup>1</sup> Mobile game addiction may be associated with Internet Gaming Disorder (IGD). IGD was introduced as a research diagnosis in DSM-5 (2013)<sup>2,3</sup> and

later as a formal diagnosis in ICD-11 (2018).<sup>4</sup> Mobile gaming addiction involves prolonged and recurrent mobile game use, resulting in adverse physical and psychological effects, including social anxiety, depression, loneliness, insomnia, eye strain, impaired self-control, headaches, and academic underperformance.<sup>1,5–7</sup>

In Taiwan, internet cafés were once highly popular among young adults for playing online games, especially with friends.<sup>8</sup> However, media outlets have depicted online gaming as hazardous, publishing accounts of sudden deaths after excessive play in internet cafés.<sup>9–11</sup> Consequently, the impact of long-term gaming on cardiovascular function has drawn attention.<sup>12</sup> An “internet-addiction hypothesis of autonomic activity” has been proposed and suggested that dysregulation of the autonomic nervous system (ANS) is related to internet addiction and characterized by shifts towards sympathetic dominance and decreased parasympathetic activity.<sup>13</sup>

## Correlation Between Gaming and Heart Rate Variability

Heart rate variability (HRV) gauges autonomic nervous system (ANS) functioning,<sup>14</sup> revealing variations in individuals with and without Internet Gaming Disorder (IGD) during resting states, suggesting potential ANS imbalances, especially in sympathetic nervous system (SNS) activity even in the presence of gaming stimuli.<sup>15</sup> The severity of IGD correlates with reduced HRV in the high-frequency (HF) range. During gaming, IGD individuals display diminished HF HRV and root mean square of successive differences (RMSSD), while the non-IGD group maintains stable HRV.<sup>16,17</sup> These findings imply heightened SNS activity and diminished parasympathetic nervous system (PNS) activity in IGD individuals during gaming,<sup>18,19</sup> as confirmed by Hsieh and Hsiao (2016).<sup>20</sup> However, comprehending the clinical implications and advantages of increased HRV during and after gaming requires further investigation. Although existing studies suggest temporary HRV elevation during gaming, the lasting clinical impact on gamers remains uncertain.<sup>13</sup>

## Correlation Between Craving and Gaming Experiences

Individuals with Internet Gaming Disorder (IGD) often experience cravings, and powerful motivational desires implicated in addictive behavior development and maintenance.<sup>21,22</sup> Such cravings can arise when encountering negative emotions<sup>23</sup> and game-related cues, stimulating cravings among those with IGD.<sup>24,25</sup> Similar patterns have been observed in alcohol use disorder, where reduced resting heart rate variability (HRV), particularly in the high-frequency (HF) range, is linked to craving and relapse.<sup>26,27</sup> Additionally, lower HRV, particularly in the HF range, has been associated with increased food cravings.<sup>28,29</sup> This interconnection between cravings, emotional regulation, and HRV in addiction suggests that interventions targeting HRV enhancement and emotional regulation may aid in addiction management. Higher HRV is associated with better emotional control, potentially reducing cravings. This understanding highlights the potential for interventions to address cravings and promote healthier emotional regulation in the context of addiction.<sup>23,30,31</sup>

Previous research has explored the connection between craving and heart rate variability (HRV) in IGD. One study found that the standard deviation of normal-to-normal intervals (SDNN) decreased during gaming, but self-reported craving scores failed to distinguish effectively between individuals with IGD and those without. This study did not, however, delve into the links between craving and ANS system metrics.<sup>32</sup> The rise of social media platforms like YouTube, Instagram, and TikTok has provided opportunities to witness the gaming experiences of others. This exposure may trigger cue-related cravings, leading to desires to continue playing or watching games despite negative consequences like fatigue, sleep deprivation or neglect of other responsibilities.

The current study investigated relationships between problematic mobile gaming (PMG), HRV, and craving to gain insight into physiological underpinnings. Specifically, we hypothesized that individuals with PMG would demonstrate increased cue-related cravings. Further, we hypothesized that craving would link to diminished HRV among individuals with PMG.

## Methods

This study was approved by the Institutional Review Board (IRB) at Tsaotun Psychiatric Center (No. 111014) and China-Medical University Hospital (CMUH109-REC2-123). Before initiating the study, all participants needed to provide signed consent. For individuals under 20 years old, a parent’s consent was also required alongside their own. Participants were given the choice to leave the study at any time.

## Participants

Ninety-three participants, recruited through digital platform advertisements (from July 2022 to December 2022), provided informed consent for their involvement in the study. A power analysis using G\*power 3.1<sup>33</sup> determined that a minimum of 66 participants was required for statistical significance in a mixed-model ANOVA test, with an assumed medium effect size ( $d = 0.5$ ) and a power of 0.8. Participants, averaging 21.75 years, included 37 males and 56 females. Based on the Problematic Mobile Gaming Questionnaire (PMGQ) thresholds, 33 were classified with problematic mobile gaming (PMG; 14 males, 19 females), and 60 as non-problematic mobile gaming (nPMG; 23 males, 37 females). Within the PMG group, 45.5% met IGD-10 criteria, indicating a relative risk of Internet Gaming Disorder. Exclusive criteria involved neurological or mental disorders, substance use, tobacco, and conditions affecting heart rate variability (HRV). Demographic details are available in Table 1.

**Table 1** Demographic Information of the Participants

Variables	nPMG (N = 60)	PMG (N = 33)		$\chi^2/F$ (p)	
		PMG <sup>-IGD</sup> (N=18)	PMG <sup>+IGD</sup> (N=15)	nPMG vs PMG	PMG <sup>-IGD</sup> vs vs PMG <sup>+IGD</sup>
Age (M $\pm$ SD)	21.77 $\pm$ 2.72	21.56 $\pm$ 2.85	21.93 $\pm$ 3.03	0.004	0.37 (0.72)
Gender (N, %)				0.36	0.20 (0.65)
Male	23 (38.3%)	7 (38.9%)	7 (46.7%)	-	
Female	37 (61.7%)	11 (61.1%)	8 (53.3%)	-	
PMGQ (M $\pm$ SD)	21.27 $\pm$ 4.91	33.89 $\pm$ 3.43	33.73 $\pm$ 3.56	84.02***	0.02 (0.90)
<b>T1-Baseline</b>					
SDNN	53.49 (21.46)	46.30 (22.15)	47.85 (14.00)	2.13 (0.15)	0.05 (0.82)
RMSSD	42.06 (23.14)	31.13 (19.88)	32.34 (20.45)	4.73 (0.03)	0.03 (0.87)
InLF	5.29 (0.90)	5.27 (1.03)	5.33 (0.89)	0.01 (0.85)	0.04 (0.85)
InHF	5.40 (1.06)	4.64 (1.12)	4.67 (1.04)	10.65 (0.002)	0.01 (0.93)
LF/HF	1.91 (5.56)	2.15 (1.25)	3.75 (5.20)	0.81 (0.37)	1.60 (0.22)
CMGSb	5.27 (5.00)	11.50 (3.43)	11.20 (4.02)	37.84 (<.00001)	0.05 (0.82)
<b>T2-Mobile gaming</b>					
SDNN	53.25 (19.49)	48.79 (20.33)	46.67 (20.49)	1.61 (0.21)	0.09 (0.77)
RMSSD	44.56 (23.48)	34.14 (23.40)	32.40 (21.93)	5.01 (0.028)	0.05 (0.83)
InLF	5.31 (0.81)	5.27 (1.03)	5.33 (0.59)	0.43 (0.52)	1.78 (0.19)
InHF	5.21 (1.00)	4.68 (1.21)	4.44 (1.20)	7.59 (0.007)	0.33 (0.57)
LF/HF	1.49 (1.19)	2.37 (1.39)	2.29 (1.52)	9.31 (0.003)	0.03 (0.88)
<b>T3-Post mobile gaming</b>					
SDNN	56.10 (19.78)	47.64 (21.71)	53.63 (20.47)	1.71 (0.19)	0.65 (0.43)
RMSSD	42.26 (21.79)	29.12 (15.41)	28.98 (15.84)	9.50 (0.003)	0.001 (0.98)
InLF	5.34 (0.93)	5.30 (1.00)	5.58 (0.95)	0.19 (0.66)	0.71 (0.41)
InHF	5.45 (0.96)	4.71 (1.03)	4.41 (0.95)	17.38 (<.00001)	0.71 (0.41)
LF/HF	1.34 (1.55)	2.19 (1.45)	4.47 (3.04)	19.72 (<.00001)	7.98 (0.008) <sup>a</sup>
CMGSpg	9.20 (4.97)	16.39 (4.57)	17.80 (4.21)	57.21 (<.0001)	0.84 (0.37)
<b>T4- (Relax)</b>					
SDNN	59.12 (21.71)	59.02 (33.09)	58.90 (25.38)	0.001 (0.98)	0.00 (0.99)
RMSSD	42.45 (19.30)	33.95 (21.79)	33.04 (16.91)	4.53 (0.04)	0.02 (0.90)
InLF	5.54 (0.93)	5.59 (0.98)	5.65 (0.87)	0.15 (0.70)	0.03 (0.86)
InHF	5.47 (0.87)	4.68 (1.02)	4.78 (0.95)	14.33 (<.0001)	0.09 (0.77)
LF/HF	1.59 (1.76)	2.76 (1.43)	2.91 (1.81)	11.12 (0.001)	0.07 (0.79)
CMGSr	4.00 (4.29)	13.61 (4.83)	13.13 (4.85)	56.05 (<.00001)	0.08 (0.78)

(Continued)

**Table 1** (Continued).

Variables	nPMG (N = 60)	PMG (N = 33)		$\chi^2/F$ (p)	
		PMG <sup>-IGD</sup> (N=18)	PMG <sup>+IGD</sup> (N=15)	nPMG vs PMG	PMG <sup>-IGD</sup> vs vs PMG <sup>+IGD</sup>
T5-Watching gaming video)					
SDNN	59.57 (21.53)	60.46 (32.76)	54.03 (20.64)	0.15 (0.70)	0.43 (0.52)
RMSSD	46.64 (23.93)	42.23 (33.50)	34.11 (23.04)	2.09 (0.15)	0.63 (0.43)
lnLF	5.63 (0.78)	5.74 (0.84)	5.37 (0.59)	0.15 (0.70)	2.05 (0.16)
lnHF	5.40 (0.91)	4.85 (1.21)	4.61 (0.96)	9.75 (0.002)	0.38 (0.54)
LF/HF	1.71 (1.66)	2.94 (1.90)	2.67 (1.63)	9.02 (0.003)	0.20 (0.66)
T6-Post watching gaming video					
SDNN	60.82 (17.65)	59.34 (24.15)	64.80 (22.95)	0.05 (0.82)	0.44 (0.51)
RMSSD	43.92 (18.96)	32.60 (19.72)	37.15 (20.47)	4.89 (0.029)	0.42 (0.52)
lnLF	5.46 (0.91)	5.62 (0.86)	5.95 (0.75)	2.68 (0.11)	1.41 (0.25)
lnHF	5.42 (0.85)	4.69 (0.82)	4.87 (0.88)	12.42 (0.001)	0.38 (0.55)
LF/HF	1.68 (2.15)	2.91 (1.65)	3.47 (1.86)	11.43 (0.001)	0.83 (0.37)
CMGSpw	7.57 (5.70)	15.61 (4.30)	16.67 (6.64)	56.05 (<.0001)	0.49 (0.49)

**Note:** <sup>a</sup>significant difference in PMG sub-groups (PMG<sup>-IGD</sup> vs PMG<sup>+IGD</sup>); \*\*\*p < 0.0001.

**Abbreviations:** PMG, Problematic mobile gaming; nPMG, non-problematic mobile gaming; CMGS, Craving for mobile gaming score; SDNN, Standard deviation of NN interval; RMSSD, Root mean square of successive differences of normal NN intervals; LF, Low-frequency power; HF, High-frequency power; LF/HF, Low Frequency/High Frequency.

## Measurements

### Problematic Mobile Gaming Questionnaire (PMGQ)

The PMGQ, developed by Pan, Chiu, and Lin (2019)<sup>34</sup> in Taiwan, was constructed based on the Smartphone Addiction Inventory (SPAI) created by Liu, Lin, Pan, and Lin (2016).<sup>35</sup> to assess symptoms of problematic mobile gaming. This 12-item scale assesses “compulsion”, “tolerance”, and “withdrawal” using a 4-point Likert scale (1=strongly disagree, 2=somewhat disagree, 3=somewhat agree, and 4=strongly agree) based on experiences in the past 3 months. A PMG is defined by a score exceeding 30 points.<sup>34,35</sup> The Cronbach’s alpha was 0.92 for this study.

### Ten-Item Internet Gaming Disorder Test (IGDT-10)

The IGDT-10 scale (Chiu et al, 2018) assesses Internet Gaming Disorder (IGD) based on DSM-5 criteria (American Psychiatric Association, 2013). It includes 10 items, with participants rating the frequency of IGD-related behaviors on a 3-point scale (0=never, 1=sometimes, and 2=often). Items 9 and 10 were combined as one criterion. A score range of 0–18 was employed, with higher scores indicating more severe IGD. A threshold of 5 was used to define IGD. The IGDT-10 demonstrated strong internal consistency, with a Cronbach’s  $\alpha$  of 0.87 in the present study.

### Craving for Internet Gaming Scale (CIGS)

The CIGS, devised by Savci and Griffiths (2019),<sup>36</sup> is a modification of the alcohol craving scale by Flannery, Volpicelli, and Petinati (1999). A Chinese adaptation, composed of 5 items rated along a 7-point Likert scale, was used here and adapted to assess craving for mobile gaming (CMGS). The Cronbach’s alpha was 0.94 for this study.

## Instruments

Electrocardiograms (ECGs) were obtained using a BioGraph Infiniti 6.0 system. ECG electrodes were positioned on participants bilaterally, beneath the midclavicular line, and at the fifth intercostal space on the left side (below the heart). Heart rate was measured in beats per minute, and subsequently, the CardioPro Infiniti HRV Analysis Module was utilized for ANS analysis. This software enabled the correction of inter-beat interval data, which were then converted into an HRV index.<sup>37,38</sup>

## Procedure

Each participant was required to fixate on the monitor while resting. ECG sensors were attached and tested to ensure that stable signals were obtained for 5 minutes while participants were in a relaxed sitting position for baseline HRV assessments. Each participant then selected their regular/favorite game and engaged in real-time gaming for 10 minutes. Next, each participant relaxed for 5 minutes and then watched their self-mobile gaming video excerpt for 10 minutes. During each experimental phase, HRV signals were measured. The experimental protocol is shown in Figure 1.

## HRV Analysis

Three ECG channels were connected to participants' chests for EEG signal acquisition. Data underwent rigorous processing to eliminate unwanted noise caused by movement, breathing, and muscle electrical activity. These steps included a third-order Butterworth high-pass filtering (0.1 Hz cutoff), a sixth-order Butterworth notch filter, and a third-order Butterworth low-pass filtering with (cutoff at 15 Hz).<sup>39</sup> The ECG signals were recorded at a 200 Hz sampling rate, and R-R intervals were automatically detected using the Pan and Tompkins method.<sup>40</sup> HRV parameters were then derived from time and frequency domains using HRV analysis software.<sup>41</sup> The time-domain analysis measured the time between R-R intervals, employing SDNN (reflecting overall HRV) and RMSSD (reflecting short-term heart rate changes and parasympathetic activity).<sup>42</sup> For frequency-domain analyses, a 20% filter removed ectopic beats, and data were interpolated at 4 Hz.<sup>43</sup> The frequency-domain parameters were transformed with an autoregressive model. The LF domain (0.04–0.15 Hz) reflects parasympathetic and sympathetic nerve activity, and the HF domain (0.15–0.4 Hz) primarily represents parasympathetic activity.<sup>44</sup> The frequency-domain HRV parameters with skewed distributions were logarithmically transformed.<sup>45</sup> The LF/HF ratio was used as an index of sympathovagal balance.<sup>46</sup>


## Statistical Analysis

Chi-square tests compared categorical data between groups while One-way analyses of variance (ANOVAs) compared demographic and HRV parameters differences between groups. Repeated measures ANOVAs compared group differences across various phases, including baseline, real-time gaming, post-gaming relaxation, self-gaming video watching, and post-watching relaxation. HRV parameters (RMSSD, SDNN, lnLF, lnHF, and LF/HF) and craving score changes were examined for significant group and phase interactions. Post-hoc comparisons identified specific phase differences. Bonferroni corrections were applied for multiple comparisons ( $p < 0.003$ ). The analysis characterized HRV patterns during mobile gaming and self-gaming video watching in PMG and nPMG participants. Pearson correlation explored links between craving score changes and HRV parameters. Statistical Package for Social Sciences version 23.0 (SPSS Inc., Chicago, IL, USA) conducted all analyses.

## Results

### Demographic Data

Approximately one-third (35.5%) of our participants had PMG, and of these, 45.45% met the criteria as IGD. The demographic data between groups are shown in Table 1. No between-group differences regarding age, sex, or educational level were observed. All HRV parameters analyzed by mixed-model ANOVA revealed a significant main effect of group and condition with no significant interaction. A main effect of period was statistically significant for the SDNN-HRV ( $F [5, 87] = 17.40, p < 0.0001$ , partial eta-squared = 0.160), the RMSSD ( $F [5, 87] = 5.75$ ,




**Figure 1** Experimental protocol.

$p < 0.00001$ , partial eta-squared = 0.06), and lnHF ( $F [5,87] = 2.32$ ,  $p = 0.04$ , partial eta-squared = 0.025). In addition, independent t-tests revealed significant differences between the two groups in the lnHF-HRV and LF/HF-HRV after real-mobile gaming and watching self-gaming videos ( $p < .0016$ ). Moreover, the PMG group displayed elevated craving scores compared to the nPMG group after real-time gaming and after watching self-mobile gaming videos, including baseline ( $ps < 0.001$ ) (Table 1).

### Correlation Between Craving Change Scores and HRV Parameters Across Different Conditions

CMGS<sub>g</sub> (Craving for mobile gaming score at the mobile gaming condition) significantly positively correlated to LF/HF<sub>g</sub> ratio ( $r = 0.43$ ,  $p < 0.0001$ ), and negatively correlated to both RMSSD<sub>g</sub> ( $r = -.31$ ,  $p = 0.003$ ) and lnHF<sub>g</sub> ( $r = -.34$ ,  $p = 0.001$ ) (see Figure 2A). A similar pattern was observed correlating between CMGS<sub>w</sub> (Craving for mobile gaming score at the watching self-mobile gameplay video condition) with HRV parameters during that viewing period ( $r_{LF/HFw} = 0.35$ ,  $p = 0.001$ ;  $r_{RMSSDw} = -.23$ ,  $p = 0.03$ , and  $r_{lnHFw} = -.38$ ,  $p < 0.0001$  respectively) (Figure 2C). Significant correlations were also found between craving scores and HRV parameters during the post-mobile-gaming condition ( $r_{LF/HFpg} = 0.42$ ,  $p < 0.0001$ ;  $r_{RMSSDpg} = -.36$ ,  $p < 0.0001$ , and  $r_{lnHFpg} = -.45$ ,  $p < 0.0001$ , respectively) and post-watching-gaming-video-excerpt condition ( $r_{LF/HFpw} = 0.29$ ,  $p = 0.005$ ; and  $r_{lnHFpw} = -.32$ ,  $p = 0.002$ ) (Figure 2B and D).



**Figure 2** Correlations between craving scores and HRV parameters during each condition. (A) mobile gaming condition; (B) following the mobile gaming condition; (C) watching self-gaming video excerpts; (D) following the watching of self-gaming video excerpts.

**Notes:** \*:  $p < 0.05$ ; \*\*:  $p < 0.003$  (Bonferroni correction); \*\*\* $p < 0.0001$ .


## Self-Regulation Among Groups Across Mobile Gaming Conditions

The results showed LF/HF significantly rose during real-time mobile gaming and self-mobile gaming video observation in both groups, followed by a subsequent decrease. Independent t-tests comparing HRV changes between conditions in the two groups revealed elevated LF/HF-HRV changes in the PMG group after gaming ( $T=-2.53$ ,  $p=0.01$ ), but no significant increase after watching self-mobile gaming video ( $T=-.84$ ,  $p=0.40$ ) compared to the nPMG group (Figure 3A). Moreover, when comparing PMG individuals who also met IGD criteria ( $PMG^{+IGD}$ ), significant differences in LF/HF changes after real-time mobile gaming emerged ( $T_{pgg} = -3.19$ ,  $p=0.003$ ). The  $PMG^{+IGD}$  subgroup displayed the most pronounced LF/HF-HRV change during mobile gaming among all groups (Figure 3B). Interestingly, this change was absent after watching self-gaming video excerpts, instead revealing increased changes in  $\ln LF$ -HRV ( $T_{pwr} = -3.79$ ,  $p=0.001$ ).

## Discussion

Our study examined heart rate variability (HRV) fluctuations in young adults engaged in active mobile gaming and self-gaming video viewing. Participants with problematic mobile gaming (PMG) exhibited significant HRV parameter variations (SDNN, RMSSD, and  $\ln HF$ ) during both real-time mobile gaming and video watching. Within the PMG group, individuals with Internet Gaming Disorder (IGD) displayed distinct longitudinal LF/HF ratio shifts compared to those without IGD, indicative of unique HRV responses during gaming. The  $PMG^{+IGD}$  subgroup demonstrated elevated LF/HF ratios, indicating differing psychophysiological responses during gaming sessions. Subjective reports revealed consistent craving score increases after both real-time gaming and self-gaming video viewing in the PMG subgroup with IGD. A negative correlation between craving scores and  $\ln HF$  HRV parameter changes suggested an impact of real-time mobile gaming on parasympathetic activity, highlighting cardiovascular responses during mobile gaming.<sup>47–49</sup> Notably, such a negative correlation was also observed after watching videos of mobile gaming experiences, the subjective craving was also induced while watching a self-mobile gameplay video. This implies that commercial advertisements for mobile games on TV may induce cardiovascular stress akin to the experience of playing a mobile game, and with a longer video and longitudinal game playing may impact more.

Previous studies have explored HRV in IGD individuals<sup>16,18,19</sup> emphasizing its relevance in the context of gaming. Our research specifically focused on the PMG group, revealing HRV alterations consistent with prior IGD findings, signifying heightened SNS activity and reduced PNS activity both at rest and in response to mobile gaming cues.<sup>16,18–20</sup>



**Figure 3** The changes in HRV parameters and LF/HF ratios between real-time gaming and watching gaming conditions among groups: **(A)** significant difference of LF/HF changes in the real-time gaming condition between nPMG and PMG group; **(B)** significant changes of LF/HF ratio in the real-time gaming condition between  $PMG^{-IGD}$  and  $PMG^{+IGD}$

Notably, we did not observe significant differences in SDNN, RMSSD, lnLF, and LF/HF between PMG and nPMG at baseline; a notable difference in lnHF was found between the PMG and nPMG, echoing previous findings in IGD.<sup>19</sup>

Furthermore, our study identified significant differences in RMSSD, lnHF, and LF/HF at rest, a departure from some prior IGD research.<sup>15,50</sup> Previous studies indicated that individuals without IGD tend to exhibit increased SNS activity during gaming, with a return to normal physiological states immediately after gaming.<sup>51</sup> In contrast, our PMG<sup>+IGD</sup> displayed altered ANS functioning after following real-time mobile gaming.

Craving may sustain addictive behaviors, while studies have examined relationships between IGD and the ANS, there is limited research investigating relationships between gaming craving and the ANS. Regarding PMG and craving, while physiological signals associated with craving have been identified, research suggests that heart rate measurements are often not robust for detecting cravings. Moreover, the study found that video game materials do not effectively stimulate the cravings of individuals with PMG with a tendency as IGD. Researchers have emphasized the importance of HRV measurements to discern game cravings in individuals with IGD, facilitating an understanding of alterations in the ANS.<sup>52</sup> The results from our study suggest that there is a significant positive correlation between gaming craving and the LF/HF ratio, and a substantial negative correlation with RMSSD and lnHF measures. The findings suggest that gaming cravings may be reflected in ANS measures in individuals engaging in PMG.

In the context of time domains, the SDNN reflects overall HRV. On the other hand, the RMSSD reflects short-term HRV and is commonly employed to estimate PNS activity.<sup>38,53</sup> Among frequency domains, HF power has been linked to the regulation of the PNS and correlated with RMSSD. Conversely, LF power has been associated with SNS regulation. However, this interpretation has been disputed, as some researchers propose that LF represents the concurrent regulation of both the SNS and PNS. The LF/HF ratio reflects balance within the ANS and is frequently used as a measure of equilibrium between the SNS and PNS. A higher LF/HF ratio indicates a predominance of the SNS within the ANS, characterized by increased SNS activity and decreased PNS activity. Conversely, a lower LF/HF ratio suggests a dominance of the PNS within the ANS, with reduced SNS activity and increased PNS activity, typically observed when an individual is engaged in restful and relaxing behaviors.<sup>38,46,53–56</sup> Prolonged dominance of the SNS can potentially elevate the risk of mortality.<sup>57,58</sup>

Subgroup differences in LF/HF ratio change post-mobile gaming indicate distinct HRV responses in PMG, more specifically in the PMG met IGD similar to previous studies in IGD.<sup>16</sup> This suggests that individuals with more severe gaming problems may have elevated cardiac risk, potentially extending to gaming disorder.<sup>59</sup> Relative to the PMG<sup>-IGD</sup> and nPMG groups, the PMG<sup>+IGD</sup> exhibited significantly decreased LF/HF after playing a mobile game, suggesting a potential mechanism by which cardiac risk could extend into resting/relaxing states. Reduced HRV implies ANS imbalance, linked to various physiological and psychological disorders. Importantly, this imbalance has associations with cardiovascular disease and mortality.<sup>58,60,61</sup>

Previous studies have suggested potential connections between gaming and ANS function, posing risks like thromboembolisms due to prolonged sedentary periods and dehydration. Sudden blood pressure spikes during gaming could lead to hemorrhages, including strokes. Additionally, factors such as short-term and long-term sleep deprivation, extreme fatigue, and stress may disrupt autonomic functions, promoting heart rhythm irregularities.<sup>10</sup> Consequently, the potential links between maladaptive gaming and cardiovascular concerns should be closely monitored, especially in more vulnerable groups like older adults.

## Limitations

There are some limitations in the current study. First, the sample size of the PMG<sup>+IGD</sup> group was small, and not all variables affecting HRV could be controlled. The participants recruited in the current study were mostly young adults, implying that this population has easier and more accessible mobile phones for gaming. Although the PMGQ was developed to evaluate the problematic mobile gaming among adolescents, young adults may have more access to mobile gaming and need more attention. In addition, the categories and types of mobile games may alter the HRV parameters with different levels of stress reaction while playing different mobile games.<sup>62,63</sup> Second, participants were recruited from online platforms, which may have biased the participants' selection. Gamers who do not often use online platforms, such as Facebook or Instagram may miss the recruitment information. Third, the gender/sex distribution was unequal,

implying the gender distribution among mobile gamers could be different. Women exhibited higher parasympathetic tone, whereas, in men, sympathetic dominance declined much later in life.<sup>64</sup> In addition, the preference for games has been reported differently between men and women. Moreover, gender-related differences in several typical CVDs have been reported.<sup>65</sup> The factor of gender and age may be considered as a factor for evaluating cardiac disease, and this could be considered a confounding factor in future studies. Future studies investigating larger samples of women and girls are needed, especially as cardiac risks may differ across sexes. Although all participants were instructed to choose and play their favorite mobile games, different games may have different physiological effects, and these should be investigated in future studies.

## Conclusion

In sum, this study represents the first to explore associations between physiological information and problematic mobile gaming experiences. In addition, viewing videos related to gameplay may also induce relevant physiological responses as game-playing does. Moreover, we first identify the psychophysiological responses between the PMG and nPMG, and further demonstrate the similarly altered psychophysiological responses at real-time gaming and viewing gaming videos. This finding implied the potential induced physiological responses while watching gaming videos. Further, we first identified possible differences between PMG with and without IGD from a psychophysiological perspective, although the sample size was small to distinguish the difference between PMG with and without IGD from such psychophysiological markers. In addition, as the development of smartphone facilities embedded several types of applications (APP) and game genres, which may increase the risk of cardiac-related diseases. Moreover, game playing is applied for leisure time and stress relief,<sup>66,67</sup> as PMG has been linked to poor physical and mental well-being among teenagers and young adults,<sup>68</sup> future investigations may examine groups with co-occurring concerns. Such efforts might consider potential interventions, such as biofeedback training. Through biofeedback, individuals may learn to increase self-awareness, observe their physiological responses, and thus help reduce risks associated with PMG.

## Acknowledgments

Shih-Ching Chin and Yun-Hsuan Chang are co-first authors of this study. The authors are thankful for Miss Han-Yu Chuang and Mr. Yu-Wei Lin's assistance in data collection as well as the contribution from all participants.

## Funding

This work was supported by the National Science and Technology Council (NSTC) of Taiwan for financial grant support (grant No. NSTC 109-2628-H-006-003-MY2, NSTC 111-2321-B-006-008, NSTC 111-2314-H-006-002-MY3, NSTC 112-2410-H-006-089-SS2 and NSTC 112-2321-B-006-013), and partly supported by China-Medical University & Asia University collaboration project (grant No. ASIA-109-CMUH-18).

## Disclosure

Professor Marc N. Potenza has a pending patent application with Yale for Novartis; a consultant for Game Day Data; reports personal fees for consulting from Boehringer Ingelheim and Baria-Tek; advisory board for Opiant, outside the submitted work. The authors report no other conflicts of interest in this work.

## References

1. Krarup K, Krarup H. The physiological and biochemical effects of gaming: a review. *Environ. Res.* 2020;184:109344.
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub; 2013.
3. Petry NM, Rehbein F, Gentile DA, et al. An international consensus for assessing internet gaming disorder using the new DSM-5 approach. *Addiction*. 2014;109(9):1399–1406.
4. World Health Organization. ICD-11 for mortality and morbidity statistics; 2023. Available from: <https://icd.who.int/browse11/l-m/en#/http%253a%252f%252fid.who.int%252fid%252fentity%252f1448597234>. Accessed April 13, 2023.
5. Laconi S, Pirès S, Chabrol H. Internet gaming disorder, motives, game genres and psychopathology. *Computers in Human Behavior*. 2017;75:652–659.
6. Sayeed MA, Rasel MSR, Habibullah AA, Hossain MM. Prevalence and underlying factors of mobile game addiction among university students in Bangladesh. *Global Mental Health*. 2021;8.

7. Wang J-L, Sheng J-R, Wang H-Z. The association between mobile game addiction and depression, social anxiety, and loneliness. *Front Public Health*. 2019;247.
8. C-S W, Cheng -F-F. Internet Café Addiction of Taiwanese Adolescents. *Cyberpsychology Behav*. 2007;10:220–225.
9. Hunt K, Ng N. Man dies in Taiwan after 3-day online gaming binge. *Cyberpsychology Behav*. 2015.
10. Kuperczko D, Kenyeres P, Darnai G, Kovacs N, Janszky J. Sudden gamer death: non-violent death cases linked to playing video games. *BMC Psychiatry*. 2022;22(1):824.
11. Parkin S. *Death by Gaming: An Investigation into the Taiwan Café Fatalities*. Eurogamer; 2012.
12. Lawley CM, Skinner JR, Turner C. Syncope Due to Ventricular Arrhythmia Triggered by Electronic Gaming. *N Engl J Med*. 2019;381(12):1180–1181.
13. Lu DW, Wang JW, Huang AC. Differentiation of Internet addiction risk level based on autonomic nervous responses: the Internet-addiction hypothesis of autonomic activity. *Cyberpsychology, Behavior and Social Networking*. 2010;13(4):371–378.
14. Montano N, Porta A, Cogliati C, et al. Heart rate variability explored in the frequency domain: a tool to investigate the link between heart and behavior. *Neurosci Biobehav Rev*. 2009;33(2):71–80.
15. Kim N, Hughes TL, Park CG, Quinn L, Kong ID. Altered autonomic functions and distressed personality traits in male adolescents with internet gaming addiction. *Cyberpsychology, Behavior and Social Networking*. 2016;19(11):667–673.
16. Lee D, Hong SJ, Jung Y-C, Park J, Kim IY, Namkoong K. Altered heart rate variability during gaming in internet gaming disorder. *Cyberpsychology, Behavior and Social Networking*. 2018;21(4):259–267.
17. Yeo M, Lim S, Yoon G. Analysis of biosignals during immersion in computer games. *J Med Systems*. 2018;42(1):3.
18. Hong SJ, Lee D, Park J, et al. Altered heart rate variability during gameplay in internet gaming disorder: the impact of situations during the game. *Frontiers in Psychiatry*. 2018;9:429.
19. Lee D, Park J, Namkoong K, Hong SJ, Kim IY, Jung Y-C. Diminished cognitive control in Internet gaming disorder: a multimodal approach with magnetic resonance imaging and real-time heart rate variability. *Prog Neuro Psychopharmacol Biol Psychiatry*. 2020;110127.
20. Hsieh D-L, Hsiao T-C Heart rate variability of internet gaming disorder addicts in emotional states. Paper presented at: 2016 International Conference on Biomedical Engineering (BME-HUST); 2016.
21. Potenza MN. Should addictive disorders include non-substance-related conditions? *Addiction*. 2006;101:142–151.
22. Tiffany ST, Wray JM. The clinical significance of drug craving. *Ann. N.Y. Acad. Sci*. 2012;1248:1.
23. Giuliani NR, Berkman ET. Craving is an Affective State and Its Regulation Can Be Understood in Terms of the Extended Process Model of Emotion Regulation. *Psychol Inq*. 2015;26(1):48–53.
24. Wegmann E, Stodt B, Brand M. Cue-induced craving in Internet-communication disorder using visual and auditory cues in a cue-reactivity paradigm. *Addict Res Theory*. 2018;26(4):306–314.
25. Zhou Y, Zhou Y, Zhou J, Shen M, Zhang M. Attentional biases and daily game craving dynamics: an ecological momentary assessment study. *J Behav Addictions*. 2022.
26. Quintana DS, Guastella AJ, McGregor IS, Hickie IB, Kemp AH. Heart rate variability predicts alcohol craving in alcohol dependent outpatients: further evidence for HRV as a psychophysiological marker of self-regulation. *Drug Alcohol Depend*. 2013;132(1–2).
27. Ralevski E, Petrakis I, Altemus M. Heart rate variability in alcohol use: a review. *Pharmacol Biochem Behav*. 2019;176:83–92.
28. Rodríguez-Ruiz S, Ruiz-Padial E, Vera N, Fernández C, Anllo-Vento L, Vila J. Effect of heart rate variability on defensive reaction and eating disorder symptomatology in chocolate cravers. *J Psychophysiol*. 2009;23(3):95–103.
29. Wu J, Pierart C, Chaplin TM, Hommer RE, Mayes LC, Crowley MJ. Getting to the heart of food craving with resting heart rate variability in adolescents. *Appetite*. 2020;155:104816.
30. Appelhans BM, Luecken LJ. Heart rate variability as an index of regulated emotional responding. *Rev General Psychol*. 2006;10(3):229–240.
31. Cosme D, Mobasser A, Zeithamova D, Berkman ET, Pfeifer JH. Choosing to regulate: does choice enhance craving regulation? *Soc Cognit Affective Neurosci*. 2018;13(3):300–309.
32. Ha J, Park S, Im C-H, Kim L. Classification of gamers using multiple physiological signals: distinguishing features of internet gaming disorder. *Frontiers in Psychology*. 2021;12:714333.
33. Ferrin DL, Bligh MC, Kohles JC. Can I Trust You to Trust Me?: A Theory of Trust, Monitoring, and Cooperation in Interpersonal and Intergroup Relationships. *Group Org Management*. 2007;32(4):465–499.
34. Pan Y-C, Chiu Y-C, Lin Y-H. Development of the problematic mobile gaming questionnaire and prevalence of mobile gaming addiction among adolescents in Taiwan. *Cyberpsychology, Behavior and Social Networking*. 2019;22(10):662–669.
35. Liu C-H, Lin S-H, Pan Y-C, Lin Y-H. Smartphone gaming and frequent use pattern associated with smartphone addiction. *Medicine*. 2016;95(28).
36. Savić M, Griffiths MD. The development of the Turkish craving for internet gaming scale (CIGS): a validation study. *Int J Ment Health Addict*. 2019;1–18.
37. Malik M. Heart rate variability: standards of measurement, physiological interpretation, and clinical use: task force of the European Society of Cardiology and the North American Society for Pacing and Electrophysiology. *Ann Noninvasive Electrocardiol*. 1996;1(2):151–181.
38. Malik M. Time-domain measurement of heart rate variability. *Cardiac Electrophysiology Review*. 1997;1(3):329–334.
39. Christov II, Daskalov IK. Filtering of electromyogram artifacts from the electrocardiogram. *Med Eng Phys*. 1999;21(10):731–736.
40. Pan J, Tompkins WJ. A real-time QRS detection algorithm. *IEEE Transactions on Bio-Med Eng*. 1985;32(3):230–236.
41. Ramshur JTJ. *Evaluation, and Application of Heart Rate Variability Analysis Software (HRVAS): Biomedical Engineering*. The University of Memphis; 2010.
42. Mietus JE, Peng CK, Henry I, Goldsmith RL, Goldberger AL. The pNNx files: re-examining a widely used heart rate variability measure. *Heart*. 2002;88(4):378–380.
43. Peretz A, Kaufman JD, Trenga CA, et al. Effects of diesel exhaust inhalation on heart rate variability in human volunteers. *Environ Res*. 2008;107(2):178–184.
44. Reyes del paso GA, Langewitz W, Mulder LJ, van Roon A, Duschek S. The utility of low frequency heart rate variability as an index of sympathetic cardiac tone: a review with emphasis on a reanalysis of previous studies. *Psychophysiology*. 2013;50(5):477–487.
45. Boardman A, Schlindwein FS, Rocha AP, Leite A. A study on the optimum order of autoregressive models for heart rate variability. *Physiol Meas*. 2002;23(2):325–336.

46. Sztajzel J. Heart rate variability: a noninvasive electrocardiographic method to measure the autonomic nervous system. *Swiss Med Weekly*. 2004;134(35–36).
47. Desai V, Gupta A, Andersen L, Ronnestrand B, Wong M. Stress-Reducing Effects of Playing a Casual Video Game among Undergraduate Students. *Trends Psychol*. 2021;29(3):563–579. doi:10.1007/s43076-021-00062-6
48. Porter AM, Goolkasian P. Video Games and Stress: how Stress Appraisals and Game Content Affect Cardiovascular and Emotion Outcomes. *Front Psychol*. 2019;10:967.
49. Thomée S. Mobile Phone Use and Mental Health. A Review of the Research That Takes a Psychological Perspective on Exposure. *Int J Environ Res Public Health*. 2018;15(12).
50. Park SM, Lee JY, Choi AR, et al. Maladaptive neurovisceral interactions in patients with Internet gaming disorder: a study of heart rate variability and functional neural connectivity using the graph theory approach. *Addict Biol*. 2020;25(4).
51. Maia DTL, Costa DR, de Vasconcelos Dantas EP, Dos Santos Maciel T, Osório RAL. Heart rate variability analysis in children submitted to electronic games. *Scientia Medica*. 2020;30(1):3.
52. Kim H, Ha J, Chang W-D, Park W, Kim L, Im C-H. Detection of craving for gaming in adolescents with internet gaming disorder using multimodal biosignals. *Sensors*. 2018;18(1):102.
53. Force T. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Heart rate variability. Standards of measurement, physiological interpretation, and clinical use. *Circulation*. 1996;93(5):1043–1065.
54. Shaffer F, Combatalade DC. Don't add or miss a beat: a guide to cleaner heart rate variability recordings. *Biofeedback*. 2013;41(3):121–130.
55. Shaffer F, Ginsberg J. An overview of heart rate variability metrics and norms. *Front Public Health*. 2017;5:258.
56. Shaffer F, McCraty R, Zerr CL. A healthy heart is not a metronome: an integrative review of the heart's anatomy and heart rate variability. *Frontiers in Psychology*. 2014;5:1040.
57. Malliani A, Pagani M, Lombardi F. Methods for assessment of sympatho-vagal balance: power spectral analysis. *Vagal Control Heart: Exp Basis Clin Implications*. 1994;433–454.
58. Thayer JF, Brosschot JF. Psychosomatics and psychopathology: looking up and down from the brain. *Psychoneuroendocrinology*. 2005;30(10):1050–1058.
59. Pilver CE, Potenza MN. Increased incidence of cardiovascular conditions among older adults with pathological gambling features in a prospective study. *J Addiction Med*. 2013;7(6):387–393.
60. Thayer JF, Åhs F, Fredrikson M, Sollers III JJ, Wager TD. A meta-analysis of heart rate variability and neuroimaging studies: implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev*. 2012;36(2):747–756.
61. Thayer JF, Yamamoto SS, Brosschot JF. The relationship of autonomic imbalance, heart rate variability and cardiovascular disease risk factors. *Int J Cardiol*. 2010;141(2):122–131.
62. Kim JY, Kim HS, Kim DJ, Im SK, Kim MS. Identification of Video Game Addiction Using Heart-Rate Variability Parameters. *Sensors*. 2021;21(14).
63. Porter AM, Goolkasian P. Video Games and Stress: how Stress Appraisals and Game Content Affect Cardiovascular and Emotion Outcomes. *Frontiers in Psychology*. 2019;10.
64. Kunikullaya UK, Kunnavil R, Vijayadas J, Prakash VS, Murthy NS. Normative data and gender differences in heart rate variability in the healthy young individuals aged 18–30 years, a South Indian cross-sectional study. *Indian Pacing Electrophysiol J*. 2021;21(2):112–119.
65. Gao Z, Chen Z, Sun A, Deng X. Gender differences in cardiovascular disease. *Med Novel Technol Devices*. 2019;4:100025.
66. Chen C. Playing mobile games for stress recovery purposes: a survey of Chinese adolescents. *Telematic Informatic*. 2021;56:101481.
67. Croissant M, Frister M. A data-driven approach for examining the demand for relaxation games on Steam during the COVID-19 pandemic. *PLoS One*. 2021;16(12).
68. Zakaria AS, Adnan WH. Youth awareness: a survey on mobile gaming addiction concerning physical health performance on young adults in Malaysia. *J Media Information Warfare*. 2022;15(1):85–98.

## Psychology Research and Behavior Management

Dovepress

### Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>