

Effect of a Point-of-Care Ultrasound-Driven vs Standard Diagnostic Pathway on 24-Hour Hospital Stay in Emergency Department Patients with Dyspnea—Protocol for A Randomized Controlled Trial

Stig Holm Ovesen^{1,2}, Søren Helbo Skaarup³, Rasmus Aagaard⁴, Hans Kirkegaard¹, Bo Løfgren^{1,5}, Michael Dan Arvig^{6–8}, Bo Martin Bibby⁹, Stefan Posth¹⁰, Christian B Laursen^{11,12}, Jesper Weile^{1,2}

¹Research Center for Emergency Medicine, Department of Clinical Medicine, Aarhus University Hospital and Aarhus University, Aarhus, Denmark; ²Emergency Department, Horsens Regional Hospital, Horsens, Denmark; ³Department of Respiratory Diseases and Allergy, Aarhus University Hospital, Aarhus, Denmark; ⁴Department of Anesthesiology, Aarhus University Hospital, Aarhus, Denmark; ⁵Department of Internal Medicine, Randers Regional Hospital, Randers, Denmark; ⁶Emergency Department, Slagelse Hospital, Slagelse, Denmark; ⁷Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark; ⁸Department of Regional Health Research, University of Southern Denmark, Odense, Denmark; ⁹Department of Biostatistics, Aarhus University, Aarhus, Denmark; ¹⁰Emergency Department, Odense University Hospital, Odense, Denmark; ¹¹Department of Respiratory Medicine, Odense University Hospital, Odense, Denmark; ¹²Odense Respiratory Research Unit (ODIN), Department of Clinical Research, University of Southern Denmark, Odense, Denmark

Correspondence: Stig Holm Ovesen, Research Center for Emergency Medicine, Aarhus University Hospital, Palle Juul-Jensens Boulevard 99, J103, Aarhus, 8200, Denmark, Email stigholm@clin.au.dk

Purpose: Point-of-care ultrasound (POCUS) helps emergency department (ED) physicians make prompt and appropriate decisions, but the optimal diagnostic integration and potential clinical benefits remain unclear. We describe the protocol and statistical analysis plan for a randomized controlled trial. The objective is to determine the effect of a POCUS-driven diagnostic pathway in adult dyspneic ED patients on the proportion of patients having a hospital stay of less than 24 hours when compared to the standard diagnostic pathway.

Patients and Methods: This is a multicenter, randomized, investigator-initiated, open-labeled, pragmatic, controlled trial. Adult ED patients with chief complaint dyspnea are eligible. Patients are randomized (1:1) to the POCUS-driven diagnostic pathway or standard diagnostic pathway, with 337 patients in each group. The primary outcome is the proportion of patients having a hospital stay (from ED arrival to hospital discharge) of less than 24 hours. Key secondary outcomes include hospital length-of-stay, 72-hour revisits, and 30-day hospital-free days.

Conclusion: Sparse evidence exists for any clinical benefit from a POCUS-integrated diagnostic pathway. The results from this trial will help clarify the promising signals for POCUS to influence patient care among ED patients with dyspnea.

Keywords: focused lung ultrasound, focused cardiac ultrasound, shortness-of-breath, diagnostic effectiveness

Introduction

Background and Rationale

Acute dyspnea is a dominating chief complaint in emergency departments (ED) worldwide.^{1–6} Nevertheless, diagnostic uncertainty is still substantial due to unspecific and overlapping symptom complexes in disorders causing dyspnea, most frequently chronic obstructive pulmonary disease, pneumonia, heart failure, pulmonary embolism, and asthma.^{7–11} The evaluation of dyspnea is highly dependent on medical history and physical examination. But despite the additional usage of electrocardiograms, chest x-rays, arterial blood gas analysis, and venous blood sample test panels, expert adjudicators post-discharge still identify 30–40% of dyspneic patients with incorrect presumptive ED diagnoses.^{12–15}

Point-of-care ultrasound (POCUS) poses as a potentially valuable examination in the diagnostic work-up of ED patients with dyspnea. The diagnostic accuracy of focused lung and cardiac ultrasound for diagnosing most disorders causing acute dyspnea is excellent.^{16–23} However, the clinical benefit and how to optimally integrate it into current diagnostic pathways are still unclear.

For ED patients with dyspnea, no studies have found evidence that a diagnostic pathway guided by POCUS increases survival. However, it seems to lead to more prompt and appropriate decision-making, likely minimizing the patient's disease burden and making the healthcare process more efficient.^{12,13,15,24,25} Recent studies have found promising results for shorter hospital admissions.^{15,26–28} Based on these results, we hypothesize that a POCUS-driven diagnostic pathway shortens the hospital length of stay for patients admitted to the ED with chief complaint dyspnea.

Objectives

The study's primary objective is to compare the number of patients having a hospital stay of less than 24 hours in the POCUS versus standard care group.

The secondary objectives are to investigate the effect of a POCUS-driven diagnostic pathway on (1) hospital length of stay; (2) use of image resources; (3) 72-hour revisits; (4) 30-day hospital-free days; (5) time to treatment; and (6) patients' experiences.

A diagnostic sub-study is planned a priori (see the [Supplemental Material](#)).

Materials and Methods

Protocol

The entire protocol is provided in the [Supplemental Material](#), and all previous versions are available from the trial website.²⁹ The protocol is developed in accordance with the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) statement.^{30,31} The trial was registered at ClinicalTrials.gov (Identifier: NCT 05674916 (date: 2023 January 9)). The Committee on Health Research Ethics in the Central Denmark Region approved the trial protocol (Identifier: 1–10-72-160-22 (date: 2022 November 24)). This trial will comply with the Declaration of Helsinki.

A preliminary version of this work was presented at the PHD DAY conference 2023 at Aarhus University, Denmark.³²

Trial Design

The trial is a multicenter, randomized, investigator-initiated, open-labeled, pragmatic, controlled trial with 1:1 allocation.

Setting

The trial is conducted at EDs in Denmark. A complete list of participating centers is provided in the protocol.

Investigators

Participating physicians were recruited based on the following parameters: 1) certified competence in POCUS 2) an experience of >50 cardiac and lung scans; 3) clinical experience corresponding to three post-graduate years; 4) be in residency training or be specialized; and 5) be working in emergency medicine.

Eligibility Criteria

Patients will be included based on the following inclusion criteria: ED visit, age ≥ 18 years, chief complaint is dyspnea, and trial physician presence.

Chief complaint dyspnea is defined through one of the following four criteria:

- a. Triaged with Dyspnea (or similar, ie, Breathlessness) as the chief complaint in the local ED OR prehospital triaging system (determined by a triaging nurse or emergency medical service (EMS) technician, respectively).
- b. Chief complaint is not implemented in local triage system AND prehospital triage not performed, BUT

- i. In the general practitioner referral note, the wording is compatible with chief complaint Dyspnea, according to a predefined systematic word search string ([Supplementary Material](#)).
- ii. In the prehospital notification OR EMS crew notes, the wording is compatible with chief complaint Dyspnea, according to a predefined systematic word search-string ([Supplementary Material](#)).
- iii. The chapter for Dyspnea (or similar, ie, Breathlessness) is used in the criteria-based EMS dispatch tool.

Exclusion criteria are: Fulfillment of coded rapid-response team criteria (ie, trauma, surgical, or medical emergencies), prior focused lung or focused cardiac ultrasound in the current ED stay, prior enrollment in the trial, unable to consent, and non-Danish-speaking.

Study Flow

Patients will be screened by the trial physician, who will look through lists of patients arriving at the ED. Patients that trial physicians screen but do not include will be documented prospectively with the reason for exclusion in a screening log using the paper Case Report Form (CRF) ([Supplementary Material](#)). During the initial assessment, and upon fulfillment of inclusion criteria, the physician will ask for informed consent, enroll, and randomize the patient. The exact time of randomization should be immediately after the consent has been signed.

A diagnostic survey will be filled out twice by the physician during the diagnostic process ([Supplementary Material](#)). After hospital discharge, patients will receive an electronic questionnaire concerning their experiences from the ED stay ([Figure 1](#)) ([Supplementary Material](#)).

Interventions

Standard Diagnostic Pathway

The standard diagnostic pathway will be based on the treating physician's discretion and diagnostic probabilities; it will likely include, but not be limited to, blood samples, blood gases, electrocardiogram, and chest x-ray. Focused lung and cardiac ultrasound cannot be performed while the patients stay in the ED unless clinical deterioration and patient safety demand it. Other POCUS modalities can be applied immediately (eg, lower extremity compression or focused abdominal ultrasound). Chest computed tomography (CT), CT pulmonary angiogram, echocardiography, and other next-line testing remain exclusively at the treating physicians' discretion, guided by national disease-specific guidelines.^{33–36}

Point-of-Care Ultrasound-Driven Diagnostic Pathway

Focused lung and cardiac ultrasound will be performed by the treating physician during the primary assessment. The final decision on next-line imaging (ie, chest x-ray, echocardiography, or CT) and further diagnostic testing remain at the treating physicians' discretion and should incorporate history, physical examinations, and POCUS. Per protocol, diagnostic decision recommendations will be provided for the treating physician ([Supplementary Material](#)).

POCUS will be performed by the treating ED physician at the discretion of timing, workflow and preference of ultrasound apparatus, transducer, and pre-set; the protocols are in accordance with international standards.^{37–39} Focused lung ultrasound will include eight zones (anterior and lateral) and evaluate pneumothorax, B-line pattern, lung consolidation, and pleural effusion ([Supplementary Material](#)). Focused cardiac ultrasound will consist of four views (subxiphoid four-chamber view, parasternal long-axis view, parasternal short-axis view, and apical four-chamber view) and evaluate pericardial fluid, right ventricle dilation, and left ventricular systolic contractility ([Supplementary Material](#)). Other POCUS modalities can be applied at the physician's discretion (eg, lower extremity compression or focused abdominal ultrasound).

Blinding

Blinding patients or clinical teams will not be attempted.

	STUDY PERIOD						
	Enrolment	Randomization	Post-randomization		Close-out		Follow-up
TIMEPOINT	-t1	0	t1 PA/ \pm POCUS	Ended SDP	ED Discharge	Discharge	30 days
ENROLMENT:							
Eligibility screen	X						
Screening log	X						X
Informed consent		X					
Randomization		X					
INTERVENTIONS:							
POCUS-driven DP			X				
SDP			X				
ASSESSMENTS:							
Case Report Form	X	X					
Diagnostic survey			X	X			
Patient Questionnaire						X	
Retrospective data retrieval (baseline characteristics, vital parameters, biochemical findings, microbiologic findings, treatments, admission metrics [timing and units], readmissions, mortality)						X	X
Manual record review (PA findings, ultrasound findings, radiologic findings)						X	X

Figure 1 SPIRIT flow diagram: Schedule of enrolment, interventions, and assessments.

Abbreviations: DP, diagnostic pathway; SDP, standard diagnostic pathway; PA, primary assessment.

Randomization

Patients will be randomized in a 1:1 ratio in blocks with random sizes of 2, 4, or 6 within each site. The randomized sequence of study identification numbers will be generated using the “blockrand” R package.^{40,41} Opaque randomization envelopes will be prepared, packed, and shipped to participating centers based on their current tally and inclusion rate by the principal investigator. The allocation sequence list and block sizes are only known by the principal investigator and steering committee statistician.

Outcomes

The primary outcome, 24-hour hospital stay, will be defined as the proportion of patients having a hospital stay (from ED arrival to hospital discharge) of less than 24 hours.

Secondary outcomes

Hospital length-of-stay, defined from ED arrival to hospital discharge, will secondarily be analyzed as a non-binary outcome.

Image resources will be quantified in numbers of the following imaging techniques performed during the current hospital stay: chest x-rays, echocardiography, CT pulmonary angiography, and chest CT.

Seventy-two-hour revisits will be defined as a composite outcome including any unplanned hospital stay within 72 hours from the previous hospital discharge, in-hospital mortality, and mortality within 72 hours from the previous hospital discharge.

Hospital-free days will be defined as the number of days within the 30-day period following hospital discharge where the patient is not hospitalized and alive.⁴² In-hospital death will be computed as zero hospital-free days. Registrations showing that a patient has left home for a hospital contact will be computed as 1 day that was not hospital-free (ie, outpatient or paraclinical follow-up visits). Long- or short-stay nursing facilities, inpatient hospice facilities, or rehabilitation facilities will be hospital-free, as will all days at home, including those with home-based medical services.⁴²

Time to treatment will be defined as time from ED arrival until administration of any of the following treatment subgroups: antibiotics, diuretics, and anticoagulants. Only ED administrations will be included. Oral and intravenous treatments will not be separated.

Patients' experiences will be defined on a 1–5 Likert scale (from “Not at all” to “Very much”) with two alternative responses (“Don't know” and “Not relevant for me”).

Sample Size Calculation

Based on a previous study with a similar population, the proportion of patients discharged within 24 hours is expected to be 24%.¹⁵ They found that 40% of POCUS group patients were discharged within 24 hours, corresponding to a risk difference of 16% (24% versus 40%). We regard a risk difference of 10% as clinically relevant. Based on a chi-squared test, assuming 5% dropouts and an alpha of 0.05, a sample size of 674 patients (337 per group) is needed to obtain 80% power.

Statistical Analysis Plan

The statistical analyses will adhere to the Consolidated Standards of Reporting Trials (CONSORT)-guidelines.^{43,44} Post-randomization exclusion from analyses will only happen from withdrawal or mislaying of patient consent or duplicate inclusion of the same individual. Apart from that, all analyses will be conducted on an intention-to-treat basis.^{45,46} All tests will be two-sided, a p-value <0.05 will be considered significant, and all confidence intervals will have 95% coverage.

The primary outcome will be presented as the absolute number and proportions in each group. Groups will be compared using risk difference and relative risk with 95% confidence intervals, and a p-value will be calculated based on a binary regression.

Hospital length-of-stay will be analyzed using time-to-event analysis and pseudo-observations. The Nelson-Aalen estimator will be applied to estimate the cumulative hazard rates in the intervention and control groups. A Cox regression

will be used to compare the overall hazards. Lastly, a generalized linear model using pseudo-observations accounting for in-hospital mortality as a competing risk will compare between-group discharge risks within 3 and 7 days (including the endpoint day, 72 and 168 hours).

Image resources and 72-hour revisits will be presented as the composite number and incidence proportion and compared by risk difference and relative risk based on a binary regression.

Thirty-day hospital-free days will be presented as raw means in the two groups with 95% confidence intervals. A Tobit regression model will be built upon a logistic model for mortality and linear regression for days alive outside the hospital within 30 days for patients discharged alive. Between-group comparison will be quantified as a mean difference from the Tobit regression.

Time to treatment will be presented as a composite outcome with a median and interquartile interval for both groups. Comparison will be performed using Cox regression analysis.

Patients' experiences will be treated continuously, comparing mean scores between groups using an unpaired *t*-test. For descriptive comparison, category distributions will be depicted by group.

Subgroup analyses will be performed on the relative scale for the primary outcome, according to: (1) ED arrival (day, evening, or night); (2) weekdays or weekends/holidays; (3) physicians' ultrasound competence categorized; (4) physicians' post-graduate years categorized; (5) prehospital presumptive diagnosis; and (6) 10-year patient-age intervals.

Data Collection and Data Management

The case report form (CRF) will be available in paper and electronic format. The paper-CRF will be filled and stored during all prospective screenings of non-included patients. The electronic CRF (eCRF) will be filled out for all included patients.

Baseline characteristics, vital parameters, biochemical and microbiologic results, treatments, admission metrics (timing and departments), readmissions, and mortality will be collected retrospectively using routinely collected patient-record data. All data will be collected longitudinally using the personal identification number as the key identifier. The primary physical assessment, ultrasound, and radiologic findings will be manually entered and coded into a REDCap[®] database based on a review of medical records.

To minimize missing data, e-mail and SMS reminders of the diagnostic survey will be sent to treating physicians at four, six, and eight hours post-randomization.

Trial physicians will additionally fill out a survey regarding their experience, including post-graduate years, specialization, ultrasound certification, and years of using ultrasound ([Supplementary Material](#)).

All data will be kept in a secure content management system (Hyland Alfresco[™], Boston, Massachusetts, USA). Data management will be done using STATA[®].⁴⁷ Ultrasound film clips will be stored securely, according to local legal regulations.

Data handling will comply with the General Data Protection Regulation (European Union 2016/679) and the Danish Act for Data Protection (ACT 502 of 23rd May 2018).

Ethical Considerations and Consent

Consent is obtained according to Danish law.

Data Sharing

Six months after the publication of the last results, all de-identified individual patient data will be made available for data sharing.⁴⁸ Procedures, including re-coding of critical variables, will be put in place to allow for complete de-identification of the data. Data will be completely anonymized according to Danish law.

Discussion

The current article describes the design of the POCUS PATHWAY trial, which tests the effect of a POCUS-driven diagnostic pathway in adult dyspneic ED patients on the proportion of patients having a hospital stay of less than 24 hours when compared to the standard diagnostic pathway.

Increasing costs, crowding, and elderly burden call for healthcare systems to provide simplifications. For that matter, in ED patients with dyspnea, POCUS holds a potential that is still not fully elucidated.

The outcomes addressed in this trial will cover important interests of patients and organizations.⁴⁹ We believe length-of-stay as the primary outcome is meaningful if it does not compromise 72-hour revisits and 30-day hospital-free days. Together, these results will help clarify the promising signals for POCUS to influence patient care.

Status

The trial started on the 25th of January 2023. To date, 490 patients have been included (1.3 per day). Assuming a similar inclusion rate, inclusion is expected to be finalized by July 2024. The primary results will be reported after 30 days of follow-up and are anticipated in late 2024.

Acknowledgments

We thank Dr. Minna Riishede for providing valuable help in the early phases of planning the trial. We also thank the conference organizers of the PHD DAY 2023 at Aarhus University for the opportunity to present an earlier version of this work, and acknowledge the feedback and comments received during this conference.

Funding

This study is funded by the Central Denmark Region (DKK 1,629,000), Frimodt-Heineke Foundation (DKK 40,000), Johan Schrøders Foundation (DKK 200,000), and Riisfort Foundation (DKK 211,858). The funding agencies have no role in designing or executing the study or interpreting, writing, or submitting the manuscript.

Disclosure

Dr. Christian B Laursen reports the following financial relationships: Royalties as author/editor from Munksgaard (Publisher); fees for giving presentations at educational events organized by AstraZeneca, Chiesi Pharma, and GSK. These financial relationships do not influence the design, conduct, or reporting of the research findings in this manuscript. The authors report no other conflicts of interest in this work.

References

1. Parshall MB, Schwartzstein RM, Adams L, et al. An official American thoracic society statement: update on the mechanisms, assessment, and management of dyspnea. *Am J Respir Crit Care Med*. 2012;185(4):435–452. doi:10.1164/rccm.201111-2042ST
2. Kelly AM, Keijzers G, Klim S, et al. An observational study of dyspnea in emergency departments: the Asia, Australia, and New Zealand Dyspnea in emergency departments study (AANZDEM). *Acad Emerg Med*. 2017;24(3):328–336. doi:10.1111/acem.13118
3. Nielsen MK, Henriksen DP, Ostergaard KB, et al. Association between presenting complaints of acutely admitted medical patients and mortality: a cohort study. *Eur J Intern Med*. 2018;54:e29–e32. doi:10.1016/j.ejim.2018.06.011
4. Sørensen SF, Ovesen SH, Lisby M, et al. Predicting mortality and readmission based on chief complaint in emergency department patients: a cohort study. *Trauma Surg Acute Care Open*. 2021;6(1):e000604. doi:10.1136/tsaco-2020-000604
5. Cairns CKK, Santo L. National hospital ambulatory medical care survey: 2018 emergency department summary tables. Available from: https://www.cdc.gov/nchs/data/nhamcs/web_tables/2018_ed_web_tables-508.pdf. Accessed March 30, 2022.
6. Pines JM, Mullins PM, Cooper JK, et al. National trends in emergency department use, care patterns, and quality of care of older adults in the United States. *J Am Geriatr Soc*. 2013;61(1):12–17. doi:10.1111/jgs.12072
7. Green SM, Martinez-Rumayor A, Gregory SA, et al. Clinical uncertainty, diagnostic accuracy, and outcomes in emergency department patients presenting with dyspnea. *Arch Intern Med*. 2008;168(7):741–748. doi:10.1001/archinte.168.7.741
8. Hale ZE, Singhal A, Hsia RY. Causes of shortness of breath in the acute patient: a national study. *Acad Emerg Med*. 2018;25:1227–1234. doi:10.1111/acem.13448
9. Hunold KM, Caterino JM. High diagnostic uncertainty and inaccuracy in adult emergency department patients with dyspnea: a national database analysis. *Acad Emerg Med*. 2018;26:267–271. doi:10.1111/acem.13553
10. Ray P, Birolleau S, Lefort Y, et al. Acute respiratory failure in the elderly: etiology, emergency diagnosis and prognosis. *Crit Care*. 2006;10(3):R82.
11. Ovesen SH, Sørensen SF, Lisby M, et al. Change in diagnosis from the emergency department to hospital discharge in dyspnoeic patients. *Dan Med J*. 2022;69(2):1.
12. Laursen CB, Sloth E, Lassen AT, et al. Point-of-care ultrasonography in patients admitted with respiratory symptoms: a single-blind, randomised controlled trial. *Lancet Respir Med*. 2014;2(8):638–646. doi:10.1016/S2213-2600(14)70135-3
13. Pivetta E, Goffi A, Nazerian P, et al. Lung ultrasound integrated with clinical assessment for the diagnosis of acute decompensated heart failure in the emergency department: a randomized controlled trial. *Eur J Heart Fail*. 2019;2019:1.
14. Becker TK, Tafoya CA, Osei-Ampofo M, et al. Cardiopulmonary ultrasound for critically ill adults improves diagnostic accuracy in a resource-limited setting: the Africa trial. *Trop Med Int Health*. 2017;22(12):1599–1608. doi:10.1111/tmi.12992

15. Riishede M, Lassen AT, Baatrup G, et al. Point-of-care ultrasound of the heart and lungs in patients with respiratory failure: a pragmatic randomized controlled multicenter trial. *Scandinavian J Trauma Resuscit Emerg Med*. 2021;29(1):60. doi:10.1186/s13049-021-00872-8
16. Gartlehner G, Wagner G, Affengruber L, et al. Point-of-care ultrasonography in patients with acute dyspnea: an evidence report for a clinical practice guideline by the American college of physicians. *Ann Intern Med*. 2021;174(7):967–976. doi:10.7326/M20-5504
17. Martindale JL, Wakai A, Collins SP, et al. Diagnosing acute heart failure in the emergency department: a systematic review and meta-analysis. *Acad Emerg Med*. 2016;23(3):223–242. doi:10.1111/acem.12878
18. Wang M, Luo X, Wang L, et al. A comparison of lung ultrasound and computed tomography in the diagnosis of patients with COVID-19: a systematic review and meta-analysis. *Diagnostics*. 2021;11(8):1351. doi:10.3390/diagnostics11081351
19. Orso D, Guglielmo N, Copetti R. Lung ultrasound in diagnosing pneumonia in the emergency department: a systematic review and meta-analysis. *Eur J Emerg Med*. 2017;2017:1.
20. Falster C, Jacobsen N, Coman KE, et al. Diagnostic accuracy of focused deep venous, lung, cardiac and multiorgan ultrasound in suspected pulmonary embolism: a systematic review and meta-analysis. *Thorax*. 2021;2021:216838.
21. Yousefifard M, Baikpour M, Ghelichkhani P, et al. Screening performance characteristic of ultrasonography and radiography in detection of pleural effusion; a meta-analysis. *Emerg*. 2016;4(1):1–10.
22. Alrajhi K, Woo MY, Vaillancourt C. Test characteristics of ultrasonography for the detection of pneumothorax. *Chest*. 2012;141(3):703–708. doi:10.1378/chest.11-0131
23. Al Deeb M, Barbic S, Featherstone R, et al. Point-of-care ultrasonography for the diagnosis of acute cardiogenic pulmonary edema in patients presenting with acute dyspnea: a systematic review and meta-analysis. *Acad Emerg Med*. 2014;21(8):843–852. doi:10.1111/acem.12435
24. Nakao S, Vaillancourt C, Taljaard M, et al. Evaluating the impact of point-of-care ultrasonography on patients with suspected acute heart failure or chronic obstructive pulmonary disease exacerbation in the emergency department: a prospective observational study. *CJEM*. 2020;22(3):342–349. doi:10.1017/cem.2019.499
25. De Carvalho H, Javaudin F, Le Bastard Q, et al. Effect of chest ultrasound on diagnostic workup in elderly patients with acute respiratory failure in the emergency department: a prospective study. *Eur J Emerg Med*. 2022;30:59–60. doi:10.1097/MEJ.0000000000000959
26. Mozzini C, Di Dio Perna M, Pesce G, et al. Lung ultrasound in internal medicine efficiently drives the management of patients with heart failure and speeds up the discharge time. *Intern Emerg Med*. 2018;13(1):27–33.
27. Zare MA, Bahmani A, Fathi M, et al. Role of point-of-care ultrasound study in early disposition of patients with undifferentiated acute dyspnea in emergency department: a multi-center prospective study. *J Ultrasound*. 2021;25:443–449. doi:10.1007/s40477-021-00582-y
28. Cid-Serra X, Royse A, Canty D, et al. Effect of a multiorgan focused clinical ultrasonography on length of stay in patients admitted with a cardiopulmonary diagnosis. *JAMA Network Open*. 2021;4(12):e2138228.
29. Website for the POCUS PATHWAY trial; 2024. Available from: <https://pocuspathway.org/>. Accessed August 21, 2024.
30. Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med*. 2013;158(3):200–207. doi:10.7326/0003-4819-158-3-201302050-00583
31. Chan AW, Tetzlaff JM, Gøtzsche PC, et al. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ*. 2013;346(15):e7586–e86. doi:10.1136/bmj.e7586
32. Abstract Book PHD DAY 2023. Aarhus University; 2023. Available from: https://phd.health.au.dk/fileadmin/phd.health.au.dk/Health_NY_filliste/About_us/Events/PhD_Day_2023/Abstract_book_PhD_Day_2023.pdf. Accessed August 21, 2024.
33. COPD exacerbation: Danish Respiratory Society 2020; 2020. Available from: <https://lungemedicin.dk/kol-exacerbation-og-niv/>. Accessed August 21, 2024.
34. Acute heart failure: Danish society of cardiology; 2023. Available from: <https://www.cardio.dk/akut-hf>. Accessed August 21, 2024.
35. Pulmonary embolism and deep venous thrombosis: Danish Society of Cardiology; 2023. Available from: <https://www.cardio.dk/lungeemboli>. Accessed August 21, 2024.
36. Adult patients hospitalized with pneumonia: Danish respiratory society and Danish infectious disease society; 2021. Available from: <https://lungemedicin.dk/pneumoni/>. Accessed August 21, 2024.
37. Via G, Hussain A, Wells M, et al. International evidence-based recommendations for focused cardiac ultrasound. *J Am Soc Echocardiogr*. 2014;27(7):683.e1–683.e33. doi:10.1016/j.echo.2014.05.001
38. Volpicelli G, Mussa A, Garofalo G, et al. Bedside lung ultrasound in the assessment of alveolar-interstitial syndrome. *Am J Emergency Med*. 2006;24(6):689–696. doi:10.1016/j.ajem.2006.02.013
39. Volpicelli G, Elbarbary M, Blaivas M, et al. International evidence-based recommendations for point-of-care lung ultrasound. *Intensive Care Med*. 2012;38(4):577–591.
40. Clinical Trials_ [program]. _blockrand: randomization for Block Random; 2020.
41. R Foundation for Statistical Computing. R: a language and environment for statistical computing [program]. Vienna, Austria: R Foundation for Statistical Computing; 2023.
42. Auriemma CL, Taylor SP, Harhay MO, et al. Hospital-free days: a pragmatic and patient-centered outcome for trials among critically and seriously ill patients. *Am J Respir Crit Care Med*. 2021;204(8):902–909. doi:10.1164/rccm.202104-1063PP
43. Zwarenstein M, Treweek S, Gagnier JJ, et al. Improving the reporting of pragmatic trials: an extension of the CONSORT statement. *BMJ*. 2008;337:a2390.
44. Schulz KF, Altman DG, Moher D, et al. Consort 2010 statement: updated guidelines for reporting parallel group randomized trials. *Ann Intern Med*. 2010;152(11):726–732.
45. Schulz KF, Grimes DA. Sample size slippages in randomised trials: exclusions and the lost and wayward. *Lancet*. 2002;359(9308):781–785.
46. Detry MA, Lewis RJ. The intention-to-treat principle. *JAMA*. 2014;312(1):85. doi:10.1001/jama.2014.7523
47. StataCorp LLC. Stata statistical software: release 18. College Station, TX: StataCorp LLC; 2023.
48. Taichman DB, Sahni P, Pinborg A, et al. Data Sharing Statements for Clinical Trials: a Requirement of the International Committee of Medical Journal Editors. *Lancet*. 2017;376(23):2277–2279.
49. Van Den Ende ES, Schouten B, Kremers MNT, et al. Understanding what matters most to patients in acute care in seven countries, using the flash mob study design. *BMC Health Serv Res*. 2021;21(1):1.

Open Access Emergency Medicine

Dovepress

Publish your work in this journal

The Open Access Emergency Medicine is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of emergency medicine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/open-access-emergency-medicine-journal>