

The Association Between Autistic Traits and Depression in College Students: The Mediating Roles of Interpersonal Emotion Regulation and Social Self-Efficacy

Caina Liu^{1,*}, Qingqing Zhang^{1,2,*}, Yiming Liu¹, Zixu Wang¹, Fangchen Chen¹, Yitian Li¹, Yami Zhao¹, Jing Zhu¹, Dandan Li^{1,3-5}, Chunyan Zhu^{1-4,6}

¹School of Mental Health and Psychological Sciences, Anhui Medical University, Hefei, 230032, People's Republic of China; ²Institute of Artificial Intelligence, Hefei Comprehensive National Science Center, Hefei, 230000, People's Republic of China; ³Anhui Province Key Laboratory of Cognition and Neuropsychiatric Disorders, Hefei, 230032, People's Republic of China; ⁴Collaborative Innovation Center of Neuropsychiatric Disorders and Mental Health, Hefei, 230032, People's Republic of China; ⁵Department of Neurology, The First Affiliated Hospital of Anhui Medical University, Hefei, 230022, People's Republic of China; ⁶Department of Psychology, The Second Affiliated Hospital of Anhui Medical University, Hefei, 230601, People's Republic of China

*These authors contributed equally to this work

Correspondence: Dandan Li; Chunyan Zhu, Email 799200213@qq.com; ayswallow@126.com

Purpose: Higher rates of depression are associated with autistic traits; however, the precise association between autistic traits and depression has yet to be fully elucidated. Good interpersonal emotional regulation and social self-efficacy are crucial for mental health; therefore, in this study, we investigated the relationships between autistic traits, interpersonal emotional regulation, social self-efficacy, and depression.

Patients and Methods: In total, 1024 participants (613 females and 411 males) aged 16 to 23 years old ($M = 19.03$, $SD = 0.95$) completed questionnaires that were designed to evaluate their autistic traits, interpersonal emotion regulation, social self-efficacy and depression. Data were analyzed by Spearman correlation and mediation effects analyses.

Results: Autistic traits were significantly and positively correlated with depression ($r = 0.39$, $p < 0.001$), and autistic traits were shown to positively predict depression ($B = 0.28$, $p < 0.001$). Interpersonal emotion regulation and social self-efficacy were found to play a serial mediating role between autistic traits and depression (indirect effect = 0.020, $p = 0.006$).

Conclusion: This study is a supplement to the mechanism of the relationship between autistic traits and depression. Interpersonal emotion regulation and social self-efficacy are important predictors of possible depression in individuals with high autistic traits. These findings suggest combining interpersonal emotional regulation training and exploring the clinical value of interpersonal emotional regulation interventions in individuals with high autistic traits and autism spectrum disorder, aiming to build social confidence, reduce negative moods, restore social functioning, and other aspects.

Keywords: autistic traits, depression, interpersonal emotion regulation, social self-efficacy, chain mediation

Introduction

People with autism spectrum disorder (ASD) are a form of neurodiversity, and the main challenges they face are negative attitudes towards them, including discrimination and exclusion, and a lack of supportive environments appropriate for ASD.¹ Based on large-scale epidemic and surveillance studies, approximately 1–2% of the general population is estimated to be autistic.²⁻⁴ Autistic traits are distributed across the general population in a normal distribution pattern.^{5,6} Numerous studies have demonstrated that individuals with high autistic traits exhibit similarities with those with ASD in various aspects, including emotions, cognition and behavior. For example, they often exhibit elevated levels

of emotional distress,⁷ limited cognitive flexibility,⁸ and poor social skills.⁹ Based on the high similarity between autistic traits and ASD, studies of autistic traits cannot only help us to understand the mechanisms underlying ASD and develop appropriate intervention strategies, but also help us to improve the mental health and social functionality of groups with high autistic traits.

Approximately 70–80% of individuals with ASD encounter mental health issues.^{10–12} Of these psychiatric concerns, anxiety, irritability, and depression are extremely common,¹³ and mood disorders often co-occur.^{14,15} Individuals with autism co-occurrence with other psychiatric disorders increases the risk of dysfunction, poor life satisfaction, and suicide.^{16,17} One study of adults with autism reported higher levels of depression, which in turn affected their loneliness.¹⁸ Similarly, among a non-clinical community or student population, participants with a higher level of autistic traits have been found to report heightened anxiety and depression,^{19,20} an increased likelihood of being bullied^{21,22} and lower relationship satisfaction.^{23,24} Collectively, these findings demonstrate a strong positive association between autistic traits and depression in both clinical and non-clinical samples. Over recent years, several studies have attempted to investigate the mechanisms underlying autistic traits and depression. A longitudinal study focusing on children with high autistic traits demonstrated that cross-neurotype interaction difficulty was a factor that increased the likelihood of depression.²² A recent study showed that neurodiverse first-semester college students who reported being less satisfied with their social connectedness tended to express more symptoms of depression and anxiety.²⁵ However, the exact pathway of autism traits and depression has yet to be elucidated. Therefore, it is necessary to further investigate the underlying mechanisms of autistic traits and depression if we are to reduce the risk of depression in individuals with high autistic traits and provide theoretical support to develop interventions to improve depression.

Interpersonal emotion regulation (IER) refers to the process by which individuals change their own or others' emotional states through interactions between individuals.²⁶ Numerous studies have shown that individuals with high autistic traits tend to exhibit a weaker ability to regulate their emotions,^{27–29} thus making these individuals more susceptible to depression.^{30,31} However, emotions are social responses.³² Thus, we focus here on emotion regulation that relies on external resources. Unlike the vast majority of individuals who only focus on the emotional regulation of their own emotional states,^{33,34} IER emphasizes the fact that individuals can change their own emotional state, or that of others, via interpersonal interactions.^{35,36} IER involves four distinct factors: soothing, enhancing positive affect, perspective taking, and social modeling.³⁷ Good IER is crucial for the improvement and maintenance of mental health.³⁸

The interpersonal relationship theory of depression³⁹ suggests that individuals who experience more rejection, conflict, and other negative social interactions, tend to develop more negative cognitive patterns and self-evaluations. Moreover, these negative social experiences can lead to the deterioration of self-awareness, ultimately increasing the risk of depression. Autistic people may feel more comfortable with less social engagement due to negative past experiences such as bullying or social exclusion.¹⁸ This may increase their risk of social withdrawal, isolation, and interpersonal relationship problems. When the mindset and social relationships are severely disrupted, individuals with high autistic traits may be unwilling to exploit others to regulate their own emotions due to social difficulties and fewer social connections.⁴⁰ In other words, individuals with high autistic traits may engage in less IER processes to regulate their negative emotions and this difficulty in regulating emotion may further contribute to depressive symptoms in individuals with high autistic traits. Empirical studies have also shown that IER was an important factor for predicting depression and that the enhancement of positive affect was inversely related to the severity of depression.^{41,42} By applying IER, individuals can reduce their own negative emotions and lower their level of depression while helping others to improve their emotions.^{43,44} Therefore, it is important to study and manipulate the process of IER in individuals with autistic traits, and further investigate whether IER mediates the relationship between autistic traits and depression. Furthermore, we are interested in which aspects of depression with high autistic traits can be prevented through effective interpersonal emotion regulation, and clarify the specific mechanisms of action involved; this research could yield suggestions for developing effective interventions.

According to the “double empathy problem” theory, autistic people face the challenge of the cross-neurotype interaction difficulty.⁴⁵ In reference to a discrepancy in reciprocity between autistic individuals and neurotypical individuals, this is likely to lead to interpersonal relationship issues and reduce confidence in social interactions for autistic individuals. Social self-efficacy refers to an individual's cognitive belief in their ability to initiate social

interactions and maintain interpersonal relationships, thus representing a form of social confidence.^{46,47} The temporal need-threat model suggests that social exclusion can threaten an individual's needs for efficacy.⁴⁸ Ren et al⁴⁹ also found that introverted individuals are more likely to exhibit withdrawal behaviors after experiencing rejection, and negative social interactions may lead to a reduction in an individual's social self-efficacy. Furthermore, those with high autistic traits are more likely to experience negative social interactions in their daily life, and also more likely to exhibit social avoidance behaviors, resulting in a series of negative emotions, including social anxiety.⁵⁰ Individuals with social anxiety tend to increasingly doubt their social skills;⁵¹ this leads to decreased levels of self-confidence and a further reduction in social self-efficacy. However, on the one hand, individuals with low social self-efficacy, despite having lower confidence in their social abilities, still have strong social needs and may be more eager to improve their social skills and expand their social circle. On the other hand, they may find it difficult to overcome challenges in real life. This conflict between interpersonal needs and practical difficulties is likely to lead to interpersonal stress and even depression in individuals with high autism traits.⁵² A strong body of evidence has demonstrated that social self-efficacy can positively predict depression.^{53,54} Therefore, we hypothesized that social self-efficacy may also act as a mediating factor between autistic traits and depression.

In addition, the regulation of interpersonal emotion is closely related to self-efficacy; Hofmann et al³⁷ explained individuals' impaired interpersonal emotion regulation, due to lower levels of self-efficacy, can lead to psychological distress. Social self-efficacy is a specific concept of self-efficacy within the interpersonal domain. Based on this, we hypothesized that individuals with high autistic traits may have difficulties in IER, which resulting in low social self-efficacy and subsequently, depressive symptoms. In summary, in the present study, we aimed to investigate the relationship and underlying mechanisms that linked autistic traits and depression, and formulated two hypotheses based on existing literature. First, we hypothesized that autistic traits were positively correlated with depression; second, and we hypothesized that IER and social self-efficacy acted as sequential mediators between autistic traits and depression.

Materials and Methods

Participants

Using convenience sampling, the study conducted a cross-sectional survey in Anhui Province, China, from September to November 2023. Through an online questionnaire platform, the reasons for the study were explained and confidentiality was guaranteed, and college students who were not diagnosed with autism, major depressive disorder, and other psychiatric disorders were surveyed. 1050 questionnaires were collected, of which 26 participants were excluded from the final sample for reasons such as not answering the questionnaire carefully (all items selected the same response), and the unusually short time to complete the survey. The validity of the final questionnaire was 97.52%.

Participants were aged between 16 and 23 years ($M = 19.03$, $SD = 0.95$). Among them, 411 were males and 613 were females. According to grade, 166 were freshmen, 689 were sophomores, and 169 were juniors. Detailed demographic information is shown in Table 1.

Table 1 Demographic Characteristics of the Participants (N = 1024)

Characteristics		Autistic Traits M±SD	Interpersonal Emotion Regulation M±SD	Social Self-efficacy M±SD	Depression M±SD
Sex	Female(n=613)	117.11±10.06	62.38±16.06	57.88±13.31	13.73±10.28
	Male(n=411)	118.66±9.96	63.58±17.20	59.02±14.99	16.52±11.23
Grade	Freshman(n=166)	114.78±9.73	61.84±15.13	60.13±13.43	14.90±9.78
	Sophomore(n=689)	118.21±10.33	63.08±17.36	58.23±14.78	14.86±10.87
	Junior(n=169)	118.67±8.62	62.96±14.27	57.05±12.45	14.73±11.24

Demographic Questionnaire

All participants were asked to fill out basic personal information, including sex, age, and grade. They were also asked for their contact information in preparation for future behavioral experiments.

Autism-Spectrum Quotient (AQ)

The AQ scale was originally developed by Baron-Cohen.⁵ It has five dimensions: social skill, attention switching, attention to detail, communication, and imagination. The AQ includes 50 four-point items (1 = completely disagree to 4 = completely agree). The higher the scores are on this scale, the higher levels of autistic traits.⁵ This scale has demonstrated good reliability and validity in a previous cohort of Chinese students.⁵⁵ The Cronbach's alpha coefficient was 0.720.

Interpersonal Emotion Regulation Questionnaire (IERQ)

The IERQ scale³⁷ was used to assess how individuals take advantage of others to regulate their own emotions. The IERQ features 20 five-point items (1 = not true for me at all to 5 = extremely true for me). The higher the scores are on this scale, the more an individual applies IER strategies.³⁷ The Chinese adaptation of this scale has demonstrated good reliability and validity.⁵⁶ The Cronbach's alpha coefficient was 0.958.

Social Self-Efficacy (SSE)

The SSE scale⁴⁷ was used to assess perceived self-confidence in social interactions. The SSE consists of a single-factor structure. The Chinese version of the SSE has demonstrated good reliability and validity⁵⁷ and consists of 18 five-point items (1 = no confidence at all to 5 = complete confidence). The higher the score are on this scale, the higher an individual's level of perceived social self-efficacy.⁴⁷ The Cronbach's alpha coefficient was 0.962.

Center for Epidemiologic Studies Depression Scale (CES-D)

The CES-D scale⁵⁸ was used to assess current depressive symptoms. The CES-D comprises four factors: depressed affect, positive affect, somatic and retarded activity, and interpersonal problems. The CES-D is a 20 four-point item instrument (0 = seldom or none of the time to 3 = most or all of the time). The higher score on the scale indicates more severe depressive symptoms.⁵⁸ The Chinese version of this scale has demonstrated good reliability and validity in college students.⁵⁹ The Cronbach's alpha coefficient was 0.927.

Statistical Analyses

A Microsoft Excel database was generated by exporting data from the completed questionnaires. After data processing, statistical analysis was conducted utilizing SPSS version 23.0 software and Mplus version 8.3. In SPSS, Harman's single factor test was used to examine whether there were significant common method biases in this study. Pearson's correlations analysis was performed on the variables in order to detect initial bivariate relationship between the variables. In Mplus, two sets of main analyses were used to examine the key hypotheses. The first set of analyses addressed the hypothesized chain mediation model, controlling for the initial levels of demographic covariates (ie, gender). Autistic traits represented the independent variable (X), depression was the dependent variable (Y), IER was the first order mediator (M1), and social self-efficacy was the second order mediator (M2). Drawing on the results of the first set of analyses, our second set of analyses investigated which dimension of depression was influenced by the variables encompassing the entire model. Specifically, four dimensions of depression (depressed affect, positive affect, somatic and retarded activity, and interpersonal problems) were added into the model as dependent variables. A 95% confidence interval (CI) that excluded zero indicates that the parameter estimates are significant.

Results

Common Method Bias

The results of Harman's single factor test indicated that the variance interpretation rate of the first component was 20.52%, which is less than the critical standard requirement of 40%.⁶⁰ Consequently, there is no significant method bias in this study.

Table 2 Descriptive Statistics of Study Variables and Bivariate Correlations Between Variables (N=1024)

	1	2	3	4	5
1.Sex	1				
2.Autistic traits	0.076*	1			
3.Interpersonal emotion regulation	0.036	-0.357***	1		
4.Social self-efficacy	0.040	-0.525***	0.504***	1	
5.Depression	0.127***	0.394***	-0.247***	-0.398***	1
M	0.40	117.73	62.86	58.34	14.85
SD	0.49	10.04	16.53	14.01	10.75


Notes: Sex: 0 = female, 1 = male. * $p < 0.05$, *** $p < 0.001$.

Descriptive Analyses

The descriptive statistical results of each variable and the detailed results of correlation analysis are presented in Table 2. Autistic traits were significantly and positively correlated with depression ($r = 0.39$, $p < 0.001$), but were significantly and negatively correlated with IER ($r = -0.36$, $p < 0.001$) and social self-efficacy ($r = -0.53$, $p < 0.001$). IER ($r = -0.25$, $p < 0.001$) and social self-efficacy ($r = -0.40$, $p < 0.001$) were significantly and negatively correlated with depression. Furthermore, IER was significantly and positively correlated with social self-efficacy ($r = 0.50$, $p < 0.001$). As sex was significantly associated with autistic traits and depression, we controlled for sex in subsequent testing of the model.

The Chain-Mediated Role of Interpersonal Emotion Regulation and Social Self-Efficacy

First, we tested the hypothesized chain mediation model. The model presented an adequate fit to the data: χ^2 (126) = 900.479, $p < 0.001$; $\chi^2/df = 7.147$; CFI = 0.928; TLI = 0.914; RMSEA = 0.077 (90% CI = 0.073–0.082); SRMR = 0.076. In Figure 1, the model's path coefficients are illustrated. Autistic traits negatively predicted IER ($B = -0.46$, $p < 0.001$) and social self-efficacy ($B = -0.59$, $p < 0.001$), but positively predicted depression ($B = 0.28$, $p < 0.001$). Although IER

**Figure 1** The chain mediating model of interpersonal emotion regulation and social self-efficacy on the relationship between autistic traits and depression.

Notes: Standardized coefficients are reported. Sex was included as a covariate in the examination but not explicitly displayed in the figure. ** $p < 0.01$, *** $p < 0.001$.

Table 3 Results Arising from Chain Mediation Analysis of Autistic Traits and Depression (N=1024)

Effect	Path	B	SE	95% CI	p
Direct effect		0.278	0.065	[0.200,0.556]	0.000
Indirect effect	Total indirect effect	0.102	0.044	[0.017,0.192]	0.021
	AT→IER→D	-0.022	0.021	[-0.064,0.016]	0.289
	AT→SSE→D	0.103	0.035	[0.039,0.178]	0.003
	AT→IER→SSE→D	0.020	0.007	[0.008,0.038]	0.006
Total		0.380	0.038	[0.303,0.450]	0.000

Notes: Bootstrap sample size = 5000.

Abbreviations: B, standardized beta (effect); SE, standard error; CI, confidence interval; AT, autistic traits; IER, interpersonal emotion regulation; SSE, social self-efficacy; D, depression.

did not directly predicted depression ($B = 0.05$, $p = 0.28$), IER positively predicted social self-efficacy ($B = 0.25$, $p < 0.001$). In addition, social self-efficacy negatively predicted depression ($B = -0.17$, $p = 0.002$).

Table 3 presents the indirect effects. Analysis indicated that the total indirect effect was significant [$B = 0.102$, $SE = 0.044$, 95% CI (0.017, 0.192), $p = 0.021$], thus accounting for 26.84% of the total effect of autistic traits on depression. The indirect effect of autistic traits through IER and social self-efficacy was also significant [$B = 0.020$, $SE = 0.007$, 95% CI (0.008, 0.038), $p = 0.006$], thus accounting for 5.26% of the total effect of autistic traits on depression. Furthermore, two specific indirect effects were identified in the model. While IER did not mediate association between autistic traits and depression alone, it did explain the variance of depression via social self-efficacy.

Second, the suggested SEM where depressed affect, positive affect, somatic and retarded activity, and interpersonal problems were used as dependent variables. The model exhibited a good fit to the data: $\chi^2(120) = 703.585$, $p < 0.001$; $\chi^2/df = 5.863$; CFI = 0.946; TLI = 0.931; RMSEA = 0.069 (90% CI = 0.064–0.074); SRMR = 0.050. In Figure 2, the model's path coefficients are illustrated. Table 4 shows the results from the four mediation analyses. The serial indirect




Figure 2 The chain mediating model of interpersonal emotion regulation and social self-efficacy on the relationship between autistic traits and depression across four dimensions.

Notes: Standardized coefficients are reported. Sex was included as a covariate in the examination but not explicitly displayed in the figure. ^aautistic traits → depressed affect, ^bautistic traits → positive affect, ^cautistic traits → somatic and retarded activity, ^dautistic traits → interpersonal problems. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Table 4 Results Arising from Chain Mediation Analysis of Autistic Traits and Depression Across Four Dimensions (N=1024)

Model		Path	B	95% CI	p
Depressed affect as the dependent variable	Direct		0.273	[0.210,0.579]	0.000
	Indirect	Total indirect	0.076	[-0.006,0.228]	0.071
		AT→IER→DA	-0.026	[-0.100,0.017]	0.211
		AT→SSE→DA	0.085	[0.032,0.224]	0.012
		AT→IER→SSE→DA	0.017	[0.006,0.047]	0.018
	Total		0.349	[0.384,0.619]	0.000
Positive affect as the dependent variable	Direct		0.234	[0.115,0.342]	0.000
	Indirect	Total indirect	0.201	[0.118,0.273]	0.000
		AT→IER→PA	0.113	[0.073,0.151]	0.000
		AT→SSE→PA	0.074	[0.009,0.136]	0.026
		AT→IER→SSE→PA	0.015	[0.002,0.029]	0.033
	Total		0.435	[0.347,0.504]	0.000
Somatic and retarded activity as the dependent variable	Direct		0.257	[0.138,0.418]	0.000
	Indirect	Total indirect	0.096	[0.013,0.193]	0.026
		AT→IER→SRA	-0.024	[-0.072,0.014]	0.227
		AT→SSE→SRA	0.100	[0.041,0.181]	0.003
		AT→IER→SSE→SRA	0.020	[0.008,0.040]	0.006
	Total		0.353	[0.298,0.472]	0.000
Interpersonal problems as the dependent variable	Direct		0.280	[0.051,0.163]	0.288
	Indirect	Total indirect	0.049	[-0.015,0.057]	0.046
		AT→IER→IP	-0.036	[-0.030,0.000]	0.062
		AT→SSE→IP	0.071	[0.001,0.057]	0.046
		AT→IER→SSE→IP	0.014	[0.000,0.012]	0.058
	total		0.330	[0.096,0.164]	0.000

Note: Bootstrap sample size = 5000.

Abbreviations: B, standardized beta (effect); CI, confidence interval; AT, autistic traits; IER, interpersonal emotion regulation; SSE, social self-efficacy; DA, depressed affect; PA, positive affect; SRA, somatic and retarded activity; IP, interpersonal problems.

effects and the indirect effects mediated by social self-efficacy were significant across all models. Specifically, in the model that used positive affect as the dependent variable, the indirect effect of IER as the mediating variable was significant [$B = 0.113$, $SE = 0.020$, 95% CI (0.073, 0.151), $p < 0.001$]; this effect was not significant in other models.

Discussion

The study investigated the relationship between autistic traits and depression, as well as the potential mechanisms that mediate this relationship. The findings indicated that IER and social self-efficacy played a serial mediating role between autistic traits and depression. This finding showed that individuals with high autistic traits may have difficulty in social interaction, thus resulting in the reduced use of IER strategies based on interpersonal interaction to cope with negative social interaction experiences. In contrast, more negative experiences will lead to a reduction in their confidence in social situations, and may even lead to more severe depressive symptoms.

First, we identified a significant positive correlation between autistic traits and depressive mood; this was consistent with previous research findings.^{7,61} On the one hand, individuals with high autistic traits had a strong need for social adaptation but had difficulty interacting with others and missed out on many opportunities to socialize, thus making it difficult for these individuals to build and maintain social relationships and support networks;⁶² this may lead to a depressive mood. On the other hand, individuals with autistic traits immersed themselves in their own world and had difficulty with attention switching.⁶³ This could explain why individuals with higher autistic traits are more likely to develop mental disorders. For example, a study investigating the relationship between autistic traits and obsessive-

compulsive disorder (OCD) showed that autistic traits in people with OCD were higher than those in healthy people and it was associated with attentional impulsivity.⁶⁴ Besides, their limited imagination and poor cognitive flexibility also affected their ability to cope effectively with life challenges; this may further exacerbate depressive mood.⁶⁵

The regulation of emotion is closely related to an individual's social competence and mental health,^{66,67} and is of great significance for social and environmental adaptation. Previous research showed that individuals with higher autistic traits tended to have a poorer ability to regulate emotion.²⁷ However, contemporary emotion regulation research emphasizes intrapersonal processes such as cognitive reappraisal and expressive suppression, but people experiencing affect commonly choose not to go it alone. An emerging literature highlighted the importance of IER,³⁶ which utilized social cues to facilitate the regulation of emotional states. Few studies have explored the relationship between autistic traits and IER. Our present findings revealed a distinct relationship between autistic traits and IER; specifically, autistic traits were negatively correlated with IER. This may be because individuals with high autistic traits are difficult to understand by others and therefore less likely to participate in interpersonal interactions.¹⁸ Previous research has shown that individuals with high autistic traits have difficulties in emotion recognition and also exhibit lower levels of empathy.^{68,69} The introspection-centric simulation theory posits that the reduced cognitive empathy associated with autistic traits may stem from difficulties in alexithymia, specifically the challenge individuals face in understanding their own emotions, which subsequently leads to difficulties in understanding the emotions of others.⁷⁰ This could also explain the difficulties in IER experienced by individuals with high autistic traits.

Although most studies demonstrated that IER was correlated with depression,^{71,72} the path from IER to depression was not significant when social self-efficacy was added to the chain mediation model of the impact of autistic traits on depression. IER can affect a regulated individual's feelings of hope, calmness, and happiness,^{38,73} while the regulator may also experience less happiness⁷⁴ and more negative emotions⁷⁵ due to the inability to achieve regulatory goals or in response to the suffering of others. However, when we incorporated the four dimensions of depression into our model, we found that autistic traits only influenced positive emotions related to depression through IER. Previous research showed that IER was associated with a reduction in feelings of happiness.^{76,77} Furthermore, research has also shown that increasing positive emotions in individuals with depression could promote their recovery of social functionality and lead to a greater sense of happiness.⁷⁸ Similarly, previous research on IER training has shown that participants who helped others regulated their emotions to a greater extent during a three-week training reported a more significant increase in subjective well-being.⁷⁹ This highlights the importance of manipulating IER to increase positive emotions in order to prevent depression in individuals with high autistic traits.

The model described herein demonstrated that social self-efficacy mediated both IER and depression. We found that IER was a positive predictor of social self-efficacy, and that social self-efficacy was a negative predictor of depression. This could be because successful communication experiences between family members and their peers could enhance an individual's evaluation of their social abilities, while individuals with high social self-efficacy could effectively use interpersonal skills to establish relationships with others to derive positive emotional experiences.^{80,81} IER is an important method by which we can maintain and develop social relationships.³⁸ Previous research demonstrated that the daily frequency of IER strategies can positively predict the quality of interpersonal relationships.⁸²

Furthermore, the ability of college students to perform IER can predict the establishment and development of their social relationships during their time in school.⁸³ The interpersonal relationship theory of depression posits that individuals who experience negative interpersonal experiences often develop more negative self-evaluations, and that the deterioration of self-awareness due to negative interpersonal experiences continues to intensify, ultimately increasing the likelihood of depression.³⁹ Our present findings suggested that successful experiences of individuals with high autistic traits in regulating negative emotions through interpersonal interactions were likely to enhance an individual's social confidence, thus protecting them from depression.

Finally, our analyses indicated that social self-efficacy mediated the relationship between autistic traits and depression. This finding concurred with the results of Camus,⁵⁴ who detected a negative correlation between social self-efficacy and depressive symptoms in individuals with autism. Individuals with high autistic traits are associated with lower levels of satisfaction with interpersonal relationships;²⁴ this is likely to result in social problems, including social anxiety and low levels of social self-efficacy. Social self-efficacy is universally important for an individual's mental health, and social self-

efficacy is closely associated with internalizing behaviors such as depression,^{84,85} loneliness^{86,87} and social anxiety.⁵¹ In general, individuals with high autistic traits are not understood and ostracized in social interactions; this may lead to long-term interpersonal problems and poor social adaptation, thus resulting in low self-esteem. The contradiction between high social adaptation needs and social exclusion makes individuals with autistic traits more prone to depression. This suggests that we should pay close attention to the social self-efficacy of individuals with high autistic traits, prompt society toward reciprocity, help them build confidence in social interactions, and actively participate in social activities.

Limitations and Implications

This study has several limitations. First, we only recruited college students in Anhui Province, China, which could result in a generalizability issue; future research should be designed using a representative national sample. Secondly, the data utilized in this study were cross-sectional, implying that our findings may suggest predictive trends rather than definitive causal linkages. To uncover actual causal relationships among these variables, future research could adopt a longitudinal design or experimental approaches. Finally, this study focused on individuals with high levels of autistic traits; future research should consider whether these results can be generalized to the ASD population. Given the genetic, cognitive, and behavioral similarities between autistic traits and ASD, the results of this study could advance our understanding of the mechanisms underlying the impact of ASD on depression and the development of appropriate interventional strategies. However, high levels of autistic traits do not equate to ASD.⁸⁸ Future research should further validate the relationships between ASD symptoms, IER, social self-efficacy, and depression within the ASD population.

Despite these limitations, this study has important implications for the prevention and intervention of depression in individuals with high autistic traits. The mediating effect of IER between autistic traits and positive affect suggests that IER may provide mental health benefits for individuals with high autistic traits. It's interesting that this benefit refers to influencing the positive mood of depression without affecting other aspects of depression. Many studies have highlighted the importance of paying attention to positive emotions in people with mental illness.^{89,90} Therefore, focusing on interpersonal emotion regulation training may be beneficial for enhancing positive emotions in individuals with higher autistic traits. Additionally, this is the first study assessing the pathway between autistic traits and depression via IER and social self-efficacy. The results of this study suggest that people with higher autistic traits are in with a higher chance of depression, which is mediated by a pathway from IER to social self-efficacy, and ultimately to depression. This study has identified potential factors for preventing depression in people with high autistic traits. Interventions targeting IER may enhance social self-efficacy and thereby reduce depressive symptoms. IER is malleable and a recent study showed that IER may be a promising avenue for effective emotion regulation in current major depressive disorder.⁹¹ One possible avenue for future research would be to develop IER training and examine whether they provide mental health benefits for autistic persons.

Conclusion

This study shows relationships between autistic traits, interpersonal emotion regulation, social self-efficacy and depression. Interpersonal emotion regulation and social self-efficacy play a mediating role between autistic traits and depression. These findings expand the underlying mechanisms between autistic traits and depression, which could promote the recovery of positive emotions in individuals with depression. In addition, these results suggest that when preventing the risk of depression in individuals with high autistic traits, appropriate interventions could focus on improving IER, enhancing social self-efficacy, and promoting positive emotions.

Data Sharing Statement

Due to reasons of data protection, the preprocessed data is available only upon reasonable request to the corresponding author (Chunyan Zhu).

Ethics Approval and Informed Consent

All the methods were performed in accordance with the Declaration of Helsinki. The study was approved by the Ethical Committee of Anhui Medical University. All the participants provided written informed consent, and for minors, consent was obtained from their legal guardian(s).

Acknowledgments

We would like to thank all participants.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This research was funded by the National Natural Science Foundation of China (grant number 32271134, 91432301), the National Key Research and Development Program (program number 2021YFC3300500) and the University Synergy Innovation Program of Anhui Province (program number GXXT-2021-003).

Disclosure

The authors declare no conflicts of interest in this work.

References

- Botha M, Chapman R, Giwa Onaiwu M, Kapp SK, Stannard Ashley A, Walker N. The neurodiversity concept was developed collectively: an overdue correction on the origins of neurodiversity theory. *Autism*. 2024;28(6):1591–1594. doi:10.1177/13623613241237871
- Christensen DL, Baio J, Van Naarden Braun K, et al. Prevalence and characteristics of autism spectrum disorder among children aged 8 years--autism and developmental disabilities monitoring network, 11 sites, United States, 2012. *MMWR Surveill Summ*. 2016;65(3):1–23. doi:10.15585/mmwr.ss6503a1
- Elsabbagh M, Divan G, Koh YJ, et al. Global prevalence of autism and other pervasive developmental disorders. *Autism Res*. 2012;5(3):160–179. doi:10.1002/aur.239
- Roman-Urrestarazu A, van Kessel R, Allison C, Matthews FE, Brayne C, Baron-Cohen S. Association of race/ethnicity and social disadvantage with autism prevalence in 7 million school children in England. *JAMA Pediatr*. 2021;175(6):e210054. doi:10.1001/jamapediatrics.2021.0054
- Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The autism-spectrum quotient (AQ): evidence from asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *J Autism Dev Disord*. 2001;31(1):5–17. doi:10.1023/a:1005653411471
- Ronald A, Happé F, Price TS, Baron-Cohen S, Plomin R. Phenotypic and genetic overlap between autistic traits at the extremes of the general population. *J Am Acad Child Adolesc Psychiatry*. 2006;45(10):1206–1214. doi:10.1097/01.chi.0000230165.54117.41
- Liew SM, Thevaraja N, Hong RY, Magiati I. The relationship between autistic traits and social anxiety, worry, obsessive-compulsive, and depressive symptoms: specific and non-specific mediators in a student sample. *J Autism Dev Disord*. 2015;45(3):858–872. doi:10.1007/s10803-014-2238-z
- Gökçen E, Petrides KV, Hudry K, Frederickson N, Smillie LD. Sub-threshold autism traits: the role of trait emotional intelligence and cognitive flexibility. *Br J Psychol*. 2014;105(2):187–199. doi:10.1111/bjop.12033
- Demizu Y, Matsumoto J, Yasuda Y, et al. Relationship between autistic traits and social functioning in healthy individuals. *Neuropsychopharmacol Rep*. 2022;42(2):226–229. doi:10.1002/npr2.12249
- Lever AG, Geurts HM. Psychiatric co-occurring symptoms and disorders in young, middle-aged, and older adults with autism spectrum disorder. *J Autism Dev Disord*. 2016;46(6):1916–1930. doi:10.1007/s10803-016-2722-8
- Roy M, Prox-Vagedes V, Ohlmeier MD, Dillo W. Beyond childhood: psychiatric comorbidities and social background of adults with Asperger syndrome. *Psychiatria Danubina*. 2015;27(1):50–59.
- Simonoff E, Pickles A, Charman T, Chandler S, Baird G, Baird G. Psychiatric disorders in children with autism spectrum disorders: prevalence, comorbidity, and associated factors in a population-derived sample. *J Am Acad Child Adolesc Psychiatry*. 2008;47(8):921–929. doi:10.1097/CHI.0b013e318179964f
- Guerrera S, Menghini D, Napoli E, Di Vara S, Valeri G, Vicari S. Assessment of psychopathological comorbidities in children and adolescents with autism spectrum disorder using the child behavior checklist. *Front Psychiatry*. 2019;10:535. doi:10.3389/fpsy.2019.00535
- Hossain MM, Khan N, Sultana A, et al. Prevalence of comorbid psychiatric disorders among people with autism spectrum disorder: an umbrella review of systematic reviews and meta-analyses. *Psychiatry Res*. 2020;287:112922. doi:10.1016/j.psychres.2020.112922
- Rosen TE, Mazefsky CA, Vasa RA, Lerner MD. Co-occurring psychiatric conditions in autism spectrum disorder. *Int Rev Psychiatry*. 2018;30(1):40–61. doi:10.1080/09540261.2018.1450229

16. Dell'Osso L, Carpita B, Muti D, et al. Mood symptoms and suicidality across the autism spectrum. *Compr Psychiatry*. 2019;91:34–38. doi:10.1016/j.comppsy.2019.03.004
17. Park SH, Song YJC, Demetriou EA, et al. Disability, functioning, and quality of life among treatment-seeking young autistic adults and its relation to depression, anxiety, and stress. *Autism*. 2019;23(7):1675–1686. doi:10.1177/1362361318823925
18. Ee D, Hwang YIJ, Reppermund S, et al. Loneliness in adults on the autism spectrum. *Autism Adulthood*. 2019;1(3):182–193. doi:10.1089/aut.2018.0038
19. Albantakis L, Brandi ML, Zillekens IC, Henco L, Schilbach L. Alexithymic and autistic traits: relevance for comorbid depression and social phobia in adults with and without autism spectrum disorder. *Autism*. 2020;24(8):136236132093602. doi:10.1177/1362361320936024
20. Kanne SM, Christ SE, Reiersen AM. Psychiatric symptoms and psychosocial difficulties in young adults with autistic traits. *J Autism Dev Disord*. 2009;39(6):827–833. doi:10.1007/s10803-008-0688-x
21. Kunihiro Y, Senju A, Dairoku H, Wakabayashi A, Hasegawa T. 'Autistic' traits in non-autistic Japanese populations: relationships with personality traits and cognitive ability. *J Autism Dev Disord*. 2006;36(4):553–566. doi:10.1007/s10803-006-0094-1
22. Rai D, Culpin I, Heuvelman H, et al. Association of autistic traits with depression from childhood to age 18 years. *J Am Med Assoc*. 2018;75(8):835–843.
23. Pollmann MM, Finkenauer C, Begeer S. Mediators of the link between autistic traits and relationship satisfaction in a non-clinical sample. *J Autism Dev Disord*. 2010;40(4):470–478. doi:10.1007/s10803-009-0888-z
24. De Groot K, Van Strien JW. Self-report and brain indicators of impaired emotion regulation in the broad autism spectrum. *J Autism Dev Disord*. 2017;47(7):2138–2152.
25. McKenney EE, Richards JK, Day TC, et al. Satisfaction with social connectedness is associated with depression and anxiety symptoms in neurodiverse first-semester college students. *Autism*. 2024;28(8):1972–1984. doi:10.1177/13623613231216879
26. Dixon-Gordon KL, Bernecker SL, Christensen K. Recent innovations in the field of interpersonal emotion regulation. *Curr Opin Psychol*. 2015;3:36–42. doi:10.1016/j.copsyc.2015.02.001
27. Liu S, Yu C, Conner BT, Wang S, Lai W, Zhang W. Autistic traits and internet gaming addiction in Chinese children: the mediating effect of emotion regulation and school connectedness. *Res Dev Disabil*. 2017;68:122–130. doi:10.1016/j.ridd.2017.07.011
28. Rinaldi LJ, Simmer J, Koursarou S, Ward J. Autistic traits, emotion regulation, and sensory sensitivities in children and adults with misophonia. *J Autism Dev Disord*. 2023;53(3):1162–1174. doi:10.1007/s10803-022-05623-x
29. Zhao X, Li X, Song Y, Li C, Shi W. Autistic traits and emotional experiences in Chinese college students: mediating role of emotional regulation and sex differences. *Res Autism Spectr Disord*. 2020;77:101607. doi:10.1016/j.rasd.2020.101607
30. Liu DY, Thompson RJ. Selection and implementation of emotion regulation strategies in major depressive disorder: an integrative review. *Clin Psychol Rev*. 2017;57:183–194. doi:10.1016/j.cpr.2017.07.004
31. Young KS, Sandman CF, Craske MG. Positive and negative emotion regulation in adolescence: links to anxiety and depression. *Brain Sci*. 2019;9(4):76. doi:10.3390/brainsci9040076
32. Parkinson B. Emotions are social. *Br J Psychol*. 2011;87.
33. Li S, Xie H, Zheng Z, et al. The causal role of the bilateral ventrolateral prefrontal cortices on emotion regulation of social feedback. *Hum Brain Mapp*. 2022;43(9):2898–2910. doi:10.1002/hbm.25824
34. Zhao J, Mo L, Bi R, et al. The VLPFC versus the DLPFC in downregulating social pain using reappraisal and distraction strategies. *J Neurosci*. 2021;41(6):1331–1339. doi:10.1523/jneurosci.1906-20.2020
35. Morawetz C, Riedel MC, Salo T, et al. Multiple large-scale neural networks underlying emotion regulation. *Neurosci Biobehav Rev*. 2020;116:382–395. doi:10.1016/j.neubiorev.2020.07.001
36. Zaki J, Williams WC. Interpersonal emotion regulation. *Emotion*. 2013;13(5):803–810. doi:10.1037/a0033839
37. Hofmann SG, Carpenter JK, Curtiss J. Interpersonal emotion regulation questionnaire (IERQ): scale development and psychometric characteristics. *Cognit Ther Res*. 2016;40(3):341–356. doi:10.1007/s10608-016-9756-2
38. Zaki J. Integrating Empathy and interpersonal emotion regulation. *Annu Rev Psychol*. 2020;71(1):517–540. doi:10.1146/annurev-psych-010419-050830
39. Anderson JC, McGee R. Child and Adolescent Mental Health. In: *Handbook of Depression in Children and Adolescents*. 2009.
40. Stice LV, Lavner JA. Social connectedness and loneliness mediate the association between autistic traits and internalizing symptoms among young adults. *J Autism Dev Disord*. 2019;49(3):1096–1110. doi:10.1007/s10803-018-3812-6
41. Koç MS, Aka BT, Doğruyol B, Curtiss J, Carpenter JK, Hofmann SG. Psychometric properties of the Turkish version of the interpersonal emotion regulation questionnaire (IERQ). *J Psychopathol Behav Assess*. 2019;41(2):294–303. doi:10.1007/s10862-019-09732-3
42. Ray-Yol E, Altan-Atalay A. Interpersonal emotion regulation and psychological distress: what is the function of negative mood regulation expectancies in this relationship? *Psychol Rep*. 2022;125(1):280–293. doi:10.1177/0033294120968086
43. Doré BP, Morris RR, Burr DA, Picard RW, Ochsner KN. Helping others regulate emotion predicts increased regulation of one's own emotions and decreased symptoms of depression. *Pers Soc Psychol Bull*. 2017;43(5):729–739. doi:10.1177/0146167217695558
44. Guendelman S, Bayer M, Prehn K, Dziobek I. Regulating negative emotions of others reduces own stress: neurobiological correlates and the role of individual differences in empathy. *Neuroimage*. 2022;254:119134. doi:10.1016/j.neuroimage.2022.119134
45. Milton DEM. On the ontological status of autism: the 'double empathy problem'. *Disabil Soc*. 2012;27(6):883–887. doi:10.1080/09687599.2012.710008
46. Alessandri G, Caprara GV, Eisenberg N, Steca P. Reciprocal relations among self-efficacy beliefs and prosociality across time. *J Pers*. 2009;77:1229–1259.
47. Smith HM, Betz NE. Development and validation of a scale of perceived social self-efficacy. *J Career Assess*. 2000;8(3):283–301. doi:10.1177/10690727000800306
48. Williams KD. Ostracism. *Annu Rev Psychol*. 2007;58(1):425–452. doi:10.1146/annurev.psych.58.110405.085641
49. Ren D, Wesselmann E, Williams KD. Evidence for another response to ostracism: solitude seeking. *Soc Psychol Pers Sci*. 2016;7:1948550615616169.
50. Wainer AL, Block N, Donnellan MB, Ingersoll B. The broader autism phenotype and friendships in non-clinical dyads. *J Autism Dev Disord*. 2013;43(10):2418–2425. doi:10.1007/s10803-013-1789-8

51. Aune T, Juul EML, Beidel DC, Nordahl HM, Dvorak RD. Mitigating adolescent social anxiety symptoms: the effects of social support and social self-efficacy in findings from the Young-HUNT 3 study. *Eur Child Adolesc Psychiatry*. 2021;30(3):441–449. doi:10.1007/s00787-020-01529-0
52. Losiewicz OM, Metts AV, Zinbarg RE, Hammen C, Craske MG. Examining the indirect contributions of irritability and chronic interpersonal stress on symptoms of anxiety and depression in adolescents. *J Affect Disord*. 2023;329:350–358. doi:10.1016/j.jad.2023.02.125
53. Wei M, Russell DW, Zakalik RA. Adult attachment, social self-efficacy, self-disclosure, loneliness, and subsequent depression for freshman college students: a longitudinal study. *J Couns Psychol*. 2005;52(4):602–614. doi:10.1037/0022-0167.52.4.602
54. Camus L, Jones K, O'Dowd E, Auyeung B, Rajendran G, Stewart ME. Autistic traits and psychosocial predictors of depressive symptoms. *J Autism Dev Disord*. 2024. doi:10.1007/s10803-024-06361-y
55. Zhang L, Sun Y, Chen F, et al. Psychometric properties of the autism-spectrum quotient in both clinical and non-clinical samples: Chinese version for mainland China. *BMC Psychiatry*. 2016;16(1):213. doi:10.1186/s12888-016-0915-5
56. Chen C, Liu X, Song J, Shen Y, Jin F, Niu Y. Reliability and validity of the interpersonal emotion regulation questionnaire in Chinese college students. *Chin J Health Psychol*. 2021;29(11):1684–1688.
57. Fan J, Meng H, Lopez F. Validation of an American social self-efficacy inventory in Chinese population. *Presented in the 21st Annual Conference of the SIOP*; 2006.
58. Radloff LS. The CES-D scale a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1(3):385–401. doi:10.1177/014662167700100306
59. Jiang L, Wang Y, Zhang Y, et al. The reliability and validity of the center for epidemiologic studies depression scale (CES-D) for Chinese university students. *Front Psychiatry*. 2019;10:315. doi:10.3389/fpsy.2019.00315
60. Podsakoff PM, MacKenzie SB, Lee JY, Podsakoff NP. Common method biases in behavioral research: a critical review of the literature and recommended remedies. *J Appl Psychol*. 2003;88(5):879–903. doi:10.1037/0021-9010.88.5.879
61. Galvin J, Richards G. The indirect effect of self-compassion in the association between autistic traits and anxiety/depression: a cross-sectional study in autistic and non-autistic adults. *Autism*. 2023;27(5):1256–1270. doi:10.1177/13623613221132109
62. Takara K, Kondo T. Comorbid atypical autistic traits as a potential risk factor for suicide attempts among adult depressed patients: a case-control study. *Ann Gen Psychiatry*. 2014;13(1):33. doi:10.1186/s12991-014-0033-z
63. Pelton MK, Cassidy SA. Are autistic traits associated with suicidality? A test of the interpersonal-psychological theory of suicide in a non-clinical young adult sample. *Autism Res*. 2017;10(11):1891–1904. doi:10.1002/aur.1828
64. Namlı Z, Tamam L, Demirkol ME, Karaytuğ MO, Sun T. The relationship among autistic traits, impulsivity, and functionality in patients with obsessive-compulsive disorder. *J Nerv Ment Dis*. 2023;211(3):195–202. doi:10.1097/nmd.0000000000001601
65. Parsi K, Elster N. A life of one's own: challenges in the transition from childhood to adulthood with autism spectrum disorder. *AMA J Ethics*. 2015;17(4):342–347. doi:10.1001/journalofethics.2015.17.4.pfor1-1504
66. Benavides-Nieto A, Romero-López M, Quesada-Conde AB, Corredor GA. Basic executive functions in early childhood education and their relationship with social competence. *Procedia*. 2017;237:471–478. doi:10.1016/j.sbspro.2017.02.092
67. Hassan R, Schmidt LA. Inhibitory control, dyadic social behavior, and mental health difficulties in preschoolers. *Child Dev*. 2022;93(3):e251–e265. doi:10.1111/cdev.13725
68. Marotta A, Aranda-Martín B, De Cono M, Ballesteros-Duperón M, Casagrande M, Lupiáñez J. Integration of facial expression and gaze direction in individuals with a high level of autistic traits. *Int J Environ Res Public Health*. 2022;19(5):2798. doi:10.3390/ijerph19052798
69. Martin R, Mckenzie K, Metcalfe D, Pollet T, Mccarty K. A preliminary investigation into the relationship between empathy, autistic like traits and emotion recognition. *Pers Individ Differ*. 2018;137:12–16. doi:10.1016/j.paid.2018.07.039
70. Goldman AI. *Simulating Minds: The Philosophy, Psychology, and Neuroscience of Mindreading*. Oxford University Press; 2006.
71. Altan-Atalay A, Saritas-Atalar D. Interpersonal emotion regulation strategies: how do they interact with negative mood regulation expectancies in explaining anxiety and depression? *Cur Psychol*. 2019;41:1–7.
72. Gökdağ C. How does interpersonal emotion regulation explain psychological distress? The roles of attachment style and social support. *Pers Individ Dif*. 2021;176:110763. doi:10.1016/j.paid.2021.110763
73. Niven K, Totterdell P, Holman D, Headley T. Does regulating others' feelings influence people's own affective well-being? *J Soc Psychol*. 2012;152(2):246–260. doi:10.1080/00224545.2011.599823
74. Henkel AP, Bromuri S, Iren D, Urovi V. Half human, half machine – augmenting service employees with AI for interpersonal emotion regulation. *J Serv Manage*. 2020;31(2):247–265. doi:10.1108/JOSM-05-2019-0160
75. Brown CL, West TV, Sanchez AH, Mendes WB. Emotional empathy in the social regulation of distress: a dyadic approach. *Pers Soc Psychol Bull*. 2020;47(5):146167220953987.
76. Cheung EO, Gardner WL, Anderson JF. Emotionships: examining people's emotion-regulation relationships and their consequences for well-being. *Soc Psychol Pers Sci*. 2015;6(4):407. doi:10.1177/1948550614564223
77. Niven K, Macdonald I, Holman D. You spin me right round: cross-relationship variability in interpersonal emotion regulation. *Front Psychol*. 2012;3:3. doi:10.3389/fpsyg.2012.00394
78. Dunn BD, Widnall E, Reed N, Owens C, Kuyken W, Kuyken W. Bringing light into darkness: a multiple baseline mixed methods case series evaluation of Augmented Depression Therapy (ADepT). *Behav Res Ther*. 2019;120(103418):103418. doi:10.1016/j.brat.2019.103418
79. Doré BP, Morris RR, Burr DA, Picard RW, Ochsner KN. Helping others regulate emotion predicts increased regulation of one's own emotions and decreased symptoms of depression. *Pers Soc Psychol Bull*. 2017;43(5):014616721769555. doi:10.1177/0146167217695558
80. Fan J, Meng H, Zhao B, Patel T. Further Validation of a U.S. Adult social self-efficacy inventory in Chinese populations. *J Career Assess*. 2012;20(4):463–478. doi:10.1177/1069072712450006
81. Minter A, Pritzker S. Measuring adolescent social and academic self-efficacy: cross-ethnic validity of the SEQ-C. *Res Soc Work Pract*. 2017;27(7):818–826. doi:10.1177/1049731515615677
82. Chan S, Rawana JS. Examining the associations between interpersonal emotion regulation and psychosocial adjustment in emerging adulthood. *Cogn Ther Res*. 2021;45(4):652–662. doi:10.1007/s10608-020-10185-2
83. Williams WC, Morelli SA, Ong DC, Zaki J. Interpersonal emotion regulation: implications for affiliation, perceived support, relationships, and well-being. *J Pers Soc Psychol*. 2018;115(2):224–254. doi:10.1037/pspi0000132

84. Niu GF, Shi XH, Yao LS, Yang WC, Jin SY, Xu L. Social exclusion and depression among undergraduate students: the mediating roles of rejection sensitivity and social self-efficacy. *Curr Psychol*. 2022;42:1–10. doi:10.1007/s12144-022-03318-1
85. Sharma PK, Kumra R. Relationship between mindfulness, depression, anxiety and stress: mediating role of self-efficacy. *Pers Individ Differ*. 2022;186:111363. doi:10.1016/j.paid.2021.111363
86. Erozkan A, Deniz S. The Influence of social self-efficacy and learned resourcefulness on loneliness. *Online J Couns Educ*. 2012;1(2):57–74.
87. Gazo AM, Mahasneh AM, Abood MH, Muhediat FA. Social self-efficacy and its relationship to loneliness and internet addiction among hashemite university students. *Int J High Educ*. 2020;9(2):144. doi:10.5430/ijhe.v9n2p144
88. Sasson NJ, Bottema-Beutel K. Studies of autistic traits in the general population are not studies of autism. *Autism*. 2022;26(4):1007–1008. doi:10.1177/13623613211058515
89. Sanislow CA, Ferrante M, Pacheco J, Rudorfer MV, Morris SE. Advancing translational research using NIMH research domain criteria and computational methods. *Neuron*. 2019;101(5):779–782. doi:10.1016/j.neuron.2019.02.024
90. Straszewski T, Siegel JT. Differential effects of high- and low-arousal positive emotions on help-seeking for depression. *Appl Psychol Health Well Being*. 2020;12(3):887–906. doi:10.1111/aphw.12214
91. Liu DY, Strube MJ, Thompson RJ. Do emotion regulation difficulties in depression extend to social context? Everyday interpersonal emotion regulation in current and remitted major depressive disorder. *J Psychopathol Clin Sci*. 2024;133(1):61–75. doi:10.1037/abn0000877

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>