



Advances in the Treatment of Rare Mutations in Non-Small Cell Lung Cancer

Yanning Sun*, Li Ma , Xiaofei Zhang*, Zhaoxia Wang 

Cancer Medical Center, The Second Affiliated Hospital of Nanjing Medical University, Nanjing, People's Republic of China

*These authors contributed equally to this work

Correspondence: Zhaoxia Wang, Cancer Medical Center, The Second Affiliated Hospital of Nanjing Medical University, Nanjing, 210011, People's Republic of China, Email wangzhaoxia@njmu.edu.cn

Abstract: Lung cancer is a malignant tumor with the highest morbidity and mortality rate worldwide, with nearly 2.5 million new cases and more than 1.8 million deaths reported globally in 2022. Lung cancer is broadly categorized into two main types: non-small cell lung cancer (NSCLC) and small cell lung cancer (SCLC), with NSCLC accounting for about 85% of all cases. Early-stage lung cancers often present without obvious symptoms, resulting in most patients being diagnosed at an advanced stage where traditional chemotherapy has limited efficacy. Recent advances in molecular biology have elucidated the pivotal role of gene mutations in tumor development, paving the way for targeted therapies that have markedly benefited patients. Beyond the well-known epidermal growth factor receptor (EGFR) mutation, an increasing number of new molecular targets have been identified, including ROS1 rearrangement, BRAF mutation, NTRK fusion, RET fusion, MET mutation, KRAS G12C mutation, HER2 mutation, ALK rearrangement, and NRG1 fusion. Some of these targeted therapies have already been approved by the Food and Drug Administration (FDA), and many others are currently undergoing clinical trials. This review summarizes recent advances in NSCLC treatment with molecular targets, highlighting progress, challenges, and their impact on patient prognosis.

Keywords: non-small cell lung cancer, rare mutations, targeted therapy, immunotherapy

Introduction

Lung cancer is the malignant tumor with the highest morbidity and mortality rates worldwide, with 2.48 million new cases and 1.82 million deaths reported globally in 2022. The overall five-year survival rate for patients with advanced lung cancer is often less than 20%.¹ Lung cancer can be classified into non-small cell lung cancer (NSCLC) and small cell lung cancer (SCLC) according to the histological type, accounting for approximately 85% and 15%, respectively. NSCLC is further subdivided into adenocarcinoma (40%), squamous cell carcinoma (25–30%), and large cell carcinoma (5–10%).²

Traditionally, lung cancer has been treated with surgery and chemotherapy, and approximately 30% of patients with NSCLC are eligible for surgery. However, the incidence of postoperative recurrence or metastasis is as high as 21.7%, even in patients with completely resected stage I.³ Similarly, chemotherapy has limited effectiveness, with a median survival of less than 9 months, a response rate of 25–35%, and a 2-year survival rate of only 10%.⁴ In 2003, the FDA approved gefitinib as a first-line treatment for metastatic NSCLC with epidermal growth factor receptor (EGFR) mutations. A Phase III trial comparing the efficacy of gefitinib and platinum-based combinations showed that the 12-month progression-free survival (PFS) rate was 24.9% in the gefitinib group, compared to 6.7% in the carboplatin-paclitaxel group. This demonstrated that targeted therapy is superior to carboplatin combined with paclitaxel as an initial treatment for patients with EGFR-positive NSCLC, while gefitinib also exhibited lower toxicity in these patients.⁵ As a result, the National Comprehensive Cancer Network (NCCN) guidelines recommend targeted therapy as a first-line treatment option for driver-positive patients receiving palliative care, marking the beginning of the targeted therapy era for advanced NSCLC.⁶ Defining molecular subtypes is essential for targeted therapy in NSCLC, and the application of next-generation sequencing (NGS) has created opportunities to identify rare gene mutations. In recent years, in-depth

research into the molecular biology of lung cancer and the development of targeted drugs have comprehensively transformed the treatment of advanced lung cancer. About 30%–50% of NSCLC patients have oncogenic driver alterations, and approximately 50%–75% of these patients can receive targeted therapy.⁴

With advances in detection technology, molecular targets beyond EGFR have been gradually identified. Oncogenic targets with an incidence of $\leq 5\%$ are classified as rare targets.⁷ Although the incidence of rare mutations is low compared to common targets like EGFR, the large population of lung cancer patients means the actual number of those with rare mutations is significant. These patients often face challenges, including the lack of targeted therapies, poor response to standard chemotherapy, poor prognosis, and increased susceptibility to recurrence and metastasis. Additionally, the low frequency of these mutations complicates the ability to conduct large-scale randomized clinical trials. Currently, targeted therapies for rare mutations primarily include small-molecule drugs and monoclonal antibodies, with kinase inhibitors representing the main class of small-molecule drugs. Kinases can be categorized into tyrosine kinases (eg, EGFR, HER2, MET, ALK, RET, ROS1), serine/threonine kinases (eg, BRAF), and dual-specific kinases (eg, MEK).⁷ Corresponding targeted therapies have undergone extensive clinical studies, with some already approved for clinical use and others still in the research phase but showing significant efficacy (Table 1). In the context of immunotherapy, immune checkpoint inhibitors (ICIs) have become the standard treatment for stage IV lung cancer without targeted mutations. However, their use in treating lung cancer with driver mutations remains controversial, with low response rates often observed in first-line therapy. As a result, it remains uncertain whether the use of molecularly targeted agents, ICIs, or chemotherapy has good efficacy in NSCLC patients with specific oncogenic driver gene mutations. This review will explore the latest advancements in therapeutic drugs for oncogenic driver mutations.

ROS1 Rearrangement

In 1987, a ROS1 gene fusion (FIG-ROS, now known as GOPC-ROS1) was first identified in the U118MG glioblastoma cell line. Subsequent studies demonstrated the oncogenic potential of this fusion in gliomagenesis. In 2007, the ROS1 gene was also found in a small subset of NSCLCs. The ROS1 gene, located on chromosome 6q22,⁶⁴ belongs to the insulin receptor family of receptor tyrosine kinases (RTKs) and promotes tumor proliferation through the activation of the PI3K-AKT-mTOR, VAV3, JAK-STAT, and MAPK-ERK pathways^{65–67} (Figure 1). ROS1 rearrangement is found in approximately 2.6% of NSCLC,⁶⁸ with a higher prevalence among females, younger individuals, and non-smokers.^{69,70} Compared to traditional chemotherapy with platinum plus pemetrexed, targeted therapy against ROS1 rearrangement has significantly improved patient prognosis and quality of life. Currently, three ROS1 inhibitors are approved for clinical use: crizotinib (approved by the FDA in 2016),⁷¹ entrectinib (approved by the FDA in 2019),⁷² and repotrectinib (approved by the FDA in 2023).⁷³

Crizotinib is a TKI against multiple targets, including ROS1, ALK and MET. It selectively binds to the ATP-binding site of tyrosine kinases, blocking their phosphorylation and subsequent activation of downstream signaling pathways, thereby inhibiting tumor cell growth and invasion.⁷⁴ In the phase I/II AcSe study, crizotinib was administered to patients with ROS1-rearrangement NSCLC, resulting in an ORR of 47.2%, a median progression-free survival (mPFS) of 5.5 months, and a median overall survival (mOS) of 17.2 months.⁸ Two other studies, EUCROSS and METROS, also demonstrated significant anti-tumor activity of crizotinib in patients with ROS1-positive advanced NSCLC.^{9,10} The predominant site of disease progression in patients with ROS1-positive NSCLC was brain metastases, likely due to crizotinib's low blood-brain barrier penetration rate, despite its strong anti-tumor activity. Furthermore, nearly 36% of patients with ROS1-positive NSCLC had baseline brain metastases. Consequently, future research should focus on developing ROS1-TKIs with improved blood-brain barrier penetration capabilities.⁶⁴

Entrectinib is an oral multi-target inhibitor of ROS1, ALK and pan-tropomyosin receptor kinase (TRK) that effectively penetrates the blood-brain barrier and demonstrates central nervous system (CNS) activity. Its anti-ROS1 activity was found to be 40-fold more potent than that of crizotinib *in vitro*.⁷⁵ A comprehensive analysis based on prospective, global, multicenter datasets (STARTRK-1, STARTRK-2, and ALKA-372-001) demonstrated that entrectinib had a favorable DCR in the CNS and other systems in patients with advanced ROS1-positive NSCLC. The data showed an mPFS of 19 months and a median effective duration of 24.6 months, surpassing the median duration of response (mDoR) of crizotinib therapy (19.7 months). In 20 patients with CNS metastases, the intracranial response rate was 55%

Table I Key Randomized Trials in Advanced Oncogene-Driven NSCLC

Rare Oncogenic Mutation	Rate (%)	Drug	Trial ID	N	Efficacy					Reference						
					ORR (%)	mOS(mo)	mPFS (mo)	DCR (%)	mDoR (mo)							
ROS1 rearrangement	2.6	Crizotinib	NCT02034981	37	47.2	17.2	5.5	83.3	NA	[8]						
			NCT02183870	30	70.0	NA	20.0	90.0	NA	[9]						
			NCT02499614	26	65.0	NA	22.8	85.0	21.4	[10]						
			NCT02097810; NCT02568267; EudraCT 2012-000148-88	53	77.0	NA	19	NA	24.6	[11]						
BRAF mutation	2.1	Lorlatinib	NCT01970865	21	62.0	NA	NA	NA	25.3	[12]						
			NCT01964157	32	62.0	NA	9.3	81.0	NA	[13]						
		Repotrectinib (TPX-0005)	NCT03093116	71	79.0	NA	35.7	NA	34.1	[14]						
			Vemurafenib	NCT01524978	54 pretreated 8 naive	37.0 37.5	15.4 NA	6.1 12.9	NA NA	NA NA	[15]					
		Vemurafenib	AcSe	96	44.8	10.0	5.2	NA	NA	[16]						
			EURAF	24	53.0	10.8	5.0	85.0	NA	[17]						
		Dabrafenib	NCT01336634	9	33.1	12.7	5.5	58.0	9.6	[18,19]						
		Sorafenib		1							NA					
		Dabrafenib		78							66.7	18.2	10.2	80.7	9.8	[19-21]
		Dabrafenib +trametinib		57							64.0	12.7	10.9	75.0	10.2	[20]
		NTRK fusion	<1%	Encorafenib+binimetinib	NCT03915951	59 naive 39 pretreated	75.0 46.0	NA NA	NA 9.3	64.0 41.0	NA 16.7	[22]				
					153	79.0	44.4	28.3	NA	35.2	[23]					
Larotrectinib	NCT02122913			15	73.0	40.7	35.4	NA	33.9	[24]						
	NCT02637687															
Larotrectinib	NCT02576431			54	57.0	NA	11.2	NA	10.4	[25]						
	NCT02576431															
Entrectinib	NCT02097810	22	63.6	NA	14.9	NA	19.9	[26]								
Entrectinib	NCT02568267															
Entrectinib	NCT02097810	31	64.5	NA	NA	NA	NA	[27]								
									NCT02568267							

(Continued)

Table 1 (Continued).

Rare Oncogenic Mutation	Rate (%)	Drug	Trial ID	N	Efficacy					Reference
					ORR (%)	mOS(mo)	mPFS (mo)	DCR (%)	mDoR (mo)	
RET fusion	1–2	Selpercatinib (Loxo-292)	NCT03157128	105 pretreated	64.0	NA	16.5	NA	17.5	[28]
				39 naive	85.0	NA	NA	NA	NA	
			NCT03157128	69 naive	84.0	NA	22.0	NA	20.2	
		Pemetrexed+ platinum-based +pembrolizumab	NCT04194944	247 pretreated	61.0	NA	24.9	NA	28.6	[29]
				83	65.0	NA	11.2	NA	11.5	
				129	84.0	NA	24.8	NA	24.2	
		Pemetrexed+ platinum-based ±pembrolizumab	NCT04194944	159	63.0	NA	11.2	NA	12.0	[30]
				102	84.0	NA	24.8	NA	24.2	
				27 naive	70.0	NA	9.1	85.0	9.0	
		Selpercatinib ±pembrolizumab	NCT03037385	87 pretreated	61.0	NA	17.1	91.0	NA	[31]
				75 naive	72.0	NA	13.0	91.0	NA	
				136 prior platinum-based chemotherapy	59.0	NA	16.5	90.0	22.3	
MET mutation	3	Tepotinib	NCT02864992	164 naive	57.3	21.3	12.6	NA	46.4	[32]
				149 pretreated	45.0	19.3	11.0	NA	12.6	
		Capmatinib	NCT02414139	28 naive	67.9	20.8	12.4	NA	12.6	[33]
				32 naive	65.6	NA	10.8	NA	NA	
				69 pretreated	40.6	13.6	5.4	NA	9.7	
		Savolitinib	NCT02897479	31 pretreated	51.6	NA	6.9	NA	8.4	[34]
				70	47.1	12.5	NA	81.4	NA	
				28	36.0	8.1	2.4	39.0	NA	
		Crizotinib	NCT02034981	26	27.0	5.4	4.4	69.0	3.7	[8]
				36	33.0	NA	6.7	58.3	NA	
		Amivantamab	NCT02499614							[10]
			NCT02609776							[35]

KRAS G12C mutation	2.8	Sotorasib	NCT03600883	59	32.2	NA	6.3	88.1	NA	[37]
				124	37.1	NA	6.8	80.6	11.1	[38]
		Sotorasib	NCT04303780	169	28.1	10.6	5.6	82.5	8.6	[39]
		Docetaxel		151	13.2	11.3	4.5	60.3	6.8	
		Adagrasib	NCT03785249	15	53.3	NA	11.1	NA	16.4	[40]
			NCT03785249	112	42.9	12.6	6.5	NA	8.5	[41]
		Nivolumab	NCT01673867	292	19.0	12.2	2.3	NA	NA	[42]
		Docetaxel		290	12.0	9.4	4.2	NA	NA	
		T-DMI	NCT02675829	18	44.0	NA	5.0	NA	4.0	[43]
			NCT02675829	49	51.0	NA	5.0	NA	4.4	[44]
HER2 mutation	2.3		JapicCTI-194620	22	38.1	8.1	2.8	52.4	3.5	[45]
		T-DXd	NCT03505710	91	55.0	17.8	8.2	92.0	9.3	[46]
		Pyrotinib	NCT02834936	60	30.0	14.4	6.9	NA	6.9	[47]
			ChiCTR1800020262	78	19.2	10.5	5.6	74.4	9.9	[48]
		Pozotinib	NCT03066206	90	27.8	NA	5.5	70.0	5.1	[49]
			NCT03318939	70	41.0	NA	5.6	73.0	5.7	[50]
		Crizotinib	NCT01154140	172	74.0	NA	10.9	NA	11.3	[51]
		Pemetrexed+ platinum-based		171	45.0	NA	7.0	NA	5.3	
		Crizotinib	NCT00932451	173	65.0	NA	7.7	NA	32.1	[52]
		Pemetrexed/docetaxel		174	20.0	NA	3.0	NA	24.4	
ALK rearrangement	3.8	Ceritinib	NCT01828099	189	72.5	NA	16.6	NA	23.9	[53]
		Pemetrexed/docetaxel		175	26.7	NA	8.1	NA	11.1	
		Ceritinib	NCT01828112	115	39.1	NA	5.4	76.5	NA	[54]
		Pemetrexed/docetaxel		116	6.9	NA	1.6	36.2	NA	
		Brigatinib	NCT02737501	137	71.0	NA	24.0	NA	33.2	[55,56]
		Crizotinib		138	60.0	NA	11.1	NA	13.8	
		Alectinib	NCT02075840	152	82.9	NA	25.7	NA	48.2	[57,58]
		Crizotinib		151	75.5	57.4	10.4	NA	23.3	
		Alectinib	NCT02604342	79	50.6	NA	10.9	NA	NA	[59]
		Pemetrexed/docetaxel		40	2.5	NA	1.4	NA	NA	
NRG1 fusion	0.1–0.4	Ensartinib	NCT02767804	143	74.0	NA	25.8	NA	NA	[60]
		Crizotinib		147	67.0	NA	12.7	NA	27.3	
		Lorlatinib	NCT01970865	198	47.0	NA	NA	NA	NA	[61]
		Lorlatinib	NCT03052608	149	76.0	NA	NA	NA	NA	[62]
		Crizotinib		147	58.0	NA	9.3	NA	11.0	
		Seribantumab	NCT04383210	12	30.0	NA	NA	90.0	NA	[63]

Abbreviations: ORR, objective response rate; mOS, median overall survival; mPFS, median progression-free survival; DCR, disease control rate; mDOR, median duration of response; mo, month; NA, not applicable.




Figure 1 Key signaling pathways of oncogenic driver genes in non-small cell lung cancer. Each of the RTKs activates several downstream signaling molecules, leading to diverse cellular responses. SHP1/SHP2 signaling leads to activation of ESYT1. VAV3 controls the RHO family GTPase via CDC42. The PI3K pathway is activated by various RTKs, leading to AKT phosphorylation and downstream effects on mTOR and NF- κ B signaling, modulated by PTEN. The RAS-RAF-MEK-ERK/MAPK cascade is initiated by several RTKs, progressing through BRAF and RAF to activate MEK and eventually MAPK. The PLC- γ pathway regulates intracellular signaling molecules such as DAG, PKC, IP3, and Ca^{2+} . The JAK-STAT pathway is another critical branch, resulting in the activation of STAT proteins. RALGDS signaling controls the RAL GTPase and its effector PLD1. This complex network coordinates various cellular processes including proliferation, survival, migration, and differentiation. Arrowheads indicate activation, while flat-head lines represent inhibitory signals.

and the median intracranial DoR was 12.9 months.⁶⁴ This gives entrectinib an advantage over crizotinib, with overall manageable and mostly grade 1 toxicity that was reversible.⁷⁴

Repotrectinib (TPX-0005) is a low molecular weight, multi-target TKI that targets ROS1, TRK and ALK. Phase I/II clinical trials showed that repotrectinib exhibited durable antitumor activity with a mDoR of 34.1 months and a mPFS of 35.7 months in patients who had not received prior ROS1 TKI therapy, regardless of prior chemotherapy. Repotrectinib was similarly effective in patients with ROS1-positive NSCLC who had received at least one prior ROS1 TKI therapy. In comparison, entrectinib had an mDoR of 20.5 months and an mPFS of 15.7 months, while crizotinib had an mDoR of 24.7 months and an mPFS of 19.3 months.¹⁴

In addition to the three approved and marketed drugs mentioned above, several other targeted drugs have shown significant efficacy in clinical studies. Lorlatinib is a potent oral inhibitor of ALK and ROS1 with activity against crizotinib- and ceritinib-resistant mutations (G2032R, D2033N, and S1986Y) and the ability to penetrate the blood-brain barrier. In a Phase II study enrolling patients with ROS1-positive NSCLC, including those who had not been treated with a TKI and those pretreated, lorlatinib achieved an mPFS of 21.0 months and an ORR of 61.5% in treatment-naïve patients, compared with an mPFS of 8.5 months and an ORR of 26.5% in patients pretreated with crizotinib. Surprisingly, in treatment-naïve patients with brain metastases, the ORR was 66.7%. Lorlatinib has been reported to be effective against resistance caused by activation of bypass signaling, though it has limited efficacy in patients with G2032R mutation.¹² Ceritinib is an ALK inhibitor with potential anti-ROS1 rearrangement activity. Regardless of prior crizotinib treatment, ceritinib has shown significant efficacy in clinical trials, with favorable intracranial responses and manageable safety profiles in patients with brain metastases.¹³ Cabozantinib is a multi-targeted TKI that targets MET, RET, and ROS1. Sun et al reported the clinical efficacy of cabozantinib in four patients with ROS1-positive NSCLC who were resistant to crizotinib and ceritinib. Patients in the cabozantinib group had a PFS of 4.9 to 13.8 months.⁷⁴ Brigatinib is a multi-targeted inhibitor of ROS1 and ALK with activity against multiple resistant ROS1 mutations. A study showed that brigatinib had treated TKI-naïve patients and crizotinib-resistant patients with ROS1-positive NSCLC had ORRs of 37% and 29% in the overall population and crizotinib-resistant subgroup, respectively.⁷⁶ In a case report, brigatinib maintained efficacy even after the patient had received multiple ROS1-TKIs.⁷⁷ According to in vitro experiments, brigatinib is ineffective against NSCLC with the G2032R mutation but shows substantial anti-tumor efficacy against those with the L2026M mutation.⁷⁴

Little is known about the effects of ICIs on ROS1-mutant NSCLC. A retrospective study showed that anti-PD-(L)1 antibody monotherapy was poorly active, with only one out of six patients achieving a short-term response rate (ORR of 17%).⁷⁸

BRAF Mutation

Ulf Rapp et al first described v-raf and its association with cancer in 1983 as a mouse retroviral oncogene with a mammalian cell homolog called CRAF.⁷⁹ In 1984–1985, BRAF was discovered to be related to CRAF.^{80,81} BRAF mutations occur in 2.1% of NSCLC cases and can be categorized into V600 and non-V600 codon mutations.⁶⁸ BRAF belongs to the RAF group of serine/threonine kinases and is involved in cell proliferation and differentiation through the MAPK signaling pathway^{82,83} (Figure 1). Precision-targeted therapies significantly enhance the prognosis of BRAF mutation-positive NSCLC patients compared to standard paclitaxel-based chemotherapy.⁸⁴ Currently, FDA-approved targeted agents for BRAF mutations include dabrafenib in combination with trametinib,⁸⁵ and encorafenib in combination with binimetinib.⁸⁶

Vemurafenib (PLX4032) is a V600-BRAF mutation inhibitor.⁸⁷ In the VE Basket trial, vemurafenib treatment yielded an ORR of 37% among BRAF V600E-mutated NSCLC patients, with consistent ORR in treatment-naïve and previously treated groups. Median overall survival (mOS) was not reached and 15.4 months, while mPFS was 12.9 and 6.1 months, respectively.¹⁵ Another study evaluating vemurafenib monotherapy in BRAF V600 and non-V600 mutated NSCLC reported an ORR of 44.8% and mPFS of 5.2 months in the V600 group, whereas no tumor response was observed in the non-V600 group, with an mPFS of 1.8 months.¹⁶ Vemurafenib monotherapy demonstrates efficacy specifically in BRAF V600-mutant NSCLC, with limited activity against non-V600 mutations. The EURAF cohort further validates the antitumor activity of vemurafenib and dabrafenib in advanced BRAF-mutant lung cancer.¹⁷

Dabrafenib is an ATP-competitive BRAF inhibitor.⁸⁸ A phase II clinical trial found that dabrafenib monotherapy in patients with metastatic BRAF V600-mutated NSCLC who had received prior systemic therapy with an ORR of 33.1% and a DCR of 58%. Dabrafenib in combination with trametinib provided substantial clinical benefit and durable responses, demonstrating comparable efficacy in both treatment-naïve (ORR=63.9%, mDoR=10.2 months) and previously treated (ORR=68.4%, mDoR=9.8 months) groups. Combination therapy with dabrafenib and trametinib significantly improved outcomes in NSCLC patients harboring BRAF V600E mutations, surpassing the efficacy of dabrafenib monotherapy irrespective of prior treatment history.^{18–20}

Despite initial effectiveness, BRAF inhibitor monotherapy is limited by acquired resistance mechanisms that reactivate the MAPK pathway. To mitigate resistance activation, BRAF monomer inhibitors are combined with downstream MEK inhibitors, enhancing efficacy and tolerability.⁸⁹ Encorafenib and binimetinib inhibit tumor proliferation by blocking ERK signaling in the MAPK pathway.⁸⁸ In a phase II clinical trial, the combination of encorafenib and binimetinib showed significant antitumor activity in BRAF-mutant NSCLC patients, achieving ORRs of 75% and 46%, mDoR of not reached and 16.7 months, and mPFS of not reached and 9.3 months in treatment-naïve and previously treated groups, respectively.²² Compared to dabrafenib combined with trametinib, the combination of encorafenib and binimetinib demonstrated significantly better efficacy in both treatment-naïve patients and those who had received prior therapy. Regarding adverse effects, nearly all patients experienced side effects of varying severity. However, those treated with encorafenib and binimetinib had significantly less pyrexia, while the rates of gastrointestinal and constitutional toxicities were similar between the two drug combinations.⁹⁰

Data on the efficacy of anti-PD-(L)1 antibodies in patients with BRAF-mutated NSCLC are limited and conflicting. One retrospective study reported that anti-PD-(L)1 antibody monotherapy exhibited low clinical activity in patients with BRAF-mutated NSCLC, with an ORR of 24% and a mPFS of 3.1 months.⁷⁸ The benefit of treatment in patients with BRAF V600-mutant tumors may be particularly limited, with one study reporting an ORR of only 9% and a time to progression on ICIs of 1.9 months.⁹¹ However, a large study involving NSCLC patients treated with anti-PD-(L)1 antibody monotherapy reported an ORR of 62% for patients with BRAF-mutated NSCLC. This study also demonstrated that patients with BRAF V600-mutant tumors had a longer mPFS and OS compared to those with non-V600 mutations (9.8 months vs 5.4 months and 20.8 months vs 14.9 months, respectively).⁹²

For patients with BRAF V600-mutated NSCLC, the optimal first-line treatment remains unclear. Combination therapy with dabrafenib and trametinib may be preferable to ICIs based on available data, but anti-PD-(L)1 antibody monotherapy (in patients with PD-L1 TPS ≥50%) or chemoimmunotherapy, with or without bevacizumab (regardless of PD-L1 status), are also reasonable options. Additionally, evidence from melanoma research supports the use of upfront immunotherapy. In a phase III

trial involving treatment-naïve patients with advanced BRAF V600-mutant melanoma, those who received dual ICI therapy (nivolumab plus ipilimumab) followed by BRAF-targeted therapy (dabrafenib plus trametinib) had better OS than those who received the therapies in the reverse sequence (72% vs 52% at two years).⁹³

NTRK Fusion

NTRK fusions were first reported in a case of colorectal cancer in 1982. Since then, over 80 different NTRK gene fusions involving various ligands have been identified across a wide range of tumors in both adults and children,⁹⁴ occurring in less than 1% of NSCLC cases.⁹⁵ NTRK genes encode the TRK proteins, pivotal in embryonic development and maintaining normal nervous system function. NTRK fusion drives various solid tumors in adults and children, promoting tumor proliferation by activating the MAPK, PI3K, and PKC pathways^{96,97} (Figure 1). Larotrectinib⁹⁸ and entrectinib⁷² have both gained FDA approval for the treatment of patients with NTRK fusion-positive tumors.

Larotrectinib, an oral small molecule highly selective for pan-TRK inhibition, was first approved in the US in November 2018 for locally advanced or metastatic solid tumors containing NTRK fusions without acquired resistance mutations.⁹⁵ A comprehensive analysis of three clinical trials (LOXO-TRK-14001, SCOUT, and NAVIGATE) found that larotrectinib was highly active in TRK fusion-positive lung cancer, achieving an ORR of 75% with notable CNS efficacy.²³ Compared to chemotherapy and immunotherapy, larotrectinib elicited rapid, significant, and durable responses.²⁴ Combined analyses from trials NCT02576431 and NCT0212291 showed an ORR of 83% in patients with TRK fusion-positive lung cancer treated with larotrectinib monotherapy and 80% in those with CNS metastases.⁹⁹

Entrectinib, an oral selective inhibitor of TRK, ROS1 and ALK tyrosine kinases, was initially approved in Japan in June 2019 for advanced or recurrent adult and pediatric NTRK fusion-positive solid tumors.⁹⁵ Pooled data from STARTRK-1, STARTRK-2 and ALKA-372-001 trials highlighted entrectinib's significant antitumor and CNS activity. Monotherapy in metastatic or locally advanced NTRK fusion-positive tumors achieved an ORR of 57%, an mDoR of 10.4 months, an mPFS of 11.2 months, and the ORR for patients with lung cancer was 70%.²⁵ In patients with baseline CNS disease (n=6), four showed intracranial responses, including two complete responses (CR) and two partial responses (PR), with one stable disease (SD).¹⁰⁰ The ORR in NSCLC patients with no prior TRK inhibitor therapy was 63.6%. The mDoR, mPFS and mOS were 19.9 months, 14.9 months and not reached, respectively.²⁶ Further analyses across various solid tumors with positive NTRK fusion reported an ORR of 61.3%, mDoR of 20.0 months, mPFS of 13.8 months, and mOS of 37.1 months. The ORR for the NSCLC subgroup was 64.5%.²⁷ In conclusion, entrectinib has clinically significant antitumor activity in NTRK fusion-positive patients, especially against CNS tumors and metastases.

RET Fusion

In 1985, Takahashi used DNA from human lymphoma cells to transform NIH3T3 cells. By analyzing the transformed daughter cells, he discovered a new gene activated by rearrangement with other genes, which he named the RET gene due to this rearrangement during the transformation process.¹⁰¹ Located on chromosome 10q11.2, RET encodes a RTK that triggers downstream MAPK, JAK-STAT, and PI3K-AKT-mTOR signaling pathways, promoting tumor cell proliferation¹⁰² (Figure 1). RET fusions occur in approximately 1–2% of NSCLC cases.²⁸ Prior to the development of specific RET inhibitors, treatment primarily involved multi-kinase inhibitors, which were less targeted and associated with significant side effects, limiting their efficacy. The FDA has approved selpercatinib¹⁰³ and pralsetinib¹⁰⁴ for the treatment of RET fusion-positive NSCLC patients.

Selpercatinib (Loxo-292) is an oral, highly selective RET inhibitor effective against wild-type and multiple mutant or fusion RET subtypes.^{105,106} In a phase I/II trial, selpercatinib monotherapy for RET fusion-positive NSCLC showed an ORR of 64% in pretreated patients and 85% in treatment-naïve patients. Among patients with measurable CNS metastases at enrollment, 91% achieved an objective intracranial response.²⁸ Updated data indicated ORRs of 84% and 61% in treatment-naïve and pretreated groups, respectively, with mPFS of 22 and 24.9 months.²⁹ A comparative study found that selpercatinib monotherapy resulted in longer mPFS and DoR compared to platinum-based combination combined with pemetrexed, with or without pembrolizumab. The intracranial ORR was 82%, suggesting favorable efficacy against CNS metastases.³⁰ This highlights the significant antitumor activity of selpercatinib in patients with RET fusion-positive lung cancer, including in first-line therapy, post-platinum-based chemotherapy, and those with brain metastases.

Pralsetinib (BLU-667) is another potent, oral and selective RET inhibitor for metastatic RET fusion-positive NSCLC. Pralsetinib was well tolerated and showed clinical activity, including intracranial response, in patients with RET fusion-positive NSCLC in the phase I/II ARROW study. The ORR was 61% in pretreated patients and 70% in treatment-naïve patients.³¹ More recently, updated data reported ORRs of 72% in treatment-naïve and 59% in pretreated patients, with mDoR not reached and 22.3 months, and mOS of 13 and 16.5 months, respectively. Tumor shrinkage was observed in all treatment-naïve patients and 97% of pretreated patients. Among those with measurable intracranial metastases, the intracranial response rate was 70%. In all patients with RET fusion-positive NSCLC, 7% discontinued due to treatment-related adverse events (TRAEs).³² In conclusion, pralsetinib has shown clinically meaningful and durable responses in advanced RET fusion-positive NSCLC in both primary and standard refractory populations, including intracranial efficacy, with a manageable safety profile.

A single retrospective study demonstrated significant efficacy of anti-PD-1 antibody monotherapy in patients with previously treated RET-fusion NSCLC, reporting an ORR of 37.5%, a mDoR of 12.1 months, and a mPFS of 7.6 months.¹⁰⁷ However, at least two other retrospective studies showed poor efficacy, with ORRs below 10% and mPFS of less than 3.5 months.^{78,108} These findings suggest that the efficacy of immunotherapy in patients with this subtype of mutation remains uncertain.

MET Mutation

The MET gene was first identified in 1984 as a MET-TPR fusion in a human osteosarcoma cell line. In 1996, Natali et al discovered that MET was overexpressed in several cancers and associated with poor prognosis. It was not until 2005 that somatic mutations in the MET gene were first reported in NSCLC.¹⁰⁹ The MET gene, also known as the hepatocyte growth factor receptor (HGFR), is present in approximately 3% of NSCLC cases.⁶⁸ It encodes a RTK for hepatocyte growth factor (HGF), which, upon binding to the MET receptor, activates several downstream signaling pathways, including RAS-RAF-MEK-MAPK, PI3K-AKT, STAT, and NF- κ B. These pathways are involved in cell proliferation, differentiation, and survival¹¹⁰ (Figure 1). C-MET mutations mainly consist of MET exon 14 skipping (METex14, 2%–4%), MET amplification (METamp, 1%–6%), MET overexpression (20%–25%) and MET fusion (0.2%–0.3%).¹¹¹ The first MET TKI to receive FDA approval was tepotinib,¹¹² followed by capmatinib.¹¹³ Savolitinib is currently approved only in China for patients who have progressed or are intolerant after platinum-based chemotherapy.¹¹⁴

Tepotinib is a highly selective MET inhibitor. The VISION phase II clinical trial evaluated tepotinib for MET-positive patients with advanced NSCLC who had not previously received a MET inhibitor. With the median follow-up time of 32.6 months, the ORR was 51.4%, the mDoR was 18 months, the mPFS was 11.2 months, and the mOS was 19.6 months. Among patients who had received prior platinum-based therapy or ICIs, the ORR was 45.0%, the mDoR was 12.6 months, the mPFS was 11.0 months, and the mOS was 19.3 months. Treatment-naïve patients showed an ORR of 57.3%, mDoR of 46.4 months, mPFS of 12.6 months, and mOS of 21.3 months.³³ The efficacy of tepotinib was consistent across different MET mutation types and concomitant MET AMP.¹¹⁵ The efficacy of tepotinib was similar regardless of prior treatments, with consistent ORR among patients who had received prior platinum-based chemotherapy, immunotherapy, and platinum-based combination immunotherapy. The mDoR was 12.4 months, the mPFS was 11.0 months in platinum-pretreated patients, and the mDoR was 9.5 months, the mPFS was 10.9 months in patients previously treated with ICIs.¹¹⁶ The intracranial response rate was 66.7% in patients with baseline brain metastases.³³ In conclusion, tepotinib showed clinically relevant outcomes in both naïve and pretreated patients with MET-mutated NSCLC. In patients with baseline brain metastases, tepotinib showed solid systemic and intracranial outcomes with comparable clinical benefit to patients without baseline brain metastases. Moreover, tepotinib was well tolerated and resulted in a low rate of TRAE discontinuation.

Capmatinib is another selective MET inhibitor. In the GEOMETRY-mono1 trial, capmatinib monotherapy in MET-positive NSCLC patients with prior first/second-line therapy resulted in an ORR of 40.6%, mDoR of 9.7 months, mPFS of 5.4 months, and mOS of 13.6 months. Among second-line therapy patients, the ORR was 51.6%, mDoR was 8.4 months, and mPFS was 6.9 months. Treatment-naïve patients had an ORR of 67.9%, mDoR of 12.6 months, mPFS of 12.4 months, and mOS of 20.8 months. An extended cohort of treatment-naïve patients showed an ORR of 65.6% and mPFS of 10.8 months.³⁴ This study found that capmatinib exhibited significant antitumor activity in advanced NSCLC

with MET mutation, particularly in treatment-naïve patients, with no significant differences between MET mutation types and concomitant MET AMP.¹¹⁷

Savolitinib is also a MET inhibitor. In a phase II clinical trial involving METex14 NSCLC patients, with a median follow-up of 28.4 months, the mOS of 12.5 months; in patients with brain metastases, the mOS was 17.7 months.¹¹⁸ The ORR was 47.6%, and the DCR was 81% in the pretreated group. In the treatment-naïve group, the ORR was 46.4%, and DCR was 82.1%.³⁵ These results indicate that savolitinib has clinically meaningful antitumor activity, regardless of prior therapy or the presence of brain metastases.

Although crizotinib is not approved for treating NSCLC patients with METex14 mutation, it is commonly used in clinical practice.¹¹⁹ The phase II AcSe trial showed modest efficacy with an ORR of 36%, mPFS of 2.4 months, and mOS of 8.1 months.⁸ In the METROS, an Italian multicenter phase II trial, patients with MET-positive NSCLC treated with crizotinib exhibited the ORR of 27%, the mPFS of 4.4 months, and the mOS of 5.4 months.¹⁰ In conclusion, these trials suggest that while crizotinib can induce tumor shrinkage in some MET-regulated NSCLC tumors, its overall impact on the clinical course of MET-mutated NSCLC patients is limited.

Amivantamab is a bispecific antibody that targets both the EGFR and MET, exhibiting a higher affinity for MET (40 pM) compared to EGFR (1.4 nM), along with immune cell-guiding activity. In the MET-2 arm of the CHRYSALIS Phase I trial, the amivantamab cohort demonstrated a response rate of 33% and a mPFS of 6.7 months. Among patients who experienced TRAEs, 12% had their dose reduced, and 5% discontinued treatment. The most common TRAEs included infusion-related reactions (69%, 5% of \geq grade 3), rash (7% of \geq grade 3), impetigo, and peripheral edema, none of which reached grade 3 severity.³⁶ Despite these encouraging results with amivantamab and manageable toxicity, the optimal treatment sequence is still under study.

Similar to other oncogenic factors, such as BRAF, evidence regarding the efficacy of ICIs in patients with MET-positive NSCLC is limited and controversial. Studies indicate that the clinical activity of ICI monotherapy for this genotype is relatively low. Although some cases have reported durable responses,^{107,120} at least three other retrospective studies have shown ORRs of less than 20% and mPFS of less than 5.0 months.^{78,92,120} Based on this data, MET TKIs may be the preferred treatment for patients with MET-positive NSCLC.

KRAS G12C Mutation

In the 1960s, Harvey and Kirsten et al discovered the mouse tumor genes HRAS and KRAS, which are carried by retroviruses similar to rous sarcoma virus (RSV). In 1982, Weinberg et al identified HRAS in human bladder cancer cells, making RAS the first human tumor gene to be discovered.¹²¹ Activated KRAS proteins activate downstream pathways, including RAF-MEK-ERK, PI3K-AKT-mTOR, and RALGDS-RAL pathways, driving cell differentiation and proliferation¹²² (Figure 1). Approximately 2.8% of NSCLC patients harbor the KRAS G12C mutation⁶⁸. Historically, treatments for KRASG12C mutant patients have shown limited efficacy with immunotherapy and chemotherapy, but KRAS G12C inhibitors have yielded promising results in NSCLC.¹²³ The FDA approved sotorasib (AMG510) in May 2021 and adagrasib (MRTX849) in December 2022 for the treatment of NSCLC with the KRAS G12C mutation.^{124,125}

Sotorasib is the first approved KRAS G12C inhibitor. A phase II study assessed its efficacy in NSCLC patients with KRAS G12C mutations, in the second/third-line settings, sotorasib had an ORR of 37.1%, mDoR of 11.1 months, and mPFS of 6.8 months.³⁸ The phase III CodeBreak 200 study compared sotorasib to docetaxel in pretreated NSCLC patients. In the docetaxel group, the mPFS was 4.5 months, ORR was 13.2%, DCR was 60.3%, mDoR was 6.8 months, and mOS was 11.3 months. In contrast, the sotorasib group showed an mPFS of 5.6 months, ORR of 28.1%, DCR of 82.5%, mDoR of 8.6 months, and mOS of 10.6 months. Although sotorasib did not significantly improve OS compared to docetaxel, its toxicity and intracranial activity suggest it is a better option.³⁹

Adagrasib is another KRAS G12C inhibitor that demonstrated its efficacy in the phase I KRYSTAL-1 trial,⁴⁰ leading to a phase II study. Adagrasib was used to treat patients with KRAS G12C mutated NSCLC, with an ORR of 42.9%, mDoR of 8.5 months, mPFS of 6.5 months, and mOS of 12.6 months.⁴¹ Updated 2023 data extended the mOS to 14.1 months with a median 2-year survival rate of 31%.¹²⁶ Additionally, the phase III KRYSTAL-12 trial is evaluating adagrasib versus docetaxel in patients with advanced NSCLC harboring KRAS mutations.¹²⁷

ICIs demonstrate clinical efficacy in patients with KRAS-mutated NSCLC. In the initial phase III trial of ICI-based immunotherapy for NSCLC, nivolumab monotherapy significantly improved OS compared to docetaxel in patients with KRAS mutations.⁴² A pooled analysis of data from 12 registrational trials indicated that patients with untreated advanced KRAS G12C mutations had an ORR of 33% when treated with ICIs alone, without chemotherapy.¹²⁸ The efficacy of ICIs in patients with KRAS-mutant NSCLC has been further supported by multiple retrospective studies.

HER2 Mutation

In 1979, Schechter et al discovered a new oncogene in rat cells, naming it *neu*. They later identified that *neu* shares structural similarities with the previously discovered *ERBB2* gene, leading to its alternative designation as *ErbB-2*. The protein encoded by the *ERBB2* gene in humans resembles the structure of the human epidermal growth factor receptor, which is why *neu* is also referred to as HER2.¹²⁹ It forms a dimer with the receptor, and activates the RAS-RAF-MEK-ERK and PI3K-AKT signaling pathways to promote tumor cell proliferation¹³⁰ (Figure 1). The incidence of HER2 mutation in NSCLC is approximately 2.3%.⁶⁸ The FDA has approved trastuzumab (T-DXd) for the treatment of adult patients with inoperable or metastatic NSCLC with HER2 mutations who have previously received systemic platinum-based therapy, marking the first approved treatment for HER2-mutated lung cancer.¹³¹

Pyrotinib is an irreversible small molecule inhibitor of EGFR, HER2 and HER4 receptors. A phase II trial showed that platinum-pretreated patients with HER2-mutated advanced NSCLC treated with pyrotinib had an ORR of 30%, an mPFS of 6.9 months, and an mOS of 14.4 months. Subgroup analysis indicated that pyrotinib benefitted all patients, irrespective of brain metastases or HER2 mutant subtypes.⁴⁷ Another phase II clinical trial using pyrotinib as first-line or later-line treatment, the ORR was 19.2%, mPFS was 5.6 months, mDoR was 9.9 months, and mOS was 10.5 months.⁴⁸ These studies suggest that pyrotinib has clinical efficacy in HER2-mutated advanced NSCLC, although its efficacy in second-line treatment remains unclear. Currently, the phase III study (NCT04447118) is ongoing to evaluate the efficacy of pyrotinib versus docetaxel in platinum-pretreated advanced non-squamous NSCLC with HER2 mutation.¹³²

Pozotinib is a novel covalent irreversible EGFR and HER2 receptor inhibitor. The phase II study ZENITH20 evaluated the efficacy of pozotinib monotherapy in HER2-mutated NSCLC. In this trial, the ORR was 27.8%, the DCR was 70%, the mPFS was 5.5 months, and the mDoR was 5.1 months. Pozotinib was efficacious regardless of prior therapy, CNS metastases, or HER2 mutation type. However, 97.8% of patients experienced TRAEs, leading to dose reduction and discontinuation in 76.7% and 13.3% of patients, respectively.⁴⁹ The ZENITH20-4 trial showed an ORR of 41%, a DCR of 73%, an mPFS of 5.6 months, and an mDoR of 5.7 months, with improved tolerance due to dose adjustments.⁵⁰

Antibody-drug conjugates (ADCs) are a class of antitumour medicines that effectively extend the therapeutic window by precisely delivering cytotoxic chemicals into cancer cells.¹³³ T-DM1 is an ADC. A clinical study evaluating T-DM1 in HER2-mutated NSCLC was terminated early due to poor efficacy, with the ORR of 6.7% and the mPFS of 2.0 months.¹³⁴ However, a phase II basket trial demonstrated better efficacy: patients (n=18) treated with T-DM1 had an ORR of 44%, an mPFS of 5.0 months, and an mDoR of 4.0 months.⁴³ In the expanded clinical trial (n=49), the ORR was 50%, the mPFS was 5.0 months and the mDoR was 4.4 months.⁴⁴ Another phase II clinical trial reported the ORR was 38.1%, while mPFS and mDoR were only 2.8 and 3.5 months, respectively, for HER2-mutated NSCLC treated with T-DM1.⁴⁵ Overall, despite some short-term efficacy, the benefit of T-DM1 in advanced NSCLC with HER2 mutations appears limited in duration.

T-DXd is a relatively new anti-HER2 ADC. The phase II DESTINY-Lung01 study evaluated the efficacy and safety of T-DXd in HER2-mutated relapsed or refractory NSCLC, with the ORR of 55%, the mPFS of 8.2 months, the mDoR of 9.3 months, and the mOS of 17.8 months. It also showed efficacy in patients with brain metastases, with an mPFS of 7.1 months and mOS of 13.8 months.⁴⁶ The phase III trial DESTINY-Lung04 is currently ongoing, which compares the efficacy and safety of T-DXd monotherapy with chemotherapy in combination with pembrolizumab as first-line treatment for HER2-mutated advanced NSCLC.¹³⁵ In conclusion, T-DXd showed unprecedented efficacy and a manageable safety profile in HER2-mutated NSCLC.

The clinical benefit of immunotherapy in patients with HER2-mutated lung cancer is limited, with ORRs ranging from 7% to 27% and a mPFS of approximately 2 months.^{78,107}

ALK Rearrangement

ALK was first identified in a subtype of anaplastic large cell lymphoma (ALCL), hence the name anaplastic lymphoma kinase (ALK). Before the discovery of ALK rearrangement in NSCLC, multiple types of ALK rearrangement have been identified in diffuse large B cell lymphoma and inflammatory myofibroblastic tumor, demonstrating that ALK is a potent oncogene.^{136,137} ALK rearrangement is observed in about 3.8% of NSCLC.⁶⁸ ALK induces malignant transformation through the activation of pathways such as MAPK, PI3K-AKT, and JAK-STAT¹³⁸ (Figure 1). Five ALK inhibitors (alectinib,¹³⁹ brigatinib,¹⁴⁰ ceritinib,¹⁴¹ crizotinib,¹⁴² and lorlatinib¹⁴³) have been approved by the FDA for ALK mutation-positive NSCLC.

Crizotinib is the first-generation ALK TKI. Two Phase III clinical trials showed the superior clinical efficacy of crizotinib compared to pemetrexed in combination with platinum-based chemotherapy in patients with advanced ALK-rearranged NSCLC.^{51,52} Among patients who had not received platinum-based therapy, the PFS was significantly longer in the crizotinib group compared to the chemotherapy group (median value of 10.9 months vs 7.0 months).⁵¹ However, resistance to crizotinib has led to the development of second/third-generation ALK TKIs that cover most secondary ALK mutations and effectively penetrate the blood-brain barrier to treat patients with brain metastases.

Ceritinib is the second-generation ALK TKI. The phase III ASCEND-04 trial showed a significant improvement in mPFS for ceritinib compared to pemetrexed combined with platinum-based chemotherapy in patients with ALK-rearranged lung adenocarcinoma (16.6 months vs 8.1 months).⁵³ The ASCEND-05 trial compared ceritinib to single-agent chemotherapy in patients with advanced ALK-rearranged NSCLC who had previously progressed following crizotinib and platinum-based doublet therapy, showing a significant improvement in mPFS compared with chemotherapy (5.4 months vs 1.6 months, respectively).⁵⁴ Comparison of crizotinib and ceritinib revealed a significant advantage for ceritinib in terms of PFS (25.2 months vs 10.8 months), but not in terms of OS.¹⁴⁴

Brigatinib is another ALK TKI. Patients treated with brigatinib had a significantly longer median PFS compared to those treated with crizotinib (24.0 months vs 11.1 months), and the intracranial response rate was significantly higher in the brigatinib group (78% vs 29%).^{55,56}

Alectinib, also a second-generation ALK TKI, significantly prolonged the mPFS in the ALEX study compared to crizotinib group (25.7 months vs 10.4 months). Alectinib also significantly reduced CNS progression events (12% vs 45%).⁵⁷ The phase III ALUR study compared the efficacy of alectinib with chemotherapy in patients with advanced metastatic ALK-rearranged NSCLC who had received prior platinum-based chemotherapy and crizotinib. Alectinib showed a mPFS of 10.9 months compared to 1.4 months in the chemotherapy group, with a CNS ORR of 66.7% in the alectinib group versus 0% in the chemotherapy group.⁵⁹

Ensartinib, another second-generation ALK TKI, was evaluated in the phase II eXalt3 trial. Patients treated with ensartinib had a significantly better PFS compared to those treated with crizotinib (25.8 months vs 12.7 months), and the intracranial response rate was higher in the ensartinib group (63.6% vs 21.1%).⁶⁰

Lorlatinib is a third-generation ALK TKI. In a phase II study, lorlatinib was used to treat patients with advanced NSCLC with ALK rearrangement, with an ORR of 47%.⁶¹ The phase III CROWN trial compared lorlatinib with crizotinib, demonstrating improvements in PFS, ORR, intracranial response rate, time to intracranial progress, and DoR in the lorlatinib group. The 3-year PFS of 64% for lorlatinib compared to 19% for crizotinib. Among patients without brain metastases, intracranial progression occurred in 1% of patients in the lorlatinib group versus 23% in the crizotinib group.¹⁴⁵

NRG1 Fusion

In 1997, the first mutation of the NRG1 gene was discovered. Two years later, it was identified as a fusion phenomenon, with the fusion partner gene of NRG1 initially confirmed as DOC4, along with the discovery of the NRG1 break site. By 2014, a series of partner genes fused with the NRG1 gene were identified in NSCLC, including CD74, SLC3A2, and VAMP2. By 2021, NRG1 gene fusions had also been found in ovarian cancer, prostate cancer, and endometrial cancer, garnering widespread attention.¹⁴⁶ NRG1 fusion occurs in non-squamous NSCLC at an incidence of about 0.1–0.4% and in more than 5% of invasive mucinous adenocarcinomas.¹⁴⁷ NRG1 proteins bind to cell surface HER3 receptor and

promote tumor cell proliferation by activating the PI3K-AKT-mTOR and MAPK signaling pathways¹⁴⁸ (Figure 1). Patients with NRG1 fusion usually do not respond well to chemotherapy, chemoimmunotherapy, or ICIs. However, treatment with HER2 inhibitors (indirect) or HER3 inhibitors (direct) may be an important way to stop tumor growth.¹⁴⁹

Zenocutuzumab,^{146,150} a HER2/HER3-targeting bispecific monoclonal antibody, and seribantumab,⁶³ an anti-HER3 monoclonal antibody, have shown encouraging efficacy in targeting NRG1-rearranged solid tumors, including NSCLC, and both are currently in phase II clinical trials (NCT02912949, NCT04383210).^{151,152}

FGFR Mutation

Fibroblast growth factor (FGF) is a family of peptide growth factors with at least 23 different members found. FGF1 and its receptor FGFR regulates key biological processes such as cell proliferation and differentiation and promotes the growth and angiogenesis of various tumors, including lung or breast cancer.¹⁵³ FGFR promotes tumor cell proliferation through the activation of PI3K-AKT, STAT, PKC, and RAS-MAPK pathways¹⁵⁴ (Figure 1). The mutation rate of FGFR in NSCLC is about 1%.¹⁵⁵ Currently erdafitinib has been approved by the FDA for patients with urothelial carcinoma with FGFR mutation,¹⁵⁶ and pemigatinib, infigratinib, and futibatinib are approved for the treatment of patients with cholangiocarcinoma with FGFR mutation.^{157–159}

Although most FGFR-targeting medications are still in preclinical testing or clinical research, numerous innovative therapies have been developed, including TKIs, selective inhibitors, and monoclonal antibodies like lenvatinib and dovitinib. The FGFR1 inhibitor ponatinib significantly inhibited the growth of primary lung cancer cells in vitro in NSCLC cell lines that expressed high levels of FGFR1. Moreover, in NSCLC patients with higher FGFR2/3 expression levels, erlotinib responded when combined with bevacizumab. A study showed that erdafitinib led apoptosis in FGF-dependent human squamous cell carcinoma NCI-H1581 and NCI-H520 cells by inhibiting FGF/FGFR.¹⁶⁰

PI3KCA Mutation

In 1985, M. Whitman and L. Cantley first discovered a phosphatidylinositol kinase capable of phosphorylating inositol cyclo-3-OH to produce phosphatidylinositol 3,4,5-trisphosphate (PIP3). They defined this new kinase as phosphatidylinositol-3-kinase (PI3K), marking the official beginning of PI3K research.¹⁶¹ Studies have demonstrated that PI3K overactivation of the downstream AKT-mTOR pathway mediates the development of a variety of malignant tumors, with PTEN being a key negative regulator of PI3KCA-activated PI3K-AKT-mTOR^{162,163} (Figure 1). The mutation rate of PI3KCA in NSCLC is about 2–5%, being more common in squamous cell lung carcinoma (33%–37%) than in adenocarcinoma (5%–6%).¹⁶⁴

Currently, many PI3K inhibitors are in different stages of preclinical studies and early clinical trials. Pan-class I PI3K inhibitors include pictilisib (GDC-0941), PX-866, buparlisib (BKM120), pilaralisib (XL-147), GNE-317, etc. Unfortunately, trials of buparlisib alone, buparlisib in combination with gefitinib, and PX-866 in combination with docetaxel have not shown any efficacy.^{165–167} Specific PI3K inhibitors have also been studied by Langer CJ et al. Taselisib, a selective PI3K inhibitor, was administered as monotherapy to patients with PI3K-positive lung cancer in the phase II LUNG-MAP trial, but the group was closed due to ineffectiveness.¹⁶⁸

DDR2 Mutation

Discoidin domain receptors (DDRs) were classified as orphan receptors in the early 20th century. In 1997, Shrivastava et al described DDR as a non-integral collagen-binding receptor that requires collagen to initiate cellular responses. DDR is activated upon collagen binding.¹⁶⁹ To date, dysregulation, mutations, and differential expression of DDRs have been reported in various cancers. DDR2 mutations occur in approximately 4% of squamous NSCLCs and have been shown to promote the growth of lung squamous cell carcinoma cells.¹⁶⁴ Miao et al demonstrated that DDR2 mutants enhance lung squamous cell carcinoma cell proliferation and increase migration and invasion in the absence of collagen stimulation.¹⁷⁰ Additionally, the V582E and L595P mutants of the DDR2 gene induce DDR2 activation and promote cell growth and tumor progression by binding to collagen ligands.¹⁷¹ While some DDR2 mutations function as oncogenic drivers, some studies suggest that DDR2 may exert inhibitory effects on cancer cells stimulated by fibrocollagen.¹⁷² Although the

physiological and pathological functions of DDRs are well documented, their specific signaling pathways and molecular mechanisms require further exploration and validation.

Several DDR kinase inhibitors have been identified, such as dasatinib, imatinib, and nilotinib, and other BCR-ABL inhibitors.¹⁷³ Clinical trials of dasatinib were terminated due to intolerable toxicity and poor efficacy.¹⁷⁴ However, the combination of dasatinib with Src kinase inhibitors has been shown to reduce toxicity and improve the efficacy of dasatinib.¹⁷⁵ Currently, the role of DDR2 as a therapeutic target is not yet fully understood. As basic research continues to advance and genetic screening and personalized tumor therapy become more widespread, it is anticipated that more DDR2 inhibitors will be developed and applied in clinical settings.

Future Directions

With advancements of tumor molecular biology, more and more novel anti-tumor drugs have emerged, leading to significant breakthroughs in the treatment of patients with NSCLC. Traditional cytotoxic drugs have only modestly improved the survival rates of NSCLC patients. However, the identification of oncogenic driver genes has profoundly impacted cancer prognosis and treatment. The FDA has approved several targeted therapies for patients with mutations in oncogenic factors such as EGFR, ALK, ROS1, BRAF, MET, RET, FGFR, and NTRK. Additionally, the development of next-generation TKIs and immunotherapeutic agents has further enhanced treatment outcomes, allowing some patients to achieve long-term survival. Most clinical studies indicate that patients with oncogene mutations may experience greater survival benefits from targeted therapy compared to those receiving chemotherapy or immunotherapy. As a result, targeted therapy has become the preferred first-line treatment option for patients with oncogenic mutations.

Despite significant advancements, several challenges remain in the targeted therapy of NSCLC that require urgent attention. First, patients receiving targeted therapy or immunotherapy often develop drug resistance, leading to disease progression or relapse. For instance, crizotinib, the first ALK TKI approved for treating ALK-positive NSCLC, initially demonstrates promising efficacy (Table 1). However, nearly all patients eventually develop resistance and experience disease progression within one year, often in the brain or other parenchymal regions. Overcoming resistance remains a critical barrier to the long-term success of targeted therapies. Therefore, it is essential to explore new therapeutic targets, investigate resistance mechanisms, and develop novel drugs or strategies to address this issue. Second, while targeted therapies can provide durable responses and prolonged survival for some patients, they may also lead to an increased incidence of adverse effects. A meta-analysis of 31 randomized controlled trials evaluating TKIs in NSCLC treatment found that 27 of the studies reported TRAEs, with the most common being hypertension, fatigue, rash, and gastrointestinal disturbances.¹⁷⁶ Addressing these side effects and enhancing drug safety are focal points of current research. Identifying the underlying mechanisms of these adverse reactions and discovering biomarkers that predict their occurrence can inform future mechanism-based treatment approaches, ultimately improving patient quality of life. Additionally, the combination of immunotherapy, targeted therapy, and chemotherapy is an emerging area of research. Determining whether oncogene-positive NSCLC patients can benefit from immunotherapy is a hot topic in lung cancer treatment. Therefore, there is still a lot of work to be done to achieve true precision medicine.

In conclusion, the identification of oncogenic driver genes in NSCLC has revolutionized the treatment landscape and significantly improved patient prognosis. Precision-targeted therapies have become the first-line treatment option for NSCLC. However, there is still a long way to go in the fight against malignant tumors. Continued exploration of new therapeutic targets and innovative treatment strategies will be essential for advancing anti-tumor therapy, making it more precise and effective.

Abbreviations

NSCLC, non-small cell lung cancer; SCLC, small cell lung cancer; EGFR, epidermal growth factor receptor; FDA, Food and Drug Administration; NCCN, National Comprehensive Cancer Network; NGS, next generation sequencing; TKI, tyrosine kinase inhibitor; PFS, progression-free survival; mPFS, median progression-free survival; ORR, objective response rate; mOS, median overall survival; DCR, disease control rate; mDOR, median duration of response; mo, month; NA, not applicable; RTK, receptor tyrosine kinase; TRK, tropomyosin receptor kinase; CNS, central nervous system; CR, complete response; PR, partial response; SD, stable disease; TRAE, treatment-related adverse event; HGFR,

hepatocyte growth factor receptor; HGF, hepatocyte growth factor; RSV, rous sarcoma virus; METex14, MET exon 14 skipping; METamp, MET amplification; ICI, immune checkpoint inhibitor; ADC, antibody-drug conjugate; ALCL, anaplastic large cell lymphoma; ALK, anaplastic lymphoma kinase; FGF, fibroblast growth factor; PIP3, phosphatidylinositol 3,4,5-trisphosphate; PI3K, phosphatidylinositol-3-kinase; DDR, discoidin domain receptor.

Consent for Publication

All authors reviewed and approved the manuscript.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by grants from the National Natural Science Foundation of China (No. 82072591 to ZW), Jiangsu Province Capability Improvement Project through Science, Technology and Education; Jiangsu Provincial Medical Key Discipline Cultivation Unit (NO. JSDW202235 to ZW), Bethune Medical Science Research Foundation Project (NO.2022-YJ-085-J-Z-ZZ-006 to ZW), “123” advantageous disciplines, core technologies of the Second Affiliated Hospital of Nanjing Medical University, Postgraduate Research & Practice Innovation Program of Jiangsu Province (No. SJCX23_0657 to LM).

Disclosure

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Bray F, Laversanne M, Sung H. et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* **2024**;74(3):229–263. doi:10.3322/caac.21834
2. Alduais Y, Zhang H, Fan F, et al. Non-small cell lung cancer (NSCLC): a review of risk factors, diagnosis, and treatment. *Medicine.* **2023**;102(8):e32899. doi:10.1097/MD.00000000000032899
3. Sun J, Wu S, Jin Z, et al. Lymph node micrometastasis in non-small cell lung cancer. *Biomed Pharmacother.* **2022**;149:112817. doi:10.1016/j.biopha.2022.112817
4. Phillips WJ, Leighl NB, Blais N, et al. Oral targeted therapy for the treatment of non-small cell lung carcinoma. *Cmaj.* **2024**;196(16):E558–e561. doi:10.1503/cmaj.231562
5. Mok TS, Wu Y-L, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med.* **2009**;361(10):947–957. doi:10.1056/NEJMoa0810699
6. Ettinger DS, Wood DE, Aisner DL, et al. NCCN Guidelines® Insights: non-Small Cell Lung Cancer, Version 2.2023. *J Natl Compr Canc Netw.* **2023**;21(4):340–350. doi:10.6004/jnccn.2023.0020
7. Harada G, Yang S-R, Cocco E, et al. Rare molecular subtypes of lung cancer. *Nat Rev Clin Oncol.* **2023**;20(4):229–249. doi:10.1038/s41571-023-00733-6
8. Moro-Sibilot D, Cozic N, Pérol M, et al. Crizotinib in c-MET- or ROS1-positive NSCLC: results of the AcSé phase II trial. *Ann Oncol.* **2019**;30(12):1985–1991. doi:10.1093/annonc/mdz407
9. Michels S, Massutí B, Schildhaus H-U, et al. Safety and Efficacy of Crizotinib in Patients With Advanced or Metastatic ROS1-Rearranged Lung Cancer (EUCROSS): a European Phase II Clinical Trial. *J Thorac Oncol.* **2019**;14(7):1266–1276. doi:10.1016/j.jtho.2019.03.020
10. Landi L, Chiari R, Tiseo M, et al. Crizotinib in MET -Deregulated or ROS1-Rearranged Pretreated Non-Small Cell Lung Cancer (METROS): a Phase II, Prospective, Multicenter, Two-Arms Trial. *Clin Cancer Res.* **2019**;25(24):7312–7319. doi:10.1158/1078-0432.CCR-19-0994
11. Drilon A, et al. Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three Phase 1–2 trials. *Lancet Oncol.* **2020**;21(2):261–270. doi:10.1016/S1470-2045(19)30690-4
12. Shaw AT, Solomon BJ, Chiari R, et al. Lorlatinib in advanced ROS1-positive non-small-cell lung cancer: a multicentre, open-label, single-arm, phase 1–2 trial. *Lancet Oncol.* **2019**;20(12):1691–1701. doi:10.1016/S1470-2045(19)30655-2
13. Lim SM, Kim HR, Lee J-S, et al. *Open-Label, Multicenter, Phase II Study of Ceritinib in Patients With Non-Small-Cell Lung Cancer Harboring ROS1 Rearrangement.* *J Clin Oncol.* **2017**;35(23):2613–2618. doi:10.1200/JCO.2016.71.3701
14. Drilon A, Camidge DR, Lin JJ, et al. Repotrectinib in ROS1 Fusion-Positive Non-Small-Cell Lung Cancer. *N Engl J Med.* **2024**;390(2):118–131. doi:10.1056/NEJMoa2302299

15. Subbiah V, et al. Efficacy of Vemurafenib in Patients With Non-Small-Cell Lung Cancer With BRAF V600 Mutation: an Open-Label, Single-Arm Cohort of the Histology-Independent VE-BASKET Study. *JCO Precis Oncol.* **2019**;3:1.
16. Mazieres J, Cropet C, Montané L, et al. Vemurafenib in non-small-cell lung cancer patients with BRAF(V600) and BRAF(nonV600) mutations. *Ann Oncol.* **2020**;31(2):289–294. doi:10.1016/j.annonc.2019.10.022
17. Gautschi O, Milia J, Cabarrou B, et al. Targeted Therapy for Patients with BRAF-Mutant Lung Cancer: results from the European EURAF Cohort. *J Thorac Oncol.* **2015**;10(10):1451–1457. doi:10.1097/JTO.0000000000000625
18. Planchard D, Kim TM, Mazieres J, et al. Dabrafenib in patients with BRAF(V600E)-positive advanced non-small-cell lung cancer: a single-arm, multicentre, open-label, Phase 2 trial. *Lancet Oncol.* **2016**;17(5):642–650. doi:10.1016/S1470-2045(16)00077-2
19. Planchard D, Besse B, Kim TM, et al. Updated survival of patients (pts) with previously treated BRAF V600E-mutant advanced non-small cell lung cancer (NSCLC) who received dabrafenib (D) or D + trametinib (T) in the phase II BR113928 study. *J Clin Oncol.* **2017**;35(15_suppl):9075. doi:10.1200/JCO.2017.35.15_suppl.9075
20. Planchard D, Besse B, Groen HJM, et al. Phase 2 Study of Dabrafenib Plus Trametinib in Patients With BRAF V600E-Mutant Metastatic NSCLC: updated 5-Year Survival Rates and Genomic Analysis. *J Thorac Oncol.* **2022**;17(1):103–115. doi:10.1016/j.jtho.2021.08.011
21. Planchard D, Besse B, Groen HJM, et al. Dabrafenib plus trametinib in patients with previously treated BRAF(V600E)-mutant metastatic non-small cell lung cancer: an open-label, multicentre phase 2 trial. *Lancet Oncol.* **2016**;17(7):984–993. doi:10.1016/S1470-2045(16)30146-2
22. Riely GJ, Smit EF, Ahn M-J, et al. Phase II, Open-Label Study of Encorafenib Plus Binimetinib in Patients With BRAF V600-Mutant Metastatic Non-Small-Cell Lung Cancer. *J Clin Oncol.* **2023**;41(21):3700–3711. doi:10.1200/JCO.23.00774
23. Hong DS, DuBois SG, Kummar S, et al. Larotrectinib in patients with TRK fusion-positive solid tumours: a pooled analysis of three phase 1/2 clinical trials. *Lancet Oncol.* **2020**;21(4):531–540. doi:10.1016/S1470-2045(19)30856-3
24. Drilon A, Tan DSW, Lassen UN, et al. Efficacy and Safety of Larotrectinib in Patients With Tropomyosin Receptor Kinase Fusion-Positive Lung Cancers. *JCO Precis Oncol.* **2022**;6(6):e2100418. doi:10.1200/PO.21.00418
25. Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1–2 trials. *Lancet Oncol.* **2020**;21(2):271–282. doi:10.1016/S1470-2045(19)30691-6
26. Demetri GD, De Braud F, Drilon A, et al. Updated Integrated Analysis of the Efficacy and Safety of Entrectinib in Patients With NTRK Fusion-Positive Solid Tumors. *Clin Cancer Res.* **2022**;28(7):1302–1312. doi:10.1158/1078-0432.CCR-21-3597
27. Krzakowski MJ, Lu S, Cousin S, et al. Updated analysis of the efficacy and safety of entrectinib in patients (pts) with locally advanced/metastatic NTRK fusion-positive (NTRK -fp) solid tumors. *J Clin Oncol.* **2022**;40(16_suppl):3099. doi:10.1200/JCO.2022.40.16_suppl.3099
28. Drilon A, Oxnard GR, Tan DSW, et al. Efficacy of Selpercatinib in RET Fusion-Positive Non-Small-Cell Lung Cancer. *N Engl J Med.* **2020**;383(9):813–824. doi:10.1056/NEJMoa2005653
29. Drilon A, Subbiah V, Gautschi O, et al. Selpercatinib in Patients With RET Fusion-Positive Non-Small-Cell Lung Cancer: updated Safety and Efficacy From the Registrational LIBRETTO-001 Phase I/II Trial. *J Clin Oncol.* **2023**;41(2):385–394. doi:10.1200/JCO.22.00393
30. Zhou C, Solomon B, Loong HH, et al. First-Line Selpercatinib or Chemotherapy and Pembrolizumab in RET Fusion-Positive NSCLC. *N Engl J Med.* **2023**;389(20):1839–1850. doi:10.1056/NEJMoa2309457
31. Gainor JF, Curigliano G, Kim D-W, et al. Pralsetinib for RET fusion-positive non-small-cell lung cancer (ARROW): a multi-cohort, open-label, phase 1/2 study. *Lancet Oncol.* **2021**;22(7):959–969. doi:10.1016/S1470-2045(21)00247-3
32. Griesinger F, Curigliano G, Thomas M, et al. Safety and efficacy of pralsetinib in RET fusion-positive non-small-cell lung cancer including as first-line therapy: update from the ARROW trial. *Ann Oncol.* **2022**;33(11):1168–1178. doi:10.1016/j.annonc.2022.08.002
33. Mazieres J, Paik PK, Garassino MC, et al. Tepotinib Treatment in Patients With MET Exon 14-Skipping Non-Small Cell Lung Cancer. *JAMA Oncol.* **2023**;9(9):1260–1266. doi:10.1001/jamaoncol.2023.1962
34. Wolf J, Garon EB, Groen HJM, et al. Capmatinib in MET exon 14-mutated, advanced NSCLC: updated results from the GEOMETRY mono-1 study. *J Clin Oncol.* **2021**;39(15_suppl):9020. doi:10.1200/JCO.2021.39.15_suppl.9020
35. Lu S, Fang J, Li X, et al. Long-Term Efficacy, Safety, and Subgroup Analysis of Savolitinib in Chinese Patients With NSCLCs Harboring MET Exon 14 Skipping Alterations. *JTO Clin Res Rep.* **2022**;3(10):100407. doi:10.1016/j.jtocrr.2022.100407
36. Krebs M, Spira AI, Cho BC, et al. Amivantamab in patients with NSCLC with MET exon 14 skipping mutation: updated results from the CHRYSALIS study. *J Clin Oncol.* **2022**;40(16_suppl):9008. doi:10.1200/JCO.2022.40.16_suppl.9008
37. Hong DS, Fakih MG, Strickler JH, et al. KRAS G12C Inhibition with Sotorasib in Advanced Solid Tumors. *N Engl J Med.* **2020**;383(13):1207–1217. doi:10.1056/NEJMoa1917239
38. Skoulidis F, Li BT, Dy GK, et al. Sotorasib for Lung Cancers with KRAS p.G12C Mutation. *N Engl J Med.* **2021**;384(25):2371–2381. doi:10.1056/NEJMoa2103695
39. de Langen AJ, Johnson ML, Mazieres J, et al. Sotorasib versus docetaxel for previously treated non-small-cell lung cancer with KRAS(G12C) mutation: a randomised, open-label, Phase 3 trial. *Lancet.* **2023**;401(10378):733–746. doi:10.1016/S0140-6736(23)00221-0
40. Ou SI, Jänne PA, Leal TA, et al. First-in-Human Phase I/IB Dose-Finding Study of Adagrasib (MRTX849) in Patients With Advanced KRAS G12C Solid Tumors (KRISTAL-1). *J Clin Oncol.* **2022**;40(23):2530–2538. doi:10.1200/JCO.21.02752
41. Jänne PA, Riely GJ, Gadgeel SM, et al. Adagrasib in Non-Small-Cell Lung Cancer Harboring a KRAS G12C Mutation. *N Engl J Med.* **2022**;387(2):120–131. doi:10.1056/NEJMoa2204619
42. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med.* **2015**;373(17):1627–1639. doi:10.1056/NEJMoa1507643
43. Li BT, Shen R, Buonocore D, et al. Ado-trastuzumab Emtansine for Patients With HER2-Mutant Lung Cancers: results From a Phase II Basket Trial. *J Clin Oncol.* **2018**;36(24):2532–2537. doi:10.1200/JCO.2018.77.9777
44. Li BT, Michelini F, Misale S, et al. HER2-Mediated Internalization of Cytotoxic Agents in ERBB2 Amplified or Mutant Lung Cancers. *Cancer Discovery.* **2020**;10(5):674–687. doi:10.1158/2159-8290.CD-20-0215
45. Iwama E, Zenke Y, Sugawara S, et al. Trastuzumab emtansine for patients with non-small cell lung cancer positive for human epidermal growth factor receptor 2 exon-20 insertion mutations. *Eur J Cancer.* **2022**;162:99–106. doi:10.1016/j.ejca.2021.11.021
46. Li BT, Smit EF, Goto Y, et al. Trastuzumab Deruxtecan in HER2-Mutant Non-Small-Cell Lung Cancer. *N Engl J Med.* **2022**;386(3):241–251. doi:10.1056/NEJMoa2112431

47. Zhou C, Li X, Wang Q, et al. Pyrotinib in HER2 -Mutant Advanced Lung Adenocarcinoma After Platinum-Based Chemotherapy: a Multicenter, Open-Label, Single-Arm, Phase II Study. *J Clin Oncol.* **2020**;38(24):2753–2761. doi:10.1200/JCO.20.00297
48. Song Z, Li Y, Chen S, et al. Efficacy and safety of pyrotinib in advanced lung adenocarcinoma with HER2 mutations: a multicenter, single-arm, phase II trial. *BMC Med.* **2022**;20(1):42. doi:10.1186/s12916-022-02245-z
49. Le X, Cornelissen R, Garassino M, et al. Pozotinib in Non–Small-Cell Lung Cancer Harboring HER2 Exon 20 Insertion Mutations After Prior Therapies: ZENITH20-2 Trial. *J Clin Oncol.* **2022**;40(7):710–718. doi:10.1200/JCO.21.01323
50. Sun S, Prelaj A, Baik C, et al. 26MO Efficacy and safety of pozotinib in treatment-naïve HER2 exon 20 insertion (ex20ins) mutated non-small cell lung cancer (NSCLC): ZENITH20-4. *Ann Oncol.* **2022**;33:S13. doi:10.1016/j.annonc.2022.01.035
51. Solomon BJ, Mok T, Kim D-W, et al. First-Line Crizotinib versus Chemotherapy in ALK-Positive Lung Cancer. *N Engl J Med.* **2014**;371(23):2167–2177. doi:10.1056/NEJMoa1408440
52. Shaw AT, Kim D-W, Nakagawa K, et al. Crizotinib versus Chemotherapy in Advanced ALK-Positive Lung Cancer. *N Engl J Med.* **2013**;368(25):2385–2394. doi:10.1056/NEJMoa1214886
53. Soria JC, Tan DSW, Chiari R, et al. First-line ceritinib versus platinum-based chemotherapy in advanced ALK-rearranged non-small-cell lung cancer (ASCEND-4): a randomised, open-label, phase 3 study. *Lancet.* **2017**;389(10072):917–929. doi:10.1016/S0140-6736(17)30123-X
54. Shaw AT, Kim TM, Crinò L, et al. Ceritinib versus chemotherapy in patients with ALK-rearranged non-small-cell lung cancer previously given chemotherapy and crizotinib (ASCEND-5): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol.* **2017**;18(7):874–886. doi:10.1016/S1470-2045(17)30339-X
55. Camidge DR, Kim HR, Ahn M-J, et al. Brigatinib versus Crizotinib in ALK-Positive Non–Small-Cell Lung Cancer. *N Engl J Med.* **2018**;379(21):2027–2039. doi:10.1056/NEJMoa1810171
56. Camidge DR, Kim HR, Ahn M-J, et al. Brigatinib Versus Crizotinib in ALK Inhibitor-Naïve Advanced ALK-Positive NSCLC: final Results of Phase 3 ALTA-1L Trial. *J Thorac Oncol.* **2021**;16(12):2091–2108. doi:10.1016/j.jtho.2021.07.035
57. Peters S, Camidge DR, Shaw AT, et al. Alectinib versus Crizotinib in Untreated ALK-Positive Non–Small-Cell Lung Cancer. *N Engl J Med.* **2017**;377(9):829–838. doi:10.1056/NEJMoa1704795
58. Peters S, Mok TSK, Gadgeel SM, et al. Updated overall survival (OS) and safety data from the randomized, phase III ALEX study of alectinib (ALC) versus crizotinib (CRZ) in untreated advanced ALK+ NSCLC. *J Clin Oncol.* **2020**;38(15_suppl):9518. doi:10.1200/JCO.2020.38.15_suppl.9518
59. Wolf J, Helland Å, Oh I-J, et al. Final efficacy and safety data, and exploratory molecular profiling from the phase III ALUR study of alectinib versus chemotherapy in crizotinib-pretreated ALK-positive non-small-cell lung cancer. *ESMO Open.* **2022**;7(1):100333. doi:10.1016/j.esmoop.2021.100333
60. Horn L, Wang Z, Wu G, et al. Ensartinib vs Crizotinib for Patients With Anaplastic Lymphoma Kinase-Positive Non-Small Cell Lung Cancer: a Randomized Clinical Trial. *JAMA Oncol.* **2021**;7(11):1617–1625. doi:10.1001/jamaoncol.2021.3523
61. Solomon BJ, Besse B, Bauer TM, et al. Lorlatinib in patients with ALK-positive non-small-cell lung cancer: results from a global phase 2 study. *Lancet Oncol.* **2018**;19(12):1654–1667. doi:10.1016/S1470-2045(18)30649-1
62. Shaw AT, Bauer TM, de Marinis F, et al. First-Line Lorlatinib or Crizotinib in Advanced ALK-Positive Lung Cancer. *N Engl J Med.* **2020**;383(21):2018–2029. doi:10.1056/NEJMoa2027187
63. Carriosa DR, Burkard ME, Elamin YY, et al. CRESTONE: initial efficacy and safety of seribantumab in solid tumors harboring NRG1 fusions. *J Clin Oncol.* **2022**;40(16_suppl):3006. doi:10.1200/JCO.2022.40.16_suppl.3006
64. Yu ZQ, Wang M, Zhou W, et al. ROS1-positive non-small cell lung cancer (NSCLC): biology, diagnostics, therapeutics and resistance. *J Drug Target.* **2022**;30(8):845–857. doi:10.1080/1061186X.2022.2085730
65. Charest A, Wilker EW, McLaughlin ME, et al. ROS fusion tyrosine kinase activates a SH2 domain-containing phosphatase-2/phosphatidylinositol 3-kinase/mammalian target of rapamycin signaling axis to form glioblastoma in mice. *Cancer Res.* **2006**;66(15):7473–7481. doi:10.1158/0008-5472.CAN-06-1193
66. Jun HJ, Johnson H, Bronson RT, et al. The oncogenic lung cancer fusion kinase CD74-ROS activates a novel invasiveness pathway through E-Syt1 phosphorylation. *Cancer Res.* **2012**;72(15):3764–3774. doi:10.1158/0008-5472.CAN-11-3990
67. Davies KD, Le AT, Theodoro MF, et al. Identifying and Targeting ROS1 Gene Fusions in Non–Small Cell Lung Cancer. *Clin Cancer Res.* **2012**;18(17):4570–4579. doi:10.1158/1078-0432.CCR-12-0550
68. Tan AC, Tan DSW. Targeted Therapies for Lung Cancer Patients With Oncogenic Driver Molecular Alterations. *J Clin Oncol.* **2022**;40(6):611–625. doi:10.1200/JCO.21.01626
69. Zhu Q, Zhan P, Zhang X, et al. Clinicopathologic characteristics of patients with ROS1 fusion gene in non-small cell lung cancer: a meta-analysis. *Transl Lung Cancer Res.* **2015**;4(3):300–309. doi:10.3978/j.issn.2218-6751.2015.05.01
70. Park S, Ahn B-C, Lim SW, et al. Characteristics and Outcome of ROS1-Positive Non-Small Cell Lung Cancer Patients in Routine Clinical Practice. *J Thorac Oncol.* **2018**;13(9):1373–1382. doi:10.1016/j.jtho.2018.05.026
71. US_FDA. FDA expands use of Xalkori to treat rare form of advanced non-small cell lung cancer | FDA. 2016 Fri, 03/11/2016-05:44, Available from: <https://www.fda.gov/news-events/press-announcements/fda-expands-use-xalkori-treat-rare-form-advanced-non-small-cell-lung-cancer>. Accessed Nov 8, 2024.
72. US_FDA. FDA approves entrectinib for NTRK solid tumors and ROS-1 NSCLC | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-entrectinib-ntrk-solid-tumors-and-ros-1-nsclc>. Accessed Nov 8, 2024.
73. US_FDA. FDA approves repotrectinib for ROS1-positive non-small cell lung cancer | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-repotrectinib-ros1-positive-non-small-cell-lung-cancer>. Accessed Nov 8, 2024.
74. D'Angelo A, Sobhani N, Chapman R, et al. Focus on ROS1-Positive Non-Small Cell Lung Cancer (NSCLC): crizotinib, Resistance Mechanisms and the Newer Generation of Targeted Therapies. *Cancers.* **2020**;12(11):3293. doi:10.3390/cancers12113293
75. Rolfo C, Ruiz R, Giovannetti E, et al. Entrectinib: a potent new TRK, ROS1, and ALK inhibitor. *Expert Opin Investig Drugs.* **2015**;24(11):1493–1500. doi:10.1517/13543784.2015.1096344
76. Dudnik E, Agbarya A, Grinberg R, et al. Clinical activity of brigatinib in ROS1-rearranged non-small cell lung cancer. *Clin Transl Oncol.* **2020**;22(12):2303–2311. doi:10.1007/s12094-020-02376-w
77. Hegde A, Hong DS, Behrang A, et al. Activity of Brigatinib in Crizotinib and Ceritinib-Resistant ROS1- Rearranged Non-Small-Cell Lung Cancer. *JCO Precis Oncol.* **2019**;3. doi:10.1200/po.18.00267

78. Mazieres J, Drilon A, Lusque A, et al. Immune checkpoint inhibitors for patients with advanced lung cancer and oncogenic driver alterations: results from the IMMUNOTARGET registry. *Ann Oncol*. 2019;30(8):1321–1328. doi:10.1093/annonc/mdz167
79. Rapp UR, Goldsborough MD, Mark GE, et al. Structure and biological activity of v-raf, a unique oncogene transduced by a retrovirus. *Proc Natl Acad Sci U S A*. 1983;80(14):4218–4222. doi:10.1073/pnas.80.14.4218
80. Bonner TI, Kerby SB, Sutcliffe P, et al. Structure and biological activity of human homologs of the raf/mil oncogene. *Mol Cell Biol*. 1985;5(6):1400–1407. doi:10.1128/mcb.5.6.1400-1407.1985
81. Bonner T, O'Brien SJ, Nash WG, et al. The Human Homologs of the raf (mil) Oncogene Are Located on Human Chromosomes 3 and 4. *Science*. 1984;223(4631):71–74. doi:10.1126/science.6691137
82. Liu F, Yang X, Geng M, et al. Targeting ERK, an Achilles' Heel of the MAPK pathway, in cancer therapy. *Acta Pharm Sin B*. 2018;8(4):552–562. doi:10.1016/j.apsb.2018.01.008
83. Frisone D, Friedlaender A, Malapelle U, et al. A BRAF new world. *Crit Rev Oncol Hematol*. 2020;152:103008. doi:10.1016/j.critrevonc.2020.103008
84. Couraud S, Barlesi F, Fontaine-Deraluelle C, et al. Clinical outcomes of non-small-cell lung cancer patients with BRAF mutations: results from the French Cooperative Thoracic Intergroup biomarkers France study. *Eur J Cancer*. 2019;116:86–97. doi:10.1016/j.ejca.2019.04.016
85. US_FDA. FDA grants accelerated approval to dabrafenib in combination with trametinib for unresectable or metastatic solid tumors with BRAF V600E mutation | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-dabrafenib-combination-trametinib-unresectable-or-metastatic-solid>. Accessed Nov 8, 2024.
86. US_FDA. FDA approves encorafenib with binimetinib for metastatic non-small cell lung cancer with a BRAF V600E mutation | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-encorafenib-binimetinib-metastatic-non-small-cell-lung-cancer-braf-v600e-mutation>. Accessed Nov 8, 2024.
87. Bollag G, Hirth P, Tsai J, et al. Clinical efficacy of a RAF inhibitor needs broad target blockade in BRAF-mutant melanoma. *Nature*. 2010;467(7315):596–599. doi:10.1038/nature09454
88. Puri M, Gawri K, Dawar R. Therapeutic strategies for BRAF mutation in non-small cell lung cancer: a review. *Front Oncol*. 2023;13:1141876. doi:10.3389/fonc.2023.1141876
89. Planchard D, Sanborn RE, Negrao MV, et al. BRAF(V600E)-mutant metastatic NSCLC: disease overview and treatment landscape. *NPJ Precis Oncol*. 2024;8(1):90. doi:10.1038/s41698-024-00552-7
90. Imyanitov EN, Mitiushkina NV, Kuligina ES, et al. Pathways and targeting avenues of BRAF in non-small cell lung cancer. *Expert Opin Ther Targets*. 2024;28(7):613–622. doi:10.1080/14728222.2024.2374742
91. Murciano-Goroff YR, Pak T, Mondaca S, et al. Immune biomarkers and response to checkpoint inhibition of BRAF(V600) and BRAF non-V600 altered lung cancers. *Br J Cancer*. 2022;126(6):889–898. doi:10.1038/s41416-021-01679-1
92. Negrao MV, Skoulidis F, Montesin M, et al. Oncogene-specific differences in tumor mutational burden, PD-L1 expression, and outcomes from immunotherapy in non-small cell lung cancer. *J Immunother Cancer*. 2021;9(8). doi:10.1136/jitc-2021-002891.
93. Otano I, Uvero AC, Zugazagoitia J, et al. At the crossroads of immunotherapy for oncogene-addicted subsets of NSCLC. *Nat Rev Clin Oncol*. 2023;20(3):143–159. doi:10.1038/s41571-022-00718-x
94. Martin-Zanca D, Hughes SH, Barbacid M. A human oncogene formed by the fusion of truncated tropomyosin and protein tyrosine kinase sequences. *Nature*. 1986;319(6056):743–748. doi:10.1038/319743a0
95. Liu F, Wei Y, Zhang H, et al. NTRK Fusion in Non-Small Cell Lung Cancer: diagnosis, Therapy, and TRK Inhibitor Resistance. *Front Oncol*. 2022;12:864666. doi:10.3389/fonc.2022.864666
96. Passiglia F, Caparica R, Giovannetti E, et al. The potential of neurotrophic tyrosine kinase (NTRK) inhibitors for treating lung cancer. *Expert Opin Investig Drugs*. 2016;25(4):385–392. doi:10.1517/13543784.2016.1152261
97. Cocco E, Scaltriti M, Drilon A. NTRK fusion-positive cancers and TRK inhibitor therapy. *Nat Rev Clin Oncol*. 2018;15(12):731–747. doi:10.1038/s41571-018-0113-0
98. US_FDA. FDA approves larotrectinib for solid tumors with NTRK gene fusions | FDA. 2024; Available from: <https://www.fda.gov/drugs/fda-approves-larotrectinib-solid-tumors-ntrk-gene-fusions-0>. Accessed Nov 8, 2024.
99. Drilon AE, Lin JJ, Kummar S, et al. Updated efficacy and safety of larotrectinib in patients with tropomyosin receptor kinase (TRK) fusion lung cancer. *J Clin Oncol*. 2022;40(16_suppl):9024. doi:10.1200/JCO.2022.40.16_suppl.9024
100. Paz-Ares L, Doebele RC, Farago AF, et al. Entrectinib in NTRK fusion-positive non-small cell lung cancer (NSCLC): integrated analysis of patients (pts) enrolled in STARTRK-2, STARTRK-1 and ALKA-372-001. *Ann Oncol*. 2019;30:ii48–ii49. doi:10.1093/annonc/mdz063.011
101. Takahashi M, Ritz J, Cooper GM. Activation of a novel human transforming gene, ret, by DNA rearrangement. *Cell*. 1985;42(2):581–588. doi:10.1016/0092-8674(85)90115-1
102. Subbiah V, Cote GJ. Advances in Targeting RET-Dependent Cancers. *Cancer Discov*. 2020;10(4):498–505. doi:10.1158/2159-8290.CD-19-1116
103. US_FDA. FDA approves selpercatinib for locally advanced or metastatic RET fusion-positive solid tumors | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-selpercatinib-locally-advanced-or-metastatic-ret-fusion-positive-solid-tumors>. Accessed Nov 8, 2024.
104. US_FDA. FDA approves pralsetinib for lung cancer with RET gene fusions | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-pralsetinib-lung-cancer-ret-gene-fusions>. Accessed Nov 8, 2024.
105. Subbiah V, Velcheti V, Tuch BB, et al. Selective RET kinase inhibition for patients with RET-altered cancers. *Ann Oncol*. 2018;29(8):1869–1876. doi:10.1093/annonc/mdy137
106. Markham A. Selpercatinib: first Approval. *Drugs*. 2020;80(11):1119–1124. doi:10.1007/s40265-020-01343-7
107. Guisier F, Dubos-Arvis C, Viñas F, et al. Efficacy and Safety of Anti-PD-1 Immunotherapy in Patients With Advanced NSCLC With BRAF, HER2, or MET Mutations or RET Translocation: GFPC 01-2018. *J Thorac Oncol*. 2020;15(4):628–636. doi:10.1016/j.jtho.2019.12.129
108. Offin M, et al. Immunophenotype and Response to Immunotherapy of RET-Rearranged Lung Cancers. *JCO Precis Oncol*. 2019;3:1.
109. Fujino T, Suda K, Mitsudomi T. Lung Cancer with MET exon 14 Skipping Mutation: genetic Feature, Current Treatments, and Future Challenges. *Lung Cancer*. 2021;12:35–50. doi:10.2147/LC.TT.S269307
110. Spagnolo CC, Ciappina G, Giovannetti E, et al. Targeting MET in Non-Small Cell Lung Cancer (NSCLC): a New Old Story? *Int J Mol Sci*. 2023;24(12):10119. doi:10.3390/ijms241210119

111. Han Y, Yu Y, Miao D, et al. Targeting MET in NSCLC: an Ever-Expanding Territory. *JTO Clin Res Rep*. 2024;5(2):100630. doi:10.1016/j.jto.2023.100630
112. US_FDA. *FDA approves tepotinib for metastatic non-small cell lung cancer* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-tepotinib-metastatic-non-small-cell-lung-cancer>. Accessed Nov 8, 2024.
113. US_FDA. *FDA grants accelerated approval to capmatinib for metastatic non-small cell lung cancer* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-capmatinib-metastatic-non-small-cell-lung-cancer>. Accessed Nov 8, 2024.
114. Markham A. Savolitinib: first Approval. *Drugs*. 2021;81(14):1665–1670. doi:10.1007/s40265-021-01584-0
115. Paik PK, Felip E, Veillon R, et al. Tepotinib in Non–Small-Cell Lung Cancer with MET Exon 14 Skipping Mutations. *N Engl J Med*. 2020;383(10):931–943. doi:10.1056/NEJMoa2004407
116. Le X, Sakai H, Felip E, et al. Tepotinib Efficacy and Safety in Patients with MET Exon 14 Skipping NSCLC: outcomes in Patient Subgroups from the VISION Study with Relevance for Clinical Practice. *Clin Cancer Res*. 2022;28(6):1117–1126. doi:10.1158/1078-0432.CCR-21-2733
117. Wolf J, Seto T, Han J-Y, et al. Capmatinib in MET Exon 14–Mutated or MET -Amplified Non–Small-Cell Lung Cancer. *N Engl J Med*. 2020;383(10):944–957. doi:10.1056/NEJMoa2002787
118. Lu S, Fang J, Li X, et al. 2MO Final OS results and subgroup analysis of savolitinib in patients with MET exon 14 skipping mutations (METex14+) NSCLC. *Ann Oncol*. 2022;33:S27. doi:10.1016/j.annonc.2022.02.011
119. Mazieres J, Vioix H, Pfeiffer BM, et al. MET Exon 14 Skipping in NSCLC: a Systematic Literature Review of Epidemiology, Clinical Characteristics, and Outcomes. *Clin Lung Cancer*. 2023;24(6):483–497. doi:10.1016/j.clcc.2023.06.008
120. Sabari JK, Leonardi GC, Shu CA, et al. PD-L1 expression, tumor mutational burden, and response to immunotherapy in patients with MET exon 14 altered lung cancers. *Ann Oncol*. 2018;29(10):2085–2091. doi:10.1093/annonc/mdy334
121. Cox AD, Der CJ. Ras history: the saga continues. *Small GTPases*. 2010;1(1):2–27. doi:10.4161/sgtp.1.1.12178
122. Shang Y, Fu S, Hao Q, et al. Multiple medicinal chemistry strategies of targeting KRAS: state-of-The art and future directions. *Bioorg Chem*. 2024;144:107092. doi:10.1016/j.bioorg.2023.107092
123. Chen QA, Lin W-H, Zhou -X-X, et al. Outcomes following KRAS(G12C) inhibitor treatment in patients with KRAS(G12C)-mutated solid tumors: a systematic review and meta-analysis. *Pharmacol Res*. 2024;200:107060. doi:10.1016/j.phrs.2024.107060
124. US_FDA. *FDA grants accelerated approval to sotorasib for KRAS G12C mutated NSCLC* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-sotorasib-kras-g12c-mutated-nsclc>. Accessed Nov 8, 2024.
125. US_FDA. *FDA grants accelerated approval to adagrasib for KRAS G12C-mutated NSCLC* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-adagrasib-kras-g12c-mutated-nsclc>. Accessed Nov 8, 2024.
126. Gadgeel S, Jänne PA, Spira AI, et al. MA06.04 KRYSTAL-1: two-Year Follow-Up of Adagrasib (MRTX849) Monotherapy in Patients with Advanced/Metastatic KRASG12C-Mutated NSCLC. *J Thorac Oncol*. 2023;18(11):S118. doi:10.1016/j.jtho.2023.09.152
127. Mok TSK, Lawler WE, Shum MK, et al. KRYSTAL-12: a randomized phase 3 study of adagrasib (MRTX849) versus docetaxel in patients (pts) with previously treated non-small-cell lung cancer (NSCLC) with KRAS G12C mutation. *J Clin Oncol*. 2021;39(15_suppl):TPS9129–TPS9129. doi:10.1200/JCO.2021.39.15_suppl.TPS9129
128. Nakajima EC, Ren Y, Vallejo JJ, et al. Outcomes of first-line immune checkpoint inhibitors with or without chemotherapy according to KRAS mutational status and PD-L1 expression in patients with advanced NSCLC: FDA pooled analysis. *J Clin Oncol*. 2022;40(16_suppl):9001. doi:10.1200/JCO.2022.40.16_suppl.9001
129. Bargmann CI, Hung MC, Weinberg RA. The neu oncogene encodes an epidermal growth factor receptor-related protein. *Nature*. 1986;319(6050):226–230. doi:10.1038/319226a0
130. Cheng H, Liu P, Ohlson C, et al. PIK3CA(H1047R)- and Her2-initiated mammary tumors escape PI3K dependency by compensatory activation of MEK-ERK signaling. *Oncogene*. 2016;35(23):2961–2970. doi:10.1038/onc.2015.377
131. US_FDA. *FDA grants accelerated approval to fam-trastuzumab deruxtecan-nxki for HER2-mutant non-small cell lung cancer*|FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-fam-trastuzumab-deruxtecan-nxki-her2-mutant-non-small-cell-lung>. Accessed Nov 8, 2024.
132. ClinicalTrials. *Phase 3 Study of Pyrotinib Versus Docetaxel in Patients With Advanced Non-squamous NSCLC Harboring a HER2 Exon 20 Mutation Who Failed Platinum Based Chemotherapy - Full Text View*. 2024; Available from: <https://clinicaltrials.gov/ct2/show/NCT04447118>. Accessed Nov 8, 2024.
133. Chau CH, Steeg PS, Figg WD. Antibody–drug conjugates for cancer. *Lancet*. 2019;394(10200):793–804. doi:10.1016/S0140-6736(19)31774-X
134. Hotta K, Aoe K, Kozuki T, et al. A Phase II Study of Trastuzumab Emtansine in HER2-Positive Non-Small Cell Lung Cancer. *J Thorac Oncol*. 2018;13(2):273–279. doi:10.1016/j.jtho.2017.10.032
135. ClinicalTrials. *A Study to Investigate the Efficacy and Safety of Trastuzumab Deruxtecan as the First Treatment Option for Unresectable, Locally Advanced/Metastatic Non-Small Cell Lung Cancer With HER2 Mutations - Full Text View*. 2024; Available from: <https://clinicaltrials.gov/ct2/show/NCT05048797>. Accessed Nov 8, 2024.
136. Fischer P, Nacheva E, Mason DY, et al. A Ki-1 (CD30)-positive human cell line (Karpas 299) established from a high-grade non-Hodgkin's lymphoma, showing a 2;5 translocation and rearrangement of the T-cell receptor beta-chain gene. *Blood*. 1988;72(1):234–240. doi:10.1182/blood.V72.1.234.234
137. Morris SW, Kirstein MN, Valentine MB, et al. Fusion of a Kinase Gene, ALK, to a Nucleolar Protein Gene, NPM, in Non-Hodgkin's Lymphoma. *Science*. 1994;263(5151):1281–1284. doi:10.1126/science.8122112
138. Parvaresh H, Roozitalab G, Golandam F, et al. Unraveling the Potential of ALK-Targeted Therapies in Non-Small Cell Lung Cancer: comprehensive Insights and Future Directions. *Biomedicine*. 2024;12(2):297. doi:10.3390/biomedicine12020297
139. US_FDA. *Alectinib approved for (ALK) positive metastatic non-small cell lung cancer (NSCLC)*|FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/alectinib-approved-alk-positive-metastatic-non-small-cell-lung-cancer-nsclc>. Accessed Nov 8, 2024.
140. US_FDA. *FDA approves brigatinib for ALK-positive metastatic NSCLC* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-brigatinib-alk-positive-metastatic-nsclc>. Accessed Nov 8, 2024.

141. US_FDA. *FDA broadens ceritinib indication to previously untreated ALK-positive metastatic NSCLC* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-broadens-ceritinib-indication-previously-untreated-alk-positive-metastatic-nsclc>. Accessed Nov 8, 2024.
142. Kazandjian D, Blumenthal GM, Chen H-Y, et al. FDA approval summary: crizotinib for the treatment of metastatic non-small cell lung cancer with anaplastic lymphoma kinase rearrangements. *Oncologist*. 2014;19(10):e5–11. doi:10.1634/theoncologist.2014-0241
143. US_FDA. *FDA approves lorlatinib for metastatic ALK-positive NSCLC* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-lorlatinib-metastatic-alk-positive-nsclc>. Accessed Nov 8, 2024.
144. Li J, Knoll S, Bocharova I, et al. Comparative efficacy of first-line ceritinib and crizotinib in advanced or metastatic anaplastic lymphoma kinase-positive non-small cell lung cancer: an adjusted indirect comparison with external controls. *Curr Med Res Opin*. 2019;35(1):105–111. doi:10.1080/03007995.2018.1541443
145. Solomon BJ, Bauer TM, Mok TSK, et al. Efficacy and safety of first-line lorlatinib versus crizotinib in patients with advanced, ALK-positive non-small-cell lung cancer: updated analysis of data from the phase 3, randomised, open-label CROWN study. *Lancet Respir Med*. 2023;11(4):354–366. doi:10.1016/S2213-2600(22)00437-4
146. Schram AM, Odintsov I, Espinosa-Cotton M, et al. Zenocutuzumab, a HER2xHER3 Bispecific Antibody, Is Effective Therapy for Tumors Driven by NRG1 Gene Rearrangements. *Cancer Discov*. 2022;12(5):1233–1247. doi:10.1158/2159-8290.CD-21-1119
147. Kazdal D, Hofman V, Christopoulos P, et al. Fusion-positive non-small cell lung carcinoma: biological principles, clinical practice, and diagnostic implications. *Genes Chromosomes Cancer*. 2022;61(5):244–260. doi:10.1002/gcc.23022
148. Fernandez-Cuesta L, Thomas RK. Molecular Pathways: targeting NRG1 Fusions in Lung Cancer. *Clin Cancer Res*. 2015;21(9):1989–1994. doi:10.1158/1078-0432.CCR-14-0854
149. Drilon A, Duruisseaux M, Han J-Y, et al. Clinicopathologic Features and Response to Therapy of NRG1 Fusion–Driven Lung Cancers: the eNRGy1 Global Multicenter Registry. *J Clin Oncol*. 2021;39(25):2791–2802. doi:10.1200/JCO.20.03307
150. Schram AM, O'Reilly EM, O'Kane GM, et al. Efficacy and safety of zenocutuzumab in advanced pancreas cancer and other solid tumors harboring NRG1 fusions. *J Clin Oncol*. 2021;39(15_suppl):3003. doi:10.1200/JCO.2021.39.15_suppl.3003
151. ClinicalTrials. *A Study of Zenocutuzumab (MCLA-128) in Patients With Solid Tumors Harboring an NRG1 Fusion (eNRGy) - Full Text View*. 2024; Available from: <https://clinicaltrials.gov/ct2/show/NCT02912949>. Accessed Nov 8, 2024.
152. ClinicalTrials. *Study of Seribantumab in Adult Patients With NRG1 Gene Fusion Positive Advanced Solid Tumors - Full Text View*. 2024; Available from: <https://clinicaltrials.gov/ct2/show/NCT04383210>. Accessed Nov 8, 2024.
153. Kharitonov A, Adams AC. Inventing new medicines: the FGF21 story. *Mol Metab*. 2014;3(3):221–229. doi:10.1016/j.molmet.2013.12.003
154. Ornitz DM, Itoh N. The Fibroblast Growth Factor signaling pathway. *WIREs Developmental Biology*. 2015;4(3):215–266. doi:10.1002/wdev.176
155. Chen J, Xu C, Lv J, et al. Clinical characteristics and targeted therapy of different gene fusions in non-small cell lung cancer: a narrative review. *Transl Lung Cancer Res*. 2023;12(4):895–908. doi:10.21037/tlcr-22-566
156. US_FDA. *FDA grants accelerated approval to erdafitinib for metastatic urothelial carcinoma* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-erdafitinib-metastatic-urothelial-carcinoma>. Accessed Nov 8, 2024.
157. US_FDA. *FDA grants accelerated approval to pemigatinib for cholangiocarcinoma with an FGFR2 rearrangement or fusion* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-pemigatinib-cholangiocarcinoma-fgfr2-rearrangement-or-fusion>. Accessed Nov 8, 2024.
158. US_FDA. *FDA grants accelerated approval to futibatinib for cholangiocarcinoma* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-futibatinib-cholangiocarcinoma>. Accessed Nov 8, 2024.
159. US_FDA. *FDA grants accelerated approval to infigratinib for metastatic cholangiocarcinoma* | FDA. 2024; Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-infigratinib-metastatic-cholangiocarcinoma>. Accessed Nov 8, 2024.
160. Barbar J, Armach M, Hodroj MH, et al. Emerging genetic biomarkers in lung adenocarcinoma. *SAGE Open Med*. 2022;10:20503121221132352. doi:10.1177/20503121221132352
161. Whitman M, Kaplan DR, Schaffhausen B, et al. Association of phosphatidylinositol kinase activity with polyoma middle-T competent for transformation. *Nature*. 1985;315(6016):239–242. doi:10.1038/315239a0
162. Courtney KD, Corcoran RB, Engelman JA. The PI3K pathway as drug target in human cancer. *J Clin Oncol*. 2010;28(6):1075–1083. doi:10.1200/JCO.2009.25.3641
163. Dienstmann R, Rodon J, Serra V, et al. Picking the point of inhibition: a comparative review of PI3K/AKT/mTOR pathway inhibitors. *Mol Cancer Ther*. 2014;13(5):1021–1031. doi:10.1158/1535-7163.MCT-13-0639
164. Guo Y, Cao R, Zhang X, et al. Recent Progress in Rare Oncogenic Drivers and Targeted Therapy For Non-Small Cell Lung Cancer. *Onco Targets Ther*. 2019;12:10343–10360. doi:10.2147/OTT.S230309
165. Levy B, Spira A, Becker D, et al. A randomized, phase 2 trial of Docetaxel with or without PX-866, an irreversible oral phosphatidylinositol 3-kinase inhibitor, in patients with relapsed or metastatic non-small-cell lung cancer. *J Thorac Oncol*. 2014;9(7):1031–1035. doi:10.1097/JTO.0000000000000183
166. Adjei AA, Bennis J, Leighl NB, et al. Safety and efficacy of buparlisib (BKM120) and chemotherapy in advanced, squamous non-small cell lung cancer (sqNSCLC): results from the phase Ib/II BASALT-2 and BASALT-3 studies. *J Clin Oncol*. 2016;34(15_suppl):e20522–e20522. doi:10.1200/JCO.2016.34.15_suppl.e20522
167. Tan DS-W, Lim KH, Tai WM, et al. A phase Ib safety and tolerability study of a pan class I PI3K inhibitor buparlisib (BKM120) and gefitinib (gef) in EGFR TKI-resistant NSCLC. *J Clin Oncol*. 2013;31(15_suppl):8107. doi:10.1200/jco.2013.31.15_suppl.8107
168. Langer CJ, Redman MW, Wade JL, et al. SWOG S1400B (NCT02785913), a Phase II Study of GDC-0032 (Taselisib) for Previously Treated PI3K-Positive Patients with Stage IV Squamous Cell Lung Cancer (Lung-MAP Sub-Study). *J Thorac Oncol*. 2019;14(10):1839–1846. doi:10.1016/j.jtho.2019.05.029
169. Fridman R, Agarwal G. New Concepts on the Interactions of Discoidin Domain Receptors with Collagen. *Biochim Biophys Acta Mol Cell Res*. 2019;1866(11):118527. doi:10.1016/j.bbamcr.2019.118527

170. Miao L, Wang Y, Zhu S, et al. Identification of novel driver mutations of the discoidin domain receptor 2 (DDR2) gene in squamous cell lung cancer of Chinese patients. *BMC Cancer*. 2014;14(1):369. doi:10.1186/1471-2407-14-369
171. Lee M-S, Jung EA, An SB, et al. Prevalence of Mutations in Discoidin Domain-Containing Receptor Tyrosine Kinase 2 (DDR2) in Squamous Cell Lung Cancers in Korean Patients. *Cancer Res Treat*. 2017;49(4):1065–1076. doi:10.4143/crt.2016.347
172. Xu X, Yu T, Wang Z. Discoidin Domain Receptor 2: a New Target in Cancer. *Oncol Res Treat*. 2022;45(4):205–215. doi:10.1159/000519645
173. Day E, Waters B, Spiegel K, et al. Inhibition of collagen-induced discoidin domain receptor 1 and 2 activation by imatinib, nilotinib and dasatinib. *Eur J Pharmacol*. 2008;599(1–3):44–53. doi:10.1016/j.ejphar.2008.10.014
174. Kelley MJ, Jha G, Shoemaker D, et al. Phase II Study of Dasatinib in Previously Treated Patients with Advanced Non-Small Cell Lung Cancer. *Cancer Invest*. 2017;35(1):32–35. doi:10.1080/07357907.2016.1253710
175. Spraker H, Price SL, Chaturvedi A, et al. The clone wars - revenge of the metastatic rogue state: the sarcoma paradigm. *Front Oncol*. 2012;2:2. doi:10.3389/fonc.2012.00002
176. Zhang K, Wang W, Zhang T, et al. Efficacy and treatment-related adverse events of multi-targeted tyrosine kinase inhibitors in advanced non-small-cell lung cancer: a meta-analysis of randomized controlled trials. *Int J Clin Pharm*. 2022;44(6):1232–1246. doi:10.1007/s11096-022-01465-w

OncoTargets and Therapy

Dovepress

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>