ORIGINAL RESEARCH

# Negative Affect and Impulsivity in Adolescents with Non-Suicidal Self-Injury: The Moderating and Mediating Roles of Perceived Social Support

Bing Pan<sup>1</sup>, Yun Gong<sup>2</sup>, Meiyi Guo<sup>3</sup>, Xiaohua Wang<sup>1</sup>, Haitao Lu<sup>1</sup>, Zheng Lin<sup>1</sup>, Bingren Zhang<sup>3</sup>

<sup>1</sup>Department of Psychiatry, Second Affiliated Hospital, Zhejiang University School of Medicine, Zhejiang University, Hangzhou, 310009, People's Republic of China; <sup>2</sup>Department of Pediatrics, Hangzhou First People's Hospital, Hangzhou, 310009, People's Republic of China; <sup>3</sup>Department of Medical Psychology, Affiliated Hospital (School of Clinical Medicine), Hangzhou Normal University, Hangzhou, 311121, People's Republic of China

Correspondence: Bingren Zhang, Department of Medical Psychology, Affiliated Hospital (School of Clinical Medicine), Hangzhou Normal University, Hangzhou, People's Republic of China, Tel +13757148402, Email bingrenz@hznu.edu.cn; Zheng Lin, Department of Psychiatry, Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, People's Republic of China, Tel +13757118261, Email 2194026@zju.edu.cn

**Background:** Negative affective states and impulsivity have been found to be closely associated with non-suicidal self-injury (NSSI), but how negative affect contributes to impulsivity in NSSI adolescents and the role of perceived social support remain unclear.

**Methods:** In Study 1, the Barratt Impulsiveness Scale (BIS-11) was administered on 225 NSSI adolescents and 225 healthy volunteers to determine the characteristics of impulsivity in the patients. In Study 2, 102 of the patients were randomly selected to further complete the Zung Self-rating Anxiety Scale (SAS), Zung Self-rating Depression Scale (SDS), and Multidimensional Scale of Perceived Social Support (MSPSS) to clarify the relationships between negative affect, perceived social support, and impulsivity in these patients.

**Results:** In Study 1, NSSI adolescents scored higher than volunteers on almost all factors of BIS-11 except for Cognitive Instability. In Study 2, Support from significant others mediated the association between SAS and BIS-11 Self-control. Moreover, Support from significant others and family moderated the prediction of SAS score on their BIS-11 Cognitive Complexity.

**Conclusion:** Compared to healthy adolescents, NSSI individuals were more impulsive, while perceived support from family and significant others prevented the anxious NSSI adolescents from lower cognitive complexity. Therefore, more social support should be available to NSSI adolescents with high anxiety to reduce their risk of self-harm due to high impulsivity.

Keywords: adolescents, impulsivity, negative affect, non-suicidal self-injury, perceived social support

### Introduction

Accepted: 23 December 2024 Published: 1 January 2025

Non-suicidal self-injury (NSSI) refers to the deliberate infliction of harm upon one's own body through self-inflicted destruction of tissues (eg, cutting or burning oneself) for socially unacceptable reasons in the absence of suicidal intent.<sup>1</sup> It is one of the strongest predictors of future suicide attempts.<sup>2</sup> Often, NSSI starts in the early teens and can persist for several years.<sup>3</sup> An earlier review showed that although the cultural context, causes and meanings of NSSI differ between individuals in developing and developed countries, the prevalence of NSSI is essentially the same.<sup>4</sup> However, a more recent meta-analysis reported a female predominance of NSSI among adolescents in North America and Europe, and comparable rates among male and female adolescents in Asia.<sup>5</sup> During 2010–2021, the global prevalence of NSSI among adolescents in a nonclinical sample was 22.0% over lifetime and 23.2% during the previous 12 months.<sup>6</sup> While in a psychiatric clinical sample, up to 40–60% of the adolescents exhibited NSSI.<sup>7</sup> In recent years, this condition has demonstrated a trend towards younger age, with 5.1–24% of the patients reporting self-injury before the age of 11–13 years<sup>8</sup>, which greatly increases the likelihood of accidental injuries and is linked to high psychiatric comorbidity.<sup>9</sup>

Evidence suggests that impulsivity plays an important role in adolescent self-harming behaviors. The Integrated Motivational-Volitional Model posits that impulsivity serves as a proximal volitional moderator of self-harming

© 2025 Pan et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (http://creativecommons.org/licenses/by-m/3.0/). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (http://www.dovepress.com/terms.php).

I

behaviors, facilitating the translation of intention into action,<sup>10</sup> this is particularly evident among adolescents<sup>11</sup> and patients diagnosed with certain affective disorders.<sup>12</sup> From a life history theory perspective, individuals who grow up in threatening environments are more likely to be impulsive, to have more distressing and provocative experiences,<sup>13</sup> and to engage in more impulsive behaviors,<sup>14</sup> including recurrent NSSIs in the context of emotional dysregulation.<sup>15</sup> Furthermore, different dimensions of impulsivity lead to variable self-harm risks. For example, a systematic study has shown that emotional impulsivity is associated with the initiation of self-harm, cognitive impulsivity is associated with the context to self-harm under negative emotional conditions.<sup>16</sup>

Further evidence indicates that individuals with heightened negative affect are more likely to engage in impulsive behavior, and the goal of relief from negative affect may drive impulsive actions for short-term gain over long-term objectives.<sup>16</sup> For instance, in individuals with anorexia nervosa, depression and anxiety were found to be related to attentional impulsivity;<sup>17</sup> Similarly, in patients with ankylosing spondylitis, depression was correlated with non-planning impulsivity.<sup>18</sup> Furthermore, research has demonstrated that negative affect, particularly depression and anxiety symptoms, exerts a significant influence on NSSI.<sup>19–22</sup>. Difficulties in emotion regulation and depression act as mediators in the relationship between the behavioral inhibition/ activation system and adolescent NSSI behaviors.<sup>23</sup> Nevertheless, it should be noted that not all adolescents with negative affect have high levels of uncontrolled impulsivity and related NSSI behaviors.

Perceived social support has been shown to play a protective role in relation to cognitive and emotional impulsivity as manifested by suicidal ideation in self-harmers<sup>24</sup> and behavioral impulsivity such as problematic internet gaming.<sup>25</sup> It also serves to safeguard individuals from traumatic events<sup>26</sup> and psychiatric symptoms,<sup>27</sup> and this function is similar in males and females.<sup>28</sup> However, individuals with NSSI typically report less perceived social support than those without NSSI,<sup>29,30</sup> no matter in a collectivist or individualist culture.<sup>31</sup> Similarly, in a study of impulsive killing among rural Chinese adolescents with suicidal behavior, high impulsivity patients had less social support compared with low impulsivity patients.<sup>32</sup> It is therefore concluded that lower perceived social support may lead to an increase in cognitive, emotional and behavioral impulsivity in NSSI adolescents.

In addition, prior research has identified a strong correlation between elevated perceived social support, particularly from parents and peers, and diminished negative cognitive and emotional strategies, depression levels, and suicidal ideation in young people with major depressive disorder.<sup>33,34</sup> Meanwhile, a positive correlation has been found between perceived informal social support from family, friends and significant others and positive affect.<sup>26</sup> Therefore, higher perceived social support may lead to lower impulsivity in NSSI adolescents by reducing their negative affect, as reported by Lee et al.<sup>35</sup> However, one might also be curious about whether negative affect leads to lower perceived social support and therefore to increased impulsivity in NSSI adolescents. The available evidence showed that patients with negative effects such as depression experienced significant social impairments, not only receiving less social support than healthy individuals, but also perceiving themselves as having impoverished social support networks,<sup>36</sup> and that those who perceived inadequate social support were less protected from impulsivity.<sup>24</sup> Moreover, it is unclear how perceived social support moderates the relationship between negative affect and impulsivity in NSSI adolescents. A study in addiction recovery demonstrated that an ecological momentary intervention targeting perceived social support was effective in reducing impulsivity and enhancing self-regulation.<sup>37</sup>

The objective of this study was to investigate the role of perceived social support in the relationship between affective states and impulsivity in adolescents with NSSI. Two studies were designed for this purpose. In Study 1, the dimensions of impulsivity were compared in detail between patients with NSSI and healthy adolescents using the 11<sup>th</sup> revision of the Barratt Impulsiveness Scale (BIS-11). It was hypothesized that, in comparison with healthy adolescents, patients with NSSI would display higher levels of cognitive, emotional, as well as behavioral impulsivity. Study 2 further examined the relationship between negative affect, perceived social support, and impulsivity in adolescents with NSSI. The following hypotheses were put forth: (1) negative affect megatively while perceived social support; (2) perceived social support acted as a mediator and moderator in the relationship between negative affect and their impulsivity.

# Methods

### Subjects

In Study 1, 225 Chinese adolescents with a history of NSSI were recruited into the NSSI group from inpatient and outpatient settings in a local top hospital from 1 January to 30 June 2024. Psychiatric diagnoses were made by two psychiatrists (BP and ZL) based on the adolescents' responses to a psychiatric interview (ie, the Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version, K-SADS-PL)<sup>38</sup> and ancillary information obtained from medical records and treatment staff. An equal number of healthy volunteers were recruited for the control group (Controls) from five middle schools in Hangzhou. They were excluded if they had a psychiatric disorder (eg, mental retardation, schizophrenia, or autism) that might interfere with the comprehension of task instructions, or if they were in state custody. All the participants completed the BIS-11 test individually in a rest room. The sociodemographic information of the participants in the two groups were showed in Table 1.

Meanwhile, all hospitalised NSSI patients among the Study 1 participants were included in Study 2, totalling 102 NSSI adolescents. They were further given questionnaires (the Zung Self-rating Anxiety Scale, the Zung Self-rating Depression Scale, and the Multidimensional Scale of Perceived Social Support) by two psychiatrists (HL and XW) to assess negative affect in the past week and perceived social support. Their socio-demographic characteristics and BIS-11 scores are shown in Table 2.

The study was conducted according to the principles of the Declaration of Helsinki. Ethical approval for this study was obtained from the Second Affiliated Hospital of Zhejiang University School of Medicine Approval (No: 20240075). After completely describing the study to the participants and their guardians, written informed consent was obtained from all of them.

### Measures

### Zung Self-Rating Anxiety Scale (SAS)

The SAS is a self-report scale with 20 items covering various symptoms of anxiety, including psychological and somatic items, and is scored on a 4-point scale ranging from 1 (none, or a little of the time) to 4 (most or all of the time). The

Sociodemographic C	haracteristics	NSSI (n, %)	Controls (n, %)	t/U	Р
Gender				24,750.00	0.567
Male		47 (20.89%)	51 (22.67%)		
Female		178 (79.11%)	174 (77.33%)		
Age	Mean ± S.D.	15.13 ±1.92	15.05 ±1.78	0.43	0.665
	Range	11–19	12–19		
Education level				23,962.50	0.243
Primary school		135 (60.00%)	147 (65.33%)		
Junior high school		90 (40.00%)	78 (34.77%)		
Family structure				23,877.00	0.266
Big family (parents, ch together)	nildren and grandparents	62 (27.56%)	53 (23.55%)		
Nuclear family (paren	ts and children)	120 (53.33%)	157 (69.78%)		
Single- parents family		27 (12.00%)	8 (3.56%)		
Children "Left-behind	,,	13 (5.78%)	6 (2.67%)		
Others		3 (1.33%)	I (0.44%)		

Table I Sociodemographic Characteristics in the NSSI Group (n = 225) and Controls (n = 225)

Abbreviation: NSSI, non-suicidal self-injury.

Sociodemographic Characteristics	n (%)	BIS-11	t/F	Р
Gender			1.54	0.225
Male	9 (8.80)	36.44± 6.6		
Female	93 (91.20)	148.15±22.24		
Age			1.99	0.497
~14	32 (31.37)	153.44±23.20		
15 ~ 19	70 (68.63)	144.23±20.95		
Education level			1.57	0.119
Primary school	49 (48.04)	150.65±21.77		
Junior high school	53 (51.96)	143.85±21.89		
Family structure			1.97	0.106
Big family (parents, children and grandparents together)	33 (32.35)	146.00±22.21		
Nuclear family (parents and children)	51 (50.00)	145.14±21.66		
Single- parents family	9 (8.73)	162.00±17.06		
Children "Left-behind"	7 (6.86)	150.00±22.92		
Others	2 (1.96)	110		

 Table 2 Sociodemographic Characteristics and Scores on BIS-11 in Non-Suicidal Self -Injury

 Adolescents (n = 102)

Abbreviation: BIS-11, the Barratt Impulsiveness Scale.

higher the score on the SAS, the more obvious the level of anxiety over the past week. The SAS has satisfactory psychometric properties.<sup>39</sup> Its Chinese version has been showed to be reliable and valid in previous studies,<sup>40</sup> including adolescents aged 13–18, with a Cronbach's alpha of 0.89.<sup>41</sup>

### Zung Self-Rating Depression Scale (SDS)

The SDS contains 20 self-report items designed to assess one's subjective depressive level. Each item is scored on a Likert scale ranging from 1 (none, or a little of the time) to 4 (most, or all of the time) according to the frequency of symptoms over the past week.<sup>42</sup> It has been proven to be reliable and valid in Chinese population,<sup>43</sup> and was used in young clinical patients.<sup>44</sup>

### Multidimensional Scale of Perceived Social Support (MSPSS)

MSPSS is a self-reported questionnaire that comprises 12 items, which measure the participants' perceived social support from three informal sources: family, friends, and significant others.<sup>45</sup> Participants rated their responses on a 7-point Likert scale (from 1 = "very strongly disagree" to 7 = "very strongly agree"). Higher scores indicate higher levels of perceived social support. Its Chinese version has been proven to be reliable and valid in a previous study.<sup>46</sup>

### The Barratt Impulsiveness Scale

The BIS-11 comprises 30 self-report items assessing impulsivity. It is scored on a Likert scale (ranging from never = 1 point to very frequently = 4 points). The scale consists of six first-order factors: Attention and Cognitive instability (measuring attentional impulsivity); Motor and Perseverance (measuring motor impulsivity); Self-controland and Cognitive complexity (measuring non-planning impulsivity).<sup>47,48</sup> The Chinese version of the scale was found to be valid and reliable in adolescents.<sup>49</sup> It has also been employed in patients with suicidal behavior in China, with Cronbach's alphas for the three BIS subscales were 0.74-0.81 for attempters.<sup>50</sup>

4

### Statistical Analyses

The data was analyzed using IBM SPSS v20.0 and StataMP 18.0. In Study 1, the Student's *t*-test and Mann–Whitney *U*-test were applied to compare the sociodemographic characteristics and the mean BIS-11 scale scores between patients and healthy controls. The effect sizes (Cohen's d values) of group comparisons were calculated.

In Study 2, Student's *t*-test and one-way ANOVA were used to compare the BIS-11 scores in patients with NSSI who differed in terms of sex, age range, educational levels, and family structures. In addition, Pearson's correlational test was applied to search for relationships among SAS/SDS, MSPSS, and BIS in the NSSI group. Taking the scores of SAS/SDS as independent variables, subscales of BIS as dependent variables, and subscales of MSPSS as mediators or moderators, the effect of perceived social support on the relationship between their negative affect and impulsivity was explored with the Process plug-in in SPSS v20.0 (for the theoretical models, see Figure 1). A p-value of less than 0.05 was considered significant. We also conducted a sensitivity analysis of the mediation model using the medsens command in StataMP 18.0, assuming that there was an unobserved continuous confounder that followed a normal distribution and was associated with the mediator and outcome, and we tested how the effects would change if the confounder was included.

# Results

### Study I: Impulsivity Characteristics in NSSI Adolescents

Sociodemographic characteristics, such as sex, age range, educational levels, and family structures, were matched between the NSSI and control groups (all Ps > 0.05) (see Table 1). Upon comparing BIS dimensional scores, patients with NSSI obtained higher scores on BIS-11 Attention (Cohen's d = 0.81), Motor (Cohen's d = 0.31), Self-control (Cohen's d = 0.47), Cognitive complexity (Cohen's d = 0.78), as well as Perseverance (Cohen's d = 0.40) than controls (see Table 3).

# Study 2: Negative Affect, Perceived Social Support, and Impulsivity in NSSI Adolescents

### Sociodemographic Characteristics and Correlations Between Factors

No significant difference was found in the BIS score among adolescents with NSSI who differed in terms of sex, age range, education levels, or family structures (all Ps > 0.05) (see Table 2).



Figure I The theoretical (A) mediation and (B) moderation models in this study. Each model consists of 36 sub-models based on the potential mediating and moderating roles of three dimensions of perceived social support between SAS/ SDS and six BIS-11 factors in NSSI adolescents, respectively.

	NSSI	Controls	t	df	Р			
BIS-11								
Attention	13.88±2.85	11.21±3.68	8.59	444	<0.001			
Cognitive Instability	6.61±1.89	6.83±1.73	-1.32	445	0.186			
Motor	14.67±4.37	13.46±3.41	3.26	421	0.001			
Perseverance	7.96±2.07	7.16±1.91	4.25	444	<0.001			
Self -control	15.53±3.26	13.83±3.90	4.99	442	<0.001			
Cognitive Complexity	12.71±2.86	10.60±2.56	8.18	442	<0.001			

**Table 3** Scale Scores (Mean  $\pm$  S.D.) of BIS-11 in the NSSI Group (n = 225) and Controls (n = 225)

Abbreviations: BIS-11, the Barratt Impulsiveness Scale; NSSI, non-suicidal self-injury.

**Table 4** Correlations Between Scores on SAS, SDS, MSPSS, and the Subscales of BIS-11 in Non-Suicidal Self -Injury Adolescents (n=102)

		SAS	SDS	MSPSS				
				Support from Family	Support from Friends	Support from Significant Others		
SAS	r	_	0.81	-0.43	-0.22	-0.31		
	Ρ	-	<0.001	<0.001	0.035	0.002		
SDS	r	-	-	-0.5 I	-0.21	-0.41		
	Ρ	-	-	<0.001	0.042	<0.001		
BIS-I I								
Attention	r	0.33	0.31	-0.15	-0.11	-0.20		
	Ρ	0.001	0.002	0.146	0.276	0.045		
Cognitive instability	r	0.07	-0.05	0.09	0.01	0.06		
	Ρ	0.507	0.645	0.376	0.934	0.595		
Motor	r	0.16	0.16	-0.09	0.01	-0.05		
	Ρ	0.107	0.107	0.365	0.936	0.663		
Perseverance	r	0.25	0.25	-0.15	-0.02	-0.13		
	Ρ	0.012	0.011	0.148	0.865	0.191		
Self-control	r	0.20	0.32	-0.23	-0.06	-0.29		
	Ρ	0.046	0.001	0.026	0.541	0.004		
Cognitive complexity	r	0.24	0.23	-0.12	-0.02	-0.10		
	Ρ	0.017	0.020	0.241	0.885	0.339		

Abbreviations: SAS, the Zung Self-rating Anxiety Scale; SDS, the Zung Self-rating Depression Scale; MSPSS, the Multidimensional Scale of Perceived Social Support; BIS-II, the Barratt Impulsiveness Scale.

The dichotomous correlations are shown in Table 4. SAS and SDS scores were positively correlated with BIS-11 Attention, Perseverance, Self-control, and Cognitive complexity (Ps < 0.05). In contrast, MSPSS Support from family was negatively correlated with BIS-11 Self-control; MSPSS Support from significant others was negatively correlated with BIS-11 Attention and Self-control (Ps<0.05). In addition, all the MSPSS subscale scores were negatively correlated with the SAS and SDS scores (Ps < 0.05).

### Mediating Role of Perceived Social Support in the Relationship Between Negative Affect and Impulsivity

Taking the scores of SAS/SDS as independent variables, the six subscales of BIS as dependent variables, and the three subscales of MSPSS as mediators respectively, the mediation analysis showed that Support from significant others played a fully mediating role in SAS and BIS-11 Self-control (a = -0.15, p < 0.01; b = -0.14, p < 0.5; c = 0.04, p > 0.05; indirect effect = 0.02, 95% CI =  $-0.02 \sim 0.09$ ). Furthermore, sensitivity analysis demonstrated that Rho = -0.24 when the average mediation effect was zero (see Figure 2).



Figure 2 Robustness analysis of intermediation effects.

# The Moderating Effect of Perceived Social Support in the Relationship Between Negative Affect and Impulsivity

As shown in Table 5, SAS total score significantly predicted score on BIS-11 Cognitive complexity (P<0.05). The interactions between SAS and MSPSS Support from family as well as Support from significant others on Cognitive complexity were significant (Ps < 0.05). These results suggest that support from significant others and family moderated the prediction of anxiety on cognitive complexity (see Figure 3).

### Discussion

The social-ecological theory indicates that individual behavior is affected by the social environment and its interaction with it, as well as the interaction among microsystem (such as physical and psychological characteristics of individuals), mesosystem (such as family, neighbors, and co-workers etc), and macrosystem (includes community, country, and culture).<sup>51</sup> Based on published literatures, the psychiatric or psychological risk factors of NSSI included behavioral disorders, personality disorders and depression or anxiety, while good family/friend relationships were most frequently identified as protective environmental or social factors.<sup>52</sup> Impulsivity has been studied as an aspect of personality and psychopathology for generations.<sup>53</sup> The study further explored the characteristics in impulsivity dimensions in adolescents with NSSI, and provided a detailed description of the relationship among negative affect, perceived social support, and their impulsivity. In this study, higher impulsivity and lower perceived social support were observed, and mediating and moderating effects of perceived social support on the relationship between anxiety and some impulsivity dimensions were noted in the NSSI group. These findings supported all our hypotheses.

**Table 5** Moderating Effect of Perceived Social Support (Indicated by MSPSS) on theRelationship Between Anxiety (by SAS), Depression (by SDS), and Impulsivity (by BIS-11) in Non-Suicidal Self -Injury Adolescents (N = 102)

Independent Variable		SE	95% CI	R <sup>2</sup>	Р
Cognitive complexity					
Anxiety		0.03	< 0.01, 0.11	0.35	0.027
Anxiety × Support from family		< 0.01	< 0.01, 0.02	0.07	0.008
Anxiety		0.02	< 0.01, 0.10	0.34	0.026
Anxiety × Support from significant others		< 0.01	< 0.01, 0.02	0.06	0.014

Abbreviations: SAS, the Zung Self-rating Anxiety Scale; SDS, the Zung Self-rating Depression Scale; MSPSS, the Multidimensional Scale of Perceived Social Support; BIS-11, the Barratt Impulsiveness Scale.



Figure 3 Interactions between anxiety symptom and perceived support from (A) family and (B) others on cognitive complexity of NSSI adolescents.

### Impulsivity Characteristics in NSSI Adolescents

In favor of our first hypothesis, adolescents with NSSI obtained higher scores in all the first-order factors of BIS-11 except for Cognitive instability, which suggested that their attentional impulsivity was mainly due to attention bias rather than cognitive instability.<sup>54,55</sup> These findings were fundamentally consistent with the recent crosssectional studies on the trend towards higher impulsivity in adolescents with NSSI,<sup>13,56</sup> and partly explained by the findings of Janis and Nock,<sup>57</sup> ie, although self-injurers reported higher impulsivity, performance-based measures of impulsivity (including more cognitive processes) failed to detect any difference between NSSI and healthy adolescents.

# Correlations Between Negative Affect, Perceived Social Support, and Impulsivity in NSSI Adolescents

Our findings that SAS and SDS scores were positively correlated with factors of BIS-11 revealed close relationships between impulsivity and individual symptoms of depression and anxiety, consistent with several previous reports that negative emotion would enhance the tendency of impulsivity.<sup>20,58,59</sup> While the subscales of the MSPSS were negatively

related to factors of the BIS-11, the results suggest that perceived social support may be a protective factor for most manifestations of impulsivity and negative affective states in adolescents with NSSI. A previous meta-analysis also reported that high levels of perceived social support prevented suicidal ideation among adolescents with NSSI and depression.<sup>60</sup> These findings also suggest that early intervention should focus on alleviating affective symptoms and increasing social support.<sup>61</sup>

## Mediating Role of Perceived Social Support in the Relationship Between Anxiety and Impulsivity

Our findings further showed mediation effect of MSPSS Support from significant others between SAS and BIS-11 Self-control, which indicated that anxiety reduced perception of support from significant others, and therefore lead to the loss of self-control in NSSI adolescents. Previous study also showed that higher anxiety symptoms were associated with more difficulties in the social domain (including difficulties in forming friendships) and more difficulties with emotional self-disclosure, which impacted significant others in an unaffiliated way and unidirectionally predicted later low social support.<sup>62</sup> And previous evidence revealed that individuals exposed to trauma with inadequate support from others were more likely to have a limited ability to exercise self-control, which subsequently contributed to impulsivity.<sup>63</sup> Our study indicated that decrease in perceived support from significant others beside support from family and friends can also lead anxiety to the risk of self-injury. Notably, the results of the sensitivity analysis suggest there's possibility that confounders affect the robustness of the mediation model. Therefore, further validation of complex mediation models with confounders such as culture, gender, socioeconomic status, education level, etc, is still needed, in consideration of their impact on adolescent NSSI behavior or social support perception.<sup>52,60,64</sup>

# The Moderating Effect of Perceived Social Support in the Relationship Between Anxiety and Impulsivity

The SAS score significantly predicted BIS-11 Cognitive complexity. We also found moderation effect of perceived support from family and significant others in the relationship between anxiety and cognitive complexity. Such results suggested that sufficient perceived support from family and significant others could lead NSSI adolescents with high anxiety levels to higher cognitive complexity, manifested by increased ability to consider multiple possibilities and solutions, which is often associate with reduce of non-planning behaviors. Previous evidence reported that patients with anxiety disorders were more likely to report impulsivity and NSSI than those without.<sup>65</sup> Another study investigating the relationship between impulsivity, comorbid anxiety and neurocognitive functioning found that anxiety predicted impulsivity and trait anxiety explained 26% of unplanned impulsivity,<sup>66</sup> which covers cognitive complexity.<sup>47</sup> There was little evidence of a moderating effect of perceived social support between anxiety and impulsivity in NSSI. However, in a study of migrant mothers only, social support was found to moderate and attenuate the relationship between internet addiction, which is associated with high anxiety, and suicidal ideation.<sup>67</sup> And perceived social support attenuated the impulsivity on anxiety and other psychiatric symptoms in women with methamphetamine use disorder.<sup>27</sup> These findings also highlight the importance of improving perceived social support in anxious NSSI patients, which reduces impulsivity and thus promotes recovery.

### Limitations of the Study and Future Directions

The study has several limitations. First, the cross-sectional design cannot explain causal relationships. Therefore, future studies are recommended to establish causal relationships between variables. Second, the sample size of this study, especially in Study 2, was relatively small due to the accessibility and cooperativeness of NSSI adolescents, which might lead to some effects not being detected. Third, it should be noted that the majority of participants in the study were female and from China. This may limit the generalisability of the findings to NSSI adolescents of different gender and

from different cultures. Finally, the study recruited mainly urban middle school students, which limited the exploration of other risk factors, such as low-income and disadvantaged children.

Therefore, a subsequent cross-cultural, multi-centre, and prospective validation with participants from different regions, ethnic groups, and socioeconomic status is recommended, and the influence of culture, gender, socioeconomic status, education level, etc. should be taken as cofounders when validating mediation models in NSSI adolescents. Furthermore, based on the results of the moderating effect of perceived social support, further research can be conducted on how to enhance the support of family and significant others for NSSI adolescents through intervention measures, as well as the long-term effects of such intervention on reducing impulsivity and preventing self-harm behaviours. Moreover, in view of the relationship between negative emotions and impulsivity found, more researches are needed upon whether this connection can be broken through methods such as emotion regulation training, thereby reducing the occurrence of self-harm behaviours.

## Conclusions

Higher anxiety led to lower perception of support from significant others in NSSI adolescents, which contributed to their loss of self-control. Whereas more perceived support from family and significant others could improve the ability of those in anxious state to perform complex cognitive activities. These findings further help to understand how micro individual psychological factors, in conjunction with socio-environmental factors (meso family and school support, macro community environment), impact on the behaviour of adolescents with NSSI. In clinical scenarios, psychological interventions such as cognitive-behavioural family therapy and motivational interviewing should be provided to NSSI adolescents with high anxiety, to improve their emotion regulation, subjective social support and reduce their risk of self-harm due to high impulsivity.

## **Data Sharing Statement**

The datasets used and/or analyzed during the current study are available from the corresponding or first author on reasonable request.

# **Ethics Approval and Consent to Participate**

Ethical approval was granted from Ethics Committee of the Second Affiliated Hospital, Zhejiang University School of Medicine (20240075).

# Acknowledgments

The authors wish to sincerely thank all of the staff participating this project. Most importantly, we wish to thank the young volunteers.

# **Author Contributions**

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

# Funding

Dr. Bingren Zhang was supported by a grant from the National Natural Science Foundation of China (No. 82101595).

# Disclosure

The authors report no conflicts of interest in this work.

- 1. Nock MK. Why do people hurt themselves? New insights into the nature and functions of self-injury. *Curr Dir Psychol Sci.* 2009;18:78–83. doi:10.1111/j.1467-8721.2009.01613.x
- 2. Bjureberg J, Sahlin H, Hellner C, et al. Emotion regulation individual therapy for adolescents with nonsuicidal self-injury disorder: a feasibility study. *BMC Psychiatry*. 2017;17(1):411. doi:10.1186/s12888-017-1527-4
- American Psychiatric Association (APA). Diagnostic and Statistical Manual of Mental Disorders: DSM-5. 5th. Washington, D.C: The Association; 2013:947.
- 4. Mannekote Thippaiah S, Shankarapura Nanjappa M, Gude JG, et al. Non-suicidal self-injury in developing countries: a review.Int. *J Soc Psychiatry*. 2021;67(5):472–482. doi:10.1177/0020764020943627
- Moloney F, Amini J, Sinyor M, Schaffer A, Lanctôt KL, Mitchell RHB. Sex differences in the global prevalence of nonsuicidal self-injury in adolescents: a meta-analysis. JAMA Network Open. 2024;7(6):e2415436. doi:10.1001/jamanetworkopen.2024.15436
- Xiao Q, Song X, Huang L, Hou D, Huang X. Global prevalence and characteristics of non-suicidal self-injury between 2010 and 2021 among a non-clinical sample of adolescents: a meta-analysis. Front Psychiatry. 2022;13:912441. doi:10.3389/fpsyt.2022.912441
- 7. Kaess M, Parzer P, Mattern M, et al. Adverse childhood experiences and their impact on frequency, severity, and the individual function of nonsuicidal self-injury in youth. *Psychiatry Res.* 2013;206:265–272. doi:10.1016/j.psychres.2012.10.012
- 8. Vega D, Sintes A, Fernández M, et al. Review and update on non-suicidal self-injury: who, how and why? Actas Esp Psiquiatr. 2018;46 (4):146–155.
- 9. Santangelo PS, Koenig J, Funke V, et al. Ecological momentary assessment of affective and interpersonal instability in adolescent non-suicidal self-injury. J Abnorm Child Psychol. 2017;45(7):1429–1438. doi:10.1007/s10802-016-0249-2
- O'Connor RC. Towards an integrated motivational-volitional model of suicidal behavior. In: O'Connor RC, Platt S, Gordon J, editors. Int Handbook of Suicide Prevention. Hoboken: Wiley-Blackwell; 2022: 181–198.
- 11. Lockwood J, Townsend E, Daley D, Sayal K. Impulsivity as a predictor of self-harm onset and maintenance in young adolescents: a longitudinal prospective study. J Affect Disord. 2020;274:583–592. doi:10.1016/j.jad.2020.05.021
- Soloff PH, Lynch KG, Malone KM KTM, Mann JJ, Mann JJ. Characteristics of suicide attempts of patients with major 383 depressive episode and borderline personality disorder: a comparative study. Ame J Psych. 2000;157(4):601–608. doi:10.1176/appi.ajp.157.4.601
- Li X, Liu XL, Wang YJ, Zhou DS, Yuan TF. The effects of childhood maltreatment on adolescent non-suicidal self-injury behavior: mediating role of impulsivity. Front Psychiatry. 2023;14:1139705. doi:10.3389/fpsyt.2023.1139705
- McLaughlin KA, Sheridan MA, Humphreys KL, Belsky J, Ellis BJ. The value of dimensional models of early experience: thinking clearly about concepts and categories. *Perspectives Psychol Sci.* 2021;16(6):1463–1472. doi:10.1177/1745691621992346
- 15. Gao Y, Liu J, Liu X, Wang Y, Qiu S. Dimensions of family stress and repetitive nonsuicidal self-injury in adolescence: examining the interactive effects of impulsivity and emotion dysregulation. *Child Abuse Negl.* 2024;152:106804. doi:10.1016/j.chiabu.2024.106804
- 16. Lockwood J, Daley D, Townsend E, Sayal K. Impulsivity and self-harm in adolescence: a systematic review. Eur Child Adolesc Psychiatry. 2017;26(4):387-402. doi:10.1007/s00787-016-0915-5
- 17. Phillipou A, Abel LA, Castle DJ, Gurvich C, Hughes ME, Rossell SL. Self-reported and behavioural impulsivity in anorexia nervosa. World J Psychiatry. 2016;6(3):345–350. doi:10.5498/wjp.v6.i3.345
- Yurdakul FG, Garip Çimen Y, Koçak Ulucaköy R, Almaz ŞE, Çalışkan Uçkun A, Bodur H. Impulsivity and ankylosing spondylitis: is there a relationship? *Turk J Phys Med Rehabil*. 2018;65(1):51–58. doi:10.5606/ffrd.2019.2580
- Xiao Q, Song X, Huang L, Hou D, Huang X. Association between life events, anxiety, depression and non-suicidal self-injury behavior in Chinese psychiatric adolescent inpatients: a cross-sectional study. *Front Psychiatry*. 2023;14:1140597. doi:10.3389/fpsyt.2023.1140597
- 20. Hack J, Martin G. Expressed emotion, shame, and non-suicidal self-injury. Int J Environ Res Public Health. 2018;15(5):890. doi:10.3390/ ijerph15050890
- Wang YJ, Li X, Ng CH, Xu DW, Hu S, Yuan TF. Risk factors for non-suicidal self-injury (NSSI) in adolescents: a meta-analysis. *EClinicalMedicine*. 2022;46:101350. doi:10.1016/j.eclinm.2022.101350
- 22. Hou S, Twayigira M, Luo X, et al. The relationship between emotional neglect and non-suicidal self-injury among middle school students in China: the mediating role of social anxiety symptoms and insomnia. *BMC Psychiatry*. 2023;23(1):248. doi:10.1186/s12888-023-04735-7
- 23. Wu RY, Huang JY, Ying JF, Gao QQ, Guo JQ, ou JN. Behavioral inhibition/approach systems and adolescent nonsuicidal self-injury: the chain mediating effects of difficulty in emotion regulation and depression. *Pers Individ Dif.* 2021;175:110718. doi:10.1016/j.paid.2021.110718
- Yaqoob N, Ahsan S. Impact of perceived social support and parental bonding in predicting suicidal intent among self-harm patients. J Pak Med Assoc. 2021;71(4):1148–1151. doi:10.47391/JPMA.1299
- Uçur Ö, Dönmez YE. Problematic internet gaming in adolescents, and its relationship with emotional regulation and perceived social support. Psychiatry Res. 2021;296:113678. doi:10.1016/j.psychres.2020.113678
- Weinberg M, Harel H, Shamani M, Or-Chen K, Ron P, Gil S. War and well-being: the association between forgiveness, social support, posttraumatic stress disorder, and well-being during and after war. Soc Work. 2017;62(4):341–348. doi:10.1093/sw/swx043
- 27. Wang S, Li J, Li Y, Xia Y, Gong Y, Mao F. The predictive role of impulsivity and perceived social support in psychiatric symptoms of women with methamphetamine use disorder. *Front Psychiatry*. 2023;14:1116650. doi:10.3389/fpsyt.2023.1116650
- 28. Fares-Otero NE, Sharp TH, Balle SR, et al. Social support and (complex) posttraumatic stress symptom severity: does gender matter? *Eur J Psychotraumatol*. 2024;15(1):2398921. doi:10.1080/20008066.2024.2398921
- Claes L, Bouman WP, Witcomb G, Thurston M, Fernandez-Aranda F, Arcelus J. Non-suicidal self-injury in trans people: associations with psychological symptoms, victimization, interpersonal functioning, and perceived social support. J Sex Med. 2015;12(1):168–179. doi:10.1111/ jsm.12711
- 30. Davey A, Arcelus J, Meyer C, Bouman WP. Self-injury among trans individuals and matched controls: prevalence and associated factors. *Health* Soc Care Community. 2016;24(4):485–494. doi:10.1111/hsc.12239
- 31. Farooq S, Tunmore J, Wajid Ali M, Ayub M. Suicide, self-harm and suicidal ideation during COVID-19: a systematic review. *Psychiatry Res.* 2021;306:114228. doi:10.1016/j.psychres.2021.114228

11

- 32. Lin L, Zhao S, Zhang J. Impulsivity kills some who did not plan to die by suicide: evidence from Chinese rural youths. *Omega*. 2019;79 (4):446–460. doi:10.1177/0030222818786417
- 33. Nam EJ, Lee JE. Mediating effects of social support on depression and suicidal ideation in older Korean adults with hypertension who live alone. J Nurs Res. 2019;27(3):e20. doi:10.1097/jnr.0000000000292
- Moller CI, Cotton SM, Badcock PB, et al. Relationships between different dimensions of social support and suicidal ideation in young people with major depressive disorder. J Affect Disord. 2021;281:714–720. doi:10.1016/j.jad.2020.11.085
- Lee TH, Liu HC, Huang YH, Sun FJ, Liu SI. How impulsivity is associated with adolescent depression: the role of substance use, gender and social support. *Psychol Res Behav Manag.* 2023;16:4959–4970 doi:10.2147/PRBM.S429923.
- 36. Park J, Lee DS, Shablack H, et al. When perceptions defy reality: the relationships between depression and actual and perceived Facebook social support. J Affect Disord. 2016;200:37–44. doi:10.1016/j.jad.2016.01.048
- 37. Runyan JD, Vermilya S, St Pierre M, Brooks NW, Fowler A, Brewer T. A mixed methods experience sampling study of a posttraumatic growth model for addiction recovery. *Sci Rep.* 2024;14(1):3511. doi:10.1038/s41598-024-53740-7
- 38. Dun Y, Li QR, Yu H, et al. Reliability and validity of the Chinese version of the kiddie-schedule for affective disorders and schizophrenia-present and lifetime version DSM-5 (K-SADS-PL-C DSM-5). J Affect Disord. 2022;317:72–78. doi:10.1016/j.jad.2022.08.062
- 39. Zung WW. A rating instrument for anxiety disorders. Psychosomatics. 1971;12:371-379. doi:10.1016/S0033-3182(71)71479-0
- 40. Wu WY. Self-Rating Anxiety Scale. In: Zhang ZJ, editor. *Behavioral Medicine Inventory Manual*. Beijing: The Chinese Medicine Electronic Audio and Video Publishing House; 2005:213–214.
- Sun J, Dunne MP. An exploratory study of the applicability of the Zung's self-rating anxiety scale (SAS) among Chinese adolescents. *Psychol Rep.* 2001;88:117–24.
- 42. Zung W. Self-Rating Depression Scale. Arch Gen Psychiatry. 1965;12:63-70. doi:10.1001/archpsyc.1965.01720310065008
- 43. Hui P, Yi Z, Ji Y, et al. Analysis of reliability and validity of Chinese version SDS scale in women of rural area. *Shanghai Med Pharm*. 2013;34 (14):20–22.
- 44. Chen SB, Hu H, Gao YS, He HY, Jin DX, Zhang CQ. Prevalence of clinical anxiety, clinical depression and associated risk factors in Chinese young and middle-aged patients with osteonecrosis of the femoral head. *PLoS One.* 2015;10(3):e0120234. doi:10.1371/journal.pone.0120234
- 45. Zimet GD, Dahlem NW, Zimet SG, Farley GK. The multidimensional scale of perceived social support. J Pers Asses. 1988;52:30–41. doi:10.1207/s15327752jpa5201\_2
- 46. Guan NC, Seng LH, Hway Ann AY, Hui KO. Factorial validity and reliability of the Malaysian simplified Chinese version of multidimensional scale of perceived social support (MSPSS-SCV) among a group of university students. Asia Pac J Public Health. 2015;27(2):225–231. doi:10.1177/ 1010539513477684
- 47. Patton JH, Stanford MS, Barratt ES. Factor structure of the Barratt impulsiveness scale. J Clin Psychol. 1995;51(6):768–774. doi:10.1002/1097-4679(199511)51:6<768::AID-JCLP2270510607>3.0.CO;2-1
- 48. Stanford MS, Mathias CW, Dougherty DM, Lake SL, Anderson NE, Patton JH. Fifty years of the Barratt Impulsiveness Scale: an update and review. *Pers Individ Dif.* 2009;47(5):385–395. doi:10.1016/j.paid.2009.04.008
- 49. Yao S, Yang H, Zhu X, et al. An examination of the psychometric properties of the Chinese version of the Barratt Impulsiveness Scale, 11th version in a sample of Chinese adolescents. *Percept Mot Skills*. 2007;104(3 Pt 2):1169–1182. doi:10.2466/pms.104.4.1169-1182
- 50. An J, Phillips MR, Conner KR. Validity of proxy-based reports of impulsivity and aggression in Chinese research on suicidal behavior. *Crisis*. 2010;31(3):137–142. doi:10.1027/0227-5910/a000019
- Chang H, Zhou J, Wang Z. Multidimensional factors affecting successful aging among empty-nesters in China based on social-ecological system theory. Int J Environ Res Public Health. 2022;19(19):11885. doi:10.3390/ijerph191911885
- 52. McEvoy D, Brannigan R, Cooke L, et al. Risk and protective factors for self-harm in adolescents and young adults: an umbrella review of systematic reviews. J Psychiatr Res. 2023;168:353–380. doi:10.1016/j.jpsychires.2023.10.017
- 53. Carver CS, Johnson SL. Impulsive reactivity to emotion and vulnerability to psychopathology. Am Psychol. 2018;73(9):1067-1078. doi:10.1037/ amp0000387
- 54. Wang MC, Deng Q, Shou Y, et al. Assessing impulsivity in Chinese: elaborating validity of BIS among male prisoners. *Criminal Justice Behav.* 2019;64(3):492–506. doi:10.1177/0093854818806033
- 55. Matera E, Margari M, Serra M, et al. Non-suicidal self-injury: an observational study in a sample of adolescents and young adults. *Brain Sci.* 2021;11(8):974. doi:10.3390/brainsci11080974
- 56. Costa RPO, Peixoto ALRP, Lucas CCA, et al. Profile of non-suicidal self-injury in adolescents: interface with impulsiveness and loneliness. *J Pediatr.* 2021;97(2):184–190. doi:10.1016/j.jped.2020.01.006
- 57. Janis IB, Nock MK. Are self-injurers impulsive?: results from two behavioral laboratory studies. *Psychiatry Res.* 2009;169(3):261–267. doi:10.1016/j.psychres.2008.06.041
- Miezah D, Mahama I, Owusu Amponsah MK, Moustafa A. Mental health states and impulsivity: role of stress, depression, and anxiety. *Psychol Neurosci Impulsivity*. 2024;2024:157–172.
- 59. Taş Torun Y, Gul H, Yaylali FH, Gul A. Intra/interpersonal functions of non-suicidal self-injury in adolescents with major depressive disorder: the role of emotion regulation, alexithymia, and childhood traumas. *Psychiatry*. 2022;85(1):86–99. doi:10.1080/00332747.2021.1989854
- 60. He H, Hong L, Jin W, et al. Heterogeneity of non-suicidal self-injury behavior in adolescents with depression: latent class analysis. *BMC Psychiatry*. 2023;23(1):301. doi:10.1186/s12888-023-04808-7
- 61. Ma Y, Li Y, Xie X, et al. The role of depressive symptoms and social support in the association of internet addiction with non-suicidal self-injury among adolescents: a cohort study in China. *BMC Psychiatry*. 2023;23(1):322. doi:10.1186/s12888-023-04754-4
- 62. Jacobson NC, Lord KA, Newman MG. Perceived emotional social support in bereaved spouses mediates the relationship between anxiety and depression. J Affect Disord. 2017;211:83–91. doi:10.1016/j.jad.2017.01.011
- Gao T, Mei S, Li M, O'Donnell K, Caron J, Meng X. Impulsivity mediates the relationship between childhood maltreatment and quality of life: does social support make it different? *Pers Individ Dif.* 2022;184:111208. doi:10.1016/j.paid.2021.111208
- 64. Gholamrezaei M, De Stefano J, Heath NL. Non suicidal self-injury across cultures and ethnic and racial minorities: a review. *Int J Psychol*. 2017;52 (4):316–326. doi:10.1002/ijop.12230

- 65. Peters EM, Bowen R, Balbuena L. Mood instability contributes to impulsivity, non-suicidal self-injury, and binge eating/purging in people with anxiety disorders. *Psychol Psychother*. 2019;92(3):422–438. doi:10.1111/papt.12192
- 66. Çörekçioğlu S, Cesur E, Devrim Balaban Ö. Relationship between impulsivity, comorbid anxiety and neurocognitive functions in bipolar disorder. *Int J Psychiatry Clin Pract.* 2021;25(1):62–72. doi:10.1080/13651501.2020.1833040
- 67. Miao S, Xu L, Gao S, Bai C, Huang Y, Peng B. Internet addiction and suicidal ideation in Chinese children with migrant parents: mediating role of anxiety and moderating role of social support. Acta Psychol. 2024;248:104427. doi:10.1016/j.actpsy.2024.104427

Neuropsychiatric Disease and Treatment



Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal

13