

# The Association Between Psychological Capital and Self-Management Behaviors in Men with Gout: A Cross-Sectional Study in Southwest China

Ying Wang<sup>1</sup>, Yanling Chen<sup>1</sup>, Qi Qi<sup>2</sup>, Yuqing Song<sup>3</sup>, Xin Guo<sup>1</sup>, Ling Ma<sup>1</sup>, Hong Chen<sup>4</sup>

<sup>1</sup>Department of Rheumatology and Immunology, West China Hospital of Sichuan University/West China School of Nursing, Sichuan University, Chengdu, Sichuan, People's Republic of China; <sup>2</sup>Department of Operating Room Nursing, West China Second University Hospital, Sichuan University/West China School of Nursing, Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, Chengdu, Sichuan, People's Republic of China; <sup>3</sup>School of Nursing, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan, People's Republic of China; <sup>4</sup>West China School of Nursing/ West China Hospital, Sichuan University, Chengdu, Sichuan, People's Republic of China

Correspondence: Hong Chen, West China School of Nursing/West China Hospital, Sichuan University, No. 37, Guoxue Alley, Wuhou District, Chengdu, Sichuan, 610041, People's Republic of China, Tel +86 18980601733, Email chong@scu.edu.cn

**Purpose:** Gout is a common, chronic inflammatory joint disease, and men are more likely to suffer from gout. Improving patient self-management behaviors is a priority in gout healthcare. Psychological capital is associated with self-management behaviors in chronic diseases and can be improved through a number of interventions. However, this topic has not been well studied in gout patients. The aim of this study was to determine the level of psychological capital among male gout patients in Southwest China and to compare differences in self-management behaviors among patients with different levels of psychological capital.

**Patients and Methods:** This was a cross-sectional study. A total of 242 male gout patients were recruited from West China Hospital of Sichuan University, and demographic characteristics, clinical characteristics, psychological capital, and behavioral variables related to patient self-management were collected. K-Means cluster analysis was used to characterize psychological capital.

**Results:** The total psychological capital score of the participants was 134.5 (SD = 21.3). Cluster analysis of the four dimensions of psychological capital yielded three clusters, namely, Cluster 1 (higher level, 29.8%), Cluster 2 (moderate level, 52.3%), and Cluster 3 (poor level, 17.9%). The differences in the self-management behaviors among the three clusters, the differences were statistically significant. Post hoc analyses revealed that cluster 1 scored higher on the self-Management behaviors and its four dimensions than either cluster 2 or cluster 3 ( $p < 0.05$ ).

**Conclusion:** The psychological capital of men with gout in Southwest China could be improved, and moderate and low levels of psychological capital are associated with suboptimal self-management behaviors. Healthcare providers may target gout patients with low or moderate levels of psychological capital as an intervention and take steps to improve their levels of psychological capital. These results may assist in decision-making for self-management behavioral interventions for gout patients.

**Keywords:** self efficacy, resilience, optimism, hope, cluster analysis

## Introduction

Gout is a common chronic inflammatory joint disease that results from persistent elevations in serum uric acid (SUA) levels and the deposition of uric acid crystals in joints, tendons, and other tissues.<sup>1</sup> Gout has impact on patients' somatic function,<sup>2</sup> social activities,<sup>2</sup> psychology and quality of life,<sup>2</sup> and it also causes significant economic loss and medical burden.<sup>3,4</sup> The global prevalence of gout ranges from 0.03% to 15.30% and is gradually increasing.<sup>1</sup> Men are more likely to suffer from gout, and the Chinese Rheumatism Data Center (CRDC) reported that the ratio of male to female gout patients in China is 15:1.<sup>5</sup> Between 1990 and 2017, the prevalence and incidence of gout in China increased by 6.88% and 6.16%,<sup>1</sup> respectively, and the prevalence and incidence of gout in men in China increased by 7.07% and 6.46%,<sup>1</sup> respectively, during this period. Thus, male patients are a priority for gout management in China.

The concept of treat-to-target (T2T) has been successfully applied to gout, and it is recommended that gout patients maintain SUA levels below 360  $\mu\text{mol/L}$  for a long period of time to promote crystal dissolution and prevent acute attacks of gout.<sup>6</sup> To achieve this therapeutic goal, gout patients should take measures such as long-term adherence to uric acid-lowering therapy, lifestyle changes, reduction of high-purine foods, and maintenance of a positive mental state, which means that self-management inevitably becomes the main mode of disease management for gout patients. Self-management is defined in the medical field as a health behavior that maintains and promotes health, wellness and management of disease, and persistent treatment of one's disease through a number of behaviors.<sup>7</sup> Self-management has now become a hot research topic in chronic disease management.<sup>8–10</sup> To the best of our knowledge, the self-management behavior of gout patients is not optimal,<sup>3</sup> the adherence to urate-lowering therapy (ULT) is only 47%,<sup>11</sup> and dietary control is also unsatisfactory.<sup>12</sup> Therefore improving patients' self-management behavior is also a focus of gout healthcare.

The implementation of self-management behavioral interventions presupposes an understanding of the factors associated with them. Previous studies have reported an association between chronic disease self-management behaviors and psychological factors.<sup>13–16</sup> Positive psychological capital, also known as psychological capital (PsyCap), comes from positive psychology and refers to the positive psychological state that emerges during an individual's growth process, including the four dimensions of self-efficacy, resilience, optimism, and hope.<sup>17</sup> Self-efficacy refers to a person's confidence in performing a challenging task. It represents a belief that the individual is capable of navigating the motivation, cognitive resources, and course of action required to successfully solve a given task.<sup>18</sup> Resilience is the ability to consistently overcome difficulties to succeed.<sup>19</sup> According to Rutter, resilience is the ability of individuals to successfully manipulate their environment to avoid negative consequences of adverse events.<sup>19</sup> Optimism refers to the positive attribution of current and possible future success. Optimism affects not only individuals' positive expectations for the future but also the coping strategies they choose.<sup>20</sup> Hope is the power to stick to a goal and adjust a path when necessary. People with high hope tend to be better at setting goals, resetting them in the face of adversity, and rationally using superior resources to achieve them.<sup>18</sup> These four concepts are independent, mutually reinforcing, and work together; in other words, individuals with high self-efficacy have hope for the future and relatively optimistic expectations for the future and are more resilient to adversity.<sup>21</sup> In recent years, PsyCap has been gradually applied in the field of health management.<sup>22,23</sup> A good positive psychology not only enables patients to face the disease correctly but also ensures that they deal with the disease correctly and make the right lifestyle choices for a rational lifestyle during the long-term progression of the disease. Previous quantitative studies have reported that PsyCap variables such as self-efficacy, resilience, optimism, and hope are associated with chronic disease self-management behaviors.<sup>16,24–26</sup> Furthermore, studies have confirmed that PsyCap can be enhanced by interventions.<sup>27–29</sup> Therefore, it can be hypothesized that understanding the PsyCap of gout patients may aid in the development of self-management intervention programs. However, worldwide research on PsyCap in gout patients has been limited. In addition, the dimensional characteristics of individual psychoanalysis are heterogeneous with different individual psychological manifestations. We need to further elucidate the potential characteristics of PsyCap in gout patients and the differences in self-management behaviors among gout patients with different psychological characteristics to quickly identify intervention targets and improve the efficiency of nursing practice. Cluster analysis, an analytical process in which sets of data objects are grouped into multiple classes composed of similar objects, is an effective method for determining the PsyCap characteristics of gout patients.

Therefore, the aim of this study was to evaluate the PsyCap levels of male gout patients in Southwest China to explore the potential characteristics of their PsyCap through cluster analysis and to analyze the differences in the self-management behaviors of gout patients with different psychological characteristics.

## Materials and Methods

### Study Design

This was a cross-sectional study using convenience sampling.

## Sampling, Recruitment and Data Collection

There are no clear recommendations for sample size estimation for cluster analysis. Some studies recommend a minimum sample size of not less than  $2^k$  ( $k$  = number of variables).<sup>30</sup> This study used the four dimensions of PsyCap as clustering variables, so the minimum sample size was  $2^4 = 16$ .

The study was conducted from February 2021 to January 2022 at West China Hospital of Sichuan University, a regional center hospital with patients mainly from the surrounding areas of Sichuan, Yunnan, and Guizhou. The inclusion criteria were (1) compliance with the 2015 American College of Rheumatology (ACR) and the European League Against Rheumatism (EULAR) diagnosis of gout by a rheumatologist,<sup>5,31</sup> (2) 18 years old and more, (3) ability to read and comprehend the questionnaire, and (4) participation. The exclusion criterion was cognitive or psychiatric abnormalities.

All participants were referred by a rheumatologist to participate in a 24-week randomized controlled trial of self-management for people with gout.<sup>32</sup> While in previous studies we have used baseline data from this program to explore the association between participants' psychosocial behaviors and quality of life,<sup>33</sup> the present study looks at participants' level of psychological capital and its association to self-management behaviors. Participants were first informed about the study and if they showed willingness, trained researchers informed participants about the purpose and the voluntary, anonymous nature of the study. All participants signed the informed consent form, were asked to independently complete a paper questionnaire, and were encouraged to seek help when needed. Submitted questionnaires were checked for completeness by the researcher. Data were manually entered by two researchers and checked.

## Ethical Considerations

The study complied with the ethical guidelines of the 1975 Declaration of Helsinki and received ethical approval from the Medical Ethics Committee of West China Hospital in 2020 (ID: 2020898). All participants signed an informed consent form before the start of the study.

## Instruments

### Demographic and Clinical Characteristics

The demographic characteristics were self-reported by the participants, included age, body mass index (BMI), marital status, education, employment status, and monthly household income; and the clinical characteristics were extracted from the hospital information system, included disease course, SUA level, visual analog scale (VAS) score of joint pain in the past 6 months, comorbidities, ULT, and tophi status.

### Psychological Capital


The Chinese version of the Positive PsyCap Questionnaire (PPQ) was used to assess participants' PsyCap.<sup>17,21</sup> Zhang et al validated a total Cronbach's alpha of 0.90.<sup>21</sup> The 26-item PPQ scores ranged from 1–7, and the total score ranged from 26–182, with higher scores indicating better PsyCap scores.<sup>17,21</sup> The PPQ consists of four dimensions: self-efficacy, resilience, optimism, and hope. In this study, Cronbach's alpha was 0.93.

### Self-Management Behavior

The Gout Patient Self-Management Assessment Scale (GPSAS) was used to measure the participants' self-management behaviors.<sup>34</sup> Yao et al developed the scale and validated its Cronbach's alpha of 0.962 and a content validity index of 0.905.<sup>34</sup> The GPSAS scores of 41 items ranged from 1–5, with a total score of 41–205, with higher scores indicating better self-management behaviors.<sup>34</sup> The GPSAS consists of four dimensions, ie, disease treatment management, diet management, lifestyle management, and psychosocial management. In this study, Cronbach's alpha was 0.92.

## Statistical Analysis

The data were analyzed using SPSS (version 25.0, IBM Corp). A one-sample K–S test was used to assess the normality of the data. Continuous variables are described using means (standard deviations) or medians (interquartile ranges) and categorical variables are expressed as frequencies and percentages.



**Figure 1** Number of clusters determined according to the elbow rule.

To categorize the data according to the four dimensions of the PPQ, we performed a k-means cluster analysis. We used the four dimensions of the PPQ as metrics, assuming a K of 2–5. The “Elbow rule” was used to review the calculation results, which showed that the optimal number of clusters was 3 (Figure 1).

Differences in demographic and clinical characteristics across PPQ clusters were analyzed by the chi-square test, one-way analysis of variance (ANOVA), the Kruskal–Wallis test or the chi-square test. Scores on the four dimensions of the PPQ and GPSAS were also compared across the three clusters using ANOVA. Post hoc Fisher’s least significant difference test (LSD-t) was used to compare variables that differed significantly between the three clusters, and  $P \leq 0.05$  was considered to indicate statistical significance.

## Results

### Demographic and Clinical Characteristics

In this study, 300 individuals were initially recruited, 242 of whom met the inclusion criteria. A total of 7 questionnaires were deleted due to the high level of duplication of all item options number; 235 questionnaires were included in the statistical analysis, and their demographic and clinical characteristics are listed in Table 1.

### Identifying the Clusters of PsyCap

The total PPQ score of the participants was 134.5 (SD = 21.3), and the self-efficacy, resilience, optimism, and hope dimensions were 36.0 (SD = 6.5), 33.9 (SD = 6.9), 32.2 (SD = 5.8), and 32.5 (SD = 6.0), respectively. Cluster analysis revealed three PPQ clusters (Table 2), and the visualization results are shown in Figure 2. The three clusters obtained were cluster 1 (N=70, 29.8%), cluster 2 (N=123, 52.3%), and cluster 3 (N=42, 17.9%), where the total PPQ scores from highest to lowest were cluster 1 (159.9, SD=11.7), cluster 2 (130.5, SD=7.5), and cluster 3 (104.1, SD=9.3). The total PPQ and four-dimensional scores of the three clusters were significantly different ( $p < 0.001$ ). Fisher’s LSD test was further used to determine significant differences between the two clusters ( $p < 0.001$ ). The demographic and clinical characteristics of the three different clusters are shown in Table 1. The differences in educational status, work status, and monthly household income of the participants were statistically significant ( $p < 0.05$ ). In terms of clinical characteristics, a statistically significant difference in disease course was observed ( $p = 0.026$ ).

### Differences in the Characteristics and Levels of GPSAS Across the Three Clusters

Table 3 shows the differences in self-management levels across the three clusters. The GPSAS score for all participants was 145.5 (SD=26.5), and the scores for the four dimensions of disease treatment management, diet management, lifestyle management, and psychosocial management were 49.2 (SD=10.7), 41.9 (SD=9.0), 28.3 (SD=7.8), and 26.1 (SD=5.7), respectively. The total GPSAS score for Cluster 1 was 161.0 (SD=21.5), that for Cluster 2 was 140.7

**Table 1** Demographic and Clinical Characteristics of the Participants and Differences Among the Three Clusters of the PPQ

Category	Range	Total (N=235)	Cluster 1 (N=70)	Cluster 2 (N=123)	Cluster 3 (N=42)	$F/\chi^2/H$	P
<b>Demographic characteristics</b>							
Age(year), Mean(SD)	18–75	40.3(12.5)	38.1(10.7)	40.5(12.6)	43.3(14.39)	2.296 <sup>a</sup>	0.103
BMI(kg/m <sup>2</sup> ), Mean(SD)	19.1–35.9	26.1(3.3)	25.6(3.3)	26.3(3.2)	26.2(3.5)	1.085 <sup>a</sup>	0.340
Marital status, N(%)							
Single/Divorced/Widowed		56(23.8)	16(22.9)	33(26.8)	7(16.7)	1.833 <sup>b</sup>	0.400
Married		179(76.2)	54(77.1)	90(73.2)	35(83.3)		
Education, N(%)							
Primary school		13(5.5)	4(5.7)	6(4.9)	3(7.1)	20.367 <sup>b</sup>	<b>0.026</b>
Middle school		38(16.2)	8(11.4)	17(13.8)	13(31.0)		
High school		37(15.7)	7(10.0)	20(16.3)	10(23.8)		
Junior college		44(18.7)	17(24.3)	21(17.1)	6(14.3)		
Undergraduate		79(33.6)	23(32.9)	49(39.8)	7(16.7)		
Postgraduate		24(10.2)	11(15.7)	10(8.1)	3(7.1)		
Employment status, N(%)							
Employed		189(80.4)	61(87.1)	100(81.3)	28(66.7)	7.117 <sup>b</sup>	<b>0.028</b>
Unemployed		46(19.6)	9(12.9)	23(18.7)	14(33.3)		
Monthly household income, (¥, CNY), N(%)							
<4000		70(29.8)	12(17.1)	38(30.9)	20(47.6)	22.83 <sup>b</sup>	<b>&lt;0.001</b>
4000–7999		74(31.5)	19(27.1)	38(30.9)	17(40.5)		
≥8000		91(38.7)	39(55.7)	47(38.2)	5(11.9)		
<b>Clinical characteristic</b>							
Disease Course(month), M(IQR)	0–490	59(96)	39(77.2)	60(88)	74(33.8)	7.331 <sup>c</sup>	<b>0.026</b>
SUA(umol/L), Mean(SD)	154–818		479.0(123.2)	498.6(128.0)	452.1(135.3)	2.160 <sup>a</sup>	0.118
Pain(VAS 0–10), Mean(SD)	0–10		5.2(2.6)	4.6(3.1)	4.7(3.3)	0.820 <sup>a</sup>	0.442
Comorbidity, N (%)							
No		124(52.8)	36(51.4)	66(53.7)	22(52.4)	0.092 <sup>b</sup>	0.955
Yes		111(47.2)	34(48.6)	57(46.3)	20(47.6)		
ULT, N (%)							
No		43(18.3)	13(18.6)	23(18.7)	7(16.7)	2.262 <sup>b</sup>	0.688
Yes		189(80.4)	55(78.6)	99(80.5)	35(83.3)		
Missing value		3(1.3)	2(2.9)	1(0.8)	0(0.0)		
Tophi, N (%)							
No		162(72.0)	49(74.2)	85(72.7)	28(66.7)	0.782 <sup>b</sup>	0.676
Yes		63(28.0)	17(25.8)	32(27.4)	14(33.3)		

**Note:**<sup>a</sup>one-way analysis of variance; <sup>b</sup>chi-square test; <sup>c</sup>Kruskal–Wallis test; the bold font:  $P \leq 0.05$ .

**Abbreviations:** PPQ, Positive PsyCap questionnaire; SD, standard deviation; M, median; IQR, interquartile range; BMI, body mass index; SUA, serum uric acid; VAS, visual analog scale; ULT, urate-lowering therapy.

(SD=25.6), and that for Cluster 3 was 133.9 (SD=25.7). Table 3 lists the four GPSAS dimension scores. The total GPSAS score and the four dimension scores for all clusters were significant ( $p < 0.05$ ). Further post hoc analysis using Fisher's LSD test showed that the scores of the GPSAS and its four dimensions in cluster 1 were greater than those in cluster 2 or cluster 3 ( $p < 0.05$ ). There was no significant difference between Cluster 2 and Cluster 3 except for the psychosocial management dimension ( $p > 0.05$ ).

**Table 2** PPQ Dimension Scores of the Three Clusters of PsyCap

PPQ Dimensions	Range	Total (N=235) Mean(SD)	Cluster 1 (N=70) Mean(SD)	Cluster 2 (N=123) Mean(SD)	Cluster 3 (N=42) Mean(SD)	Overall P	Cluster 1 VS Cluster 2 P	Cluster 1 VS Cluster 3 P	Cluster 2 VS Cluster 3 P
PPQ (total)	76–182	134.5(21.3)	159.9(11.7)	130.5(7.5)	104.1(9.3)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Self efficacy	18–49	36.0(6.5)	42.2(4.9)	35.3(4.0)	27.7(3.6)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Resilience	20–49	33.9(6.9)	40.3(6.1)	32.2(5.1)	27.9(4.2)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Optimism	14–42	32.2(5.8)	38.6(3.0)	31.2(3.3)	24.7(3.9)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Hope	16–42	32.5(6.0)	38.8(3.2)	31.8(3.2)	23.9(4.0)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>

**Note:** the bold font:  $P \leq 0.05$ .


**Abbreviations:** PPQ, Positive PsyCap questionnaire; PsyCap, psychological capital; SD, standard deviation.

## Discussion

Gout is a chronic and lifelong condition and maintaining positive self-management behaviors is critical. Gout affects patients in a variety of ways and is often accompanied by adverse emotional experiences.<sup>2–4</sup> PsyCap is a concept of human strength and positive aspects, and individuals with higher PsyCap are more likely to overcome negative emotions and produce positive behaviors.<sup>35–37</sup> Therefore, we sought to determine the level of PsyCap in gout patients and whether it is associated with self-management behaviors.

We found that the PsyCap of gout patients was 134.5 (SD=21.3). Most of the previous PsyCap-related studies used quantitative research methods and could not consider the stratification of PsyCap levels. Through cluster analysis, we identified three PPQ clusters, cluster 1 (higher level), cluster 2 (moderate level), and cluster 3 (poor level). The total PPQ scores and dimensions of the study participants in the three clusters were significantly different, suggesting that the three clusters could distinguish between the three levels of PsyCap. This study also revealed that the majority of participants (70.2%) had poor or moderate levels of PsyCap, suggesting that the PsyCap levels of male gout patients in Southwest China need to be improved and that healthcare providers need to develop interventions to improve the PsyCap of gout patients.

We also found statistically significant differences in total GPSAS scores among the three clusters, with Cluster 1 scoring higher than Cluster 2 and Cluster 3, and similar results were observed across the four dimensions of GPSAS scores. This finding suggested that moderate and low levels of PsyCap may be associated with suboptimal self-management behaviors in gout patients. Previous studies have shown that one or more of the concepts in PsyCap are associated with self-management behaviors in patients with a number of chronic diseases such as diabetes, chronic kidney disease, and multiple sclerosis.<sup>13–16,38</sup> PsyCap can provide positive psychological support and can be reflected in personal behavior.<sup>39</sup> Those with a higher PsyCap may see it from a more positive perspective and have better resilience than those with a lower PsyCap. Luthans and others suggested that a number of models including main,



**Figure 2** Visualization of the three clusters for the four subscales of PPQ.



**Table 3** Level of Gout Self-Management Behavior Based on the Three Clusters of PPQ

Gout Patient Self-Management	Range	Total (N=235) Mean(SD)	Cluster 1 (N=70) Mean(SD)	Cluster 2 (N=123) Mean(SD)	Cluster 3 (N=42) Mean(SD)	Overall P	Cluster 1 VS Cluster 2 P	Cluster 1 VS Cluster 3 P	Cluster 2 VS Cluster 3 P
GPSAS(total)	56–201	145.5(26.5)	161.0 (21.5)	140.7(25.6)	133.9(25.7)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.120
Disease treatment management	15–65	49.2(10.7)	54.1 (9.6)	47.7(9.9)	45.0(11.9)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.141
Diet management	14–60	41.9(9.0)	45.8 (7.8)	40.7(9.0)	39.1(8.8)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.293
Lifestyle management	9–45	28.3(7.8)	30.7 (7.5)	27.2(8.1)	27.5(6.7)	<b>0.008</b>	<b>0.003</b>	<b>0.031</b>	0.856
Psychosocial management	7–35	26.1(5.7)	30.4 (4.2)	25.0(5.5)	22.3(4.0)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>0.002</b>

**Note:** the bold font:  $P \leq 0.05$ .

**Abbreviations:** PPQ, Positive PsyCap questionnaire; GPSAS, gout patient self-management assessment scale; SD, standard deviation.

buffer, middle effects and dynamic effects models can influence outcomes and that these effects may be direct or indirect.<sup>40–42</sup> Our study also confirmed that PsyCap is associated with self-management behaviors in gout patients, which means that healthcare providers can use increasing PsyCap as a strategy to improve self-management behaviors in gout patients.

PsyCap, which includes self-efficacy, resilience, optimism, and hope,<sup>17</sup> can be a positive psychological resource for adhering to self-management behaviors. Previous studies have emphasized the relevance of self-efficacy and self-management in patients with chronic diseases.<sup>13,38,43,44</sup> In this study, the self-management behaviors of cluster 1 were significantly better than those of the other two clusters, clearly demonstrating the value of using self-efficacy as an intervention target. Individuals with chronic conditions who have high resilience are more likely to adopt positive self-management behaviors, a finding confirmed by our study.<sup>15,44</sup> Therefore, increasing the level of resilience may be beneficial for promoting self-management behaviors in gout patients. Wilski found that higher levels of optimism corresponded to greater self-management behavior.<sup>16</sup> This study also suggested that increasing the level of optimism in gout patients may be a way to improve their self-management behavior. Hope levels directly or indirectly influence self-management of chronic illness.<sup>14,26</sup> The present study also revealed that high hope was associated with high self-management behaviors in gout patients, suggesting that interventions targeting hope may help improve self-management behaviors in gout patients.

We also found that participants in the three groups differed in terms of educational status, employment status, income level and disease course. A study of patients with ischemic stroke have similar findings.<sup>45</sup> Future studies should explore the mechanisms by which these factors are associated with psychological capital in gout patients in order to develop interventions to improve their psychological capital.

This study has several limitations. First, these participants were recruited through convenience sampling. Therefore, the findings should be generalized with caution, and further multicenter studies are needed for validation. Second, this was a cross-sectional study, and the findings do not establish causal evidence. Longitudinal studies are needed to examine the role of PsyCap in self-management in gout patients. Third, this study used patient self-reported data, and the results are susceptible to recall bias.

Despite these limitations, the significance of this study is to compensate for the lack of attention to the psychological capital of gout patients in previous studies. We analyzed the PsyCap status of patients with gout and, through cluster analysis, classified PsyCap with multidimensional characteristics into three clusters and determined that low, medium, and intermediate levels of PsyCap were associated with suboptimal self-management behaviors.

## Conclusion

In this study, we found that the PsyCap of gout patients still needs to be improved, and we divided the PsyCap of gout patients into three subgroups, in which moderate and lower levels of PsyCap corresponded to suboptimal self-management behaviors. Therefore, healthcare providers may intervene in gout patients with low, medium, and PsyCap levels and take certain measures to improve their PsyCap levels. These results may assist in decision-making for self-management behavioral interventions for gout patients.

## Acknowledgments

The authors acknowledge gout patients who were involved in this study for providing feedback and data.

## Funding

This work was supported by the SCST (Science and Technology of Sichuan Province) (Grant No.2018FZ0110 and No.2023JDR0251).

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Tang YM, Zhang L, Zhu SZ, et al. Gout in China, 1990-2017: the global burden of disease study 2017. *Public Health*. 2021;191:33–38. doi:10.1016/j.puhe.2020.06.029
2. Coulshed A, Nguyen AD, Stocker SL, Day RO. Australian patient perspectives on the impact of gout. *Int J Rheum Dis*. 2020;23(10):1372–1378. doi:10.1111/1756-185X.13934
3. Chua CKT, Cheung PP, Santosa A, Lim AYN, Teng GG. Burden and management of gout in a multi-ethnic Asian cohort. *Rheumatol Int*. 2020;40(7):1029–1035. doi:10.1007/s00296-019-04475-6
4. Flores NM, Nuevo J, Klein AB, Baumgartner S, Morlock R. The economic burden of uncontrolled gout: how controlling gout reduces cost. *J Med Econ*. 2019;22(1):1–6. doi:10.1080/13696998.2018.1532904
5. Chinese Rheumatology Association. 2016 China gout clinical practice guideline. *Chin J Intern Med*. 2016;55(11):892–899. doi: 10.3760/cma.j.issn.0578-1426.2016.11.019
6. Kiltz U, Smolen J, Bardin T, et al. Treat-to-target (T2T) recommendations for gout. *Ann Rheum Dis*. 2017;76(4):632–638. doi:10.1136/annrheum-dis-2016-209467
7. Lorig KR, Holman H. Self-management education: history, definition, outcomes, and mechanisms. *Ann Behav Med*. 2003;26(1):1–7. doi:10.1207/S15324796ABM2601\_01
8. Nikiphorou E, Santos EJF, Marques A, et al. 2021 EULAR recommendations for the implementation of self-management strategies in patients with inflammatory arthritis. *Ann Rheum Dis*. 2021;80(10):1278–1285. doi:10.1136/annrheumdis-2021-220249
9. Safari R, Jackson J, Sheffield D. Digital self-management interventions for people with osteoarthritis: systematic review with meta-analysis. *J Med Internet Res*. 2020;22(7):e15365. doi:10.2196/15365
10. Damgaard AJ, Primdahl J, Esbensen BA, Latocha KM, Bremander A. Self-management support needs of patients with inflammatory arthritis and the content of self-management interventions: a scoping review. *Semin Arthritis Rheu*. 2023;60:152203. doi:10.1016/j.semarthrit.2023.152203
11. Yin R, Li L, Zhang G, et al. Rate of adherence to urate-lowering therapy among patients with gout: a systematic review and meta-analysis. *BMJ Open*. 2018;8(4):e017542. doi:10.1136/bmjopen-2017-017542
12. Liddle J, Richardson JC, Hider SL, et al. “It’s just a great muddle when it comes to food”: a qualitative exploration of patient decision-making around diet and gout. *Rheumatol Adv Pract*. 2021;5(3):rkab055. doi:10.1093/rap/rkab055
13. King DK, Glasgow RE, Toobert DJ, et al. Self-efficacy, problem solving, and social-environmental support are associated with diabetes self-management behaviors. *Diabetes Care*. 2010;33(4):751–753. doi:10.2337/dc09-1746
14. Liu T, Chen DH, Jia QM, et al. Effect of hope on self-efficacy and self-management in patients with chronic kidney disease(stages 1-3). *Zhongguo Yi Xue Ke Xue Yuan Xue Bao*. 2019;41(3):367–372. doi:10.3881/j.issn.1000-503X.10680
15. Jia J, Jenkins AJ, Quintiliani LM, Truong V, Lasser KE. Resilience and diabetes self-management among African-American men receiving primary care at an urban safety-net hospital: a cross-sectional survey. *Ethnic Health*. 2022;27(5):1178–1187. doi:10.1080/13557858.2020.1849566
16. Wilski M, Kocur P, Broła W, Tasiemski T. Psychological factors associated with self-management in multiple sclerosis. *Acta Neurol Scand*. 2020;142(1):50–57. doi:10.1111/ane.13236
17. Luthans F, Avolio BJ, Avey JB, Norman SM. Positive psychological capital: measurement and relationship with performance and satisfaction. *Personnel Psychol*. 2007;60(3):541–572. doi:10.1111/j.1744-6570.2007.00083.x
18. Broad JD, Luthans F. Positive resources for psychiatry in the fourth industrial revolution: building patient and family focused psychological capital (PsyCap). *Int Rev Psychiatry*. 2020;32(7–8):542–554. doi:10.1080/09540261.2020.1796600
19. Rutter M. Resilience in the face of adversity. protective factors and resistance to psychiatric disorder. *Br J Psychiatry*. 1985;147:598–611. doi:10.1192/bjp.147.6.598
20. Scheier MF, Weintraub JK, Carver CS. Coping with stress: divergent strategies of optimists and pessimists. *J Pers Soc Psychol*. 1986;51(6):1257–1264. doi:10.1037//0022-3514.51.6.1257



21. Zhang K, Zhang S, Dong YH. Positive psychological capital: measurement and relationship with mental health. *Stud Psych Behav.* 2010;8(1):58–64.
22. Zeng K, Li Y, Yang R. The mediation role of psychological capital between family relationship and antenatal depressive symptoms among women with advanced maternal age: a cross sectional study. *BMC Pregnancy Childbirth.* 2022;22(1):488. doi:10.1186/s12884-022-04811-y
23. Cao S, Zhu Y, Li P, Zhang W, Ding C, Yang D. Age difference in roles of perceived social support and psychological capital on mental health during COVID-19. *Front Psychol.* 2022;13:801241. doi:10.3389/fpsyg.2022.801241
24. Dong N, Chen WT, Bao M, Lu Y, Qian Y, Lu H. Self-Management Behaviors Among Patients With Liver Cirrhosis in Shanghai, China: a cross-sectional study. *Clin Nurs Res.* 2020;29(7):448–459. doi:10.1177/1054773818777914
25. Chang EM, Chen LS, Li YT, Chen CT. Associations between self-management behaviors and psychological resilience in patients with COPD. *Resp Care.* 2023;68(4):511–519. doi:10.4187/respcare.10416
26. Zhang D, Zhang N, Chang H, et al. Mediating role of hope between social support and self-management among Chinese liver transplant recipients: a multi-center cross-sectional study. *Clin Nurs Res.* 2023;32(4):776–784. doi:10.1177/10547738221078897
27. Corbu A, Peláez Zuberbühler MJ, Salanova M. Positive psychology micro-coaching intervention: effects on psychological capital and goal-related self-efficacy. *Front Psychol.* 2021;12:566293. doi:10.3389/fpsyg.2021.566293
28. Da S, He Y, Zhang X. Effectiveness of psychological capital intervention and its influence on work-related attitudes: daily online self-learning method and randomized controlled trial design. *Int J Environ Res Public Health.* 2020;17(23):8754. doi:10.3390/ijerph17238754
29. Song R, Sun N, Song X. The Efficacy of Psychological Capital Intervention (PCI) for depression from the perspective of positive psychology: a pilot study. *Front Psychol.* 2019;10:1816. doi:10.3389/fpsyg.2019.01816
30. Im Y, Jung S. Family functioning according to clusters of family management styles in Korean families of children with chronic atopic disease: a cross-sectional study. *Int J Nurs Stud.* 2020;109:103674. doi:10.1016/j.ijnurstu.2020.103674
31. Neogi T, Jansen TLTA, Dalbeth N, et al. 2015 Gout classification criteria: an American College of Rheumatology/European league against rheumatism collaborative initiative. *Ann Rheum Dis.* 2015;74(10):1789–1798. doi:10.1136/annrheumdis-2015-208237
32. Wang Y, Chen Y, Song Y, et al. The impact of mHealth-based continuous care on disease knowledge, treatment compliance, and serum uric acid levels in Chinese patients with gout: randomized controlled trial. *JMIR mHealth uHealth.* 2024;12:e47012. doi:10.2196/47012
33. Wang Y, Guo X, Chen B, et al. The relationship between psychosocial behavior and the quality of life of male gout patients in Southwest China: a cross-sectional study based on an information-motivation-behavioral skills model. *Patient Prefer Adher.* 2023;17:3503–3514. doi:10.2147/PPA.S434875
34. Yao XY, Liu T, Li Y, et al. Development and psychometric testing of a gout patient self-management assessment scale. *Chin J Nurs.* 2020;320(55):261–265. doi:10.3761/j.issn.0254-1769.2020.02.018
35. Luthans F. The need for and meaning of positive organizational behavior. *J Organ Behavior.* 2002;23(6):695–706. doi:10.1002/job.165
36. Yan D, Wen F, Li X, Zhang Y. The relationship between psychological capital and innovation behaviour in Chinese nurses. *J Nurs Manage.* 2020;28(3):471–479. doi:10.1111/jonm.12926
37. An analysis of positive psychological capital influencing emotional reactions of college students under COVID-19-all databases. Available from: <https://webofscience.clarivate.cn/wos/alldb/full-record/KJD:ART002659170>. Accessed December 2, 2023.
38. Wilski M, Tasiemski T. Illness perception, treatment beliefs, self-esteem, and self-efficacy as correlates of self-management in multiple sclerosis. *Acta Neurol Scand.* 2016;133(5):338–345. doi:10.1111/ane.12465
39. Abbas M, Raja U. Impact of psychological capital on innovative performance and job stress. *Can J Adm Sci.* 2015;32(2):128–138. doi:10.1002/cjas.1314
40. Luthans F, Youssef CM, Avolio BJ. *Psychological Capital: Developing the Human Competitive Edge*. New York, NY: Oxford University Press; 2007.
41. Cole K, Daly A, Mak A. Good for the soul: the relationship between work, wellbeing and psychological capital. *J Socio Econ.* 2009;38(3):464–474. doi:10.1016/j.socsc.2008.10.004
42. Wang Y, Zhu Y. Advances in psychological capital theory and related research. *Econs Mgmt.* 2007;05:32–39.
43. Papadakos J, Barnsley J, Berta W, Rowlands G, Samoil D, Howell D. The association of self-efficacy and health literacy to chemotherapy self-management behaviors and health service utilization. *Support Care Cancer.* 2022;30(1):603–613. doi:10.1007/s00520-021-06466-5
44. Wang RH, Chen SY, Lee CM, Lu CH, Hsu HC. Resilience, self-efficacy and diabetes distress on self-management behaviours in patients newly diagnosed with type 2 diabetes: a moderated mediation analysis. *J Adv Nurs.* 2023;79(1):215–222. doi:10.1111/jan.15483
45. Zhang Y, Liu Z, Wang X, Gu Y. Influencing factors associated with psychological capital among Ischemic stroke patients: a latent profile analysis. *Psychol Res Behav Manag.* 2024;17:4043–4052. doi:10.2147/PRBM.S482943

## Patient Preference and Adherence

### Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>

**Dovepress**  
Taylor & Francis Group